Caring for a Spouse with ADL Dependency: Perceived Social Support and Stress Level among Elderly Spousal Caregivers by Rahmi Setiyani

Submission date: 06-Oct-2022 11:32AM (UTC+0700) Submission ID: 1917978989 File name: AOTMPH_1.pdf (889.1K) Word count: 3250 Character count: 17547 See discussions, stats, and author profiles for this publication at: https://www.researchgate.net/publication/354399962

Caring for a Spouse with ADL Dependency: Perceived Social Support and Stress Level among Elderly Spousal Caregivers

Article in Annals of Tropical Medicine and Public Health · September 2021 DOI: 10.36295/AOTMPH.2021.7116 CITATIONS READS 75 0 3 authors: Tri Harningsih Rahmi Setiyani . Universitas Jenderal Soedirman Universitas Jenderal Soedirman 1 PUBLICATION 0 CITATIONS 18 PUBLICATIONS 24 CITATIONS SEE PROFILE SEE PROFILE Koernia Nanda Pratama Universitas Jenderal Soedirman 8 PUBLICATIONS 0 CITATIONS SEE PROFILE Some of the authors of this publication are also working on these related projects: 4 Improving diarrheal preventive behavior through therapeutic sociodramatic play in school-aged children View project

All content following this page was uploaded by Rahmi Setiyanion 07 September 2021.

The user has requested enhancement of the downloaded file.

Annals of Tropical Medicine and Public Health Volume 7, Issue 1 Print ISSN: 1755-6783 Online ISSN: 0974-6005



Caring for a Spouse with ADL Dependency: Perceived Social Support and Stress Level among Elderly Spousal Caregivers

Tri Harningsih^{1,} Rahmi Setiyani², Koernia Nanda Pratama²

¹Student of Department of Nursing, Faculty of Health Sciences, Jenderal Soedirman University ²Department of Nursing, Faculty of Health Sciences, Jenderal Soedirman University

*Corresponding author: Rahmi Setiyani. Jl. Dr. Soeparno, Karang Wangkal, Purwokerto, Central Java, Indonesia, rahmi.setiyani@unsoed.ac.id

Abstract

Background: Taking care for an individual with Activity of Daily Living (ADL) dependence is often very demanding and stressful. With an increasing in life expectancy, an elderly spouse often becomes a caregiver for an aging or ill loved one. However, changing in family structure, shrinking social circle and aging process itself might influence caregiver's social support and their stress level. This study aimed to examine perceived stress and social support among spousal caregivers of elderly with ADL dependence and correlation between these two variables.

Methods: This is a descriptive quantitative study with cross sectional design. A convenience samples of 30 caregivers aged 60 years or over who taking care for spouse with ADL dependence. Perceived social support and care giving-related stress were measured using social support questionnaire and the King stone Caregiver Stress Scale (KCSS) respectively. Data was analyzed using descriptive statistics and Somer's D test.

Results: Result showed that a half of respondents were lack of social support and the remaining only had moderate level of support (46.7%). Very few of them had good social support (3.3%). Most respondents experienced moderate stress level (56.7%), even severe (16.7%). Statistical analysis showed no relationship between social support and stress level (p>0.05).

Conclusion: Elderly spousal caregiver was lack of social support and experienced moderate stress level.

Keyword: Activity of daily living; elderly; spousal Caregiver; social support; stress

Page: 100

http://doi.org/10.36295/AOTMPH.2021.7116

Introduction

As people age, their body functions, including immune function are decreased. As a result, they become more susceptible to diseases and disability (1). About 26% of eldery perform worldwide was estimated to live with disability in activities of daily living (ADL) (2). The number is even bigger in institutional care setting. A previous study found that 72% of the nursing home's residents were partially dependent and 14% of them are totally dependent (3). When individuals gradually lose their independence, they definitely need assistance from others. Caregiver is an individual who provides care and assistance to those who have ADL disability or dependence (4). Family members such as spouse, children, or parents most often have to play a role as an informal caregiver. With an increasing in life expectancy, elderly often become caregivers for their aging or ill spouse. A spouse is considered to be an ideal caregiver because they can provide both physical care and emotional support to their partners (5-7). Putra on his study found that 23.3% of elderly became caregivers for their ill spouse (7). The long process of taking care for individual with ADL dependency undoubtedly becomes a stressor for the caregiver.

The stress experienced by caregiver might relate to physical, psychological, social, and financial burdens. Compare to younger caregiver, the older one tend to feel depressed and are at greater risk of experiencing physical and psychological problem if it left untreated (8,9). Elderly are more susceptible to have stress because they usually have to deal with other stressors, not only the care giving responsibility. For example most eldelry caregiver find themselves struggling with physical and mental health issues, economics hardship because of retirement, and also changes in social network, and family structure (10,11). People have different appraisal process of potential stressor. As a result, they deal with stress in different ways. Some people deal with it just fine and others have a hard time. According to the Buffering Hypothesis Theory by Cohen and Wills, stress appraisal can be influenced by social support (12). Social support also plays a role in reappraisal process, inhibition of maladaptive responses, and facilitation of adaptive ones which are important to counter the negative effects of stress. Social support for caregivers can be from children, siblings, spouse, other family members, neighbors, friends and health workers (13).

However, differ from younger caregiver, most elderly caregivers may find their social circles shrinking. Most often, spouse becomes the most significant person for them.

However, spouse can be turnout to be a significant stressor, for example when they suffers from debilitating diseases (14). The issue of care giving-related stress and social support have been commonly studied among younger caregivers (15).

However, little is known about that of elderly caregivers who caring for spouse with ADL dependence. This study is undertaken to examine perceived social support and stress level among spousal caregivers of elderly with ADL dependence and correlation between these two variables.

Method

This is a descriptive quantitative study with cross sectional design. This study was conducted in 7 villages in Baturraden District, Purwoker to in 2019. A total of 30 caregivers who met the study criteria was participated in this study. The inclusion criteria were aged 60 years or over, taking care for spouse with ADL dependence, have been in care giving role ≥ 6 months and willing to participate in this study. The studied variables were social support and care giving stress level. Social support was measured using an adapted social support questionnaire from previous research (16).

The instrument consists of 20 items using 4 point rating scale (1=never to 4=always).

The total score is ranging from 20-80 and then categorized into three levels of social supports: low (20-40), moderate (41-60) and high (61-80). Meanwhile, caregiving-related stress were measured using the King stone Caregiver Stress Scale (KCSS). There are 10 items with 5 point rating scale (1=feeling no stress to 5=extreme stress).

The total score is ranging from 10-50 which then categorized into three levels of care giving stress: low (10-15), moderate (16-24), and high (25-50).

Data were analyzed using descriptive stors stors and Somer's D test. Prior to data collection, respondents were given explanation about aim and nature of this study and then signed informed consent if they were agreed to participate.

and anonimity were maintained throughout the study. Ethical approval was gained from the Health Research Ethics Committee of Faculty of Health Sciences, graduate from primary school (80%) or even never taken Jenderal Soedirman University 070.1/1162/IX/2019).

Ethical principles, including autonomy, confidentiality with different level of ADL dependency, but majority was mild one (76.7%). In regards to education, most of caregivers had a low level of education either just (No. any formal education (10%). All caregivers in this study also had low social economic status and majority of them had no paid job (70%).

Results

Table 1. Characteristics of respondents (n=30)

Category		
Age (Mean, SD)	71	8.836
Education Level (n, %)		
Uneducated	3	10.0
Primary School	24	80.0
Junior High School	1	3.3
Senior High school	2	6.7
College	0	0
Employment Status (n, %)		
Employed	9	30.0
Unemployed	21	70.0
Caregiver's Status (n, %)		
Wife	14	46.7
Husband	16	53.3
Social Economics Status (Income)		
Under standard (≤1,750,000)	30	100
Above standard (> 1,750,000)	0	0
ADL status of the treated spouse		
"Mild" dependency	23	76.7
"Moderate" dependency	2	6.7
"Severe" dependency	1	3.3
"Total" dependency	4	13.3

Table 1 shows that mean age of caregivers was 71 years old. Caregivers were either husband or wife of the ill one (53.3% and 46.7% respectively). They treated for spouse with different level of ADL dependency, but majority was mild one (76.7%). In regards to education, most of caregivers had a low level of education either just graduate from primary school (80%) or even never taken any formal education (10%). All caregivers in this study also had low social economic status and majority of them had no paid job (70%).

Table 2. Caregiver's Stress Level (n = 30)

Category	Frequency	Percentage
Low	8	26.6
Moderate	17	56.7
High	5	16.7

Table 2 shows that most caregivers in this study experienced significant stress when caring for their ill spouses. This was indicated by more than half of them experiencing moderate level of stress (56.7%), and some caregivers even experiencing a severe stress (16.7%).

Category	Frequency	Percentage
High	1	3.3
Moderate	14	46.7
Low	15	50.0

Table 3. Perceived Social Support (n = 30)

Table 3 shows that most caregivers tended to not receive significant amount of support. Half of the caregivers received low social support and just very few of them had the high one. The remaining had moderate level of social support (46.7%)

Table 4. Relationship between social support and stress level (n = 30)

Stress				
Low f(%)	Moderate High f(%) f(%)	High	Total	P Value
		f(%)	f(%)	
0(0)	0 (0)	1(100)	1(100)	
3(21.4)	9(64.3)	2(14.3)	14(100)	0.289
5(33.3)	8(53.3)	2(13.4)	15(100)	
	Low f(%) 0(0) 3(21.4)	Low Moderate f(%) f(%) 0(0) 0 (0) 3(21.4) 9(64.3)	Low Moderate High f(%) f(%) f(%) 0(0) 0 (0) 1(100) 3(21.4) 9(64.3) 2(14.3)	Low Moderate High Total f(%) f(%) f(%) f(%) 0(0) 0 (0) 1(100) 1(100) 3(21.4) 9(64.3) 2(14.3) 14(100)

Table 4 shows a cross tabulation between social support and stress level. However, Somer's d test showed that there was no significant relationship between those two variables (p>0.05).

Discussion

This study described perceived social support and stress level among elderly spouse in regards to caregiving and analysed relationship between those two variables. Results showed that most respondents experienced moderate and severe level of stress. This finding was similar to a previous study that found caregivers tend to experience moderate and severe level of stress while caring for the elderly (17). Caregiver felt a significant amount of stress due to a variety of factors. Caring for spouse with ADL dependence, caregivers often feel overwhelmed by the caregiving responsibility. Taking care for elderly with ADL dependence is most often very time consuming (17). This issue is significant especially for those who have other responsibilities than just caregiving. They have to spend time caring for their ill loved one and at the same times doing household chores or job responsibilities.

Besides, care giving process itself seems never ending due to the nature of diseases. Caregivers in this study had been in care giving role at least for six months. They didn't know how long they had to be in that role because their spouses were suffering from chronic debilitating diseases such as stroke and diabetes melitus. Moreover, elderly caregivers most often also had to deal with other situations than can worsen the stress, including decrease in body functions, psychological issues, and changes in social and economic functions (10,11).

Caregiving-related stress that continues without relief can lead to physical problems such as fatique and sleep problem, psychological problems such as anxiety and depression, and social problems such as impaired social interaction and isolation (18). It was interesting that some caregivers in this study only experienced low level of stress. Possible explanation for this finding was that caregiving seen as expression of love and devotion to the loved one. Thus, they were likely to appraise the carevigiving experience in a positive way which can hinder them from extreme stress. In regards to social support, findings showed that caregivers in this study were more likely to have less social support. A study by Andriyani, Anggreny, & Utami demonstrated that majority of elderly had a lower level of support (66,7%) (19). The low level of social support received by the caregivers can be caused by the fact that the elderly caregivers have different conditions than the younger one. Husband or wife is considered to be the greatest source of support for a married couple. Thus, when their loved ones suffer from debilitating disease, they most often lose their main source of support (14). Family members particularly adult children also becomes the main source of support. Thus, when their adult children get married and start to live independently, they lose source of support (20). However, the availability of adult children in the family doesn't guarantee the availability

Page: 103

of support. It is very possible that they could not take part in care giving because of work responsibility. Meanwhile, married children are usually busy taking care for their own family. Friends and neighbours can also be source of social support. However, the elderly caregivers usually have entered retirement age so they might lose friends from workplace (21). Because of demanded caregiving tasks, many caregivers also left their social activates and had less social interaction with their neighbours. Besides, changes in social norms might cause people to mind their own bussiness only and neglect others (22). Those might cause caregivers were likely to be lack of social support from their neighbours. This study demonstrated that social support did not associate to caregiving-related stress. This finding seems not to support the theory of The Buffering Hypothesis by Cohen & Wills, which assume that the presence of a social support system helps buffer or shield, an individual from the negative impact of stressful events (12). Possible explanation for this finding was elderly caregivers might have specific characteristics that differ to other age groups which possibly influence their mechanisme to encounter stressfull life events. Elderly people are less likely to rely on external support, rather more on themselves, their most significant person, or even a superior being. The latter is known as spirituality. The stage of human development affects a person's spiritual level. As people get old, they reaches maturity in thinking and behaving (23). Spirituality can also give meaning to life, comfort and strength during difficult times, so extreme stress can be avoided (24). Even though majority of elderly in Indonesia is still living in a three generational type family, the present study has provided an insight about the issue around eldelry care giving and old age support. This issue is expected to increase along with the changing in preferred living arrangement. A previous study found that about 11% of adult children wanted their parents to live independently either alone or only with spouse in their old age (25).

Few limitations were identified in this study. First, the number of participants were considered small for a quantitative study. Second, type of living arrangement of participants were not considered.

Conclusions

Elderly spousal caregiver was likely to experience caregiving-related stress and lack of social support. Because social support was found not to relate to stress level, elderly caregivers are suggested to develop mechanisme to encounter stress which more focuse on self or a supreme being. Further researches which involving larger number of study participants with specified criteria are needed to produce a more conclusive findings.

Annals of Tropical Medicine and Public Health

References

 Dewi SR. Buku Ajar Keperawatan Gerontik. Yogyakarta: Deepublish; 2014.

2. United Nations Population Fund (UNFPA). UNFPA Monographs Series No.1: Indonesia on the Threshold of

Population Ageing. Indonesia: UNFPA; 2014.

3. Rohaedi S, Putri ST, Karimah AD. Tingkat Annendirian Lansia dalam Activities Daily Living di Panti Sosial Tresna Wredha Senja Rawi. J Pendidik Keperawatan Indones. 2016;2(1).

4. Given BA, Given CW, Sherwood P. The Challene of Quality Cancer Care for Family Caregivers. Semin Oncol Nurs. 2012;28(4):205–12.

5. Ayuningputri N, Maulana H. Persepsi Tekanan Terhadap Kesejahteraan Psikologis pada Pasangan Suami-Istri dengan Stroke. J Psikol Integr. 2014;2(2):27–34.

6. Diana CA, Sukarlan AD, Pohan LD. Hubungan antara Caregiver Strain dan Kepuasan Pernikahan pada Istri sebagai Spouse Caregiver dari Penderita Stroke. INSAN. 2012;14(03):171–8.

7. Putra DA. Gambaran Strategi Koping Family Caregiver pada Lansia dengan Hambatan Activity of Daily Living di Dusun Pringgading Guwosari Pajangan Bantul Yogyakarta. Sekolah Tinggi Ilmu Kesehatan Jenderal Achmad Yani; 2017.

 Tania E. Depresi pada Lansia yang Menjadi Caregiver Pasien Pasca-Stroke. J Kedokt Meditek. 2015;21(56):1– 15.

9. Gaol NTL. Teori Stres: Stimulus, Respons, dan Transaksional. Bul Psikol. 2016;24(1):1–11.

 Indriana Y, Kristiana IF, Sonda AA, Intanirian A. Tingkat Stres Lansia Di Panti Wredha "Pucang Gading" Semarang. J Psikol Undip. 2010;8(2):87–96.

11. Parasari GAT, Lestari MD. Hubungan Dukungan Sosial Keluarga dengan Tingkat Depresi pada Lansia di Kelurahan Sading. J Psikol Udayana. 2015;2(1):68–77.

12. Cohen S, Wills TA. Stress, Social Support, and the Buffering Hypothesis. 1985;

13. Sujatmiko EP. Dukungan Sosial pada Caregiver Pasangan Penderita Diabetes Mellitus Tipe 2. Universitas Muhammadiyah Surakarta; 2016.

14. Isnawati D, Suhariadi PHF. Hubungan antara Dukungan Sosial dengan Penyesuaian Diri Masa

Page: 104

http://doi.org/10.36295/AOTMPH.2021.7116

Persiapan Pensiun Pada Karyawan PT Pupuk Kaltim. J Psikol Ind dan Organ 2013;02(1):1–6.

15. Wulandari YY, Herawati, Setyowati A. Dukungan Sosial Keluarga dengan Tingkat Stres Anggota Keluarga yang Merawat Pasien Gangguan Jiwa. Dunia Keperawatan. 2016;4(2):133–7.

 Setiyani R, Sumarwati M, Ramawati D. Attitude Towards Future Elderly Support: A Study Among Indonesian Young Adults. Int J Res Med Sci. 2015;3(1):74–8.

17. Mubin MF, Livana, Mahmudah AR. Gambaran Tingkat Stres Keluarga Lansia. J Keperawatan. 2018;6(2):128–33.

18. Purdani KS, Agusman \overline{F} , Rahma N. Studi Deskriptif Beban Caregiver Insan Pasca Stroke Di Rumah Sakit Panti Wilasa dr . Cipto Semarang. J Ilm Keperawatan Dan Kesehat. 2016;2(1):8–13.

19. Andriyani R, Anggreny Y, Utami A. Hubungan Dukungan Keluarga Terhadap Depresi Dan Interaksi Sosial Pada Lansia. J Kesehat Masy. 2019;4(3):105–11. 20. Hernilawati. Konsep dan Proses Keperawatan Keluarga. Takalar: Pustaka As Salam; 2013.

21. Wiliyanarti PF. Lanjut Usia Perspektif dan Masalah. Surabaya: UMSurabaya Publishing; 2018.

22. Effendi TN. Budaya Gotong-Royong Masyarakat dalam Perubahan Sosial Saat Ini. J Pemikir Sosiol. 2013:2(1).

23. Syam A. Hubungan Antara Kesehatan Spiritual Dengan Kesehatan Jiwa Pada Lansia Muslim Di Sasana Tresna Werdha Kbrp Jakarta Timur. Universitas Indonesia; 2010.

24. Destarina V, Agrina, Dewi YI. Gambaran Spiritualitas Lansia Di Panti Sosial Tresna Werdha Khusnul Khotimah Pekanbaru. JOM Psik. 2014;1(2):1– 8.

25. Setiyani R, Sumarwati M, Ramawati D. Future Living Arrangement for Aging Parents and Its Associated Factors. J Keperawatan Soedirman [Internet].
12 9;14(3):210–7. Available from: http://jks.fikes.unsoed.ac.id/index.php/jks/article/view/1 196

Page: 105

Caring for a Spouse with ADL Dependency: Perceived Social Support and Stress Level among Elderly Spousal Caregivers

ORIGINALITY REPORT 5% 5% SIMILARITY INDEX PUBLICATIONS **INTERNET SOURCES** STUDENT PAPERS **PRIMARY SOURCES** Submitted to Universitas Jenderal Soedirman 2% Student Paper www.scielo.br **í** % 2 Internet Source repository.uhamka.ac.id 1% 3 Internet Source nursing.ui.ac.id 1% 4 Internet Source es.scribd.com 1% 5 Internet Source pdfs.semanticscholar.org **1**% 6 Internet Source repositori.usu.ac.id 1% Internet Source scholar.unand.ac.id % 8 Internet Source repository.unjaya.ac.id 9 % Internet Source

10	revalidatiegeneeskunde.nl	1%
11	Sunardi Sunardi, Renidayati Renidayati. "Peningkatan Kemandirian Lansia Melalui Okupasi Terapi dan Group Psychotheraphy di PSTW Sabai Nan Aluih Sicincin Padang Pariaman", Jurnal Sehat Mandiri, 2020 Publication	1 %
12	journal.poltekkes-mks.ac.id	1 %
13	repository.unsoed.ac.id	1%
14	repository.unika.ac.id	1%

Exclude quotes	On	Exclude matches	< 1%
Exclude bibliography	Off		