



# A concise review of the potential utilization based on bioactivity and pharmacological properties of the genus *Gelidium* (Gelidiales, Rhodophyta)

Maria Dyah Nur Meinita<sup>1,2</sup> · Dicky Harwanto<sup>3</sup> · Amron<sup>1</sup> · Md. Abdul Hannan<sup>4</sup> · Gwi-Taek Jeong<sup>5</sup> · Il Soo Moon<sup>6</sup> · Jae-Suk Choi<sup>7</sup>

Received: 4 October 2022 / Revised: 13 March 2023 / Accepted: 15 March 2023  
© The Author(s), under exclusive licence to Springer Nature B.V. 2023

## Abstract

Being an agarophyte, *Gelidium* is extensively exploited for its agar-producing capacity. However, the food value and medicinal importance of this red alga are not to be underestimated. The pharmacological properties of *Gelidium* species have potential for the pharmaceutical, nutraceutical and cosmeceutical industries. This review collects, identifies and analyzes comprehensively the studies that deal with the bioactive properties of *Gelidium* species in the last decade. The principal bioactive compounds of *Gelidium* include R-phycoerythrin, R-phycoyanin, alkaloids, terpenoids, tannins, flavonoids, saponins, coumarins, cardiac glycosides, and steroids. *Gelidium* in the form of extracts or isolated compounds have been reported to show antibacterial, antioxidant, anticancer, anti-inflammatory, anti-obesity, immunomodulatory, neuroprotective, and antidiabetic properties. Most of the evidence has been reported in various in vitro models. Therefore, further experiments using appropriate animal and human subjects are necessary to develop the preclinical findings into clinical use. The main challenge in developing bioactive compounds on a commercial scale is the sustainable supply of *Gelidium* biomass that requires integrated seaweed aquaculture through employing biotechnological approaches and effective utilization of industrial byproducts.

**Keywords** Seaweed · Macroalgae · Bioactive · Pharmacological · Agarophyte · *Gelidium* · Rhodophyta

## Introduction

Macroalgae are one of the most potential marine resources because of their polysaccharide and bioactive compounds (Zemke-White and Ohno 1999; Bixler and Porse 2011; Ale and Meyer 2013; Andrade et al. 2013). *Gelidium* is one of the agar-producing red macroalgae and has been used as sea vegetable, human food and as traditional herb medicine in East Asian countries. *Gelidium* has been used as a food ingredient for centuries by boiling it in water to make edible agar (McHugh 1991). It contains polysaccharides in the form of agar, which consist of agarose and a small mixture of agaropectin (Qi et al. 2008; Bixler and Porse 2011). Agarose is a neutral agar that consisted of a linear polymer of the disaccharide agarobiose consisting of 1,3-linked-D-galactose and 1,4-linked 3,6-anhydrous-L-galactose, while a sulphated galactan with some anionic structure such as small anionic amounts of pyruvic, D-glucuronic acid and acid ester sulfates build up agaropectin (Qi et al. 2008; Park et al. 2011). Polysaccharides from agarophytes have been used in industry for a long time (Bixler and Porse 2011). Agar production

✉ Maria Dyah Nur Meinita  
maria.meinita@unsoed.ac.id

<sup>1</sup> Faculty of Fisheries and Marine Science, Jenderal Soedirman University, Purwokerto 53123, Indonesia

<sup>2</sup> Center for Maritime Bioscience Studies, Jenderal Soedirman University, Purwokerto 53123, Indonesia

<sup>3</sup> Faculty of Fisheries and Marine Science, Diponegoro University, Semarang 50275, Indonesia

<sup>4</sup> Department of Biochemistry and Molecular Biology, Bangladesh Agricultural University, Mymensingh 2202, Bangladesh

<sup>5</sup> Department of Biotechnology, Pukyong National University, Busan 48513, Korea

<sup>6</sup> Department of Anatomy, College of Medicine, Dongguk University, Gyeongju 38066, Republic of Korea

<sup>7</sup> Department of Seafood Science and Technology, The Institute of Marine Industry, Gyeongsang National University, 38 Cheondaegukchi-Gil, Tongyeong-Si 53064, Gyeongsangnam-Do, Korea

increased by 7% from 2009 to 2015 with an increase in sales value of 6% (Porse and Rudolph 2017). *Gracilaria* and *Gelidium* species are the prominent agar-producing species in the world. The agar produced by *Gelidium* shows stronger gel strength compared to the agar produced by *Gracilaria*. Hence, *Gelidium* is widely used commercially and industrially as a reagent for the manufacture of bacto agar and agarose (Porse and Rudolph 2017). *Gelidium* also produces some bioactive compounds that potentially can be applied and developed in the pharmaceutical, food, and nutraceutical industries. These compounds include R-phycoerythrin, R-phycoyanin, alkaloids, terpenoids, tannins, flavonoids, saponins, coumarins, cardiac glycosides and steroids (Mohy El-Din and Alagawany 2019; Sukwong et al. 2019). The bioactivity of *Gelidium* bioactive compounds includes antibacterial, antioxidant, anticancer, antitumor, anti-inflammatory, immune-promoting, neuroprotective, and antidiabetic activities (Bouhlal et al. 2010; Rhimou et al. 2010a; Hannan et al. 2013; Oumaskour et al. 2013; Wang et al. 2013; Kang et al. 2016; Alghazeer et al. 2018; Saeed et al. 2020).

A comprehensive review of agarophyte macroalgae from the genus *Gracilaria* has been done by Torres et al. (2019). Previous reviews on the commercial utilization, cultivation, and harvest methods of the genus *Gelidium* were conducted by McHugh (1991), Santelices (1991), Melo (1998), Friedlander (2008), Porse and Rudolph (2017) and Santos and Melo (2018). However, there is no up-to-date comprehensive review on the bioactivity and pharmacological properties of *Gelidium*. The main aim of this review is to analyse the potential and challenge of the current utilization of *Gelidium* as a source of bioactive compounds. This study was done systematically by collecting, identifying, screening, and analysing scientific articles published during 2001–2021. This review gives updated information on *Gelidium* potential utilization and application in the industry. Hence, we can find the research gaps and future directions on the utilisation of *Gelidium*.

## Distribution

*Gelidium* belongs to the division Rhodophyta, class Floridophyceae, order Gelidiales, and family Gelidiaceae. *Gelidium* can be found both in the tropics and subtropics, with the highest diversity concentrated in the tropics. The highest producers of *Gelidium* species are Morocco, Japan, Republic of Korea, and Indonesia which produced 9,500 t dry weight of *Gelidium* or 85% of world total production (Porse and Rudolph 2017). *Gelidium* species also have been reported in other countries including Madagascar (Boo et al. 2022), Brazil (Iha et al. 2015), China (Fei and Huang 1991), India (Santelices 1991), Argentina (Croce et al. 2015) Portugal (Melo 1998), Spain (Melo 1998), South Africa (Melo 1998),


Australia (Kim and Boo 2012) and other locations as shown in Fig. 1.

## Morphology

Morphologically *Gelidium* can be identified based on the characteristics of the thallus including thallus tips, branching pattern, cell size, cystocarp, nutritive filament, pericarp, holdfast and apical cell. The genus *Gelidium* is one of the most diverse among the Gelidiaceae having high plasticity. Furthermore, within its wide geographical range and environmental conditions, *Gelidium* shows physiological and morphological adaptations (Santelices 1991). The tropical *Gelidium* species have a higher optimum temperature for growth compared to the subtropical species. Morphologically, *Gelidium* species have a cartilaginous thallus, red to deep purple or blackish color, arising from cylindrical or compressed, branched or unbranched creeping axes with numerous short haptera extending as individual axes or forming massive disc-like holdfasts disconnected from the interior of the cortical cells, are distichously, feathery or irregularly branched, with translucent rhizoidal filaments in the cortex and medulla, horned globose cystocarps and cruciate tetrasporangia arranged in sori (Santelices 2007; Kim et al. 2011; Guiry and Guiry 2022).

## Taxonomy

Currently, 144 *Gelidium* species are accepted in Algaebase, as well as 1 accepted subspecies, 16 accepted varieties, and 8 accepted formae (Guiry and Guiry 2022). *Gelidium* species show high plasticity of morphological features and characteristics, hence it is difficult to identify the species solely based on morphological features. Molecular approaches are useful tools to identify and analyse the phylogenetic relationship of the *Gelidium* species which will give information regarding taxonomy and evolutionary pattern. The use of DNA barcoding to identify *Gelidium* species is increasing. The plastid *RuBisCO* (*rbcL*), mitochondrial cytochrome *c* oxidase (*cox1/COI*) and the chloroplast universal plastid amplicon (UPA) are commonly used for identification and phylogenetic analysis of *Gelidium* species (Freshwater et al. 2010; Kim et al. 2012; Iha et al. 2015; López et al. 2017; Boo et al. 2022). Freshwater et al. (2010) compared *COI* and *rbcL* and verified the barcoding gap among six *Gelidium* species. Kim and Boo (2012) identified and analysed the phylogenetic relationship of two *Gelidium* species, namely *G. crinale* and *G. pusillum*, using mitochondrial *cox1* and plastid *rbcL* markers. They found that both species were different based on the dataset of mitochondrial *cox1* and plastid *rbcL* markers. The phylogenetic relationship of *G. crinale*



**Fig. 1** Geographic distribution of *Gelidium* in the world based on (A) top producers of *Gelidium* and (B) global distribution of *Gelidium* (Porse and Rudolph 2017; Guiry & Guiry 2022)

showed a kinship with *G. capense* and *G. coulteri*, while *G. pusillum*'s relationship and infraspecific classification were still not determined. Based on the morphological characteristic and DNA barcoding using mitochondrial *cox1* and plastid *rbcL*, Boo et al. (2022) observed the diversity of *Gelidium* species in southern Madagascar and identified some new records of *Gelidium* species, namely *G. usmanghanii*, *G. leptum* and *G. sclerophyllum*. Iha et al. (2015) used DNA barcoding of some Brazilian specimens of Gelidiales and to confirm three *Gelidium* species namely *G. crinale*, *G. floridanum*, *G. microdonticum*. Moreover, next-generation

genome sequencing has been used to obtain the complete mitochondrial genomes of *Gelidium* species (Yang et al. 2014; Boo et al. 2016).

## Cultivation

To date, the supply of *Gelidium* species is collected from the wild stock. Several cultivation methods of *Gelidium* have been developed to overcome the dependency on wild stock. The development of *Gelidium* cultivation for providing seedstock

using vegetative propagation and spore culture have been developed in laboratory, sea, pond, and tank culture systems (Fei and Huang 1991; Melo et al. 1991; Titlyanov and Titlyanova 2006; Titlyanov et al. 2006; Boulus et al. 2007; Friedlander 2008; Otaíza et al. 2018, 2019; Alemañ et al. 2019). Unfortunately, most of the cultivation technology of *Gelidium* is still at the experimental level and cannot fulfill the commercial demand. Plantlets of *Gelidium* can be also regenerated from callus and protoplasts through cell culture. However, there is a limited update on the cell culture of *Gelidium*. Cultivation experiments of *Gelidium* are commonly based on the fragmentation of thalli and spore. Otaíza et al. (2018) observed the fragmentation of *G. lingulatum* and investigated the optimum cultivation condition including the substrate, irradiance, and temperature. The effect of physical, chemical, and biological factors was also investigated in fragment thalli cultivation (Friedlander 2008; Otaíza et al. 2019). The cultivation from *Gelidium* fragment thalli showed a slow growth rate reaching 1.2% day<sup>-1</sup> (Otaíza et al. 2019), 2% day<sup>-1</sup> (Melo et al. 1991), 4.6% day<sup>-1</sup> (Fei and Huang 1991), and 11–13% day<sup>-1</sup> (Titlyanov and Titlyanova 2006). Besides the slow growth, the cultivation of thallus fragments is not environmentally friendly compared to spore culture. However, spore cultivation also has some limitations and drawbacks. Further studies on spore culture are needed before it can be applied. Studies on genetic engineering and efficient cultivation technology are needed to overcome the slow growth of *Gelidium*.

## Nutrient content

*Gelidium* species contain carbohydrate (40.64–67.3% dw), protein (11.31–15.6% dw), lipid (0.0–2.16% dw) and ash (5.5–24.74% dw) (Park et al. 2011; Siddique et al. 2013; Ashoush et al. 2017). Major minerals of *Gelidium* species are Ca (0.091–0.87% dw), Mg (0.045–0.23% dw), K (0.054–0.56% dw), Na (0.128–0.53% dw) (Panayotova and Stancheva 2013; Rubio et al. 2017). The monosaccharide composition of *Gelidium* species mainly consists of galactose, glucose, xylose, galacturonic acid and the sulfated polysaccharide (Park et al. 2012; Meinita et al. 2017; Cui et al. 2019).

## Agar characteristics

*Gelidium* has been used extensively to extract high-quality agar for biotechnological applications. Agar yield and characteristics such as gel strength, gelling temperature, and melting point vary depending on the extraction techniques, species, country of origin, and seaweed harvesting period (Table 1). In general, *Gelidium* species yield a better-quality agar in terms of gel strength compared to other agarophytes, however, the yield is relatively low. Although the quality of

*Gelidium* agar extracted by hot water extraction is adequate, several techniques, including alkaline pretreatment, enzymatic hydrolysis, ultrasonication, stirring, and autoclaving have been employed to increase agar yield and extract high-quality agar. Among seaweed species used for agar extraction, *Gelidium amansii* yields the highest quantity of agar (49.10% by sonication and autoclaving). Gómez Barrio et al. (2022) have shown that ultrasound-assisted extraction can improve yield and gel strength compared to conventional extraction. Alkali pretreatment followed by enzymatic hydrolysis or ultrasonication with *Gelidium corneum* (formerly *Gelidium sesquipedale*) can result in higher yield of agar and stronger gel strength compared to alkali pretreatment alone (Li et al. 2021). Again, higher yield of agar and stronger gel strength were observed in sonicated *G. amansii* when the autoclave method was applied rather than direct heating (Din et al. 2019).

## Utilization

*Gelidium* has been widely used, especially in East Asian countries including China, Japan and Korea. The utilization of *Gelidium* species has developed from the traditional use by the community to the modern utilization. Traditionally, macroalgae have been widely used by people in East Asian countries both for food and herbal medicine. People in South East Asian countries have long experience using *Gelidium* as food ingredient by boiling it in water to make edible agar and also by consuming *Gelidium* in the form of conventional strip and square (McHugh 1991; Armisen 1995). *Gelidium* is one of the macroalgae which has been commonly used in China as herbal medicine since two thousand years ago for stomach ailments, hemorrhoids, and anal fistulas (Chengkui et al. 1984).

In modern times, *Gelidium* is mainly used in food technology, pharmaceutical and bioenergy fields based on their primary and secondary metabolites. In bioenergy field, several studies have reported biohydrogen, bioethanol, biodiesel and platform chemical production from *Gelidium* (Wi et al. 2009; Afify et al. 2010; Jeong and Park 2010; Park et al. 2011; Meinita et al. 2013, 2021; Ra et al. 2015).

The research on bioactivity of *Gelidium* extracted compounds is also increasing during 2001–2021 as shown in Fig. 2a. We found 63 research articles focused on the bioactivities of *Gelidium* species. *Gelidium* has been widely studied for antibacterial, antioxidant, anticancer, antitumor, anti-inflammatory, immune-promoting, neuroprotective, antidiabetic activities (Fig. 2b).


Sixteen *Gelidium* species have been studied during 2001–2021 (Fig. 3). *Gelidium amansii* (formerly *Gelidium elegans*) was the most well-studied species (34.8%) in the genus *Gelidium*, followed by *Gelidium spinosum* (formerly *Gelidium latifolium*) (10.1%), *Gelidium corneum* (10.1%) and *Gelidium pristoides* (6.7%). The pharmacological

**Table 1** Yield, gel strength, gelling point, and melting point of agar extracted from various species of *Gelidium*

<i>Gelidium</i> species	Agar extraction technique	Agar yield (%)	Gel strength (g cm <sup>-2</sup> )	Gelling point	Melting point	Country	Ref
<i>Gelidium corneum</i>	Conventional extraction and ultrasound-assisted extraction	1.50–15.85	377 ± 136 and 438 ± 47	n.d.	n.d.	Ireland	(Gómez Barrio et al. 2022)
<i>Gelidium corneum</i>	Traditional extraction conditions (95 °C, 3 h)	18.4	~750	n.d.	n.d.	Portugal	(Gomes-Dias et al. 2022)
<i>Gelidium</i> sp.	Not mentioned	8.21	674	31.5 °C	80 °C	Indonesia	(Subaryono and Sinurat 2021)
<i>Gelidium corneum</i>	0.5% NaOH plus enzymatic hydrolysis and/or ultrasonication	3.3–18.2	119–670	n.d.	n.d.	Ireland	(Li et al. 2021)
<i>Gelidium corneum</i> ; <i>Gelidium microdon</i>	i) Alkali pretreatment using 10% NaOH solution and heated in a water bath at 90 °C for 2 h ii) Alkali pretreatment using 0.5% Na <sub>2</sub> CO <sub>3</sub> solution at 90 °C for 30 min	i) 6.20–20.50 ii) 12.23–17.73	528.55 ± 11.08 and 489.00 ± 19.41	34–39 °C	85.65–86.71 °C, 86.32–87.13 °C	Morocco	(Belattmania et al. 2021)
<i>Gelidium spinosum</i>	Not mentioned	32.79 ± 0.78 – 39.26 ± 0.37	206.83 ± 5.36 to 591.85 ± 2.42	47.46 ± 0.20 °C	93.86 ± 0.27 °C	Turkey	(Öğretmen and Kaya 2019)
<i>Gelidium amansii</i>	i) Sonication pretreatment using autoclaving method and ii) Direct heating	Autoclaving 49.10 and Direct heating 13.26	840	26.9 to 25.2 °C	51.4 °C to 68.0 °C	Malaysia	(Din et al. 2019)
<i>Gelidium spinosum</i>	Water bath plus magnetic stirrer extraction method	45.79 ± 0.85 and 37.29 ± 0.76	n.d.	39.06 ± 0.26 °C	94.8 ± 0.29 °C	Turkey	(Öğretmen and Duyar 2018)
<i>Gelidium amansii</i>	Not mentioned	3.5–4.8	742	25.80.1 °C	76.80.1 °C	Malaysia	(Chew et al. 2018)
<i>Gelidium</i> spp.	Alkaline treatment with 6% alkali at 80 °C for 1.5 h	23.14	1068.15	28.1–43.2 °C	96.4–99.6 °C	China	(Wang et al. 2017)
<i>Gelidium robustum</i>	0.1 M phosphate buffer, pH 6.3, for 12 h	35% to 37	205 ± 5 – 444 ± 25	33.3 °C–35.3 °C	91.9–95.2 °C	Mexico	(Hurtado et al. 2011)
<i>Gelidium canariense</i>	0.5% (w/v) solution of Na <sub>2</sub> CO <sub>3</sub> at 85–90 °C for 30 min	18 to 27.8	n.d.	n.d.	n.d.	Algeria	(Freile-Pelegrín et al. 1995)
<i>Gelidium spinosum</i>	Hot water extraction	26% — 42.5	400–800	n.d.	n.d.	France	(Mouradi-Givernaud et al. 1992)

\* n.d., not determined

**Fig. 2** The number of publications on *Gelidium* (A) based on the year and (B) based on the bioactivities



properties of *G. amansii* encompassed their bioactivity as anticancer, antimicrobial, anti-inflammatory, immunomodulatory, neuroprotective, antidiabetic, anti-photoaging, anti-obesity and other bioactivities. Most of studies focused on the antimicrobial (26.8%), antioxidant (15.9%) and neuroprotective (12.2%) activities of *Gelidium* species.

## Bioactive compounds

Macroalgae produces secondary metabolites in response to environmental stress. Macroalgae as sessile organisms produce secondary metabolites to survive from

environmental pressures. Macroalgae that live in different environmental conditions may also produce different types of bioactives. The types of bioactive compounds in *Gelidium* can be seen in Table 2.

The chemical structure of some bioactive compounds found in *Gelidium* can be seen in Fig. 4. Red algae contain photosynthetic pigments namely, phycoerythrin and phycocyanin (Ficner et al. 1992; Kohata et al. 2010). Phycocyanin and phycoerythrin are pigments in *Gelidium* that show bright colors (Pangestuti and Kim 2011). Phycocyanin is a blue pigment while phycoerythrin is a red pigment, and both are water-soluble compounds

(Sukwong et al. 2019). These compounds have remarkable antioxidant activity (Pan-utai and Iamtham 2019).

Mycosporine-like amino acids (MAAs) are small water soluble molecules categorised as secondary metabolites synthesized by macroalgae including *Gelidium* species that have bioactivity as photoprotectors and antioxidants (Pandey et al. 2017). MAAs such as shinorine and porphyra-334 isolated from *Gelidium* sp. have been investigated in the pharmaceutical field. Their immunomodulatory effects have been investigated in human myelomocytic cell lines and they have anti-inflammatory activity (Becker et al. 2016). Shinorine and porphyra-334 are MAAs which can be found in organism that live in the high intensity of sun exposure. Hence, they can be developed as UV protectors and cell proliferation activator in cosmetic products because they show a protective response on human fibroblast cells from UV induction (Oyamada et al. 2008; Lalegerie et al. 2019; Sen and Mallick 2021).

Cholesterol and lumichrome isolated from *G. microdon* have been shown to have antibacterial properties (Silva et al. 2013). Other compounds found in *Gelidium* are alkaloids, terpenoids, tannins, flavonoids, saponins, coumarins, cardiac glycosides, and steroids (Mohy El-Din and Alagawany 2019). Alkaloid compounds in macroalgae contain many biological amines and contain halogenated cyclic nitrogen-containing compounds (Güven et al. 2010). Bioactive compounds such as flavonoids, alkaloids, tannins, terpenoids, saponins, and steroids in macroalgae have the potential to be used as antioxidants and antibacterials (Taleb-Contini et al. 2003; Shan et al. 2007; Güven et al. 2010). Terpenoids play a role in the development of new pesticides and may become prototype agrochemical agents (Born et al. 2012). Coumarin is a natural product with pharmacological properties such as anti-inflammatory, antibacterial, antioxidant and neuroprotective properties (Kontogiorgis and Hadjipavlou-Litina 2005; Venugopala et al. 2013). Cardiac glycosides have been extensively studied for their anticancer activity (Kepp et al. 2012).

## Pharmacological properties

### Anticancer activity

*Gelidium* spp. extracts have shown anticancer properties (Table 3) and can be used as new chemotherapeutic agents or as a source of chemical leads for new agents (Rocha et al. 2018; Meinita et al. 2022). For example, an ethanol extract of *G. latifolium* showed inhibited human colorectal carcinoma (Caco-2) cell lines with a substantial inhibition (85%) at a concentration of 200  $\mu\text{g mL}^{-1}$ . However, the extract also had a cytotoxic effect on normal cells, namely Human Corneal Epithelial Cells (HCEC) with inhibition not exceeding

24% at a concentration of 200  $\mu\text{g mL}^{-1}$ . The  $\text{IC}_{50}$  against Caco-2 cell line was 120  $\mu\text{g mL}^{-1}$  which is classified as moderate cytotoxic activity (Alghazeer et al. 2018). Furthermore, polysaccharide extract from *G. corneum* showed potential cytotoxicity against a number of cancer cell lines (Abdala Díaz et al. 2019).


Chen et al. (2004) examined the effects of freeze-dried PBS and DMSO extracts from *G. amansii* agar which were tested on three cancer cell lines. The results showed that both extracts can inhibit Hepa-1 and NIH-3T3 cells, but not the growth of HL-60 cells. This study was confirmed by Xu et al. (2004) who reported that *G. amansii* extract had  $\text{IC}_{50}$  values greater than 50  $\mu\text{g mL}^{-1}$  against three cancer cell lines and one normal cell line. In addition, Silva et al. (2013) stated that crude methanol extracts from *G. microdon* showed moderate inhibition against MCF-7, NCI-H460 and A375-C5 cell lines with an  $\text{IC}_{50}$  ranging from 36 to 75  $\mu\text{g mL}^{-1}$ .

As shown above, extracts of several *Gelidium* species showed moderate and potential cytotoxicity against multiple cancer cell lines. However, the cytotoxicity of *Gelidium* extracts is not limited to cancer cells, but also affects normal cells. Thus further research is needed on the potential of using *Gelidium* species as a source of anticancer compounds.

### Antioxidant activity

Antioxidants play a role in the potential prevention of several diseases related to oxidative stress, occurring when the balance between antioxidants and reactive oxygen species (ROS) is disturbed (Corsetto et al. 2020). Table 4 summarizes the antioxidant activity of *Gelidium*. The antioxidant activity of *Gelidium* was mostly determining by evaluating DPPH (2,2-diphenyl-1-picrylhydrazyl) degradation. Based on research carried out, the degradation rate of *Gelidium* extract against DPPH is medium but its antioxidant activity is low (Alghazeer et al. 2018; Abdala Díaz et al. 2019; Nursid et al. 2020; Saeed et al. 2020). The antioxidant parameters based on  $\text{IC}_{50}$  are very strong (50 ppm), strong (50–100 ppm), moderate (100–150 ppm), and weak (> 200 ppm) (Molyneux 2004; Juszczak et al. 2019). Mohy El-Din and Alagawany (2019) showed that *G. crinale* extract had the lowest antioxidant effect compared to the other macroalgae using total antioxidant capacity (TAC) evaluation. This is consistent with recent findings (Lekameera et al. 2008) that brown algae have stronger antioxidant activity than red or green algae. *Gelidium* sp. showed a low antioxidant effect based on DPPH assay (Ishakani et al. 2016), FRAP method (Ortiz-Viedma et al. 2021), and ORAC assay (Pacheco et al. 2020) compared to other algae tested. On the other hand, *G. foliaceum*, *G. microdon*, and *G. pristoides* demonstrated the highest free radical scavenging activity of the macroalgae tested (Rengasamy et al. 2015;

**Fig. 3** The number of publications on *Gelidium* based on the species and their bioactivities




Paiva et al. 2016; Olasehinde et al. 2019b). In other studies of hydromethanolic, chloroform, diethyl ether, ethyl acetate, and n-butanol fractions of *G. corneum* possessed low antioxidant activity compared to synthetic antioxidants

BHT,  $\alpha$ -tocopherol and ascorbic acid (Metidji et al. 2015). Amongst the five extracts, the ethyl acetate fraction had the maximum activity (87.02%), followed by the diethyl ether fraction (83.52%), however, there was a substantial

**Table 2** Bioactive compounds of *Gelidium*

Species	Compound	Reference
<i>G. amansii</i>	R-phycoerythrin (R-PE)	(Baghel et al. 2015; Sukwong et al. 2019)
<i>G. amansii</i>	R-phycoyanin (R-PC)	(Baghel et al. 2015; Sukwong et al. 2019)
<i>Gelidium sp.</i>	Shinorine	(Pandey et al. 2017)
<i>Gelidium sp.</i>	Porphyra-334	(Pandey et al. 2017)
<i>G. microdon</i>	Cholesterol	(Silva et al. 2013)
<i>G. microdon</i>	Lumichrome	(Silva et al. 2013)
<i>G. crinale</i>	Alkaloids	(Metidji et al. 2015; Mohy El-Din and Alagawany 2019)
<i>G. crinale</i>	Terpenoids	(Mohy El-Din and Alagawany 2019)
<i>G. crinale</i>	Tannins	(Mohy El-Din and Alagawany 2019)
<i>G. crinale</i>	Flavonoids	(Mohy El-Din and Alagawany 2019)
<i>G. crinale</i> and <i>G. corneum</i>	Saponins	(Metidji et al. 2015; Mohy El-Din and Alagawany 2019)
<i>G. crinale</i>	Coumarins	(Mohy El-Din and Alagawany 2019)
<i>G. crinale</i>	Cardiac Glycosides	(Mohy El-Din and Alagawany 2019)
<i>G. crinale</i>	Steroids	(Mohy El-Din and Alagawany 2019)
<i>G. corneum</i>	Anthocyanine	(Metidji et al. 2015)

**Fig. 4** Structures of R-phycoerythrin (A), Phycocyanobilin (B) Shinorine (C) Porphyrin-334 (D) Cholesterol (E) Lumi-chrome (F) isolated from *Gelidium*



difference when compared to the positive control in the  $\beta$ -carotene-linoleate assay. Furthermore, MAAs extract namely asterina 330 plus palythine (AS-330 + PNE) from *G. corneum* revealed a significant activity of  $\beta$ -carotene oxidation inhibition referring to superoxide radical scavenging and vitamin E (de la Coba et al. 2009). Based on  $\beta$ -carotene-linoleate assay, *Gelidium* extracts have potential as an antioxidant agent.

### Antimicrobial activity

Antimicrobial activity against bacteria, fungi, and viruses has been studied (Table 5). Antibacterial activity was tested using the disc diffusion technique (Bouhlal et al. 2010; Rhimou et al. 2010a; Oumaskour et al. 2013; Saeed et al. 2020). *Gelidium* extracts show high inhibitory activity against both

Gram-positive (Rhimou et al. 2010a; Oumaskour et al. 2013) and Gram-negative bacteria (Bouhlal et al. 2010; Rhimou et al. 2010a). *Gelidium pusillum* showed antibacterial activity against all strains of bacteria tested and the highest activity was exhibited by *G. microdon* (as *G. spinulosum*) against *Staphylococcus aureus*. Among five *Gelidium* spp. tested, *G. attenuatum* possessed highest antibacterial activity, while *G. corneum* exhibited weak inhibition against all of the bacteria strains (Bouhlal et al. 2013). On the contrary, Oumaskour et al. (2013) observed that *G. corneum* showed highest inhibition and *G. spinosum* (formerly *G. latifolium*) possessed moderate inhibition. *Gelidium spinosum* showed weak and moderate inhibition against the bacteria tested but higher inhibition against *P. mirabilis* compared to positive control (Saeed et al. 2020). *Gelidium microdon* showed higher antibacterial activity after

**Table 3** Anticancer activities of *Gelidium* extracts

Species	Experimental Models	Extract or Constituent	Study Type	Optimum Dose ( $\mu\text{g mL}^{-1}$ )	IC <sub>50</sub> ( $\mu\text{g mL}^{-1}$ )	Ref
<i>G. spinosum</i>	Human colorectal carcinoma (Caco2) cell lines	Methanol extract	In vitro	200	120	(Alghazeer et al. 2018)
<i>G. corneum</i>	Human leukemia cell line (U-937), human colon cancer cell line (HTC-116), and human malignant melanoma (G-361)	Sulfated polysaccharides	In vitro	-	60.01 $\pm$ 4.84	(Abdala Díaz et al. 2019)
<i>G. amansii</i>	Murine hepatoma cells (Hepa-1c1c7), human leukemia cells (HL-60), and normal murine embryo fibroblast cells (NIH-3T3)	Phosphate-buffered saline (PBS) and dimethyl sulfoxide (DMSO) extract	In vitro	150	-	(Chen et al. 2004)
<i>G. microdon</i>	Breast adenocarcinoma (MCF-7), non-small cell lung cancer (NCI-H460) and melanoma (A375-C5)	Methanol extract	In vitro	-	36.3 $\pm$ 8.0 to $\geq$ 200	(Silva et al. 2013)
<i>G. amansii</i>	KB, hepatocellular carcinoma Bel7402 and lung cancer A549,	Methanol extract	In vitro	-	> 50	(Xu et al. 2004)

IC<sub>50</sub> < 100  $\mu\text{g mL}^{-1}$ , potential cytotoxicity; 100 < IC<sub>50</sub> < 1000  $\mu\text{g mL}^{-1}$ , moderate cytotoxicity; IC<sub>50</sub> > 1000  $\mu\text{g mL}^{-1}$ , non-toxic (Prayong et al. 2008)

removal of chlorophylls against methicillin resistant *S. aureus* (MRSA) compared to remaining bacteria (Silva et al. 2013).

Antifungal activity has been found in extracts of *G. microdon*, *G. spinosum* and *G. amansii*. *Gelidium microdon* showed higher antifungal activity after removal of chlorophylls against *Trichophyton rubrum* and *Microsporium canis* with minimal inhibitory concentration (MIC) and minimal lethal concentration (MLC) values ranging from 64.0 to 128.0  $\mu\text{g mL}^{-1}$  (Silva et al. 2013). Saeed et al. (2020) reported that *G. spinosum* extract showed weak inhibition against *Aspergillus fumigatus* and *Aspergillus niger* compared to ketoconazole. In addition, among four fungi tested, *G. amansii* exhibited highest inhibition against *Fusarium oxysporum* (Yi et al. 2001). Besides being able to show antibacterial and antifungal activities, *Gelidium* extracts also show antiviral. The methanol extract from *G. attenuatum* showed high antiviral activity against Herpes simplex virus type 1 with an EC<sub>50</sub> value of 9.3  $\mu\text{g mL}^{-1}$  (Rhimou et al. 2010b) and the *G. spinosum* n-butanol extract inhibited avian and human influenza virus replication (Kamenarska et al. 2009).

### Anti-inflammatory activity

The anti-inflammatory properties of *Gelidium* potentially can play a role in treating inflammation. Inflammation is natural defense mechanism of vascular tissue to antagonism agents which attack and weaken the immune system (de Cássia da Silveira e Sá et al. 2013).

Anti-inflammatory activity of *Gelidium* has been studied in vitro and in vivo (Table 6). In vitro studies by Cui et al. (2019) showed that a *G. pacificum* sulfated polysaccharide with a molecular weight of 28,807 Da protected human monocytic (THP-1) cells from LPS-stimulated cytotoxicity. A concentration of 5  $\mu\text{g mL}^{-1}$  suppressed mRNA and protein expression of the mRNA and protein expression of toll-like receptor-4 (TLR-4), myeloid differentiation factor (MyD88) and tumor necrosis factor receptor-associated factor-6 (TRAF-6). In another in vitro study, the dichloromethane/methanol (50:50) extract of *G. sesquipedale* inhibited phospholipase A2 by 100% and elastase by 95 (Oumaskour et al. 2013).

In vivo assay of sulfated galactan of *G. crinale* showed that in rat paw edema, there was an involvement of arachidonic acid metabolism (de Sousa et al. 2013). Agar free-*G. amansii* extracts also increased the production of anti-inflammatory cytokines (IL-10 antibodies) in diet-induced obese C57BL/6 J mice (Lee et al. 2018).

### Immunomodulatory effect

Immune response regulation is closely related to various physiological and pathological conditions (Wang et al. 2013). Immunomodulation has complex mechanisms that

**Table 4** Antioxidant activities of *Gelidium* extracts

Species	Extract or Constituent	Antioxidant Assay	DPPH	Effect	Ref
<i>G. corneum</i>	Sulfated polysaccharides	DPPH assay	40.21 ± 0.44%	Low antioxidant effect	(Abdala Díaz et al. 2019)
<i>G. spinosum</i>	Ethanol extract	DPPH assay	46.68%	Low antioxidant effect (IC <sub>50</sub> = 300 µg mL <sup>-1</sup> )	(Nursid et al. 2020)
<i>G. spinosum</i>	Methanol extract	DPPH assay	nd	Low antioxidant effect (IC <sub>50</sub> = 300 µg mL <sup>-1</sup> )	(Alghazeer et al. 2018)
<i>G. spinosum</i>	Ethanol, methanol, ethyl acetate, hexane, chloroform and acetone extract	DPPH assay	56.55 ± 1.20%	Low antioxidant effect (IC <sub>50</sub> = 8.78 mg mL <sup>-1</sup> )	(Saeed et al. 2020)
<i>G. chilense</i>	Phenolic extract	DPPH assay and FRAP method	51.2 ± 9.1%	Low antioxidant effect	(Ortiz-Viedma et al. 2021)
<i>G. pusillum</i>	Methanol extract	DPPH assay	11.50 + 0.43 mM eq ascorbic acid g <sup>-1</sup> DW	Low antioxidant effect	(Ishakani et al. 2016)
<i>G. foliaceum</i>	Methanol extract	DPPH assay	52.24 ± 0.86 mg mL <sup>-1</sup>	High free radical scavenging activity	(Rengasamy et al. 2015)
<i>G. corneum</i>	Asterina-330 plus palythine (AS-330 + PNE)	β-carotene/ linoleate bleaching and pyrogallol autooxidation method	80.0 to 85.0%	High activity β-carotene oxidation inhibition (IC <sub>50</sub> = 10 µM)	(de la Coba et al. 2009)
<i>Gelidium</i> sp.	Phenolic and acetone extracts	DPPH assay and ORAC assay	4.8 ± 0.4% and 4.7 ± 0.2%	Low antioxidant effect	(Pacheco et al. 2020)
<i>G. microdon</i>	Phenolic extract	DPPH assay	47.73%	High free radical scavenging activity	(Paiva et al. 2016)
<i>G. crinale</i>	Methanol extract	TAC evaluation	0.923 mgAAE g <sup>-1</sup>	Low antioxidant effect	(Mohy El-Din and Alagawany 2019)
<i>G. pristoides</i>	Phenolic extract	DPPH assay	153.22 µg mL <sup>-1</sup>	The scavenging activity higher than remaining seaweed tested	(Olasehinde et al. 2019b)
<i>G. corneum</i>	Hydromethanolic crude extract and four fractions (chloroform, diethyl ether, ethyl acetate, n-butanol)	DPPH assay and β-carotene/ linoleate bleaching	61.25%	Low antioxidant effect	(Metidji et al. 2015)

nd, not determined

regulate the pathophysiology and pathogenesis of various diseases that affect the immune system (Yoo et al. 2019). The bioactive compounds in *Gelidium* have been investigated to play a role in immune promoting. Table 7 summarises the immunomodulatory activity of *Gelidium*.

A number of inflammatory mediators such as tumor necrosis factor-α (TNF-α), prostaglandin and E2 (PGE2), interleukin (IL)-1b, and IL-6 and nitric oxide (NO), are produced from macrophages (Yun et al. 2008). *Gelidium* as an immune-promoter showed that *Gelidium* gel phosphate buffered saline (PBS) extract can activate RAW 264.7 macrophages by increasing the production of immune mediators and these inflammatory mediators (Wang et al. 2013). Furthermore, hot-water extract of *G. amansii* increased immune ability in white shrimp *Litopenaeus vannamei*. In contrast, sulfated polysaccharides of *G. corneum* showed low immunomodulatory activity compared to other species (Abdala Díaz et al. 2019).

## Neuroprotective activity

Neuroprotective activity is closely related to antioxidant activity, which plays a role in the defence system in the brain (Takahashi et al. 2003). Antioxidants play a role in defence while neuroprotective capacity plays a role in repairing neuronal damage in the brain (Hannan et al. 2020b). Marine algae, including *Gelidium* have been reported to have neuroprotective potential in various preclinical studies (Hannan et al. 2020a). Table 8 summarises the neuroprotective activity of *Gelidium*. A series of studies by Hannan and his team from 2012 to 2020 showed that *Gelidium* has neurotrophic and neuroprotective activities. Preliminary research suggested that *Gelidium* had highest neurotropic potential (Hannan et al. 2013). In subsequent studies, *Gelidium* has been shown to promote the morphological and functional maturation of neurons with the potential for prevention

Table 5 Antimicrobial activities of *Gelidium* extracts

Species	Extract or Constituent	Microorganism	Effect	Ref
<b>Antibacterial</b>				
<i>G. attenuatum</i>	Methanol extract	Gram positive: • <i>Staphylococcus aureus</i> • <i>Enterococcus faecalis</i> ATCC 29,212 • <i>Enterococcus faecalis</i> ATCC 29,213	Highest antibacterial activity with DIZ ranging from 15.0 to 35.0 mm	(Rhimou et al. 2010a)
<i>G. spinosum</i>		Gram negative: • <i>Escherichia coli</i> • <i>Klebsella pneumoniae</i>		
<i>G. pulchellum</i>		• <i>Escherichia coli</i> • <i>Salmonella typhi</i> <i>Klebsiella</i> sp. • <i>Pseudomonas aeruginosa</i>	Highest antibacterial activity with DIZ ranging from 15.0 to 34.0 mm	(Alghazeer et al. 2013)
<i>G. pusillum</i>		• <i>Bacillus</i> sp. • <i>Bacillus subtilis</i>		
<i>G. corneum</i>		• <i>Staphylococcus aureus</i> • <i>Staphylococcus epidermidis</i> • <i>Staphylococcus epidermidis</i> • <i>Bacillus</i> sp.	DIZ ranging from 7.0 to 19.5 mm	(Bouhlal et al. 2013)
<i>G. microdon</i>		• <i>Staphylococcus aureus</i> • <i>Staphylococcus epidermidis</i> • <i>Staphylococcus haemolyticus</i> • <i>Staphylococcus hominis</i> • <i>Staphylococcus simulans</i> • <i>Bacillus</i> sp. • <i>Cytophaga</i> sp. • <i>Micrococcus 5J6</i>		
<i>G. spinosum</i>	Methanol extract	• <i>Streptococcus faecalis</i> • <i>Bacillus cereus</i> • <i>Staphylococcus aureus</i> • <i>Staphylococcus aureus ssp aureus</i> • <i>Bacillus subtilis</i>	DIZ ranging from 9.0 to 12.0 mm	(Saeed et al. 2020)
<i>G. latifolium</i>	Dichloromethane-Methanol	• <i>Klebsella pneumoniae</i> • <i>Proteus mirabilis</i>	<i>G. latifolium</i> showed moderate inhibition while <i>G. sesquipedale</i> showed highest inhibition	(Oumaskour et al. 2013)
<i>G. corneum</i>				
<i>G. microdon</i>	Methanol extract	• <i>Staphylococcus aureus</i> ATCC 25,923 • <i>Methicillin Resistant Staphylococcus aureus</i> (MRSA) • <i>Staphylococcus aureus</i>	Higher antibacterial activity after removal of chlorophylls	(Silva et al. 2013)
<i>G. spinosum</i>	Butanol extract, water extract, and volatile compound	• <i>Escherichia coli</i> ATCC 25,922 • <i>Pseudomonas aeruginosa</i> ATCC 27,853	Volatile compound showed highest antibacterial activity	(Kamenarska et al. 2009)
<b>Antifungal</b>				
<i>G. microdon</i>	Methanol extract	<i>Epidermophyton floccosum</i> FF9, <i>Microsporium canis</i> FF1, <i>Microsporium gypseum</i> FF3, <i>Trichophyton mentagrophytes</i> FF7, and <i>Trichophyton rubrum</i> FF5	Higher antifungal activity after removal of chlorophylls	(Silva et al. 2013)
<i>G. spinosum</i>	Methanol extract	<i>Aspergillus fumigatus</i> and <i>Aspergillus niger</i>	Lower than positive control	(Saeed et al. 2020)
<i>G. amansii</i>	Ethanol extract	<i>Penicillium citrinum</i> , <i>Aspergillus niger</i> , <i>Fusarium oxysporum</i> , and <i>Alternaria dianthi</i>	Highest inhibition against <i>Fusarium oxysporum</i>	(Yi et al. 2001)
<b>Antiviral</b>				

Table 5 (continued)

Species	Extract or Constituent	Microorganism	Effect	Ref
<ul style="list-style-type: none"> <li>• <i>G. attenuatum</i></li> <li>• <i>G. spinosum</i></li> <li>• <i>G. pulchellum</i></li> <li>• <i>G. pusillum</i></li> <li>• <i>G. corneum</i></li> <li>• <i>G. microdon</i></li> </ul>	Methanol extract	<i>Herpes simplex virus type 1 (HSV-1)</i>	Methanol extract showed highest antiviral activity	(Rhimou et al. 2010b)
<i>G. spinosum</i>	Butanol extract and water extract	<i>Avian influenza virus A/Germany/27, strain Weybridge (A/Wey-bridge) and Human influenza virus A/PR8/34 (H1N1) (A/PR8)</i>	n-butanol extract inhibited influenza virus replication	(Kamenarska et al. 2009)

and treatment of neurodegenerative diseases (Hannan et al. 2014). Then, a proteomic study of the chloroform (CHCl<sub>3</sub>) subfraction and the ethyl acetate (EtOAc) fractions of *G. amansii* ethanol extract found that neurite development was enhanced by up-regulation of several proteins such as profilin 2a, septin 7, cdc42, protein phosphatase 2A, DA11, eukaryotic translation initiation factor 5A-1, and enolase (Hannan et al. 2017). In a hypoxia/reoxygenation-induced oxidative injury model of primary hippocampal neurons, *G. amansii* protected against oxidative damage through suppressing GluN2B expression (Hannan et al. 2020c).

Inhibition of acetylcholinesterase (AChE) activity is one of the commonly available treatment strategies against Alzheimer's disease (AD) and *G. pristoides* possessed moderate acetylcholinesterase (AChE) and butyrylcholinesterase (BChE) inhibitory activity with IC<sub>50</sub> values of 52.70 ± 1.85 and 58.28 ± 1.79 mg mL<sup>-1</sup> respectively (Olasehinde et al. 2019b). Rengasamy et al. (2015) showed that *G. foliaceum* had the lowest AChE inhibitory activity with IC<sub>50</sub> value of 0.16 ± 0.02 mg mL<sup>-1</sup> among the macroalgae tested. *Gelidium* and its bioactives can protect against Zn-induced neuronal damage in HT-22 cells through inhibiting apoptosis and oxidative damage, and regulating redox imbalance (Olasehinde et al. 2019c, 2020). Extracts of *G. pristoides* also inhibit β-amyloid aggregation and β-secretase and cholinesterase activities (Olasehinde et al. 2019a; d). Based on these data, *Gelidium* can be a potential source of bioactive metabolites with therapeutic significance against oxidative stress-related neurodegeneration, including ischemic stroke and neurodegenerative diseases (Hannan et al. 2020c).

### Antidiabetic activity

Natural antidiabetics are needed for the treatment of diabetes mellitus. Diabetes treatment uses insulin level regulation, such as diet modification and oral hypoglycemic drugs (Marles and Farnsworth 1995; Jung et al. 2006). Macroalgae have bioactive peptides that can play a role in antidiabetic activity (Admassu et al. 2018). Research on the potential antidiabetic activity of *Gelidium* is summarized in Table 9. Several in vitro and in vivo studies conducted on *Gelidium* as antidiabetic agent have been reported. The development of type II diabetes and the complications associated with the disease, including macro-vascular and micro-vascular diseases is strongly influenced by postprandial hyperglycemia (Ko et al. 2011). Inhibition of carbohydrate hydrolyzing enzymes in the digestive organs, such as α-glucosidase and α-amylase, is one therapeutic method for reducing postprandial hyperglycemia (Clissold and Edward 1988). Acetone extracts of *Gelidium* sp. decreased the α-amylase activity to 77.9 ± 2.1% and did not show significant α-glucosidase inhibition at concentrations of

**Table 6** Anti-inflammatory activities of *Gelidium* extracts

Species	Extract or Constituent	Experimental Models	Study Type	Optimum Dose	Effects		Ref
					Molecular	Cellular	
<i>G. sesquipedale</i>	Methanol, acetone, chloroform, hexane, dichloromethane/methanol extracts	PLA inhibition and elastase assay	In vitro	nd	↓ phospholipase enzyme A2 (100%) and elastase enzyme (95%)		(Oumaskour et al. 2013)
<i>G. pacificum</i>	Sulfated polysaccharide	Human monocytic (THP-1) cells	In vitro	5 µg mL <sup>-1</sup>	↓ LPS-stimulated cytotoxicity		(Cui et al. 2019)
<i>G. crinale</i>	Sulfated galactan	Rat paw edema	In vivo	1 µg mL <sup>-1</sup>		↓ histamine (49%) and phospholipase A2 (44%)	(de Sousa et al. 2013)
<i>G. amansii</i>	Cellulose nanocrystal	Human keratinocytes and mice skin	In vitro and in vivo	100 µg mL <sup>-1</sup>	↓ c-Jun-induced UVB phosphorylation		(So et al. 2021)
<i>G. amansi</i>	Ethanol extract	Diet-induced obese (DIO) C57BL/6 J mice	In vivo	500 µg mL <sup>-1</sup>		↑ production of anti-inflammatory cytokines (IL-10 antibodies)	(Lee et al. 2018)
<i>G. amansii</i>	Phosphate-buffered saline (PBS) or ethanol extract	RAW 264.7 macrophages	In vitro	nd	↓ cytokine production in LPS-stimulated RAW 264.7 macrophages		(Wang et al. 2013)

(1, 10, 100, 1000 µg dry extract mL<sup>-1</sup>) (Pacheco et al. 2020). Furthermore, *G. amansii* exhibited greater inhibition against α-amylase and α-glucosidase compared to acarbose, an oral hypoglycemic agent. *Gelidium amansii* extract administered to STZ-induced diabetic mice can lower the level of postprandial blood glucose (Park et al. 2017a). Ko et al. (2011) also found that *G. amansii* extract exhibited high α-glucosidase inhibitory effects (96.47%) using high throughput screening. *Gelidium elegans* extract also regulated glucose homeostasis and significantly lowered blood glucose level in ICR mice administered (Choi et al. 2017a, 2018). Furthermore, the *G. elegans* extract activated insulin receptor substrate-1 (IRS-1) and phosphoinositide 3-kinase (PI3K), thus enhancing the expression of glucose transporter type 4 (GLUT4). The extract also lowered mitogen-activated protein kinase (MAPK) activity. These findings demonstrated that upregulating insulin pathway and downregulating MAPK pathway by *Gelidium* extract potentially can be used for blood metabolism.

### Anti-photoaging effect

UV radiation is one of the most damaging environmental factors, causing a kind of skin damage known as photoaging. Photoaging produced by persistent UV exposure results in premature aging of the skin, which is indicated by dryness, coarse and fine wrinkle formation, laxity, and pigmentation (Xiang et al. 2011). Studies conducted on the anti-photoaging effect of *Gelidium* spp. have been reported (Table 10).

Tyrosinase inhibitors are important in cosmetics for depigmentation and may be clinically effective for the treatment of various dermatological conditions linked to melanin hyperpigmentation (Cabanes et al. 1994; Shiino et al. 2001). Previous studies have reported that *Gelidium* extracts have potential as a tyrosinase inhibitor. Nursid et al. (2020) found that *G. spinosum* extract showed low inhibition of tyrosinase (< 20%) similar to Khatulistiwa et al. (2020) who observed 38.54% tyrosinase inhibition. However, Cha et al. (2011) observed higher tyrosinase inhibition (73.87%) at 20 °C. The differences may be due to different environmental conditions the algae grew in, resulting in variation in biochemical and physiological traits of *Gelidium* species (Véliz et al. 2019).

Kim et al. (2014) conducted research on a fermented *G. amansii* and *Cirsium japonicum* (Angiosperm) extract mixture (FGCM) on UVB-induced skin photoaging on hairless mice. FGCM showed anti-photoaging activity by inhibiting UVB-induced matrix metalloproteinase-1 (MMP-1) expression and increased type I procollagen in vitro. Meanwhile, hairless mice administered orally with FGCM showed strong activity in decreasing the number and total depth of wrinkles on dorsal skin.

**Table 7** Immunomodulatory activity of *Gelidium* extracts

Species	Extract or Constituent	Experimental Models	Study Type	Optimum Dose	Effects		Ref
					Molecular	Cellular	
<i>G. amansii</i>	Phosphate-buffered saline (PBS) or ethanol extract	RAW 264.7 macrophages	In vitro	150 µg mL <sup>-1</sup>		↑production of immune mediators	(Wang et al. 2013)
<i>G. corneum</i>	Sulfated polysaccharides	human colon cancer cell line (HCT-116), human malignant melanoma (G-361), breast adenocarcinoma human (MCF-7), human leukemia cell line (U-937), and murine macrophage cell line (RAW 264.7)	In vitro	-	Low immunomodulatory activity		(Abdala Díaz et al. 2019)
<i>G. amansii</i>	Hot-water extract	White shrimp <i>Litopenaeus vannamei</i>	In vivo	1.0 - 2.0 g kg <sup>-1</sup> in the diet		↑ immune ability	(Fu et al. 2007)

According to So et al. (2021) *G. amansii* extract inhibited UVB-induced AP-1 and COX-2 gene and protein expression in HaCaT cells. UVB irradiation activated *G. amansii* extract to suppress c-Jun translocation from cytosol to nucleus. In HaCaT cells, *G. amansii* inhibited UVB-induced phosphorylation of extracellular signal-regulated kinases (ERKs1)/2/MEK/2/B-Raf, c-Jun N-terminal kinase (JNK1)/2/MKK4/7, Akt, and epidermal growth factor receptor (EGFR).

### Anti-obesity activity

Obesity is a global health care issue that is associated with metabolic diseases such as type II diabetes and hyperlipidemia (Aubin et al. 2008; Crawford et al. 2010). Obesity is caused by body fat accumulation in adipose tissue due to abnormalities in energy metabolism (Choi et al. 2017b). Fat accumulation is caused by two different types of white adipocyte development processes, hyperplasia (adipocyte number increase) and hypertrophy (adipocyte size increase) in white adipocyte tissue (Guilherme et al. 2008). Recent studies reported that *Gelidium* can be used as anti-obesity by reducing lipid accumulation and adipogenesis (Table 11). *Gelidium amansii* extract showed inhibited adipogenesis in 3T3-L1 preadipocytes (Lee et al. 2011), and prevent weight gain through modulation of the AMPK, PRDM16, and UCP-1 pathways in a mice model (Choi et al. 2017b). Moreover, *G. amansii* extract decreased blood glucose and serum insulin levels when administered on mice induced by obesity diet (Kang et al. 2016). This finding is in line with other reported research that *G. amansii* ethanol extracts can reduce

fat accumulation in a 3T3-L1 cell model (Seo et al. 2012), and reduce body and epididymal fat weights, as well as serum total cholesterol (TC) and triglyceride (TG) levels in an obese mice model (Kang et al. 2017; Park et al. 2017b). On the other hand, hot-water extracts of *G. amansii* (GHE), can lower plasma and liver lipids in an animal model by decreasing cholesterol absorption and enhancing bile acid and fecal fat excretion (Yang et al. 2019). The water soluble fibres in *G. amansii* (GHE) may help to reduce lipid accumulation in the liver and adipose tissues. Furthermore, *G. elegans* extract at concentration of 100 µg mL<sup>-1</sup> inhibited mRNA expression of PPARγ (50%) and C/EBPα (90%), thus enhancing SOD1, SOD2, Gpx, and GR levels significantly in 3T3-L1 preadipocytes and RAW 264.7 cells (Jeon et al. 2014). The extract also inhibited the expression of C/EBPβ and increased C/EBP homologous protein 10 (Choi et al. 2016). These results suggested that *G. amansii* extract has the ability to inhibit adipogenesis and fat accumulation in 3T3-L1 preadipocytes and RAW 264.7 cells. The extract reduced weight gain of high-fat diet-induced obese mice and hamster through upregulating AMPK phosphorylation, PPARα, PRDM16, UCP-1, and UCP-2 pathways. However, further clinical studies are urgently needed to determine the effect of *Gelidium* treatment on obese patients in the future.

### Miscellaneous effects

Besides the above bioactivities, *Gelidium* also shows other biological activities (Table 12). Kim et al. (2016) evaluated the genotoxic effect of *G. elegans* in vitro and in vivo using five different strains of bacteria, CHL/IU cell lines, and ICR mice. In

**Table 8** Neuroprotective activities of *Gelidium* extract

Species	Extractor/Constituent	Experimental Models	Study Type	Optimum Dose	Effects		Ref
					Molecular	Cellular	
<i>G. amansii</i>	Ethanol extract	Rat hippocampal neurons	In vitro	15 $\mu\text{g mL}^{-1}$	-	↑ early neuronal differentiation ↑ the axonal and dendritic development and maturation index of neurons	(Hannan et al. 2013)
<i>G. amansii</i>	Ethanol extract	Rat hippocampal neurons	In vitro	15 $\mu\text{g mL}^{-1}$	-	↑ number of filopodia and dendritic spines ↑ synaptogenesis, modulate NMDA receptor recruitment	(Hannan et al. 2014)
<i>G. amansii</i>	Ethanol extract	Rat hippocampal neurons	In vitro	7.5 $\mu\text{g mL}^{-1}$	-	↑ neurite development by up- or down-regulating several key proteins	(Hannan et al. 2017)
<i>G. amansii</i>	Ethanol extract	Rat hippocampal neurons	In vitro	15 $\mu\text{g mL}^{-1}$	-	potential source of oxidative stress-related neurodegeneration	(Hannan et al. 2020c)
<i>G. foliaceum</i>	Methanol extract	-	In vitro	1.0 mg $\text{mL}^{-1}$	Low inhibition of AChE	-	(Rengasamy et al. 2015)
<i>G. pristoides</i>	Water extract	-	In vitro	21 to 84 $\mu\text{g mL}^{-1}$	AChE and BChE inhibitory	-	(Olasehinde et al. 2019b)
<i>G. pristoides</i>	Sulfated polysaccharide	Hippocampal neuron cell line cells (HT-22)	In vitro	-	↓ apoptosis, oxidative damage and AChE	-	(Olasehinde et al. 2020)
<i>G. pristoides</i>	Aquaeosus-ethanol extract	-	In vitro	-	↑ antioxidant enzyme activities and GSH levels ↑ in malondialdehyde and nitric oxide levels	-	(Olasehinde et al. 2019c)
<i>G. pristoides</i>	Aquaeosus-ethanol extract	-	In vitro	-	↓ $\beta$ -amyloid aggregation, $\beta$ -secretase, and cholinesterases	-	(Olasehinde et al. 2019d)
<i>G. pristoides</i>	Sulfated polysaccharide	-	In vitro	-	↓ $\beta$ -amyloid aggregation, $\beta$ -secretase, and cholinesterases	-	(Olasehinde et al. 2019a)

the in vitro experiment, there was no increase of mutation and chromosome aberration of bacteria at a concentration of 5000  $\mu\text{g}$  per plate. Furthermore, in vivo study showed that *G. elegans* did not affect the frequency of micronucleated bone marrow

polychromatic erythrocytes (MNPCE). These results suggest that *G. elegans* extract could be considered a safe dietary ingredient.

The coagulation of blood as a result of haemostasis failure causes clogging of the vascular system and is occasionally

**Table 9** Antidiabetic activities of *Gelidium* extract

Species	Extract or Constituent	Experimental Models	Study Type	Optimum Dose	Effects		Ref
					Molecular	Cellular	
<i>Gelidium</i> sp.	Ethanol/ water hot pressurized liquid extraction (HPLC) and Acetone extract	Human colon carcinoma (HT-29) cell line	In vitro	1–1000 µg dry extract mL <sup>-1</sup>	↓ α-amylase activity No effect in α-glucosidase activity		(Pacheco et al. 2020)
<i>G. amansii</i>	Ethanol extract	ICR mice	In vivo			↓ α-amylase and α-glucosidase activities ↓ postprandial blood glucose levels	(Park et al. 2017a)
<i>G. elegans</i>	Industrial extract (Newtree co., Ltd)	ICR mice	In vivo	200 mg kg <sup>-1</sup>		↓ blood glucose levels ↑ glucose homeostasis	(Choi et al. 2017a)
<i>G. amansii</i>	Methanol extract	-	In vitro	-	↓ α-glucosidase activity		(Ko et al. 2011)
<i>G. elegans</i>	Industrial extract (Newtree co., Ltd)	C57BL/KsJ-db/db mice and C57BL/KsJ m +/+ db mice	In vivo	50 mg kg <sup>-1</sup> day <sup>-1</sup>		↓ blood glucose ↑ activated IRS-1 ↓ activation of MAPK pathways	(Choi et al. 2018)

fatal. Activated partial thromboplastin time (APTT) and prothrombin time (PT) experiments revealed that *G. crinale* extracts had anticoagulant properties (Mohy El-Din and Alagawany 2019). This study suggested that the presence of phenols or their derivatives promote the anticoagulant effect. Angiotensin I-converting enzyme (ACE) is an enzyme that regulates blood pressure through the rennin-angiotensin

system, which is found in the vascular endothelium lining of the lungs (Cha et al. 2006). In humans ACE inhibitors have been widely utilized to treat essential hypertension and heart failure. Both the aqueous extract and the methanol extract of *G. amansii* was shown to have low ACE inhibitory activity (9.0–18.0%) compared to other macroalgae tested.

**Table 10** Anti-photoaging effects of *Gelidium* extract

Species	Extract or Constituent	Experimental Models	Study Type	Optimum Dose	Effects		Ref
					Molecular	Cellular	
<i>G. spinosum</i>	Ethanol extract	Tyrosinase enzyme	In vitro	1000 ppm	Low inhibition of tyrosinase	-	(Nursid et al. 2020)
<i>G. amansii</i>	Aqueous extract	Murine B16F10 melanocyte	In vitro	-	High inhibition of tyrosinase	-	(Cha et al. 2011)
<i>G. amansii</i>	Ethanol extract	SKH-1 hairless mice and Hs68 human dermal fibroblasts	In vitro and in vivo	200 µL of PBS + 500 mg FGCM per kg body weight	↑ type I procollagen levels ↓ UVB-induced MMP-1 expression	↓ number and total depth of wrinkles	(Kim et al. 2014)
<i>Gelidium</i> sp.	Water extract	Tyrosinase enzyme	In vitro	-	Low inhibition of tyrosinase	-	(Khatulistiwa et al. 2020)
<i>G. amansii</i>	Water extract	Human keratinocyte HaCaT, Beas-2B, and Raw 264.7 cells	In vitro	100 µg mL <sup>-1</sup>	↓ COX-2 expression and UVB-induced AP-1	-	(So et al. 2021)

PBS phosphate-buffered saline

**Table 11** Anti-obesity effects of *Gelidium* extract

Species	Extract or Constituent	Experimental Models	Study Type	Optimum Dose	Effects		Ref
					Molecular	Cellular	
<i>G. elegans</i>	Ethanol extract	3T3-L1 preadipocytes and RAW264.7 cells	In vitro	100 µg mL <sup>-1</sup>	↓ lipid accumulation ↓ mRNA expression of PPARγ and C/EBPα ↓ ROS production and NO level ↑ SOD1, SOD2, Gpx, and GR levels	-	(Jeon et al. 2014)
<i>G. amansii</i>	Ethanol extract	3T3-L1 preadipocytes and C57BL/6 mice	In vitro and in vivo	100 µg mL <sup>-1</sup>	↓ differentiation of 3T3-L1	↓ blood glucose and serum insulin levels	(Kang et al. 2016)
<i>G. amansii</i>	Water extract	Syrian hamsters	In vivo	200 mg kg <sup>-1</sup> diet		↓ triglyceride and total cholesterol ↑ AMPK phosphorylation, PPARα and UCP-2	(Yang et al. 2019) (Lee et al. 2011)
<i>G. amansii</i>	Ethanol extract	3T3-L1 adipocytes	In vitro	100 µg mL <sup>-1</sup>	↓ fat accumulation ↓ differentiation of 3T3-L1		(Choi et al. 2017b)
<i>G. amansii</i>	Industrial extract (Newtree co., Ltd)	ICR mice	In vivo	50, 200 mg kg <sup>-1</sup> day <sup>-1</sup>		↑ AMPK, PRDM16, and UCP-1	(Choi et al. 2017b)
<i>G. amansii</i>	Ethanol extract	Epididymal adipose tissue and liver; C57BL/6 J mice	In vitro and in vivo	0.75 to 2.00 g per 100 g diet	↓ FAS, SREBP-1c, PPARγ, and C/EBPα ↑ HSL and phospho-AMPKα (p-AMPK)	↓ body and epididymal fat weights, TC and TG levels	(Kang et al. 2017)
<i>G. amansii</i>	Ethanol extract	C57BL/6 J lean mice and C57BL/6 J-ob/ob mice	In vivo	0.5%		↓ body weight, epididymal adipose tissue weight, plasma triglycerides, and hepatic lipid accumulation	(Park et al. 2017b)
<i>G. amansii</i>	Industrial extract (Newtree co., Ltd)	3T3-L1 pre-adipocytes	In vitro	100 µg mL <sup>-1</sup>	↑ C/EBP homologous protein 10 ↓ the expression of C/EBPβ		(Choi et al. 2016)
<i>G. amansii</i>	Ethanol extract	3T3-L1 pre-adipocytes	In vitro	40 µg mL <sup>-1</sup>	↓ PPARγ and aP2 ↓ mRNA levels of a ROS-generator, NOX4 ↑ levels of antioxidant enzymes including SOD1/2, Gpx, and GR		(Seo et al. 2012)

Table 12 Miscellaneous effects of *Gelidium* extract

Bioactivity	Species	Extract or Constituent	Experimental Models	Study Type	Optimum Dose	Effects		Ref
						Molecular	Cellular	
Genotoxic effect	<i>G. elegans</i>	Industrial extract (Newtree co., Ltd)	<i>S. typhimurium</i> TA98, TA100, TA1535, and TA1537, <i>E. coli</i> WP2uvrA, and hamster lung cell line (CHL/IU); ICR mice	In vitro and in vivo	5000 µg plate <sup>-1</sup>	No increase in reverse mutation and chromosome aberration	No effect on MNPCE	(Kim et al. 2016)
Anticoagulant	<i>G. crinale</i>	Sulphated polysaccharide extract	Normal plasma	In vitro	50 µL	Second highest anticoagulant effect after <i>S. hornschurchii</i>		(Mohy El-Din and Alagawany 2019)
Hypertension effect	<i>G. amansii</i>	Methanol extract	-	In vitro	200 µg mL <sup>-1</sup>	Low ACE inhibitory activity (9.0–18.0%)		(Cha et al. 2006)
Antibiotic	<i>G. pacificum</i>	Water and ethanol extract	C57BL/6 mice	In vivo	-		Promoted antibiotic effect on mice with antibiotic-associated diarrhea (AAD)	(Cui et al. 2020)


In clinical treatment, antibiotics are used to treat various bacterial infections and infectious diseases. According to Cui et al. (2020) the sulfated polysaccharide of *G. pacificum* had beneficial effects on mice with antibiotic-associated diarrhea (AAD). This study showed that there was an increase in the richness and diversity of the gut microbiome and mucosal barrier function. *Gelidium pacificum* also can reverse metabolic disorders, inhibited the levels of cytokines inflammation and upregulated the content of SCFAs.

## Prospects and Challenges

The various species of *Gelidium* are a substantial source of biomaterials, including hydrocolloids and bioactives with profound industrial and medicinal uses (Fig. 5). However, several factors need to be considered when developing *Gelidium* as a source of biomaterials at a larger scale. In the hydrocolloid industry, *Gelidium* species together with *Gracilaria* are two main agarophyte species that are used as raw materials in the agar industry. In 2015, the production of *Gelidium* species was 11,100 dry t and total production of *Gracilaria* was 114,100 dry t. These two agarophytes contribute 246 million US\$ in total sales values of agar (Porse and Rudolph 2017). Agar extracted from *Gelidium* has higher gel strength and lower gelling temperature compared to *Gracilaria* (McHugh 1991; Bixler and Porse 2011). Hence, *Gelidium* is mostly chosen as agar source for bacteriological and pharmaceutical purposes. Although *Gelidium* produces high-quality agar, there are some obstacles to the large-scale production of *Gelidium*. Moreover, the agar production from *Gelidium* is steadily declining as it relies on natural stock; hence the supply of *Gelidium* biomass remains the main issue.

The first challenge is to adopt cultivation technology for *Gelidium* species. To develop an efficient cultivation method, we need to investigate the life history of *Gelidium* species. Most of the life history, reproductive features and complexity of *Gelidium* are still unknown. Rico and Guiry (1997) observed the characteristics and structure of tetrasporophytes and gametophytes of *Gelidium* and also observed the cystocarp structure and the bisexual nature of the gametophyte. The complexity and uniqueness of *Gelidium*'s life history as well as sexual gametophytes and meiotic sporophyte is challenging to be studied. Environmental parameters, including photoperiod, irradiance levels, temperature, and nutrients of *Gelidium* are also important factors in developing the optimum cultivation method (McHugh 1991).

The second challenge is to identify the potential candidate species of *Gelidium* to produce the desired bioactive compound. Out of 144 *Gelidium* species, which have been submitted to AlgaeBase (<https://www.algaebase.org/>), only a few species have been confirmed taxonomically (Guiry and



**Fig. 5** The potential use, bioactive compounds and pharmacological properties of *Gelidium*

Guiry 2022). In addition, genetic studies should be done to identify the species and also to investigate the genome diversity of *Gelidium* species. Study of genome sequencing, strain development, and genome editing technologies are needed to resolve the limitation of *Gelidium* species in

producing bioactive compounds and other valuable products. Advanced tools for genetic engineering potentially can be used to enhance the capability of *Gelidium* in producing desired bioactive compounds.

The third challenge is to develop efficient and sustainable extraction and isolation techniques for bioactive compounds to fulfill industrial needs. Pharmaceutical, nutraceutical and cosmeceutical industries require efficient, affordable, high yields and sustainable extraction techniques. Advanced extraction techniques such as solvent, ultrasound-assisted (UAE), microwave-assisted (MAE), enzyme-assisted (EAE) and pressurized liquid (PLE) extractions (Nigam et al. 2021) need to be used. These advanced extraction techniques can enhance and improve the yield and productivity of bioactive compounds from *Gelidium* compared to conventional extraction techniques. Once these challenges are overcome, sustainable and productive bioactive production from *Gelidium* can be feasible.

## Conclusion

*Gelidium* spp. contain a wide variety of bioactive compounds including R-phycoerythrin, phycocyanobilin, shinorine, porphyra-334, cholesterol and lumichrome which have potential applications in pharmaceutical and nutraceutical industries. Evidence shows that *Gelidium* spp. and their bioactive compounds have pharmacological properties, including anticancer, antioxidant, antimicrobial, anti-inflammatory, immunomodulatory, neuroprotective, antidiabetic, anti-photoaging, anti-obesity and many other bioactivities, which suggest that these marine natural products can be utilized in the development of therapeutic agents against chronic human diseases and can be incorporated as functional food ingredients to sustain healthy living. As highlighted in the Prospects and Challenges, the biotechnological approaches need to be exploited towards a sustainable supply of *Gelidium* biomass to ensure efficient industrial application of *Gelidium*.

**Acknowledgements** The authors thank Pukyong National University, Silla University and Dongguk University, South Korea for the international collaboration.

**Author contributions** M.D.N.M: Investigation, Data curation, Analysis, Validation, Visualization, Writing—original draft; A, D.H and M.A.H: Investigation, Data curation, Writing—review & editing; J.-S.C: Conceptualization, Supervision, Methodology, Writing—review & editing; G.-T.J and I.-S.M: Investigation, Data curation, Supervision, Methodology, Writing—review & editing.

**Funding** This research was funded by Indonesian Ministry of Education, Culture, Research and Technology under Scheme for Academic Mobility and Exchange (SAME) number 109/D2.3/KK.0403.

**Data Availability** Data supporting reported results are available upon request.

## Declarations

**Competing interests** The authors declare no competing interests.

**Conflict of interest** The authors declare no conflict of interest. The authors agree to the submission of the manuscript for peer review and publication in Journal of Applied Phycology.

## References

- Abdala Díaz RT, Casas Arrojo V, Arrojo Agudo MA, Cárdenas C, Dobretsov S, Figueroa FL (2019) Immunomodulatory and antioxidant activities of sulfated polysaccharides from *Laminaria ochroleuca*, *Porphyra umbilicalis*, and *Gelidium corneum*. *Mar Biotechnol* 21:577–587
- Admassu H, Gasmalla MAA, Yang R, Zhao W (2018) Bioactive peptides derived from seaweed protein and their health benefits: anti-hypertensive, antioxidant, and antidiabetic properties. *J Food Sci* 83:6–16
- Afify AEMMR, Shalaby EA, Shanab SMM (2010) Enhancement of biodiesel production from different species of algae. *Grasas Aceites* 61:416–422
- Ale MT, Meyer AS (2013) Fucoidans from brown seaweeds: An update on structures, extraction techniques and use of enzymes as tools for structural elucidation. *RSC Adv* 3:8131–8141
- Alemañ AE, Robledo D, Hayashi L (2019) Development of seaweed cultivation in Latin America: current trends and future prospects. *Phycologia* 58:462–471
- Alghazeer R, Howell NK, El-Naili MB, Awayn N (2018) Anticancer and antioxidant activities of some algae from western Libyan coast. *Nat Sci* 10:232–246
- Andrade PB, Barbosa M, Matos RP, Lopes G, Vinholes J, Mouga T, Valentao P (2013) Valuable compounds in macroalgae extracts. *Food Chem* 138:1819–1828
- Armisen R (1995) World-wide use and importance of *Gracilaria*. *J Appl Phycol* 7:231–243
- Ashoush YA, El-sayed SM, Farid HE, Abd-elwahab MA (2017) Biochemical Studies on red algae *Gelidium* sp. grown in Egypt. *Chem Res J* 2:334–341
- Aubin M-C, Lajoie C, Clément R, Gosselin H, Calderone A, Perrault LP (2008) Female rats fed a high-fat diet were associated with vascular dysfunction and cardiac fibrosis in the absence of overt obesity and hyperlipidemia: Therapeutic potential of resveratrol. *J Pharmacol Exp Ther* 325:961–968
- Baghel RS, Trivedi N, Gupta V, Neori A, Reddy CRK, Lali A, Jha B (2015) Biorefining of marine macroalgal biomass for production of biofuel and commodity chemicals. *Green Chem* 17:2436–2443
- Becker K, Hartmann A, Ganzera M, Fuchs D, Gostner JM (2016) Immunomodulatory effects of the mycosporine-like amino acids shinorine and porphyra-334. *Mar Drugs* 14:229
- Belattmania Z, Bentiss F, Jama C, Nadri A, Reani A, Sabour B (2021) Spectroscopic characterization and gel properties of agar from two *Gelidium* species from the Atlantic coast of Morocco. *Biointerface Res Appl Chem* 11:12642–12652
- Bixler HJ, Porse H (2011) A decade of change in the seaweed hydrocolloids industry. *J Appl Phycol* 23:321–335
- Boo GH, Hughey JR, Miller KA, Boo SM (2016) Mitogenomes from type specimens, a genotyping tool for morphologically simple species: Ten genomes of agar-producing red algae. *Sci Rep* 6:35337
- Boo GH, Le Gall L, Hwang IK, Rousseau F, Yoon HS (2022) Species diversity of *Gelidium* from southern Madagascar evaluated by an integrative taxonomic approach. *Diversity* 14:826
- Born FS, Bianco ÉM, Da Camara CAG (2012) Acaricidal and repellent activity of terpenoids from seaweeds collected in Pernambuco, Brazil. *Nat Prod Commun* 7:463–466

- Bouhhal R, Riadi H, Bourgougnon N (2010) Antibacterial activity of the extracts of Rhodophyceae from the Atlantic and the Mediterranean coasts of Morocco. *Afri J Biotechnol* 9:7968–7975
- Bouhhal R, Riadi H, Bourgougnon N (2013) Antibacterial activity of the extracts of Rhodophyceae from the Atlantic and the Mediterranean coasts of Morocco. *J Microbiol Biotechnol Food Sci* 2:2431–2439
- Boulus A, Spaneir E, Friedlander M (2007) Effect of outdoor conditions on growth rate and chemical composition of *Gelidium crinale* in culture. *J Appl Phycol* 19:471–478
- Cabanes J, Chazarra S, Garcia-Carmona F (1994) Kojic acid, a cosmetic skin whitening agent, is a slow-binding inhibitor of catecholase activity of tyrosinase. *J Pharm Pharmacol* 46:982–985
- Cha S-H, Lee K-W, Jeon Y-J (2006) Screening of extracts from red algae in Jeju for potentials marine angiotensin-I converting enzyme (ACE) inhibitory activity. *Algae* 21:343–348
- Cha SH, Ko SC, Kim D, Jeon YJ (2011) Screening of marine algae for potential tyrosinase inhibitor: Those inhibitors reduced tyrosinase activity and melanin synthesis in zebrafish. *J Dermatol* 38:354–363
- Chen YH, Tu CJ, Wu HT (2004) Growth-inhibitory effects of the red alga *Gelidium amansii* on cultured cells. *Biol Pharm Bull* 27:180–184
- Chengkui Z, Tseng CK, Junfu Z, Chang CF (1984) Chinese seaweeds in herbal medicine. *Hydrobiologia* 116–117:152–154
- Chew KW, Show PL, Yap YJ, Juan JC, Phang SM, Ling TC, Chang J-S (2018) Sonication and grinding pre-treatments on *Gelidium amansii* seaweed for the extraction and characterization of Agarose. *Front Environ Sci Eng* 12:2
- Choi J, Kim K, Koh E, Lee B (2016) Altered *Gelidium elegans* extract-stimulated beige-like phenotype attenuates adipogenesis in 3T3-L1 Cells. *J Food Nutr Res* 4:448–453
- Choi J, Kim KJ, Koh EJ, Seo YJ, Lee BY (2017a) *Gelidium elegans* regulates blood glucose homeostasis in ICR mice. *J Food Nutr Res* 5:274–280
- Choi J, Kim KJ, Koh EJ, Lee BY (2018) *Gelidium elegans* extract ameliorates type 2 diabetes via regulation of MAPK and PI3K/Akt signaling. *Nutrients* 10:51
- Choi J, Kim KJ, Koh EJ, Lee BY (2017b) *Gelidium elegans* regulates the AMPK-PRDM16-UCP-1 pathway and has a synergistic effect with orlistat on obesity-associated features in mice fed a high-fat diet. *Nutrients* 9:342
- Clissold SP, Edward C (1988) Acarbose. A preliminary review of its pharmacodynamic and pharmacokinetic properties, and therapeutic potential. *Drugs* 35:214–243
- Corsetto PA, Montorfano G, Zava S, Colombo I, Ingadottir B, Jonsdottir R, Sveinsdottir K, Rizzo AM (2020) Characterization of antioxidant potential of seaweed extracts for enrichment of convenience food. *Antioxidants* 9:249
- Crawford AG, Cote C, Couto J, Daskiran M, Gunnarsson C, Haas K, Haas S, Nigam SC, Schuette R (2010) Prevalence of obesity, type II diabetes mellitus, hyperlipidemia, and hypertension in the United States: Findings from the GE Centricity electronic medical record database. *Popul Health Manag* 13:151–161
- Croce ME, Villar MA, Parodi ER (2015) Assessment of alternative sources of seaweed polysaccharides in Argentina: potentials of the agarophyte *Gelidium crinale* (Hare ex Turner) Gaillon (Rhodophyta, Gelidiales). *J Appl Phycol* 27:2099–2110
- Cui M, Wu J, Wang S, Shu H, Zhang M, Liu K, Liu K (2019) Characterization and anti-inflammatory effects of sulfated polysaccharide from the red seaweed *Gelidium pacificum* Okamura. *Int J Biol Macromol* 129:377–385
- Cui M, Zhou R, Wang Y, Zhang M, Liu K, Ma C (2020) Beneficial effects of sulfated polysaccharides from the red seaweed: *Gelidium pacificum* Okamura on mice with antibiotic-associated diarrhea. *Food Funct* 11:4625–4637
- de Cássia da Silveira e Sá R, Andrade LN, de Sousa DP (2013) A review on anti-inflammatory activity of monoterpenes. *Molecules* 18:1227–1254
- de la Coba F, Aguilera J, Figueroa FL, de Gálvez MV, Herrera E (2009) Antioxidant activity of mycosporine-like amino acids isolated from three red macroalgae and one marine lichen. *J Appl Phycol* 21:161–169
- de Sousa AA, Benevides NM, de Freitas PA, Fiúza FP, Queiroz MG, Morais TM, Pereira MG, Assreuy AM (2013) A report of a galactan from marine alga *Gelidium crinale* with in vivo anti-inflammatory and antinociceptive effects. *Fundam Clin Pharmacol* 27:173–180
- Din SS, Chew KW, Chang YK, Show PL, Phang SM, Juan JC (2019) Extraction of agar from *Euचेuma cottonii* and *Gelidium amansii* seaweeds with sonication pretreatment using autoclaving method. *J Oceanol Limnol* 37:871–880
- Fei XG, Huang LJ (1991) Artificial sporeling and field cultivation of *Gelidium* in China. *Hydrobiologia* 221:119–124
- Ficner R, Lobeck K, Schmidt G, Huber R (1992) Isolation, crystallization, crystal structure analysis and refinement of B-phycoerythrin from the red alga *Porphyridium sordidum* at 2.2 Å resolution. *J Mol Biol* 228:935–950
- Freile-Pelegrín Y, Robledo DR, García-Reina G (1995) Seasonal agar yield and quality in *Gelidium canariensis* (Grunow) Seoane-Camba (Gelidiales, Rhodophyta) from Gran Canaria, Spain. *J Appl Phycol* 7:141–144
- Freshwater DW, Tudor K, O'Shaughnessy K, Wysor B (2010) DNA barcoding in the red algal order Gelidiales: comparison of COI with *rbcL* and verification of the “barcoding gap.” *Cryptogam Algal* 31:435–449
- Friedlander M (2008) Advances in cultivation of Gelidiales. *J Appl Phycol* 20:451–456
- Fu YW, Hou WY, Yeh ST, Li CH, Chen JC (2007) The immunostimulatory effects of hot-water extract of *Gelidium amansii* via immersion, injection and dietary administrations on white shrimp *Litopenaeus vannamei* and its resistance against *Vibrio alginolyticus*. *Fish Shellfish Immunol* 22:673–685
- Gomes-Dias JS, Pereira SG, Teixeira JA, Rocha CMR (2022) Hydrothermal treatments – A quick and efficient alternative for agar extraction from *Gelidium sesquipedale*. *Food Hydrocoll* 132:107898
- Gómez Barrio LP, Cabral EM, Zhao M, Álvarez García C, Senthamarakannan R, Padamati RB, Tiwari U, Curtin JF, Tiwari BK (2022) Comparison study of an optimized ultrasound-based method versus an optimized conventional method for agar extraction, and protein co-extraction, from *Gelidium sesquipedale*. *Foods* 11:805
- Guilherme A, Virbasius JV, Puri V, Czech MP (2008) Adipocyte dysfunctions linking obesity to insulin resistance and type 2 diabetes. *Nat Rev Mol Cell Biol* 9:367–377
- Guiry, M.D. & Guiry GM (2022) AlgaeBase. World-wide electronic publication, National University of Ireland, Galway. <https://www.algaebase.org>. Accessed 23 Dec 2022
- Güven KC, Percot A, Sezik E (2010) Alkaloids in marine algae. *Mar Drugs* 8:269–284
- Hannan MA, Dash R, Haque MN, Mohibullah M, Sohag AAM, Rahman MA, Uddin MJ, Alam M, Moon IS (2020a) Neuroprotective potentials of marine algae and their bioactive metabolites: pharmacological insights and therapeutic advances. *Mar Drugs* 18:347
- Hannan MA, Dash R, Sohag AAM, Haque MN, Moon IS (2020b) Neuroprotection against oxidative stress: Phytochemicals targeting TrkB signaling and the Nrf2-ARE antioxidant system. *Front Mol Neurosci* 13:116
- Hannan MA, Haque MN, Mohibullah M, Dash R, Hong YK, Moon IS (2020c) *Gelidium amansii* attenuates hypoxia/

- reoxygenation-induced oxidative injury in primary hippocampal neurons through suppressing GluN2B expression. *Antioxidants* 9:223
- Hannan A, Kang JY, Hong YK, Lee H, Choi JS, Choi IS, Moon IS (2013) The marine alga *Gelidium amansii* promotes the development and complexity of neuronal cytoarchitecture. *Phyther Res* 27:21–29
- Hannan MA, Mohibullah M, Hong YK, Nam JH, Moon IS (2014) *Gelidium amansii* promotes dendritic spine morphology and synaptogenesis, and modulates NMDA receptor-mediated postsynaptic current. *Vitr Cell Dev Biol Anim* 50:445–452
- Hannan MA, Mohibullah M, Hong YK, Moon IS (2017) Proteomic analysis of the neurotrophic effect of *Gelidium amansii* in primary cultured neurons. *J Med Food* 20:279–287
- Hurtado MÁ, Manzano-Sarabia M, Hernández-Garibay E, Pancho-Ruiz I, Zertuche-González (2011) Latitudinal variations of the yield and quality of agar from *Gelidium robustum* (Gelidiales, Rhodophyta) from the main commercial harvest beds along the western coast of the Baja California Peninsula, Mexico. *J Appl Phycol* 23:727–734
- Iha C, Milstein D, Guimarães SMPB, Freshwater DW, Oliveira MC (2015) DNA barcoding reveals high diversity in the Gelidiales of the Brazilian southeast coast. *Bot Mar* 58:295–305
- Ishakani AH, Joshi NH, Ayaz M, Sumara K, Vadher KH (2016) Antioxidant potential, polyphenols and diphenol content of seaweed available at Veraval coast, Saurashtra region - Gujarat. *Indian J Sci Technol* 9:1–6
- Jeon HJ, Seo MJ, Choi HS, Lee OH, Lee BY (2014) *Gelidium elegans*, an edible red seaweed, and hesperidin inhibit lipid accumulation and production of reactive oxygen species and reactive nitrogen species in 3T3-L1 and RAW264.7 cells. *Phyther Res* 28:1701–1709
- Jeong GT, Park DH (2010) Production of sugars and levulinic acid from marine biomass *Gelidium amansii*. *Appl Biochem Biotechnol* 161:41–52
- Jung M, Park M, Lee H, Kang YH, Kang ES, Kim SK (2006) Antidiabetic agents from medicinal plants. *Curr Med Chem* 13:1203–1218
- Juszczak A, Ramos P, Szczolko W, Pilawa B, Stanisz BJ (2019) Can angiotensin-converting enzyme inhibitors interfere with the free radicals? Measurement of antioxidant capacity using DPPH radical reduction examined by UV-Vis method. *Acta Pol Pharm - Drug Res* 76:233–239
- Kamenarska Z, Serkedjieva J, Najdenski H, Stefanov K, Tsvtkova I, Dimitrova-Konaklieva S, popov S (2009) Antibacterial, antiviral, and cytotoxic activities of some red and brown seaweeds from the Black Sea. *Bot Mar* 52:80–86
- Kang JH, Lee HA, Kim HJ, Han JS (2017) *Gelidium amansii* extract ameliorates obesity by down-regulating adipogenic transcription factors in diet-induced obese mice. *Nutr Res Pract* 11:17–24
- Kang MC, Kang N, Kim SY, Lima IS, Ko SC, Kim YT, Kim YB, Jeung HD, Choi KS, Jeon YJ (2016) Popular edible seaweed, *Gelidium amansii* prevents against diet-induced obesity. *Food Chem Toxicol* 90:181–187
- Kepp O, Menger L, Vacchelli E, Adjemian S, Martins I, Ma Y, Sukkurwala AQ, Michaud M, Galluzzi L, Zitvogel L, Kroemer G (2012) Anticancer activity of cardiac glycosides: At the frontier between cell-autonomous and immunological effects. *Oncimmunology* 1:1640–1642
- Khatulistiani TS, Noviendri D, Munifah I and Melanie S (2020) IOP Conf Ser: Earth Environ Sci 404:012065
- Kim HM, Lee DE, Park SD, Kim YT, Kim YJ, Jeong JW, Lee JH, Jang SS, Chung DK, Sim JH, Hu CS (2014) Preventive effect of fermented *Gelidium amansii* and *Cirsium japonicum* extract mixture against UVB-induced skin photoaging in hairless mice. *Food Sci Biotechnol* 23:623–631
- Kim K-M, Boo S-M (2012) Phylogenetic relationships and distribution of *Gelidium crinale* and *G. pusillum* (Gelidiales, Rhodophyta) using *cox1* and *rbcl* sequences. *Algae* 27:83–94
- Kim K, Choi J, Lee B (2016) Evaluation of the genotoxicity of a *Gelidium elegans* extract in vitro and in vivo. *J Food Nutr Res* 4:653–657
- Kim KM, Hoarau GG, Boo SM (2012) Genetic structure and distribution of *Gelidium elegans* (Gelidiales, Rhodophyta) in Korea based on mitochondrial *cox1* sequence data. *Aquat Bot* 98:27–33
- Kim KM, Hwang IK, Park JK, Boo SM (2011) A new agarophyte species, *Gelidium eucorneum* sp. nov. (Gelidiales, Rhodophyta), based on molecular and morphological data. *J Phycol* 47:904–910
- Ko SC, Lee SH, Kang SM, Ahn G, Cha SH, Jeon YJ (2011) Evaluation of  $\alpha$ -glucosidase inhibitory activity of Jeju seaweeds using high throughput screening (HTS) technique. *J Mar Biosci Biotechnol* 5:33–39
- Kohata S, Matsunaga N, Hamabe Y, Yumihara K, Sumi T (2010) Photo-stability of mixture of violet pigments phycoerythrin and phycocyanin extracted without separation from discolored nori seaweed. *Food Sci Technol Res* 16:617–620
- Kontogiorgis CA, Hadjipavlou-Litina DJ (2005) Synthesis and anti-inflammatory activity of coumarin derivatives. *J Med Chem* 48:6400–6408
- Lalegerie F, Lajili S, Bedoux G, Taupin L, Stiger-Pouvreau V, Connan S (2019) Photo-protective compounds in red macroalgae from Brittany: Considerable diversity in mycosporine-like amino acids (MAAs). *Mar Environ Res* 147:37–48
- Lee OH, Yoon KY, Kim KJ, You SG, Lee BY (2011) Seaweed extracts as a potential tool for the attenuation of oxidative damage in obesity-related pathologies. *J Phycol* 47:548–556
- Lee Y, Oh H, Lee M (2018) Anti-inflammatory effects of agar free-*Gelidium amansii* (GA) extracts in high-fat diet-induced obese mice. *Nutr Res Pract* 12:479–485
- Lekameera R, Vijayabaskar P, Somasundaram ST (2008) Evaluating antioxidant property of brown alga *Colpomenia sinuosa* (DERB. ET SOL). *Afr J Food Sci* 2:126–130
- Li Y, Zhao M, Gomez LP, Senthamaraikannan R, Padmanti RB, O'Donnell CP, Tiwari BK (2021) Investigation of enzyme-assisted methods combined with ultrasonication under a controlled alkali pretreatment for agar extraction from *Gelidium sesquipedale*. *Food Hydrocoll* 120:106905
- López BA, Tellier F, Retamal-Alarcón JC, Pérez-Araneda K, Fierro AO, Mcaya EC, Tala F, Thiel M (2017) Phylogeography of two intertidal seaweeds, *Gelidium linguatum* and *G. rex* (Rhodophyta: Gelidiales), along the South East Pacific: patterns explained by rafting dispersal? *Mar Biol* 164:188
- Marles RJ, Farnsworth NR (1995) Antidiabetic plants and their active constituents. *Phytomedicine* 2:137–189
- McHugh DJ (1991) Worldwide distribution of commercial resources of seaweeds including *Gelidium*. *Hydrobiologia* 221:19–29
- Meinita MDN, Amron A, Trianto A, Harwanto D, Caesarendra W, Jeong G-T, Choi J-S (2021) Levulinic acid production from macroalgae: Production and promising potential in industry. *Sustainability* 13:13919
- Meinita MDN, Harwanto D, Choi JS (2022) Seaweed exhibits therapeutic properties against chronic diseases: An overview. *Appl Sci* 12: 2638
- Meinita MDN, Marhaeni B, Hong YK, Jeong GT (2017) Enzymatic saccharification of agar waste from *Gracilaria verrucosa* and *Gelidium latifolium* for bioethanol production. *J Appl Phycol* 29:3201–3209
- Meinita MDN, Marhaeni B, Winanto T, Jeong GT, Kahn MNK, Hong YK (2013) Comparison of agarophytes (*Gelidium*, *Gracilaria*, and *Gracilariopsis*) as potential resources for bioethanol production. *J Appl Phycol* 25:1957–1961

- Melo RA (1998) *Gelidium* commercial exploitation: Natural resources and cultivation. *J Appl Phycol* 10:303–314
- Melo RA, Harger BWW, Neushul M (1991) *Gelidium* cultivation in the sea. *Hydrobiologia* 221:91–106
- Metidji H, Dob T, Toumi M, Krimat S, Ksouri A, Nouasri A (2015) In vitro screening of secondary metabolites and evaluation of antioxidant, antimicrobial and cytotoxic properties of *Gelidium sesquipedale* Thuret et Bornet red seaweed from Algeria. *J Mater Environ Sci* 6:3184–3196
- Mohy El-Din SM, Alagawany NI (2019) Phytochemical constituents and anticoagulation property of marine algae *Gelidium crinale*, *Sargassum hornschurchii* and *Ulva linza*. *Thalassas* 35:381–397
- Mouradi-Givernaud A, Givernaud T, Cosson J, Morvan H (1992) Agar from *Gelidium latifolium* (Rhodophyceae, Gelidiales): Biochemical composition and seasonal variations. *Bot Mar* 35:153–160
- Nigam S, Singh R, Bhardwaj SK, Sami R, Nikolova MP, Chavali M, Sinha S (2021) Perspective on the therapeutic applications of algal polysaccharides. *J Polym Environ* 30:785–809
- Nursid M, Khatulistiani TS, Novlendi D, Hapsari F, Hadiyati T (2020) Total phenolic content, antioxidant activity and tyrosinase inhibitor from marine red algae extract collected from Kupang, East Nusa Tenggara. *IOP Conf Ser Earth Environ Sci* 493:012013
- Öğretmen ÖY, Duyar HA (2018) The effect of different extraction methods and pre-treatments on agar yield and physico-chemical properties of *Gelidium latifolium* (Gelidiaceae, Rhodophyta) from Sinop Peninsula Coast of Black Sea, Turkey. *J Appl Phycol* 30:1355–1360
- Öğretmen ÖY, Kaya Y (2019) Seasonal changes in the yield and gel properties of agar extracted from *Gelidium latifolium* (Rhodophyta). *J Appl Phycol* 31:3091–3100
- Olasehinde TA, Mabinya LV, Olaniran AO, Okoh AI (2019a) Chemical characterization, antioxidant properties, cholinesterase inhibitory and anti-amyloidogenic activities of sulfated polysaccharides from some seaweeds. *Bioact Carbohydrates Diet Fibre* 18:100182
- Olasehinde TA, Olaniran AO, Okoh AI (2019b) Phenolic composition, antioxidant activity, anticholinesterase potential and modulatory effects of aqueous extracts of some seaweeds on  $\beta$ -amyloid aggregation and disaggregation. *Pharm Biol* 57:460–469
- Olasehinde TA, Olaniran AO, Okoh AI (2020) Sulfated polysaccharides of some seaweeds exhibit neuroprotection via mitigation of oxidative stress, cholinergic dysfunction and inhibition of Zn-induced neuronal damage in HT-22 cells. *BMC Complement Med Ther* 20:251
- Olasehinde TA, Olaniran AO, Okoh AI (2019c) Neuroprotective effects of some seaweeds against Zn-induced neuronal damage in HT-22 cells via modulation of redox imbalance, inhibition of apoptosis and acetylcholinesterase activity. *Metab Brain Dis* 34:1615–1627
- Olasehinde TA, Olaniran AO, Okoh AI (2019d) Aqueous-ethanol extracts of some South African seaweeds inhibit beta-amyloid aggregation, cholinesterases, and beta-secretase activities in vitro. *J Food Biochem* 43:e12870
- Ortiz-Viedma J, Aguilera JM, Flores M, Lemus-Mondaca R, Larrazabal MJ, Miranda JM, Aubourg SP (2021) Protective effect of red algae (Rhodophyta) extracts on essential dietary components of heat-treated salmon. *Antioxidants* 10:1108
- Otaíza RD, Cáceres JH, Rodríguez CY, Sanhueza ÁG (2019) Seeding of fragments of the agarophyte *Gelidium lingulatum* (Rhodophyta, Gelidiales) for the repopulation of lower levels of wave-exposed, intertidal rocky shores. *J Appl Phycol* 31:2133–2143
- Otaíza RD, Rodríguez CY, Cáceres JH, Sanhueza ÁG (2018) Fragmentation of thalli and secondary attachment of fragments of the agarophyte *Gelidium lingulatum* (Rhodophyta, Gelidiales). *J Appl Phycol* 30:1921–1931
- Oumaskour K, Boujaber N, Etahiri S, Assobhei O (2013) Anti-inflammatory and antimicrobial activities of twenty-three marine red algae from the coast of Sidi Bouzid (El Jadida-Morocco). *Int J Pharm Pharm Sci* 5:145–149
- Oyamada C, Kaneniwa M, Ebitani K, Murata M, Ishihara K (2008) Mycosporine-like amino acids extracted from scallop (*Patinopecten yessoensis*) ovaries: UV protection and growth stimulation activities on human cells. *Mar Biotechnol* 10:141–150
- Pacheco LV, Parada J, Pérez-Correa JR, Mariotti-Celis MS, Erpel F, Zambrano A, Palacios M (2020) Bioactive polyphenols from southern Chile seaweed as inhibitors of enzymes for starch digestion. *Mar Drugs* 18:353
- Paiva L, Lima E, Neto AI, Marcone M, Baptista J (2016) Health-promoting ingredients from four selected Azorean macroalgae. *Food Res Int* 89:432–438
- Pan-utai W, Iamtham S (2019) Extraction, purification and antioxidant activity of phycobiliprotein from *Arthrospira platensis*. *Process Biochem* 82:189–198
- Panayotova V, Stancheva M (2013) Mineral composition of marine macroalgae from the Bulgarian Black Sea coast. *Scr Sci Medica* 45:42–45
- Pandey A, Pandey S, Rajneesh PJ, Ahmed H, Singh V, Singh SP, Sinha RP (2017) Mycosporine-like amino acids (MAAs) Profile of two marine red macroalgae, *Gelidium* sp. and *Ceramium* sp. *Int J Appl Sci Biotechnol* 5:12–21
- Pangestuti R, Kim SK (2011) Biological activities and health benefit effects of natural pigments derived from marine algae. *J Funct Foods* 3:255–266
- Park J, Kim J, Han J (2017a) *Gelidium amansii* Extract, a potent  $\alpha$ -glucosidase and  $\alpha$ -amylase inhibitor, alleviates postprandial hyperglycemia in diabetic mice. *J Life Sci* 27:1052–1058
- Park JH, Hong JY, Jang HC, Oh SG, Kim SH, Yoon JJ, Kim YJ (2012) Use of *Gelidium amansii* as a promising resource for bioethanol: A practical approach for continuous dilute-acid hydrolysis and fermentation. *Bioresour Technol* 108:83–88
- Park JH, Yoon JJ, Park HD, Kim YJ, Lim DJ, Kim SH (2011) Feasibility of biohydrogen production from *Gelidium amansii*. *Int J Hydrogen Energy* 36:13997–14003
- Park MH, Kang JH, Kim HJ, Han JS (2017b) *Gelidium amansii* ethanol extract suppresses fat accumulation by down-regulating adipogenic transcription factors in ob/ob mice model. *Food Sci Biotechnol* 26:207–212
- Molyneux P (2004) The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity. *Songklanakarin J Sci Technol* 26:211–219
- Porse H, Rudolph B (2017) The seaweed hydrocolloid industry: 2016 updates, requirements, and outlook. *J Appl Phycol* 29:2187–2200
- Prayong P, Barusrux S, Weerapreeyakul N (2008) Cytotoxic activity screening of some indigenous Thai plants. *Fitoterapia* 79:598–601
- Qi H, Li D, Zhang J, Liu L, Zhang Q (2008) Study on extraction of agaropectin from *Gelidium amansii* and its anticoagulant activity. *Chin J Oceanol Limnol* 26:186–189
- Ra CH, Jung JH, Sunwoo IY, Jeong GT, Kim SK (2015) Enhanced ethanol production by fermentation of *Gelidium amansii* hydrolysate using a detoxification process and yeasts acclimated to high-salt concentration. *Bioprocess Biosyst Eng* 38:1201–1207
- Rengasamy KRR, Amoo SO, Aremu AO, Stirk WA, Gruz J, Šubratová M, Doležal K, Van Staden J (2015) Phenolic profiles, antioxidant capacity, and acetylcholinesterase inhibitory activity of eight South African seaweeds. *J Appl Phycol* 27:1599–1605
- Rhimou B, Hassane R, José M, Nathalie B (2010a) The antibacterial potential of the seaweeds (Rhodophyceae) of the Strait of Gibraltar and the Mediterranean coast of Morocco. *Afr J Biotechnol* 9:6365–6372

- Rhimou B, Hassane R, Nathalie B (2010b) Antiviral activity of the extracts of Rhodophyceae from Morocco. *Afr J Biotechnol* 9:7968–7975
- Rico J, Guiry M (1997) Life history and reproduction of *Gelidium maggsiae* sp. nov. (rhodophyta, gelidiales) from Ireland. *Eur J Phycol* 32:267–277
- Rocha DHA, Seca AML, Pinto DCGA (2018) Seaweed secondary metabolites in vitro and in vivo anticancer activity. *Mar Drugs* 16:410
- Rubio C, Napoleone G, Luis-González G, Gutiérrez AJ, González-Weller D, Hardisson A, Revert C (2017) Metals in edible seaweed. *Chemosphere* 173:572–579
- Saeed AM, Abotaleb SI, Alam NG, El Mehalawy AA, Gheda SF (2020) In vitro assessment of antimicrobial, antioxidant and anticancer activities of some marine macroalgae. *Egypt J Bot* 60:81–96
- Santelices B (1991) Production ecology of *Gelidium*. *Hydrobiologia* 221:31–44
- Santelices B (2007) Testing the usefulness of attachment structures in the taxonomy of small-sized gelidioids. *Phycologia* 46:293–299
- Santos R, Melo RA (2018) Global shortage of technical agars: back to basics (resource management). *J Appl Phycol* 30:2463–2473
- Sen S, Mallick N (2021) Mycosporine-like amino acids: Algal metabolites shaping the safety and sustainability profiles of commercial sunscreens. *Algal Res* 58:102425
- Seo MJ, Lee OH, Choi HS, Lee BY (2012) Extract from edible red seaweed (*Gelidium amansii*) inhibits lipid accumulation and ROS production during differentiation in 3T3-L1 cells. *Prev Nutr Food Sci* 17:129–135
- Shan B, Cai YZ, Brooks JD, Corke H (2007) The in vitro antibacterial activity of dietary spice and medicinal herb extracts. *Int J Food Microbiol* 117:112–119
- Shiino M, Watanabe Y, Umezawa K (2001) Synthesis of N-substituted N-nitrosohydroxylamines as inhibitors of mushroom tyrosinase. *Bioorg Med Chem* 9:1233–1240
- Siddique MAM, Khan MSK, Bhuiyan MKA (2013) Nutritional composition and amino acid profile of a sub-tropical red seaweed *Gelidium pusillum* collected from St. Martin's Island, Bangladesh. *Int Food Res J* 20:2287–2292
- Silva M, Vieira LMM, Almeida AP, Silva AMS, Seca AML, Brreto MC, Neto AI, Pedro M, Pinto E, Kijoa A (2013) Chemical study and biological activity evaluation of two Azorean macroalgae: *Ulva rigida* and *Gelidium microdon*. *Oceanography* 1:1000102
- So BR, Yeo HJ, Lee JJ, Jung YH, Jung SK (2021) Cellulose nanocrystal preparation from *Gelidium amansii* and analysis of its anti-inflammatory effect on the skin in vitro and in vivo. *Carbohydr Polym* 254:117315
- Subaryono, Sinurat E (2021) Characteristics of *Gelidium* sp seaweed and its agar from Lampung Waters. *IOP Conf Ser Earth Environ Sci* 860:012074
- Sukwong P, Sunwoo IY, Nguyen TH, Jeong GT, Kim SK (2019) R-phycoerythrin, R-phycoyanin and ABE production from *Gelidium amansii* by *Clostridium acetobutylicum*. *Process Biochem* 81:139–147
- Takahashi H, Kosaka M, Watanabe Y, Nakade K, Fukuyama Y (2003) Synthesis and neuroprotective activity of bergenin derivatives with antioxidant activity. *Bioorg Med Chem* 11:1781–1788
- Taleb-Contini SH, Salvador MJ, Watanabe E, Ito IY, Rodrigues de Oliveira DC (2003) Antimicrobial activity of flavonoids and steroids isolated from two *Chromolaena* species. *Rev Bras Cienc Farm* 39:403–408
- Titlyanov EA, Titlyanova TV (2006) Production of plantlets of the red alga of *Gelidium* genus (Rhodophyta) from thallus fragments. *Russ J Mar Biol* 32:284–288
- Titlyanov EA, Titlyanova TV, Kadel P, Lüning K (2006) Obtaining plantlets from apical meristem of the red alga *Gelidium* sp. *J Appl Phycol* 18:167–174
- Torres P, Santos JP, Chow F, dos Santos DYAC (2019) A comprehensive review of traditional uses, bioactivity potential, and chemical diversity of the genus *Gracilaria* (Gracilariales, Rhodophyta). *Algal Res* 37:288–306
- Véliz K, Chandía N, Karsten U, Lara C, Thiel M (2019) Geographic variation in biochemical and physiological traits of the red seaweeds *Chondracanthus chamissoi* and *Gelidium lingulatum* from the south east Pacific coast. *J Appl Phycol* 31:665–682
- Venugopala KN, Rashmi V, Odhav B (2013) Review on natural coumarin lead compounds for their pharmacological activity. *Biomater Res Int* 2013:963248
- Wang L, Shen Z, Mu H, Lin Y, Zhang J (2017) Impact of alkali pretreatment on yield, physico-chemical and gelling properties of high quality agar from *Gracilaria tenuistipitata*. *Food Hydrocoll* 70:356–362
- Wang ML, Hou YY, Chiu YS, Chen YH (2013) Immunomodulatory activities of *Gelidium amansii* gel extracts on murine RAW 264.7 macrophages. *J Food Drug Anal* 21:397–403
- Wi SG, Kim HJ, Mahadevan SA, Yang DJ, Bae HJ (2009) The potential value of the seaweed Ceylon moss (*Gelidium amansii*) as an alternative bioenergy resource. *Bioresour Technol* 100:6658–6660
- Xiang Y, Liu G, Yang L, Zhong JL (2011) UVA-induced protection of skin through the induction of heme oxygenase-1. *Biosci Trends* 5:239–244
- Xu N, Fan X, Yan X, Tseng CK (2004) Screening marine algae from China for their antitumor activities. *J Appl Phycol* 16:451–456
- Yang EC, Kim KM, Boo GH, Lee JH, Boo SM, Yoon HS (2014) Complete mitochondrial genome of the agarophyte red alga *Gelidium vagum* (Gelidiales). *Mitochondrial DNA* 25:267–268
- Yang TH, Chiu CY, Lu TJ, Liu SH, Chiang MT (2019) The anti-obesity effect of polysaccharide-rich red high-fat diet-induced obese hamsters. *Mar Drugs* 17:532
- Yi Z, Yin-Shan C, Hai-Sheng L (2001) Screening for antibacterial and antifungal activities in some marine algae from the Fujian coast of China with three different solvents. *Chin J Oceanol Limnol* 19:327–331
- Yoo HJ, You DJ, Lee KW (2019) Characterization and immunomodulatory effects of high molecular weight fucoidan fraction from the sporophyll of *Undaria pinnatifida* in cyclophosphamide-induced immunosuppressed mice. *Mar Drugs* 17:447
- Yun KJ, Koh DJ, Kim SH, Park SJ, Ryu JH, Kim DG, Lee JY, Lee KT (2008) Anti-inflammatory effects of sinapic acid through the suppression of inducible nitric oxide synthase, cyclooxygenase-2, and proinflammatory cytokines expressions via nuclear factor- $\kappa$ B inactivation. *J Agric Food Chem* 56:10265–10272
- Zemke-White WL, Ohno M (1999) World seaweed utilisation: An end of century summary. *J Appl Phycol* 11:369–376

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.