




KnE Life Sciences

3rd International Conference on Advance & Scientific Innovation (ICASI) – Life Sciences Chapter

**20–21 Julv 2021
Medan, Indonesia**

ISSN 2413-0877



www.KnEpublishing.com

PUBLISHED

- 2022-09-13

[KnE Life Sciences](#) / The 3rd International Conference on Advance & Scientific Innovation (ICASI) – Life Sciences Chapter

This collection contains selected papers from the **3rd International Conference on Advance & Scientific Innovation (ICASI) – Life Sciences Chapter** held in Medan, Indonesia on the 20th–21st of July 2021. This event was hosted by the Forum Kerjasama Pendidikan Tinggi.

The purpose of this conference was to discuss the epidemiology of some of the major public health issues and diseases prevalent in Indonesia.

This *KnE Life Sciences* collection contains 76 peer-reviewed articles containing ground breaking research on biology, healthcare, and pharmacology topics. Key themes include new studies into illnesses such as leprosy, tuberculosis and hypertension, and new insights into medication efficacy in combatting disease. In addition, this collection offers a unique overview of the disparate diseases which predominate in different Indonesian provinces.

These papers will be of interest to academics, students and professionals researching or working in the fields of healthcare and pharmaceutical and biological sciences.

Conference date: 20–21 July 2021

Location: Medan, Indonesia

Editor: Robbi Rahim

Organizer: Forum Kerjasama Pendidikan Tinggi

Published: 13 September 2022

ISSN: 2413-0877

[ORGANISATION](#)

[PUBLICATION ETHICS](#)

[THINK CHECK
ATTEND](#)

[COPYRIGHT](#)

[ARCHIVING POLICY](#)

[MEET THE TEAM](#)

ABOUT KNE PUBLISHING

We help you publish your journals and conference proceedings on our platform while making sure that the highest standards of quality and ethics are adhered to.

CONTACT US

ORGANISATION

KnE Publishing is the publishing arm of Knowledge E.

Knowledge E's mission is to disseminate regional academic journal articles and conference papers to the broader global research community thereby creating a permanent, discoverable, archived, and open access record of high quality scholarly content. All articles published by Knowledge E are made freely accessible online immediately upon publication. All content on the KnE Publishing Platform is published under the [Creative Commons Attribution License \(CC-BY\)](#).

Knowledge E takes a responsible approach to peer review insisting on the application of peer-review processes by journal and conference editors together with strict ethical policies and standards to ensure the addition of high-quality scientific content to the field of scholarly publication. Knowledge E adheres to the COPE [Code of Conduct](#) and [Best Practice Guidelines](#).

Knowledge E is a member of several industry organisations, including the Association of Learned and Professional Society Publishers ([ALPSP](#)), the Publishers International Linking Association ([Crossref](#)), and the Open Access Scholarly Publishing Association ([OASPA](#)), and is a signatory to the United Nations Sustainable Development Goals [Publishers Compact](#). It is also a member of the Open Researcher and Contributor ID ([ORCID](#)) and is its Institutional Ambassador in the Middle East.

All of Knowledge E's content is archived in [Portico](#) a not-for-profit organization with a mission and singular focus to provide a permanent archive of electronic scholarly content.

[Read more about Knowledge E on our corporate webpage](#)

PUBLICATION ETHICS

We make every effort to ensure high quality scholarship in all published articles and conference papers published by us.

[Back to top](#)

[ORGANISATION](#)

[PUBLICATION ETHICS](#)

[THINK CHECK
ATTEND](#)

[COPYRIGHT](#)

[ARCHIVING POLICY](#)

[MEET THE TEAM](#)

This means that we consider best practices such as **peer review, anti-plagiarism control**, and adherence to **industry standard ethical practices and guidelines** to be matter of course, and therefore mandatory.

KnE Publishing adheres to the [Committee on Publication Ethics \(COPE\) Core practices.](#)

[Read our full publication ethics statement](#)

THINK CHECK ATTEND

Think. Check. Attend. (www.thinkcheckattend.org) is an initiative that aims to guide and assist researchers and scholars to judge the legitimacy and academic credentials of conferences in order to help them decide whether to or not attend the same. Knowledge E adheres to the Core Principles of the Committee On Publication Ethics (COPE). As the initiator of the Think. Check. Attend. initiative, Knowledge E ensures all conference proceedings published on its platform come from conferences fitting the guidelines stipulated in the initiative.

[Read more about Think Check Attend](#)

COPYRIGHT

We publish all content under the [Creative Commons Attribution \(CC-BY\) license.](#)

In short, under this license, authors retain ownership of the copyright for their content, but it also allows anyone to download, reuse, reprint, modify, distribute, and/or copy the content **as long as the original authors and source are cited.**

We have chosen this license because we have evaluated it as being fair to both authors and readers.

[Read the full copyright policy](#)

ARCHIVING POLICY

To preserve scholarly literature published in electronic form and to ensure that these materials remain available to future generations of scholars and researchers, Knowledge E content is archived in [Portico](#), a not-for-profit organization with a mission and singular focus to provide a permanent archive of electronic scholarly content. Portico representative offices are located in New York, New Jersey, and Michigan.

MEET THE TEAM

We are the people behind the scenes doing our best to get you edited, published and indexed as quickly as possible. Feel free to contact us for anything!



[Back to top](#)

[ORGANISATION](#)

[PUBLICATION ETHICS](#)

[THINK CHECK
ATTEND](#)

[COPYRIGHT](#)

[ARCHIVING POLICY](#)

[MEET THE TEAM](#)

KAMRAN KARDAN
CEO and Founder



EMILY CHOYNOWSKI
Head of Publishing



MOHAMAD MOSTAFA
Publishing Editor

orcid.org/0000-0003-0768-6642



DEEPTI THAPA
Production & Copy Editor



LEILA ELIZABETH KARDAN
Publishing Executive



ABIGAIL ENOCH
Editorial Assistant



THERESE BILL
Head of Operations

[Back to top](#)

[ORGANISATION](#)

[PUBLICATION ETHICS](#)

[THINK CHECK
ATTEND](#)

[COPYRIGHT](#)

[ARCHIVING POLICY](#)

[MEET THE TEAM](#)



CARL FRY

BDM for Government and Projects



SUSHANT VIRDI

IT & Client Support Manager



SOMIN GAJJAR

Accountant and Financial Controller

EXTERNAL ADVISER



PIPPA SMART

Publishing Adviser

orcid.org/0000-0002-5528-4704

Contact

Email publishing@knowledgee.com
Phone +971 4 4227043
Address Knowledge E
Office 4401-04
Jumeirah Bay X2 Tower
Jumeirah Lakes Towers (JLT)
P.O. Box 488239
Dubai, UAE

Follow us

 [@KnePublishing](#)

 facebook.com/KnowledgeEGlobal

 [Youtube](#)

 [KnowledgeEGlobal](#)

[Find out more about
Knowledge E from
our corporate page](#)



[KnE Life Sciences](#) / Indexing

Indexing





KnE Life Sciences is listed with the following indexes and bibliometric databases:

- Dimensions
- Directory of Open Access Resources (ROAD)
- EBSCO
- Google Scholar
- Indonesia OneSearch
- Microsoft Academic Research
- MyScienceWork
- Neliti
- OpenAIRE
- Researcher App
- Scilit
- Semantic Scholar
- TheOneRepo
- Ulrichsweb
- UnPaywall
- WorldCat Discovery Service
- WorldWideScience

Contact

Email publishing@knowledgee.com
Phone +971 4 4227043
Address Knowledge E
Office 4401-04
Jumeirah Bay X2 Tower
Jumeirah Lakes Towers (JLT)
P.O. Box 488239
Dubai, UAE

Follow us

 [@KnePublishing](#)
 facebook.com/KnowledgeEGlobal
 [Youtube](#)
 [KnowledgeEGlobal](#)

[Find out more about Knowledge E from our corporate page](#)



Growth Characteristic and the Study of Polymorphism Growth Hormone Genes of Sentul Chicken

0

[KnE Life Sciences / The 3rd International Conference on Advance & Scientific Innovation \(ICASI\) – Life Sciences Chapter / Pages 481–488](#)

[Download fulltext](#)

[HTML](#)

[How to Cite](#)

<https://doi.org/10.18502/kls.v0i0.11834>

STATISTICS

100 Abstract Views

82 PDF Views

0 HTML Views

AUTHORS

[Ismoyowati.](#)

[S Mugiyono](#)

[E Tugiyanti](#)

[IH Sulistyawan](#)

[Rosidi.](#)

PUBLISHED DATE

Sep 13, 2022

Abstract

The research was conducted to study the characteristics of growth, carcass production and growth hormone gene polymorphisms in various male sentul chickens. The research was conducted using experimental methods with a completely randomized design. The research material was 100 male day old chickens of Sentul. The treatment was a fixed factor, namely the variation of the color of the feathers of various Sentul chickens consisting of: “Abu” Sentul, “Emas” Sentul, “Geni” Sentul, “Debu” Sentul, “Batu” Sentul.. Each experimental unit consisted of 5 chickens and 4 replications. The variables measured included: hatching weight, body weight, body weight gain and the percentage of carcass produced at the age of 8 weeks. Identification of growth hormone gene polymorphisms used the primary design Gallus gallus haplotype GH-h22 growth hormone (GH) gene, complete cds, GenBank: JN675393.1 with forward primer / Sequence: AGGTGGTTCGGTTTTCACTG and reverse primer / Sequence: TCCCTTCTTCCAGGTCCTTT. Characteristics of growth and carcass production data were analyzed by analysis of variance; while to determine the presence of GH gene, polymorphisms were analyzed using the bioedit program. Analysis of variance showed that there were no significant differences ($P > 0.05$) between hatching weight, body weight, body weight gain and carcass percentage of various Sentul chickens. The results of GH Gallus gallus gene sequencing at base length 80 bp showed a mutation from adinine to cytosine, when comparing sentul chickens and Gallus gallus data in GenBank. The GH gene were present in various sentul chickens with monomorphic characteristics with homozygous genotypes of CC. The results of this study can be concluded that the characteristics of growth and carcass production in various Sentul chickens are relatively the same, as well as the monomorphic GH gene.

Keywords: Sentul chickens, body weight, carcass percentage, growth hormone gene, monomorphic.

References

- [1] Nataamijaya AG. 2000. The native chicken of Indonesia. Buletin Plasma Nutfah 6(1):1–6.
- [2] Zhou H, AD Mitchell, JP McMurtry, CM Ashwell and S Lamont. 2005. Insulin-Like Growth Factor-1 Gene Polymorphism Association with Chicken Body

Composition, Skeleton Integrity and Metabolic Traits in Chickens. *Poult Sci.* 84:212-219.

[3] Li H, W Zhu, K Chen, X Wu, Q Tang and Y Gao. 2008. Associations between GHR and IGF-1 Gene Polymorphisms, and Reproductive Traits in Wenchang Chickens. *Turk. J. Vet. Anim. Sci.* 32:281-285.

[4] Anh NTL, S Kunhareang, M Duangjinda. (2015). Association of chicken growth hormones and insulin-like growth factor gene polymorphisms with growth performance and carcass traits in Thai broilers. *Asian-Australasian journal of animal sciences.* 28:1686.

[5] Pichner F. 1981. Population genetic and animal breeding. WH. Freeman and Co. San Fransisco.

[6] Nei M. 1978. Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics.* 89:583-590.

[7] Sartika T, S Iskandar, LH Prasetyo, Takahashi and M Minezawa. 2004. Kekerabatan genetik ayam Kampung, Pelung, Sentul dan Kedu Hitam dengan menggunakan penanda DNA Mikrosatelit I: Grup pemetaan pada makro kromosom. *J. Ilmu Ternak and Veteriner* 9 (2):81-86.

[8] Rizzi C. 2018. Plumage colour in Padovana chicken breed: growth performance and carcass quality. *Italian Journal of Animal Science* 17 (3):797–803. doi.org/10.1080/1828051X.2017.1413598.

[9] Dewi GMAK, IM Nuriyasa and IW Wijana. 2017. Production of Chicken carcass and non carcass of kampung chicken who received rations skin Dragon fruit flour (*Hylocereus polyrhizus*) Fermented. In: Proc. The International Seminar on Tropical Animal Production. Contribution of Livestock production on food sovereignty in Tropical countries. September 12-14, 2017. Yogyakarta. Indonesia.

[10] Kazemi H, M Rezaei, H Hafezian, GR Mianji and M Najafi. Genetic Analysis of SNPs in GH, GHR, IGF-I and IGFBP II Genes and their Association with Some Productive and Reproductive Traits in Native Breeder Hens. *Gene Technol* 7 (1). 1-7. DOI: 10.4172/2329-6682.1000145

[11] Yulianty.2013. Keragaman Gen GH (Growth Hormone) Pada Populasi Kambing Kacang Di Kabupaten Jenoponto. Skripsi. Program Study Produksi Ternak Jurusan Produksi Ternak Fakultas Peternakan Universitas Hasanudin Makasar.

[12] Khaerunnisaa I, Jakaria, Il Arief, C Budiman and C Sumantri. 2017. The Associations of GH and GHR Genes with Carcass Components in Indonesian Kampung and Broiler Chicken Cross. *Media Peternakan.* 40 (2):78-87. DOI: https://doi.org/10.5398/medpet.2017.40.2.78.

[13] Ge W, ME Davis, HC Hines and KM Irvin. 2000. Single nucleotide polymorphisma detected in exon 10 of the bovine growth hormone receptor gene. *J. Anim. Sci.* 78:2229-2230.

[14] Beauchemin, VR, MG Thomas, DE Frank and GA Silver. 2006. Evaluation of DNA polymorphisms involving growth hormone relative to growth and carcass characteristics in Brahman steers. *Genet. Mol. Res.* 5:438-447.

[15] Thomas MG, RM Enns, KL Shirley, MD Garcia, A.J Garrett and G.A Silver. 2007. Associations of DNA polymorphisms in growth hormone and its

transcriptional regulators with growth and carcass traits in two populations of Brangus bulls. Genet. Mol. Res. 6:222-237.

[16] Sumantri, C, A Farajallah, and JF Salamena. 2010. Keragaman genetik DNA mikrosatelit dan hubungannya dengan performa bobot badan pada domba lokal. Media Peternakan. 31(1):1-13.


Contact


Email publishing@knowledgee.com


Phone +971 4 4227043


Address Knowledge E
Office 4401-04
Jumeirah Bay X2 Tower
Jumeirah Lakes Towers (JLT)
P.O. Box 488239
Dubai, UAE

Follow us

 [@KnePublishing](https://twitter.com/KnePublishing)

 facebook.com/KnowledgeEGlobal

 [Youtube](https://www.youtube.com/KnowledgeEGlobal)

 [KnowledgeEGlobal](https://www.linkedin.com/company/KnowledgeEGlobal)

[Find out more about Knowledge E from our corporate page](#)



Conference Paper

Growth Characteristic and the Study of Polymorphism Growth Hormone Genes of Sentul Chicken

Ismoyowati*, S Mugiyono, E Tugiyanti, IH Sulistyawan and Rosidi

Faculty of Animal Science, Universitas Jenderal Soedirman, Purwokerto, Indonesia

ORCID

<https://orcid.org/0000-0002-7920-3681>

Abstract.

The research was conducted to study the characteristics of growth, carcass production and growth hormone gene polymorphisms in various male sentul chickens. The research was conducted using experimental methods with a completely randomized design. The research material was 100 male day old chickens of Sentul. The treatment was a fixed factor, namely the variation of the color of the feathers of various Sentul chickens consisting of: "Abu" Sentul, "Emas" Sentul, "Geni" Sentul, "Debu" Sentul, "Batu" Sentul.. Each experimental unit consisted of 5 chickens and 4 replications. The variables measured included: hatching weight, body weight, body weight gain and the percentage of carcass produced at the age of 8 weeks. Identification of growth hormone gene polymorphisms used the primary design Gallus gallus haplotype GH-h22 growth hormone (GH) gene, complete cds, GenBank: JN675393.1 with forward primer / Sequence: AGGTGGTTCGGTTTTCACTG and reverse primer / Sequence: TCCCTTCTTCCAGGTCCTTT. Characteristics of growth and carcass production data were analyzed by analysis of variance; while to determine the presence of GH gene, polymorphisms were analyzed using the bioedit program. Analysis of variance showed that there were no significant differences ($P > 0.05$) between hatching weight, body weight, body weight gain and carcass percentage of various Sentul chickens. The results of GH Gallus gallus gene sequencing at base length 80 bp showed a mutation from adinine to cytosine, when comparing sentul chickens and Gallus gallus data in GenBank. The GH gene were present in various sentul chickens with monomorphic characteristics with homozygous genotypes of CC. The results of this study can be concluded that the characteristics of growth and carcass production in various Sentul chickens are relatively the same, as well as the monomorphic GH gene.

Keywords: Sentul chickens, body weight, carcass percentage, growth hormone gene, monomorphic.

Corresponding Author:

Ismoyowati; email:

ismoyowati@unsoed.ac.id

Published: 13 September 2022

Publishing services provided by
Knowledge E

© Ismoyowati et al. This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Selection and Peer-review under the responsibility of the ICASI Conference Committee.

OPEN ACCESS

1. Introduction

Indonesia is a country that has a wealth of biodiversity, one of which is a local chicken. The existence of local breed chickens in Indonesia is still very diverse genetically and phenotypically, and most of them are still maintained semi-intensively or even traditionally. There are 31 breeds of local Indonesian chickens that have distinctive characteristics, including: Pelung chicken, Sentul chicken, Nunukan chicken, Sedayu chicken, Sentul chicken, Gaok chicken and others [1]. One type of local poultry with a large enough potential is the Sentul chicken. Sentul chicken is a local chicken that grows in Ciamis Regency, and is one of the original genetic resources and has been designated as a leading commodity in West Java.

The advantages of Sentul Chicken include relatively fast growth and high egg production compared to other local chickens. With these advantages, Sentul Chicken can be used as a community industrial commodity or to be further developed into superior local chickens. Based on the feather color pattern, Sentul Chicken can be divided into five types, namely Sentul Abu, Sentul Emas, Sentul Geni, Sentul Debu and Sentul Batu.

The high genetic diversity in Sentul chickens made it possible to improve the genetic quality through selection. Molecular markers associated with one or more quantitative characteristic can help increase production and health simultaneously [2]. The use of selection with the aid of molecular markers has been shown to be efficient and leads to improved production performance in livestock [3]. Previous studies reported candidate genes could result in higher efficiency for detecting traits of desired economic value, to improve production and reproductive performance in poultry breeding programs. One of the genes that correlate with growth is GH, is considered as the candidate gene that has the most role in growth performance and carcass quality traits in chickens [4]. This study aims to study the characteristics of growth, carcass production and growth hormone gene polymorphisms in various male sentul chickens.

2. Materials and Methods

The research material used 100 male Sentul day old chickens. The research material was taken from the Farmers Group Breeders of the Ciung Wanara, Ciamis District, Ciamis Regency. Materials used include: complete feed broiler starter crumble for chicks aged one day-3 weeks, containing nutrients: 22% crude protein, 12% moisture content, 5% fat, 5% crude fiber, 1% Ca, 0.5% P and ME 3000 kcal / kg. Complete Par S feed for chickens aged 4-8 weeks contain nutrients: 19% crude protein, 12% moisture content,

3% fat, 7% crude fiber, 1% Ca, 0.45% P and 2800 kcal ME / kg. The materials used to assess the polymorphism of the GH gene consist of materials for DNA isolation and PCR. Identification of growth hormone gene polymorphisms using the primary design Gallus gallus haplotype GH-h22 growth hormone (GH) gene, complete cds, GenBank: JN675393.1 with forward primer / Sequence: AGGTGGTTCGGTTTTCACTG and reverse primer / Sequence: TCCCTTCTTCCAGGTCCTTT.

The research was conducted by using experimental methods with a completely randomized design. The treatments consisted of varieties of Sentul Chicken including: "Abu" Sentul, "Emas" Sentul, "Geni" Sentul, "Debu" Sentul, and "Batu" Sentul. Each experimental unit was filled with 4 chickens and 5 replications, so that it involved 100 chickens. Chicken rearing is carried out for 8 weeks. Feeding is done in a measured adlibitum. Weighing the body weight is done once a week.

Blood samples were taken at the age of 7 weeks individually through the axillary veins of each Sentul Chicken as much as 1 ml using a syringe and put into a tube containing EDTA for DNA isolation. DNA isolation using a DNA kit to obtain pure DNA, then electrophoresis using 1% agarose gel.

The PCR amplification was carried out by targeting the Growth Hormone gene in a total reaction volume of 25 µl consisting of 12.5 µl of KAPA2G Fast Ready mix PCR kit (Kapa Biosystems), 9.5 µl of dH₂O, 1 µl of GH Gallus gallus primers, and finally 1 µl of genomic DNA. The amplification stage includes pre-denaturation carried out for 5 minutes at 94°C, then denaturation for 30 seconds at 94°C, and followed by annealing for 45 seconds at 55°C. The next step is elongation at 72°C for 1 minute. The last stage is post-elongation, namely the step to complete DNA elongation and is carried out for 5 minutes at a temperature of 72°C. The PCR reaction was repeated 35 cycles to get maximum results, the PCR results were then electrophoresed on 1% agarose gel and visualized using Ultra Violet light to determine the success of PCR. PCR Products are then sequenced using a sequencer machine.

Data analysis to determine differences in the growth performance of various Sentul chickens used analysis of variance. Analysis of genetic diversity based Gallus gallus GH gene is done by calculating the genotype and gene frequencies. Genotypes and gene frequencies were calculated by the formula Pirchner (1981) [5]:

$$\text{Formula: } FAn = \frac{\sum GH \text{ A gene}}{\sum GH \text{ A gene} + \sum GH \text{ n gene}}$$

Note: FAn = frequency of gene A at the nth locus

Genetic diversity is determined using the heterozygosity formula based on Nei (1978) [6] with the formula: $h = 1 - \sum_{i=1}^m x_i^2$

Note:

m = number of alleles,

X_i = frequency of the i th gene

3. Results and Discussion

3.1. Growth performance and carcass production

Sentul chicken is an Indonesian local chicken that has genetic diversity. The genetic diversity of Sentul Chickens can be seen from the different plumage colors. The genetic differences of each individual are thought to occur because of differences in the inherited traits of each parent. Individual traits, both qualitative traits (plumage color, comb shape) and quantitative traits (body weight, egg production) are determined by genes and alleles arranged in DNA pairs found in the cell nucleus.

The results of the analysis of variance, the growth performance and carcass production of various Sentul chickens, showed that the fix factor of various Sentul chickens had no significant effect ($P > 0.05$) on all the characteristics measured. These results indicate that the diversity of plumage color does not affect the growth characteristics and carcass production. Differences in breed in local chickens are only reflected in the appearance of their phenotypes, whereas based on their genetics, selection still needs to be done [7]. Based on the color of the feathers, Sentul chickens are divided into 6 types, namely “Batu” Sentul, “Abu” Sentul, “Debu” Sentul, “Emas” Sentul, “Geni” Sentul and “Jambe” Sentul. The results of this study are different from previous studies, which reported the process of feather formation can be a factor affecting the growth rate of poultry, so that chickens with different plumage colors of the same breed can have different viability and growth performance. Chickens with black plumage have lower growth and higher mortality compared to white plumage birds [8].

The results of the analysis of variance of consumption and feed conversion ratio on various Sentul chickens in the initial and growth period, showed that various Sentul chickens had no significant effect ($P > 0.05$) on the feed consumption and feed conversion ratio of chicken feed in a growth period. These results support measurements of body weight and growth. Various Sentul chickens consume relatively the same feed and the weight gain is relatively the same, so the feed conversion ratio is also the same (Table 1).

The results of the analysis of variance showed that carcass production in various Sentul chickens was not significantly different ($P < 0.05$). Carcass production is closely

TABLE 1: Growth performance and carcass production in various male sentul chickens.

Sentul Chickens	DOC weight (g)	Body weight at 8 weeks of age (g)	Weight gain in 8 weeks (g)	Feed consumption (g)	Feed conversion ratio	Carcass weight (g)	Carcass percentage (%)
"Abu"	29.15 ± 1.01	687.00 ± 37.74	657.85 ± 37.60	1710.14 ± 63.74	2.603 ± 0.107	541.2 ± 78.7	66.47 ± 0.88
"Emas"	29.74 ± 1.54	699.87 ± 52.32	669.97 ± 51.28	1812.83 ± 61.07	2.714 ± 0.143	466.8 ± 51.1	63.54 ± 2.14
"Geni"	28.80 ± 1.06	681.35 ± 28.82	652.62 ± 29.09	1741.80 ± 72.51	2.671 ± 0.108	494.2 ± 79.0	66.09 ± 3.24
"Debu"	30.64 ± 1.88	673.04 ± 57.90	642.41 ± 59.64	1729.60 ± 49.04	2.707 ± 0.205	512.4 ± 32.0	64.43 ± 6.38
"Batu"	29.79 ± 2.61	698.22 ± 55.37	668.44 ± 55.93	1771.67 ± 29.64	2.666 ± 0.235	510.8 ± 69.2	67.53 ± 6.58
Average	29.62 ± 1.70	687.90 ± 44.83	658.26 ± 45.15	1753.21 ± 63.75	2.672 ± 0.158	505.1 ± 64.0	65.61 ± 4.88

related to DOC weight, growth, body weight and feed consumption. The results of this study indicate that DOC weight, growth, body weight, and feed consumption are relatively the same, resulting in insignificant differences in carcass production (Table 1). The carcass percentage of Sentul chicken in this study ($65.61 \pm 4.88\%$), is higher than in the study the carcass percentage of native chicken (56.08-58.02%) [9].

3.2. Identification of the GH *Gallus gallus* gene polymorphism

Growth hormone (GH) is responsible for several metabolic pathways, such as growth, reproduction, puberty, immune response and regulating cell organization functions through its receptors. The growth hormone gene in chickens (cGH) is located on chromosome 1 with a length of 4098 bp containing 5 exons and 4 introns [10]. Growth hormone in growing animals plays a role in increasing the efficiency of feed use, organ growth and bone growth. This relationship makes growth hormone a candidate gene that can be used as a genetic marker in selection programs [11]. The results of the GH *Gallus gallus* gene sequencing in Sentul chickens are presented in Figure 1.



Figure 1: The change in the position of the 80 bp SNP was c.80A> C.

Based on the results of the GH *Gallus gallus* gene sequencing at a base length of 80 bp, it shows a mutation from adenine (A) to cytosine (C) (Figure 1). There is only one genotype of the GH *Gallus gallus* gene in all various of sentul chickens, namely CC. These results indicate that the GH *Gallus gallus* gene is monomorphic. The genotypes of various sentul chickens based on the GH gene are the same, because there is no base variation or polymorphism. Polymorphism in a gene (single nucleotide polymorphism) can occur due to a mutation process in one or more of the bases that are composed of the nucleotides.

The genotype and gene frequency in various sentul chickens, based on the GH *Gallus gallus* gene, is one, because all genotypes are homozygous CC. As a result, there was no genetic diversity based on various sentul chickens based on the identification of the GH *Gallus gallus* gene. The results of this study were different from previous studies, which reported that there were polymorphisms of the GH gene in chickens from crosses between native chickens and broilers, but there was no association with chicken growth, so it could not be used as a candidate gene [12].

This has an impact on body weight, growth and carcass production produced in various Sentul chickens which are relatively the same. Genetic and environmental factors affect the growth rate of Sentul chicken body weight. The results of this study are different from previous studies. Previous studies have shown that the GH gene plays a major role in body weight gain reported in several species such as cattle [13]. The GH gene is very influential on growth and metabolism through interactions with the specific receptor (GHR) on the surface of the target cell, therefore GHR is a candidate gene that influences production [14]. GH and Pit-1 genes affect growth and carcass traits. Pit-1 GG genotypes have a higher percentage of intramuscular fat than AA or AG, and GH and CG genotypes have faster body weight gain than CC and GG genotypes [15]. Genes that control metabolism and energy distribution have a very important economic impact on the livestock world. These genes include growth hormone receptor (GHR), IGF2, and GH. The AG genotype on GHR affects body weight and feed efficiency [16].

4. Conclusions

Based on the results of the study, it can be concluded that there is no genetic diversity in the GH *Gallus gallus* gene, so that the characteristics of growth and carcass production in various Sentul chickens are relatively the same.

5. Acknowledgement

Thanks to the Rector Universitas Jenderal Soedirman and the Directorate General of Higher Education on Applied research funding.

References

- [1] Nataamijaya AG. 2000. The native chicken of Indonesia. *Buletin Plasma Nutfah* **6**(1):1–6.
- [2] Zhou H, AD Mitchell, JP McMurtry, CM Ashwell and S Lamont. 2005. Insulin-Like Growth Factor-1 Gene Polymorphism Association with Chicken Body Composition, Skeleton Integrity and Metabolic Traits in Chickens. *Poult Sci.* **84**:212-219.
- [3] Li H, W Zhu, K Chen, X Wu, Q Tang and Y Gao. 2008. Associations between GHR and IGF-1 Gene Polymorphisms, and Reproductive Traits in Wenchang Chickens. *Turk. J. Vet. Anim. Sci.* **32**:281-285.
- [4] Anh NTL, S Kunhareang, M Duangjinda. (2015). Association of chicken growth hormones and insulin-like growth factor gene polymorphisms with growth performance and carcass traits in Thai broilers. *Asian-Australasian journal of animal sciences.* **28**:1686.
- [5] Pichner F. 1981. *Population genetic and animal breeding*. WH. Freeman and Co. San Fransisco.
- [6] Nei M. 1978. Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics.* **89**:583-590.
- [7] Sartika T, S Iskandar, LH Prasetyo, Takahashi and M Minezawa. 2004. Kekerabatan genetik ayam Kampung, Pelung, Sentul dan Kedu Hitam dengan menggunakan penanda DNA Mikrosatelit I: Grup pemetaan pada makro kromosom. *J. Ilmu Ternak and Veteriner* **9** (2):81-86.
- [8] Rizzi C. 2018. Plumage colour in Padovana chicken breed: growth performance and carcass quality. *Italian Journal of Animal Science* **17** (3):797–803. doi.org/10.1080/1828051X.2017.1413598.
- [9] Dewi GMAK, IM Nuriyasa and IW Wijana. 2017. Production of Chicken carcass and non carcass of kampung chicken who received rations skin Dragon fruit flour (*Hylocereus polyrhizus*) Fermented. In: Proc. The International Seminar on Tropical Animal Production. Contribution of Livestock production on food sovereignty in Tropical countries. September 12-14, 2017. Yogyakarta. Indonesia.

- [10] Kazemi H, M Rezaei, H Hafezian, GR Mianji and M Najafi. Genetic Analysis of SNPs in GH, GHR, IGF-I and IGFBP II Genes and their Association with Some Productive and Reproductive Traits in Native Breeder Hens. *Gene Technol* **7** (1). 1-7. DOI: 10.4172/2329-6682.1000145
- [11] Yulianty.2013. Keragaman Gen GH (Growth Hormone) Pada Populasi Kambing Kacang Di Kabupaten Jenoponto. *Skripsi*. Program Study Produksi Ternak Jurusan Produksi Ternak Fakultas Peternakan Universitas Hasanudin Makasar.
- [12] Khaerunnisaa I, Jakaria, II Arief, C Budiman and C Sumantri. 2017. The Associations of GH and GHR Genes with Carcass Components in Indonesian Kampung and Broiler Chicken Cross. *Media Peternakan*. **40** (2):78-87. DOI: <https://doi.org/10.5398/medpet.2017.40.2.78>.
- [13] Ge W, ME Davis, HC Hines and KM Irvin. 2000. Single nucleotide polymorphism detected in exon 10 of the bovine growth hormone receptor gene. *J. Anim. Sci.* **78**:2229-2230.
- [14] Beauchemin, VR, MG Thomas, DE Frank and GA Silver. 2006. Evaluation of DNA polymorphisms involving growth hormone relative to growth and carcass characteristics in Brahman steers. *Genet. Mol. Res.* **5**:438-447.
- [15] Thomas MG, RM Enns, KL Shirley, MD Garcia, A.J Garrett and G.A Silver. 2007. Associations of DNA polymorphisms in growth hormone and its transcriptional regulators with growth and carcass traits in two populations of Brangus bulls. *Genet. Mol. Res.* **6**:222-237.
- [16] Sumantri, C, A Farajallah, and JF Salamena. 2010. Keragaman genetik DNA mikrosatelit dan hubungannya dengan performa bobot badan pada domba lokal. *Media Peternakan*. **31**(1):1-13.