

PAPER • OPEN ACCESS

Preface

To cite this article: 2022 *IOP Conf. Ser.: Earth Environ. Sci.* **1001** 011001

View the [article online](#) for updates and enhancements.

You may also like

- [PREFACE](#)
- [Optimize use of icare based student worksheet \(ICARE-BSW\) in physics learning at the introduction level](#)
Jurubahasa Sinuraya, Ida Wahyuni, Deo Demonta Panggabean et al.
- [The needs analysis of expansion blended learning using icare's model in history learning for tenth grade in senior high school level](#)
R Andriani, N Umamah and M Na'im

A promotional banner for 'Free the Science Week 2023' with a dark blue background and a futuristic circular interface. A hand is shown interacting with the interface, which features a central padlock icon. The text 'Free the Science Week 2023' is in large, light blue font, followed by 'April 2-9' in white. Below this, 'Accelerating discovery through open access!' is written, with 'open access!' in light blue. At the bottom left is the ECS logo and the website 'www.ecsdl.org'. At the bottom right is a blue button with the text 'Discover more!'.

Free the Science Week 2023 April 2-9

Accelerating discovery through
open access!

 www.ecsdl.org [Discover more!](#)

Preface

International Conference on Animal Research for Eco-Friendly Livestock Industry (ICARELI) 2021 was held in Surakarta 23 November 2021. The scientific conference was performed by Department of Animal Science, Faculty of Agriculture, Universitas Sebelas Maret, Indonesia in virtual platform due to COVID-19 pandemic. This international conference aims to explore recent researches on eco-friendly livestock industry which is in line with sustainable development goals.

The conference consisted of plenary and panel sessions. During plenary session, all of the invited speakers from Japan, USA, South Korea, Germany and Indonesia were able to present live for discussions although there was time difference between continents. Professor Siti Nur Hidayati from Middle Tennessee State University, USA gave talk about impact of climate change to tropical biodiversity. Professor Heekwon Ahn from Chungnam National University, South Korea discussed about mitigation of livestock emission. Prof. Masato Yayota from Gifu University, Japan talked about eco-friendly animal feed resources. Professor Carlos Hermosilla from Justus Liebig University Giessen, Germany made a talk about environmental effect on host-parasite interaction. Prof. Sutrisno Hadi Purnomo from Universitas Sebelas Maret, Surakarta delivered a speech about crops-livestock integration for zero-waste farming.

The Organizing Committee received 123 manuscripts. There 107 participants were able to give presentation during conference. The participants presented their scientific results live by zoom platform provided during conference. All talks and discussions during conference were well recorded. There were 7 breakout rooms of zoom provided according to conference scopes i.e.: eco-friendly animal feed resources; nutrition, forage and pasture, sustainable animal production system; animal biodiversity, ecology, and conservation, organic farming; eco-friendly animal products processing and technology; food safety and security of animal resources; animal waste management and bioenergy; socio-economic studies on sustainable animal farming; and animal health and welfare. The 55 presented manuscripts out of 107 total received were screened for IOP: EES publication and went through a double-blind review in the Morressier platform.

The ICARELI 2021 was successfully held by all support from the university president, dean, students volunteering in the event, and all of third parties involved. We are very glad that although this conference held as the first time and during Covid-19 pandemics, the event was joined by all of the participants with enthusiasms. The editorial team, reviewers and participants were also able to learn fast in the Morressier platform during reviewing process with the responsive work of Morressier team. Therefore, the proceedings of ICARELI 2021 can be published and hopefully able to give values to the community.

Editors

Dr.med.vet. Penny Humaidah Hamid

Dr.agr. Muhammad Cahyadi

Scientific reviewers of proceeding

Farouq Heidar Barido, Ph.D.

(Kangwon National University, South Korea)

Nurul Huda, Ph.D.

(Universiti Malaysia Sabah, Malaysia)

Prof. Dr. Endang Tri Margawati

(The National Research and Innovation Agency, Indonesia)

Dr. Muhsin Al Anas

(University of Arkansas, USA)

Prof. Dr.sc.ETH. Anuraga Jayanegara

(IPB University, Indonesia)

Prof. Sutrisno Hadi Purnomo, Ph.D.

(Universitas Sebelas Maret, Indonesia)

Agus Setiadi, Ph.D.

(Diponegoro University, Indonesia)

April Hari Wardhana, Ph.D.

(The National Research and Innovation Agency, Indonesia)

Dr. Cahya Utama

(Diponegoro University, Indonesia)

Dicky Tri Utama, Ph.D.

(Universitas Padjadjaran, Indonesia)

Dimas Hand Vidya Paradhita, Ph.D.

(Universitas Gadjah Mada, Indonesia)

Dian Wahyu Harjanti, Ph.D.

(Diponegoro University, Indonesia)

Widi Nugroho, Ph.D.

(Universitas Brawijaya, Indonesia)



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Daud Samsudewa, Ph.D.	(Diponegoro University, Indonesia)
Lilik Retna Kartikasari, Ph.D.	(Universitas Sebelas Maret, Indonesia)
Muhlisin, Ph.D.	(Universitas Gadjah Mada, Indonesia)
Siwi Gayatri, Ph.D.	(Diponegoro University, Indonesia)
Sutaryo, Ph.D.	(Diponegoro University, Indonesia)
Yuli Yanti, Ph.D.	(Universitas Sebelas Maret, Indonesia)

Scientific committee of conference

Prof. Dr. Carlos Hermosilla	(Justus Liebig Universitaet Giesen, Germany)
Prof. Dr. Heekwon Ahn	(Chungnam National University, South Korea)
Prof. Dr. Masato Yayota	(Gifu University, Japan)
Prof. Dr. Sudibya	(Universitas Sebelas Maret, Indonesia)
Prof. Dr. Rini Widayanti	(Universitas Gadjah Mada, Indonesia)
Prof. Richard Chinomona	(Walter Sisulu University, South Africa)
Dr.med.vet. Tamara Munoz Caro	(Universidad Santo Tomás Talca, Chile)
Dr. Chifumi Takaghi	(National Chung Hsing University, Taiwan)
Abu Musa Md. Talimur Reza, Ph.D.	(Polish Academy of Sciences, Poland)
Amali Alahakoon, Ph.D.	(University of Sri Jayewardenepura, Sri Lanka)
Dewanto Harjunowibowo, Ph.D.	(Universitas Sebelas Maret, Indonesia)

Organizing committee

Dr. Winny Swastike	(Chairman)
Dian Meididewi Nuraini, DVM.	(Secretary)
Endang Tri Rahayu, DVM.	(Finance)
Wari Pawestri, DVM.	(Finance)
Ari Kusuma Wati, M.Sc.	(Publication)
Bayu Setya Hertanto, M.Sc.	(Public Relation and Promotion)
Dr. Ahmad Pramono	(Event)
Aqni Hanifa, M.Si.	(Event)

Table of contents

Volume 1001

2022

◀ Previous issue Next issue ▶

1st International Conference on Animal Research for Eco-Friendly Livestock Industry 23/11/2021 - 23/11/2021 Online

Accepted papers received: 22 February 2022

Published online: 21 March 2022

Open all abstracts

Preface

OPEN ACCESS	011001
Preface	

✚ Open abstract  View article  PDF

OPEN ACCESS	011002
Peer Review Statement	

✚ Open abstract  View article  PDF

Eco-friendly Animal Feed Resources, Nutrition, Forage and Pasture

OPEN ACCESS	012001
Evaluation of gas production kinetics from phyllode and acacia plant (<i>Acacia mangium</i>) pod through fermentation by <i>in vitro</i> gas test	

J W Kusuma, I N Tuti, E Handayanta, A Hanifah, Sudiyono and R F Hadi

✚ Open abstract  View article  PDF

OPEN ACCESS	012002
A study of agricultural waste potential to support the expansion of cattle and buffalo populations in Aceh Barat District	

Y Yusriani, S Y Hayanti, H Saputra, N Qomariyah, F Fitriawaty, M A Da Costa, S Surya and S Asmaireicen

✚ Open abstract  View article  PDF

OPEN ACCESS	012003
Impact of ammoniation rice straw treatment with direct-fed microbials and <i>Hibiscus tiliaceus</i> leaf meal supplemented in concentrate on local sheep performances	

M Bata, S Rahayu, E A Rimbawanto and M Fita

✚ Open abstract  View article  PDF

OPEN ACCESS	012004
Increasing linseed (<i>Linum usitatissimum</i>) meal potential as feed ingredients and source of omega-3 and -6 fatty acids for broilers through processing technology	

O Sjoftjan, D N Adli, N Febrianingrum, H M Ghoezi, M F A Yudatama, M Faalah, M H Natsir and Y F Nuningtyas

✚ Open abstract  View article  PDF

OPEN ACCESS	012005
Effect of nucleotides and turmeric extract on blood protein and body weight of broiler kept in open cages	

E Tugiyanti, E Susanti, I Ismoyowati and R Rosidi

✚ Open abstract  View article  PDF

OPEN ACCESS	012006
The effect of explants and light conditions on callus induction of srikandi putih maize (<i>Zea mays</i> L.)	

E Satrio, M Kurniawan, R Nuzuliyah, N Susebudi and A Y Oulaga



PAPER • OPEN ACCESS

Effect of nucleotides and turmeric extract on blood protein and body weight of broiler kept in open cages

To cite this article: E Tugiyanti *et al* 2022 *IOP Conf. Ser.: Earth Environ. Sci.* **1001** 012005

View the [article online](#) for updates and enhancements.

You may also like

- [The potential of turmeric \(*Curcuma xanthorrhiza*\) in agroforestry system based on silk tree \(*Albizia chinensis*\)](#)
D Purnomo, M S Budiastuti, A T Sakya et al.
- [Addition of turmeric in feed on growth and survival rate of Nilasa red tilapia \(*Oreochromis sp.*\)](#)
R Cahyani, W H Satyantini, D D Nindarwi et al.
- [Potency of Turmeric in Reducing Motile *Aeromonas Septicaemia* \(Mas\) in *Pangasius hypophthalmus*](#)
M Riauaty, T E Y Sari and A Adelina

A promotional banner for 'Free the Science Week 2023' with a dark blue background and a futuristic digital interface. A hand is shown interacting with a circular control panel that has a padlock icon in the center. The text 'Free the Science Week 2023' is in a light blue font, followed by 'April 2-9' in white. Below this, 'Accelerating discovery through' is in white and 'open access!' is in light blue. At the bottom left is the ECS logo and the website 'www.ecsdl.org'. At the bottom right is a blue button with the text 'Discover more!' in white.

Free the Science Week 2023 April 2-9

Accelerating discovery through
open access!

 www.ecsdl.org [Discover more!](#)

Effect of nucleotides and turmeric extract on blood protein and body weight of broiler kept in open cages

E Tugiyanti*, E Susanti, I Ismoyowati, R Rosidi

Faculty of Animal Science, University of Jenderal Soedirman, Indonesia

*E-mail: elly.tugiyanti@unsoed.ac.id

Abstract. Indonesia's wet tropical climate is a condition that does not support the development chickens. Therefore, this study aims to analyze the effect of nucleotides and turmeric extract in basal feed on broilers' blood protein and body weight. Data was collected using a Completely Randomized Design (CRD) with 7 treatments that comprised of negative and positive control. Meanwhile, the negative aspects include basal feed + plus Zink Bactitracin 0.1 g, while the positive control consists of N0K1, N1K0, N1K1, N2K0, and N2K1. The results showed that nucleotide supplementation and turmeric extract in the feed had no significant effect on blood PCV, TPP, albumin levels, and broiler body weights. Furthermore, blood protein increases due to the high levels of nucleotides and turmeric extract. The 5 g of nucleotide supplementation and turmeric extract produce the same broilers blood protein and body weight.

1. Introduction

Indonesia's wet tropical climate is a condition that does not support the development of chickens. Therefore, broilers reared above the thermoneutral zone are vulnerable to environmental heat stress and exhibit behavioral, as well as physiological changes. This study result showed that 75% of the chicken industry was controlled by large breeders and integrator, while the remaining 25% is for smallholder farmers who uses open cages. However, entrepreneurs play an important role in trade dynamics and the country's economy because livestock production facilities such as breed, feed, vitamins, and cage equipment are not available for selling without their presence. These smallholder farmers often face the problems of broiler growth that takes a long time of about 37-42 days before it reaches the target body weight. In Indonesia, nucleotide synthesis is insufficient because the tropical climate conditions cause these chickens to experience heat stress. Nucleotide deficiency tends to reduce the performance of immune organs[1]. Therefore, it is necessary to find safe feed additives including nucleotides and turmeric extract for broilers and the people that consume them.

Turmeric contains curcuminoid compounds consisting of curcumin, desmethoxycumin, bisdemethoxycurcumin, and essential oils, 1-3% fat, 3% carbohydrates, 30% protein, 8% starch, 45-55% vitamin C, as well as mineral salts including iron, phosphorus, and calcium[2]. The curcuminoids and turmeric are efficacious just like antihepatotoxic, anthelmintic, antiedemic, analgesic, anti-inflammatory, and antioxidant[3]. Therefore, turmeric helps to increase appetite, work digestive organs, absorb nutrients in the body, and livestock endurance. This situation affects the ability of broilers to digest and absorb protein.

Nucleotides are semi-essential nutrients because they code and translate genetic information, contribute to energy metabolism, participate in coenzyme structures, regulate lymphocyte maturation,



activate phagocytosis and macrophages, as well as enhance immunological responses[4]. Previous studies indicated that adding nucleotides to feed tends to increase body weight gain and reduce FCR [5]. Moreover, 0.1% nucleotide supplementation help to decrease the effect of infection and *Clostridium perfringens* [6]. This supplementation did not significantly affect the immune organs' performance at different environmental temperatures. Therefore, it needs to be combined with turmeric extract which is antimicrobial and antioxidant to enable chicken growth to remain fast. This study examines the effect of nucleotide supplementation and turmeric extract on broilers' blood protein and body weight.

2. Materials and methods

2.1. Animals, diet, and experimental design

A total of 168 Day-old chic were placed in an open cage with 30 plots measuring 1.5 x 1.5 x 1 m. The starter period basal diet contains 21% protein and EM 3100 kcal/kg. Table 1 shows the nutrient content of the feed. However, BioNutrend® nucleotide Feed Grade produced by Wuhan Sunhy Biology Co. Ltd., China was used at the level of 0.25 g and 0.5 g [7] because the extra turmeric used is 0.6g [8].

Table 1. Nutrient content of the treatment.

Treatment	DM(%)	% DM				
		CP	EE	CF	Ash	NFE
NC	89.10	14.54	4.92	7.97	15.11	57.46
PC	89.33	13.98	5.52	7.56	15.86	57.08
N0K1	90.31	14.79	5.66	7.79	16.59	55.17
N1K0	90.08	14.02	5.80	7.00	15.18	58.01
N1K1	90.07	14.70	5.60	10.60	13.29	55.82
N2K0	90.45	14.89	5.60	11.16	15.68	52.66
N2K1	90.51	18.07	5.62	11.95	16.23	48.13

Table 1 consists of DM, CP, EE, CF, NFE, NC, PC, N0K1, N1K0, N1K1, N2K0, and N2K1.

Data were collected using a randomized design with 7 treatments that are comprised of negative and positive control. Meanwhile, the negative aspects consist of basal feed+zinc batriacin 0.1g, while the positive contains N0K1, N1K0, N1K1, N2K0, and N2K1. Turmeric extract is calculated based on the percentage of dry matter. Therefore, there are 28 experimental units because treatment was repeated 4 times.

2.2. Research procedure

Disease control was performed by giving NCD vaccines through eye drops while the chickens were 4 days old. Meanwhile, the vitamin was given a day before the broilers were vaccinated, but the weighing was performed weekly. The PCV measurement is performed by taking a specimen in venous or capillary blood from the brachial vein. Blood was put into 2/3 or 3/4 of two microhematocrit tubes where one part was closed using clay or a micro burner. However, the speed of centrifuge for 5 minutes is 11,000-16,000 rpm using a microhematocrit calculator. The result tends to be re-examined if it has a difference of $\pm 2\%$. Meanwhile, TPP was measured using the biuret method with Automated Chemical Analyzer Diasys Response 920, while albumin was calculated by centrifuging 3-5 ml to obtain serum or plasma. The reagents and specimens were allowed to stand at room temperature.

2.3. Statistical analysis

Data were analyzed using variance with the help of SPSS IBM version 26.

3. Results and discussion

High ambient temperature negatively affects broilers' physiological condition and productivity. Therefore, chickens especially those above three weeks old find it difficult to adapt to changes in

environmental temperatures. These physiological conditions are indicated through the protein profile of the broilers' blood and growth.

Table 2. Effect of nucleotide supplementation and turmeric extract on PVC, TPP, albumin, and broiler body weight.

Parameter	Treatment							Sig.
	NC	PC	N0K1	N1K0	N1K1	N2K0	N2K1	
PCV (%)	19.5 ± 0.3	21.25 ± 0.6	20.75 ± 0.9	23.0 ± 1.3	21.8 ± 0.9	23.5 ± 0.9	24.3 ± 1.2	ns
	2.35 ± 0.1	2.7 ± 0.1	2.5 ± 0.1	2.7 ± 0.1	2.5 ± 0.1	2.65 ± 0.2	3.1 ± 0.2	
TPP (g/dl)	40.5 ± 2.0	46.6 ± 1.7	43.1 ± 2.2	46.6 ± 2.1	43.1 ± 2.2	45.7 ± 3.9	53.4 ± 4.3	ns
	824.8 ± 36.7	830 ± 63.4	919 ± 27.6	863 ± 22.2	868.8 ± 26.0	840.5 ± 30.1	868.3 ± 22.4	

Table 2 comprises NC, PC, N0K1, N1K0, N1K1, N2K0, N2K1, PCV, TPP, and BW.

3.1. Blood protein

The comfort zone environment tends to affect chickens' feed and consumption, blood protein, as well as productivity including growth and body weight [18]. Meanwhile, high temperatures cause broilers to experience heat stress and dehydration which damage blood viscosity and proteins [19].

Table 2 shows the average blood protein including packed cell volume, total plasma protein, and albumin levels. The analysis result showed that nucleotide supplementation and turmeric extract did not significantly affect broilers' PCV, TPP, and albumin levels in tropical climates ($P < 0.05$). However, the highest pack cell volume in the N2K1 treatment was 24.3 ± 1.2 %. Table 2 shows that adding 0.6 g of turmeric extract on N0K1 led to a lower PCV than the positive control. This is in line with the study of Hartati [9] that the chickens' pack cell volume of 100 gr/kg BW was lower than PC. The addition of 0.25 g of nucleotides in the N1K1 treatment leads to a higher PCV value. Although, the increase in nucleotide is not significant but was directly proportional to the development of pack cell volume. Meanwhile, heat stress conditions do not allow chickens to meet nucleotides adequacy through the de novo and salvage pathway mechanisms [20].

The PCV indicates the blood's ability to transport oxygen and nutrients [10] because its high blood viscosity reduces blood flow and improves heart performance. Meanwhile, a low pack cell volume shows that nutrient absorption is inhibited and lead to erythrocyte damages [10], while normal PCV range from 22-35% [11]. In this study, the PCV value was normal at N1K0, N2K0, and N2K1 but was low at other treatments. This is possible because de novo synthesis does not meet the needs of erythrocyte growth and differentiation.

The same thing was indicated in TPP and albumin with the highest N2K1 treatment of 3.1 ± 0.2 g/dl and 53.4 ± 4.3 g/dl respectively. However, the combination of 0.5 g nucleotide supplementation and 0.6 g turmeric extract lead to higher TPP and albumin values. This is because nucleotides play a role in the growth and replacement of damaged cells. Also, the content of curcumin in turmeric extract acts as an anti-inflammatory to protect hepatocyte cells and prevents inflammation in the liver to enable protein synthesis to occur optimally [12].

3.2. Body weight

Under tropical climates, broiler production is encumbered by several constraints which make it difficult in attaining their genetic potential. Therefore, the harsh environmental conditions with thermal stress become one of the challenges that hinder chicken optimal growth.

Table 2 shows the average body weight of broilers. The analysis result showed that nucleotide supplementation and turmeric extract do not significantly affect chickens' bodyweight in tropical climates ($P < 0.05$). This is indicated by using nucleotides from *Saccharomyces cerevisiae* in broiler [13].

Although, the increase in nucleotides is not significant but leads to the development of body weight. Meanwhile, the availability of nucleotides helps to improve intestinal performance by enhancing intestinal villi and gut mucosal immunity [7,14,15]. The nucleotides failed to increase feed intake despite its development in body weight [5]. Therefore, the increase in nucleotide without turmeric extract leads to low body weight. This shows that additional nucleotides are needed to optimize de novo synthesis at a high temperature [16].

The highest body weight was 919 ± 27.6 g in the N0K1 treatment because turmeric extract contains antioxidants and antimicrobials that reduce lipid peroxidation and improve intestinal morphology and ecology, as well as the immune response and absorption of body nutrients [3]. In heat stress, 0.5% turmeric extract supplementation had better body weight gain and feed intake [17]. Therefore, the combination of the two results is higher than the control with 868.3 ± 22.4 grams.

4. Conclusions

In a wet tropical environment, a total of 5 g of nucleotides and turmeric extract produce the same blood protein and body weight

Conflict of interest

The author certifies that there is no conflict of interest with any financial organization regarding this study

Acknowledgments

The author thanks the Chancellor of Jenderal Soedirman University, and Wuhan Sunhy Biology co. Ltd.

References

- [1] Hakim R L, Mahfudz L D and Muryani R 2021 *J. Sain Peternak. Indones.* **16** 164
- [2] Simanjuntak P 2012 *AGRIUM: Jurnal Ilmu Pertanian* **17** 103
- [3] Lagua E B and Ampode K M B 2021 *J. Anim. Health Prod.* **9** 243
- [4] Sauer N, Mosenthin R and Bauer E 2011 *Nutr. Res. Rev.* **24** 46
- [5] Salah M, Suprijatna E, Djauhari M L and Y B I Vitus D 2021 *Livest. Res. Rural Dev.* **31** 174
- [6] Mohamed F F, Hady M M, Kamel N F and Ragaa N M 2020 *Vet. Anim. Sci.* **10** 100130
- [7] Sakinah S, Djauhari L M and Sunarti D 2019 *Pros. Sem. Nas. (Magelang: Indonesia)* p 1
- [8] Kafi A, Uddin M N, Uddin M J, Khan M M H and Haque M E 2017 *Int. J. Poult. Sci.* **16** 257
- [9] Hartati S, Widiyono I, Raharjo S, Purnamaningsih H, Nururrozi A and Fitriana I 2021 *Jurnal Ilmu Peternakan dan Veteriner Tropis* **2** 6
- [10] Alghazali F, Suharyati S and Santosa P E 2018 *Jurnal Riset dan Inovasi Peternakan* **2** 1
- [11] Jain N C 1993 *Essentials of Veterinary Hematology* (New York: New York University)
- [12] Widhyari S D, Esfandiari A and Herlina 2011 *Jurnal Ilmu Pertanakan. Indonesia* **16** 179
- [13] Safaei M and Hassanabadi A 2020 *Poult. Sci. J.* **8** 135
- [14] Wu C, Yang Z, Song C, Liang C, Li H, Chen W, Lin W and Xie Q 2018 *Poult. Sci.* **97** 3837
- [15] Jung B and Batal A B 2012 *Br. Poult. Sci.* **53** 98
- [16] de Souza C, Eyng C, Viott A M, de Avila A S, Pacheco W J, Junior N R, Kohler T L, Tenorio K I, Cirilo E H and Nunes R V. 2021 *Livest. Sci.* **251** 104659
- [17] Kanani B, Daneshyar P and Najafi M 2016 *Poult. Sci. J.* **4** 47
- [18] Ibrahim A A, Aliyu J, Abdu M I, Hassan A M 2012 *World Applied Sciences Journal* **16** 433
- [19] Jackson M L 2007 *Veterinary Clinical Pathology: An Introduction* (Iowa: Blackwell Publishing) p 25
- [20] Kruger D and Werf M V D 2018 *Benefits of Nucleotide Supplementation in Poultry* (Ohly Application Note: Ohly.com) p 1