

Journal of Bioscience and Bioengineering



Journal of
Bioscience and
Bioengineering



April 2023 • Volume 17(2) • Issue 1

Journal of Bioscience and Bioengineering

Volume 17(2) • April 2023
ISSN: 1547-4326
CODEN: JBIO-D



ScienceDirect®

Journal of Bioscience and Bioengineering

5.4

CiteScore

3.185

Impact Factor

[Submit your article](#)[Guide for authors](#)[Menu](#)[Search in this journal](#)[> Aims and scope](#)[> **Editorial board**](#)[> Abstracting & indexing](#)[> News](#)[> Conferences](#)[> Cover Gallery](#)

Editorial board by country/region

36 editors and editorial board members in 11 countries/regions

1 Japan (21)

2 Germany (3)

3 China (2)

[> See more editors by country/region](#)[FEEDBACK](#)

Editorial board

Editor-in-Chief



Noriho Kamiya

National University Corporation Kyushu University, Fukuoka, Japan

[View full biography](#)

Editorial Board



Yoshiteru Aoi

Hiroshima University, Higashihiroshima, Japan

[View full biography](#)

Atsushi Arakaki

Tokyo University of Agriculture and Technology, Fuchu, Japan

Virendra Swarup Bisaria

Indian Institute of Technology Delhi, New Delhi, India



Lars M. Blank, PhD

RWTH Aachen University Institute of Applied Microbiology, Aachen, Germany

[View full biography](#)



Jo-Shu Chang, Ph.D.

National Cheng Kung University, Tainan, Taiwan


[View full biography](#)

Howard Chase

University of Cambridge, Cambridge, United Kingdom

I-Ming Chu

National Tsing-Hua University, Department of Chemical Engineering, Hsinchu, Taiwan



Toshiaki Fukui

Tokyo Institute of Technology School of Life Science and Technology, Yokohama, Japan

[View full biography](#)



Taiki Futagami

Kagoshima University, Kagoshima, Japan

[View full biography](#)



Hiroyuki Futamata

Shizuoka University, Shizuoka, Japan

Kiyotaka Hara

University of Shizuoka, Shizuoka, Japan



Hidehiko Hirakawa

University of Tsukuba, Tsukuba, Japan

[View full biography](#)



Tomoyuki Hori

National Institute of Advanced Industrial Science and Technology Tsukuba Center Tsukuba Central,
Tsukuba, Japan

Wei-Shou Hu

University of Minnesota Twin Cities, Minneapolis, Minnesota, United States of America

Shu Ishikawa

Kobe University Graduate School of Science, Technology and Innovation, Kobe, Japan

Yoshinori Kawabe

Kyushu University, Faculty of Engineering Graduate School of Engineering, Department of Chemical
Engineering, Fukuoka, Japan



Mee-Hae Kim

Osaka University, Suita, Japan

[View full biography](#)

Takaaki Kojima

Meijo University, Nagoya, Japan

Masaaki Konishi, PhD

Kitami Institute of Technology, Kitami, Japan



Sang Yup Lee, PhD/Distinguished Professor

Korea Advanced Institute of Science and Technology, Daejeon, South Korea

[View full biography](#)



Jun-ichi Maruyama

The University of Tokyo, Bunkyo-Ku, Japan

[View full biography](#)

Kazuo Masaki

National Research Institute of Brewing, Hiroshima, Japan

Seiichi Ohta

The University of Tokyo Graduate School of Engineering, Faculty of Engineering, Department of Technology Management for Innovation, Bunkyo-Ku, Japan



Kenji Okano

Kansai University, Suita, Japan

Watanalai Panbangred

Mahidol University Departement of Biotechnology, Bangkok, Thailand

Chit-Laa Poh

Sunway University, Bandar Sunway, Malaysia

Andreas Schmid

Helmholtz-Centre for Environmental Research - UFZ, Leipzig, Germany

Francesco Secundo

Institute of Chemistry of Molecular Recognition National Research Council, Milano, Italy

Seiji Takahashi

Tohoku University, Sendai, Japan

Yukihiro Tashiro

National University Corporation Kyushu University, Fukuoka, Japan

Alisa Vangnai

Chulalongkorn University, Bangkok, Thailand



Kazuya Yamanaka

Kansai University, Suita, Japan

[View full biography](#)

Oskar Zelder

BASF SE, Ludwigshafen, Germany

Chong Zhang

Tsinghua University, Department of Chemical Engineering, Beijing, China

Jian-Jiang Zhong, PhD

Shanghai Jiao Tong University, Shanghai, China

All members of the Editorial Board have identified their affiliated institutions or organizations, along with the corresponding country or geographic region. Elsevier remains neutral with regard to any jurisdictional claims.



Copyright © 2023 Elsevier B.V. or its licensors or contributors.
ScienceDirect® is a registered trademark of Elsevier B.V.

 RELX™



ScienceDirect®

Journal of Bioscience and Bioengineering

5.4

CiteScore

3.185

Impact Factor

[Submit your article](#)[Guide for authors](#)[Menu](#)[Search in this journal](#)

Volume 101, Issue 3

Pages 203-286 (March 2006)

[Download full issue](#)[Previous vol/issue](#)[Next vol/issue](#)

Receive an update when the latest issues in this journal are published

[Sign in to set up alerts](#)

Research article ○ Abstract only

Immunogenic and allergenic potentials of natural and recombinant innocuous proteins

Tsukasa Matsuda, Takeshi Matsubara, Shingo Hino

Pages 203-211

Article preview ✓

Research article ○ Abstract only

FEEDBACK

N-linked glycan structures of a mouse monoclonal antibody produced from tobacco BY2 suspension-cultured cells

Kazuhito Fujiyama, Ryo Misaki, Atsuya Katsura, Tomoko Tanaka, ... Tatsuji Seki

Pages 212-218

Article preview 

Research article ☐ Abstract only

Bioconversion of D-psicose to D-tagatose and D-talitol by *Mucoraceae* fungi

Kazutoshi Yoshihara, Yoshihiro Shinohara, Takahiro Hirotsu, Ken Izumori

Pages 219-222

Article preview 

Research article ☐ Abstract only

Effect of salt concentration on intracellular accumulation of lipids and triacylglyceride in marine microalgae *Dunaliella* cells

Mutsumi Takagi, Karseno, Toshiomi Yoshida

Pages 223-226

Article preview 

Research article ☐ Abstract only

Effects of methanol feeding methods on chimeric α -amylase expression in continuous culture of *Pichia pastoris*

Atsushi Nakano, Chun Yeon Lee, Arei Yoshida, Takehiro Matsumoto, ... Shigeo Katoh

Pages 227-231

Article preview 

Research article ☐ Abstract only

Effects of water-soluble carbohydrate content on silage fermentation of wheat straw

Hong Yan Yang, Xiao Fen Wang, Jian Bin Liu, Li Juan Gao, ... Zong Jun Cui

Pages 232-237

Article preview 

Research article ○ Abstract only

Morphology of spheroidal hepatocytes within injectable, biodegradable, and thermosensitive poly(organophosphazene) hydrogel as cell delivery vehicle

Keun-Hong Park, Soo Chang Song

Pages 238-242

Article preview 

Research article ○ Abstract only

Gene expression property of high-density three-dimensional tissue of HepG2 cells formed in radial-flow bioreactor

Tomokatsu Hongo, Mariko Kajikawa, Seiichi Ishida, Shogo Ozawa, ... Hiroyuki Honda

Pages 243-250

Article preview 

Research article ○ Abstract only

Production of Fab fragment corresponding to surface protein antigen of *Streptococcus mutans* serotype c-derived peptide by *Escherichia coli* and cultured tobacco cells

Akira Yano, Masataka Takekoshi, Eiko Morita, Susumu Imai, ... Nobuhiro Hanada

Pages 251-256

Article preview 

Research article ○ Abstract only

Fate of transforming bacterial genome following incorporation into competent cells of *Bacillus subtilis*: a continuous length of incorporated DNA

Yukiko Saito, Hisataka Taguchi, Takashi Akamatsu

Pages 257-262

Article preview 

Research article ○ Abstract only

Cloning and characterization of *nif* structural and regulatory genes in the purple sulfur bacterium, *Halorhodospira halophila*

Hisayoshi Tsuihiji, Yoichi Yamazaki, Hironari Kamikubo, Yasushi Imamoto, Mikio Kataoka

Pages 263-270

Article preview 

Research article  Abstract only

Microbial community in methanogenic packed-bed reactor successfully operating at short hydraulic retention time

Kengo Sasaki, Shin Haruta, Masahiro Tatara, Akira Yamazawa, ... Yasuo Igarashi

Pages 271-273

Article preview 

Research article  Abstract only

Robustness of cascade pH and dissolved oxygen control in symbiotic nisin production process system of *Lactococcus lactis* and *Kluyveromyces marxianus*

Agustin K. Wardani, Sunao Egawa, Keisuke Nagahisa, Hiroshi Shimizu, Suteaki Shioya

Pages 274-276

Article preview 

Research article  Abstract only

Quantitative expression analysis of inulinase gene cluster of *Penicillium* sp. strain TN-88

Satoshi Moriyama, Michio Muguruma, Kazuyoshi Ohta

Pages 277-279

Article preview 

Research article  Abstract only

Synthesis and structure analysis of novel disaccharides containing D-psicose produced by endo-1,4- β -D-xylanase from *Aspergillus sojae*

Hisaka Oshima, Isao Kimura, Ken Izumori

Pages 280-283

Article preview 

Research article  Abstract only

Generation of monoclonal antibodies using simplified single-cell reverse transcription-polymerase chain reaction and cell-free protein synthesis

Muhamad Ali, Kiyotaka Hitomi, Hideo Nakano

Pages 284-286

Article preview 

[< Previous vol/issue](#)

[Next vol/issue >](#)

ISSN: 1389-1723

Copyright © 2023 The Society for Biotechnology, Japan. All rights reserved



Copyright © 2023 Elsevier B.V. or its licensors or contributors.
ScienceDirect® is a registered trademark of Elsevier B.V.



Rp 133,0 JT	Rp 108,0 JT	Rp 170,0 JT	Rp 145,0 JT	
Beli Mobil Pasti Dapat Garansi				
214 Indonesia				

Journal of Bioscience and Bioengineering

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
Netherlands  Universities and research institutions in Netherlands  Media Ranking in Netherlands	Biochemistry, Genetics and Molecular Biology Biotechnology Chemical Engineering Bioengineering Immunology and Microbiology Applied Microbiology and Biotechnology	Elsevier	116
PUBLICATION TYPE	ISSN	COVERAGE	INFORMATION
Journals	13474421, 13891723	1990-1991, 1993-1994, 1999-2021	Homepage How to publish in this journal jbb@sbj.or.jp



SCOPE

The Journal of Bioscience and Bioengineering is a research journal publishing original full-length research papers, reviews, and Letters to the Editor. The Journal is devoted to the advancement and dissemination of knowledge concerning fermentation technology, biochemical engineering, food technology and microbiology.

Join the conversation about this journal

Online paper Publication

DOI for Published papers

Indexing :- Google Scholar, Index Copernicus, Scribd, Academia and more

ijisrt.com

OPEN

Truly International Journal

Refereed Journal

200 + Reviewer from 70 countries.

iosrjournals.org

OPEN

FIND SIMILAR JOURNALS ?

1

Applied Microbiology and
Biotechnology

DEU

72%

similarity

2

Bioresources and
Bioprocessing

DEU

71%

similarity

3

Critical Reviews in
Biotechnology

GBR

67%

similarity


4



AMB Express

DEU

6

s





Journal of Bioscience and Bioengineering

Q2 Applied Microbiology and Biotechnology best quartile

SJR 2021 0.61

powered by scimagojr.com


← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com" data-bbox="634 301 784 316">
```

SCImago Graphica

Explore, visually communicate and make sense of data with our [new data visualization tool](#).



Online paper Publication - DOI for Published papers

Indexing :- Google Scholar, Index Copernicus, Scribd, Academia and more ijsrt.com

Metrics based on Scopus® data as of April 2022

A **Andi Rohandi** 3 years ago

good

reply

Leave a comment

▼

☐ I'm not a robot

reCAPTCHA
Privacy - Terms

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2022. Data Source: Scopus®

EST MODUS IN REBUS

Horatio (Satire 1.1.106)

[Cookie settings](#)

[Cookie policy](#)



Effect of Salt Concentration on Intracellular Accumulation of Lipids and Triacylglyceride in Marine Microalgae *Dunaliella* Cells

Mutsumi Takagi,¹* Karseno,¹ and Toshiomi Yoshida¹

International Center for Biotechnology, Osaka University, 2-1 Yamada-oka, Suita, Osaka 565-0871, Japan¹

Received 14 September 2005/Accepted 6 December 2005

In order to get the high liquefaction yield from marine algae cell mass to fuel oil, the effect of salt stress on the accumulation of lipids and triacylglyceride in *Dunaliella* cells was investigated. Although initial NaCl concentration higher than 1.5 M markedly inhibited cell growth, increase of initial NaCl concentration from 0.5 (equal to sea water) to 1.0 M resulted in a higher intracellular lipid content (67%) in comparison with 60% for the salt concentration of 0.5 M. Addition of 0.5 or 1.0 M NaCl at mid-log phase or the end of log phase during cultivation with initial NaCl concentration of 1.0 M further increased the lipid content (70%).

[Key words: algae, salt stress, liquefaction, lipid]

Since atmospheric CO₂ accumulation has a serious effect on the global environment, the control of total CO₂ emission into the atmosphere is considered to be an important issue related to the biosphere. Marine microalgae are expected to play an important role in resolving this problem because they have a high capability for photosynthesis and grow well in the sea which solubilizes a high amount of CO₂ and which accounts for 70% of the surface area of the earth. There are two major approaches to large-scale CO₂ fixation by marine microalgae.

One is the production of non-carbon energy sources such as hydrogen gas from the algal biomass. The other is the conversion of algal biomass into liquid fuel oil by thermochemical liquefaction reactions at a high temperature and pressure (1). The latter process is expected to stimulate CO₂ circulation on earth, and is considered to be a simpler and more appropriate process for worldwide implementation than the former.

One of the most important criteria in assessing the performance of thermochemical liquefaction is the oil yield from the organic materials in the algal biomass. A report on liquefaction using various model substrates revealed that a high liquefaction yield was obtained from cell components containing hydrophobic compounds such as lipids, fatty acids and fatty acid esters (2). Among marine microalgae species, cells of the genus *Nannochloris* are known to contain a large amount of intracellular lipids (3, 4). It was reported that the intracellular triacylglyceride content increased after nitrate depletion during a cultivation of *Nannochloris* sp. UTEX LB1999 cells in modified NORO medium containing 9.9 mM KNO₃ even though the intracellular lipid con-

tent did not increase and the oil yield following liquefaction of the cells increased in proportion to the intracellular triacylglyceride content (5). When 0.9 mM nitrate was intermittently fed 10 times during the log phase in addition to initial nitrate feed (0.9 mM), the concentration of *Nannochloris* sp. UTEX LB1999 reached almost the same (2.16 g/l) and the contents of intracellular lipids and the percentage of triacylglycerides in the lipids were respectively increased from 31.0% to 50.9% and 26.0% to 47.6%, compared with those of cells cultured in a modified NORO medium containing 9.9 mM KNO₃ without additional nitrate feeding (6).

The composition of intracellular lipid of microalgae was reported to change in response to environmental salinity. Increase of NaCl concentration from 0.4 M to 4 M increased saturated and monounsaturated fatty acids in *Dunaliella* cells isolated from an Antractic hypersaline lake (7), while polyunsaturated fatty acid decreased. The fatty acid composition of polar lipid in *Dunaliella salina* Teodoresco was affected significantly by the change in NaCl concentration (8). The percentage of saturated fatty acid decreased as the concentration of NaCl increased, while the percentage of highly unsaturated fatty acid increased (9). *Dunaliella tertiolecta* cells contain large amount of lipid and are highly salt tolerant (10), which might be appropriate for the large scale outdoor cultivation.

In this study, loading of salt stress during cultivation of *Dunaliella* cells for highly intracellular accumulation of lipids and triacylglycerides was investigated, because the algae cell mass having high contents of lipids and triacylglycerides should result in a high liquefaction yield.

MATERIALS AND METHODS

Algal strain and media *D. tertiolecta* ATCC 30929 were used in this study as a marine CO₂-fixing microalgae strains and grown in modified NORO medium which had the following composition (per liter): NaCl, 29.2 g; KNO₃, 1.0 g (9.9 mM); MgCl₂·H₂O, 1.5 g;

* Corresponding author. e-mail: takagi-m@eng.hokudai.ac.jp
phone/fax: +81-(0)11-706-6567

[§] Present address: Division of Biotechnology and Macromolecular Chemistry, Graduate School of Engineering, Hokkaido University, Sapporo 060-8628, Japan.

MgSO₄·7H₂O, 0.5 g; KCl, 0.2 g; CaCl₂, 0.2 g; K₂HPO₄, 0.045 g; tris(hydroxymethyl)aminomethane, 2.45 g; EDTA·2Na, 1.89 mg; ZnSO₄·7H₂O, 0.087 mg; H₃BO₃, 0.61 mg; CoCl₂·6H₂O, 0.015 mg; CuSO₄·5H₂O, 0.06 mg; MnCl₂, 0.23 mg; (NH₄)₆Mo₇O₂₄·4H₂O, 0.38 mg; Fe(III)-EDTA, 3.64 mg; the pH was adjusted to 8.0 with 1 N HCl.

Cultivation Twenty ml of the modified NORO medium in a 100-ml Erlenmeyer flask was inoculated with the cells (OD₆₈₀ 0.05) and incubated at 28°C with reciprocal shaking (60 rpm) for 6 d. The light intensity on the surface of the flask was adjusted to 65 μmol·s⁻¹·m⁻² using fluorescent lamps. The flask cultivation was transfer to 500 ml of the same fresh medium having salt concentration 0.5 or 1.0 M in a Roux bottle to obtain OD₆₈₀ of 0.05. The temperature, light intensity, and aeration conditions were 30°C, 150 μmol·s⁻¹·m⁻² and 250 ml/min CO₂-enriched air (3% CO₂), respectively. Concentrated solution of NaCl (5.0 M) was fed at log phase, the end of log phase, or stationary phase during cultivation. Besides, test-tube static cultivation (10 ml) was performed.

Determination of cell and nitrate concentrations The cell concentration was determined by measuring OD₆₈₀ and converted to ash-free dry cell weight employing the coefficient of OD₆₈₀ to cell concentration of 0.01 (g·l⁻¹·UOD⁻¹). The precipitate obtained by centrifugation (3000 rpm, 15 min) of a 10-ml culture was heated overnight at 105°C and weighed (w1 g). After carbonization of the precipitate for 3 min on a Bunsen burner, it was weighed again (w2 g). The ash-free dry cell weight (g/l) was determined by dividing the difference between w1 and w2 by the culture volume (0.01 l). The culture supernatant was diluted with the medium without KNO₃ and the nitrate concentration was determined from the difference between its absorbance values at 221.4 and 232.0 nm.

Determination of osmolality The osmolality of culture supernatant was determined using an osmometer (model OM-801; Vogel, Berlin, Germany).

Determination of the contents of lipids and triacylglycerides A culture sample containing about 30 mg cell was centrifuged and the resulting precipitate was washed with 1% NaCl. After extraction of lipids from the precipitate with methanol-chloroform (2:1), chloroform and 1% NaCl solution was added to adjust the ratio of methanol, chloroform and water to 2:2:1. All the chloroform layer collected three times were evaporated, dried in a desiccator, and weighed as the total lipid (3). The percentage of triacylglyceride (TG) in the total lipids solubilized in isopropyl alcohol was determined by the glycerol-3-phosphate oxidase-*p*-chlorophenol method (triacylglyceride G-test Wako; Wako Pure Chemical, Osaka), with triolein as the standard.

RESULTS

Effect of initial NaCl concentration on the cell growth

To clarify the effect of initial NaCl concentration on cell growth of *D. tertiolecta* ATCC 30929, test tube cultivations were performed with several initial NaCl concentrations (Fig. 1). Cell concentration decreased as NaCl concentration increased from 1.0 M to 2.0 M, while there was no decrease in the range of NaCl concentration less than 1.0 M. There should be marked growth inhibition by NaCl concentration higher than 1.0 M. Consequently, initial NaCl concentration less than 1.0 M was considered to be appropriate to achieve high cell concentration.

Effect of initial NaCl concentration on the intracellular lipid content To confirm the effect of initial NaCl concentration less than 1.0 M on intracellular lipid content, Roux bottle cultivations were performed employing two




FIG. 1. Effect of initial NaCl concentration on cell growth. Test tube cultivations were performed with several initial NaCl concentrations. The average of duplicate cultivations was shown.




FIG. 2. Cell growth with high initial NaCl concentration. During cultivations in Roux bottle with initial NaCl concentrations of 0.5 (circles) or 1.0 M (triangles), OD₆₈₀ (open symbols) and nitrate concentration (closed symbols) were measured. The average of duplicate cultures was shown.

kinds of initial NaCl concentrations (0.5 and 1.0 M). Cell growth with initial NaCl concentration of 1.0 M was almost similar to the culture with initial NaCl concentration of 0.5 M, which was equivalent to that of sea water (Fig. 2). There was no apparent difference in the course of nitrate concentration during these cultures. The intracellular contents of lipids and triacylglyceride in lipid of cells harvested at the end of cultivation with initial NaCl concentration 1.0 M (67% and 56%) were markedly higher than those in the culture with 0.5 M NaCl (60% and 40%) as shown in Table 1.

Effect of NaCl addition during culture on cell growth and lipid content

To investigate the effect of NaCl addi-

TABLE 1. Effect of initial NaCl concentration on intracellular contents of lipid and TG in lipid

Initial NaCl (M)	Osmolality (Osm/kg)	Cell (g/l)	Lipid (%)	TG in lipid (%)
0.5	1.10±0.05	1.00±0.10	60.6±0.5	40.8±2.9
1.0	1.87±0.15	1.03±0.05	67.8±0.7	56.6±1.8

The intracellular content of lipid and TG content in lipid of cells harvested at the end of Roux bottle cultivations with initial NaCl concentrations of 0.5 and 1.0 M shown in Fig. 2 were determined. Osmolality of culture supernatant was determined at the end of cultures. The average of duplicate cultures and the deviation were shown.




FIG. 3. Addition of NaCl during the culture with initial NaCl concentration of 1.0 M. During cultivation in Roux bottle with initial NaCl concentration of 1.0 M, NaCl was added at mid-log phase (triangles, 0.5 M) or the end of log phase (reverse triangles, 0.5 M; squares, 1.0 M). The data without NaCl addition during the culture was shown with circles. The addition times are indicated by arrows. OD₆₈₀ (open symbols) and nitrate concentration (closed symbols) were measured. The average of duplicate cultures was shown.

tion on cell growth and lipid content, several modes of NaCl addition were performed during the culture with initial NaCl concentration of 1.0 M. Namely, (i) addition of 0.5 or 1.0 M NaCl at the end of log-phase, (ii) addition of 0.5 M NaCl at mid-log phase. The intracellular contents of lipid and TG in lipids were measured at the end of cultivation. There was not marked difference in the course of OD₆₈₀ and nitrate concentration except for the short time lag of growth after the addition 1.0 M NaCl at the end of log phase (Fig. 3). The final volume of culture with the addition was higher than that without the addition because of the addition. Although the final cell concentration and cell mass decreased by the NaCl addition during culture, the influence was smaller than the marked inhibition by initial high concentration of NaCl shown in Fig. 1. Lipid contents in the culture with NaCl addition (70.6–71.4%, Table 2) were apparently higher than that without addition (63.5%). Consequently, the addition of NaCl (0.5 or 1.0 M) during culture with initial NaCl concentration of 1.0 M could increase lipid content with a slight decrease in cell mass.

DISCUSSION

Increase of initial NaCl concentration from 0.5 M (equal to Sea water) to 1.0 M did not decrease cell concentration, which might due to salt tolerant character of *Dunaliella* cells (8, 11). However, initial NaCl concentration higher than 1.0 M markedly inhibit cell growth. Consequently, ini-

tial NaCl concentration less than 1.0 M was considered to be appropriate to achieve high cell concentration.

Although the cultivations with initial NaCl concentration of 0.5 and 1.0 M showed similar time course of cell and nitrate concentrations, lipid content of cells cultivated with 1.0 M initial NaCl (67%) was higher than that with 0.5 M NaCl (60%) as shown in Table 1. Consequently, the reason for higher lipid content might be not nitrate limitation but high NaCl concentration. However, further increase in initial NaCl concentration should not be good strategy, because marked decrease in cell concentration by further increase in initial NaCl concentration was anticipated from the results of Fig. 1.

Addition of 0.5 and 1.0 M NaCl at mid-log phase or the end of log phase during cultivation with initial NaCl concentration of 1.0 M gave the significant increase in lipid content (Fig. 3). Increase of initial NaCl concentration from 1.0 M to 1.5 and 2.0 M resulted in the 42% and 86% decrease in cell concentration (Fig. 1). However, the decrease in final cell mass by the addition of 0.5 and 1.0 M NaCl were only 6–10% and 17%, respectively (Fig. 3), although cell concentration at the time of NaCl addition was less than half of final cell concentration. Consequently, there might be the possibility of cell adaptation to high NaCl concentration by gradual increase in NaCl concentration, while mechanism is not clear.

The reversible change of cell size in response to water potential was reported (12). Microscopic observation in this study revealed that the size of most algae cells decrease after the addition of NaCl (data not shown). Ratio of smaller cells after 2.0 M addition was apparently higher than that with 1.0 M addition. Cell size recovered within 48 h after the additions. *Dunaliella* cells was reported to secrete glycerol in response to increase of NaCl concentration (13). The increase in lipid content may correlate with the adaptive response to high NaCl concentration such as cell volume change and glycerol production. However, the mechanism of lipid content increase by high NaCl concentration is not clear.

The amount of cell mass harvested from culture may decrease somewhat by NaCl addition during culture as shown in Table 2. The final objective of this process is achievement of not higher lipid and TG productivity's but higher oil productivity from cell mass by liquefaction reaction, which is the product of cell concentration and the oil yield. Therefore, the quantitative correlation between oil yield and cell composition such as lipid content and TG content in lipid should be studied in future, in order to find the optimum operation condition about the time and amount of NaCl addi-

TABLE 2. Effect of NaCl addition on the contents of lipid and TG in lipid

Culture	NaCl addition		Cell (mg/ml)	Volume (ml)	Cell (mg)	Lipid (%)	TG in lipid (%)
	Amount (M)	Phase					
1	—	—	0.92±0.02	360	331	63.5±1.0	30.4±3.5
2	1.0	End of log	0.68±0.04	400	272	70.6±3.9	34.5±2.2
3	0.5	Middle of log	0.76±0.06	390	296	71.4±2.3	34.3±3.7
4	0.5	End of log	0.72±0.05	430	309	70.6±1.5	31.8±6.0

Several amounts of NaCl was added at several culture phase during the culture with initial NaCl concentration of 1.0 M as shown in Fig. 3. The intracellular contents of lipid and TG in lipid of cells harvested at the end of each culture were determined. The average of duplicate cultures and deviation were shown.

tion to the culture with initial NaCl concentration of 1.0 M.

In conclusion, the cultivation of *Dunaliella* cells with the high initial NaCl concentration of 1.0 M and the addition of 1.0 M NaCl during culture resulted in high intracellular lipid content and high percentage of triacylglyceride in the lipid. This cultivation technique should contribute to the increase of fuel oil productivity in the combined process of marine microalgae culture and thermo-chemical liquefaction.

REFERENCES

1. Kishimoto, M., Okakura, T., Nagashima, H., Minowa, T., Yokoyama, S., and Yamaberi, K.: CO₂ fixation and oil production using micro-algae. *J. Ferment. Bioeng.*, **6**, 479–482 (1994).
2. Evans, R. J. and Felbeck, G. T., Jr.: High temperature simulation of petroleum formation III. Effect of organic starting material structure on hydrocarbon formation. *Org. Geochem.*, **4**, 153–160 (1983).
3. Ben-Amozi, A. and Tornabene, T. G.: Chemical profile of selected species of microalgae with emphasis on lipid. *J. Phycol.*, **21**, 77–81 (1985).
4. Negoro, M., Shioji, N., Miyamoto, K., and Miura, Y.: Growth of microalgae in high CO₂ gas and effects of SO_x. *Appl. Biochem. Biotechnol.*, **28/29**, 877–886 (1991).
5. Yamaberi, K., Takagi, M., and Yoshida, T.: Nitrogen depletion for intracellular triacylglyceride accumulation to enhance liquefaction yield of marine microalgal cells into a fuel oil. *J. Mar. Biotechnol.*, **6**, 44–48 (1998).
6. Takagi, M., Watanabe, K., Yamaberi, K., and Yoshida, T.: Limited feeding of potassium nitrate for intracellular lipid and triacylglyceride accumulation of *Nanochloris* sp. UTEX LB1999. *Appl. Microbiol. Biotechnol.*, **54**, 112–117 (2000).
7. Xu, X.-Q. and Berdall, J.: Effect of salinity on fatty acid composition of a green microalgae from an Antarctic hypersaline lake. *Phytochemistry*, **45**, 655–658 (1997).
8. Peeler, T. C., Stephenson, M. B., Einsphar, K. J., and Thompson, G. A., Jr.: Lipid characterization of enriched plasma membrane fraction of *Dunaliella salina* grown in media of varying salinity. *Plant Physiol.*, **89**, 970–976 (1989).
9. Fujii, S., Uenaka, M., Nakayama, S., Yamamoto, R., and Mantani, S.: Effect of sodium chloride on the fatty acids composition in *Boekelovia hooglandii* (Ochromonadales, Chrysophyceae). *Phycol. Res.*, **49**, 73–77 (2001).
10. Elenkov, I., Stefanov, K., Konaklieva, D. S., and Popov, S.: Effect of salinity on lipid composition of *Cladophora vagabunda*. *Phytochemistry*, **42**, 39–44 (1996).
11. Borowitzka, L. J., Kessly, D. S., and Brown, A. D.: The salt relation of *Dunaliella*. Further observation on glycerol production and its regulation. *Arch. Microbiol.*, **13**, 131–138 (1977).
12. Galinski, E. A.: Osmoadaptation in bacteria. *Adv. Microbiol. Physiol.*, **37**, 273–328 (1995).
13. Sadka, A., Lers, A., Zamir, A., and Avron, M.: A critical examination of the role de novo protein synthesis in the osmotic adaptation of the halotolerant alga *Dunaliella*. *FEBS Lett.*, **244**, 93–98 (1989).