

FOOD RESEARCH

OPEN ACCESS





FOOD RESEARCH

Volume 5, Issue 3

June 2021

Mini Review

***Vibrio parahaemolyticus*: a review on the pathogenicity, antibiotic resistance, foodborne outbreaks, and detection methods**

Naziahsalam Kehinde, A., Tang, J.Y.H. and Nakaguchi, Y.

Available Online: 14 MARCH 2021

Naziahsalam *et al.* reviewed the pathogenicity, antibiotic resistance, foodborne outbreaks and detection methods of *Vibrio parahaemolyticus*.



PDF (406KB)

[Download File](#)

Issues related to animal blood into food products: a review paper

Siti Jamilah M.S., Nurulhidayah, A.F., Azura, A., Mat Jubri, S.M., Abdul

Rohman., Nur Azira, T., Arieff Salleh, R. and Rashidi, O.

Available Online: 14 MARCH 2021

Siti Jamilah *et al.* reviewed the issues related to animal blood produced into food products.



PDF (402KB)

[Download File](#)

Phytic acid in legumes: a review of nutritional importance and hydrothermal processing effect on underutilised species

Ojo, M.A.

Available Online: 31 DECEMBER 2020

Ojo reviewed on the nutritional importance and hydrothermal processing effect of phytic acid in legumes.



PDF (356KB)

[Download File](#)

Properties and potential of agarwood hydrosol as a drink: a review

Mohamad Kahar, E.E., Talip, B.A., Mohd Fauzi, N.A., Kamarulzaman, S.N., Zakaria, F., Muhammad, N., Rajat Mamat, T.N.A. and Basri, H



PDF (352KB)
Download File

Available Online: 28 MARCH 2021

The properties and potential of agarwood hydrosol as a drink were reviewed by Mohamad Kahar *et al.*

Full Paper

Determination of antiradical activity and phenolic and flavonoid contents of extracts and fractions of jackfruit (*Artocarpus heterophyllus* Lamk) seeds



PDF (427KB)
Download File

Zubaydah, W.O.S., Sahumena, M.H., Fatimah, W.O.N., Sabarudin, Arba, M. and Yamin

Available Online: 18 APRIL 2021

Zubadyah *et al.* determined the antiradical activity and phenolic and flavonoid contents of extracts and fractions of jackfruit (*Artocarpus heterophyllus* Lamk) seeds.

Extraction and characterisation of cassava starch cultivated in different locations in Sabah, Malaysia



PDF (562KB)
Download File

Hasmadi, M., Harlina, L., Jau-Shya, Lee, Mansoor, A.H., Jahurul, M.H A. and Zainol, M.K.

Available Online: 2 MAY 2021

The extraction and characterisation of cassava starch cultivated in different locations in Sabah, Malaysia was studied by Hasmadi *et al.*

Effervescent formulation based on variation of nanocapsules matrix type and roselle (*Hibiscus sabdariffa*) nanocapsules percentage



PDF (334KB)
Download File

Purbowati, I.S.M., Maksum, A. and Anggriawan, R.

Available Online: 10 MAY 2021

The effervescent formulation based on the variation of nanocapsules matrix type and roselle (*Hibiscus sabdariffa*) nanocapsules percentage was studied by Purbowati *et al.*

Composition of amino acids and fatty acids on luwak coffee processing



PDF (345KB)
Download File

Fitri, Laga, A., Dwyana, Z. and Tawali, A.B.

Available Online: 10 MAY 2021

The composition of amino acids and fatty acids on luwak coffee processing was studied by Fitri *et al.*

Potential sources of anti-obesity compound – hydroxycitric acid

among some of the underutilized fruits in the Philippines

Bagabaldo, P.A., Hurtada, W.A., Laurena, A.C. and Atienza, L.M.

Available Online: 10 MAY 2021

Bagabaldo *et al.* studied on the potentail sources of anti-obesity compound-hydroxycitric acid among some of the underutilized fruits in the Phillippines.



PDF (682KB)

[Download File](#)

Antioxidant increase by response surface optimization and Bayesian neural network modelling of pumpkin (*Cucurbita moschata* Duch) freezing

Kristianto, Y., Wignyanto, W., Argo, B.D. and Santoso, I.

Available Online: 10 MAY 2021

Kristianto *et al.* modelled the antioxidant increase by response surface optimization and Bayesian neural network modelling of pumpkin (*Cucurbita moschata* Duch) freezing.



PDF (918KB)

[Download File](#)

Antimicrobial stability of *Cosmos caudatus* extract at varies pH and temperature, and compounds identification for application as food sanitizer

Yusoff, N.A.H., Rukayadi, Y., Abas, F., Khatib, A. and Hassan, M.

Available Online: 16 MAY 2021

The antimicrobial stability of *Cosmos caudatus* extract at various pH and temperature was studied by Yusoff *et al.* and the compounds were identified for the possible food sanitizer application.



PDF (474KB)

[Download File](#)

Yeast dynamics in the black table olives processing using fermented brine as starter

Ciafardini, G., Venditti, G. and Zullo, B.A.

Available Online: 16 MAY 2021

The yeast dynamics in the processing of black table olives using fermented brine as starter was studied by Ciafardini *et al.*



PDF (712KB)

[Download File](#)

Bioactive compounds and antioxidant activity of black and green tea available in Bangladesh

Rahman, M., Jahan, I.A., Ahmed, S., Ahmed, K.S., Roy, M., Zzaman, W. and Ahmad, I.

Available Online: 16 MAY 2021

Rahman *et al.* studied on the bioactive compounds and antioxidant activity of black and green tea available in Bangladesh.



PDF (379KB)

[Download File](#)

Factors associated with household expenditure on oil and fat products in Malaysia: application of quantile regression



PDF (366KB)
Download File

Cheah, Y.K., Abdul Adzis, A., Abu Bakar, J. and Applanaidu, S.D.

Available Online: 23 MAY 2021

Cheah *et al.* evaluated the factors associated with household expenditure on oil and fat products in Malaysia.

Tempeh extract reduces cellular ROS levels and upregulates the expression of antioxidant enzymes



PDF (484KB)
Download File

Surya, R., Romulo, A. and Suryani, Y.

Available Online: 23 MAY 2021

Surya *et al.* studied on the tempeh extract to reduce cellular ROS levels and upregulates the expression of antioxidant enzymes.

The physicochemical properties of white sorghum (*Sorghum bicolor* L.) flour in various particle sizes by soaking the seeds before and after dehulling



PDF (723KB)
Download File

Antarlina, S.S., Estiasih, T., Zubaidah, E. and Harijono

Available Online: 23 MAY 2021

The physicochemical properties of white sorghum (*Sorghum bicolor* L.) flour in various particle sizes by soaking the seeds before and after dehulling were studied by Antarlina *et al.*

Identification of bacteriological quality and antimicrobial resistance of microorganisms isolated from animal foods collected from the abattoir, butcher shops and local seafood market



PDF (506KB)
Download File

Vijayalakshmi, P., Rajani Chowdary, A., Vidyullatha, P. and Sharon Sushma, M.

Available Online: 23 MAY 2021

Vijayalakshmi *et al.* studied on the identification of bacteriological quality and antimicrobial resistance of microorganisms isolated from animal foods collected from the abattoir, butcher shops and local seafood markets.

Consumption of jelly dessert containing porang (*Amorphophallus oncophyllus*) glucomannan and inulin along with low-calorie diet contributes to glycemic control of obese adults: a randomized clinical trial



PDF (583KB)
Download File

Utami, N.N., Lestari, L.A., Nurliyani and Harmayani, E.

Available Online: 23 MAY 2021

Utami *et al.* performed a randomized clinical trial on obese adults on the effect of consumption of jelly dessert containing porang (*Amorphophallus oncophyllus*) glucomannan and inulin along with low-calorie diet.

Synergistic of antimicrobial activities of lactic acid bacteria in fermented *Tilapia nilotica* incorporated with selected spices



**PDF (572KB)
Download File**

Ismail, A., Lani, M.N., Zakeri, H.A., Hasim, N.N., Alias, R. and Mansor, A.

Available Online: 30 MAY 2021

The synergistic of antimicrobial activities of lactic acid bacteria in fermented *Tilapia nilotica* with selected spices was studied by Ismail *et al.*

Could food or food contact surfaces be the favourable hideouts for *Listeria monocytogenes* in Perak, Malaysia?



**PDF (404KB)
Download File**

Chen, S.N., Yap, M.L., Kuan, C.H., Son, R. and Saw, S.H.

Available Online: 30 MAY 2021

Chen *et al.* evaluated the possibility of food contact surfaces as the favourable hideouts for *Listeria monocytogenes* in Perak, Malaysia

Quantification of bioactive compounds in guava at different ripening stages



**PDF (506KB)
Download File**

Shukla, S., Kushwaha, R., Singh, M., Saroj, R., Puranik, V., Agarwal, R. and Kaur, D.

Available Online: 16 MAY 2021

The quantification of bioactive compounds in guava at different ripening stages was studied by Shukla *et al.*

Effect of three types of oils and their level of incorporation on sensory quality of sorghum cookies



**PDF (814KB)
Download File**

Goubgou, M., Songré-Ouattara, L.T., Bationo, F., Banhoro, O., Traoré Y. and Savadogo, A.

Available Online: 30 MAY 2021

Goubgou *et al.* studied the sensorial quality of developed sorghum cookies made three types of oil.

The influence of Gum Arabic on the physicochemical and antimicrobial activity of the microencapsulated Mahkota Dewa (*Phaleria macrocarpa*)



**PDF (1139KB)
Download File**

leaves

Mohd Zin, Z., Razman, N.H., Hasmadi, M., Abd Manap, M.N. and Zainol, M.K.

Available Online: 6 JUNE 2021

The influence of Gum Arabic on the physicochemical and antimicrobial activity of the microencapsulated Mahkota Dewa (*Phaleria macrocarpa*) leaves was studied by Mohd Zin *et al.*

Isolation and characterization of acid and pepsin soluble collagen extracted from sharpnose stingray (*Dasyatis zugei*) skin



PDF (645KB)

[Download File](#)

Ong, T.Y., Shaik, M.I. and Sarbon, N.M.

Available Online: 20 DECEMBER 2021

Ong *et al.* isolated and characterized acid and pepsin soluble collagen extracted from sharpnose stingray (*Dasyatis zugei*) skin.

Effect of fish protein hydrolysate on physicochemical properties and oxidative stability of shortfin scad (*Decapterus macrosoma*) emulsion sausage



PDF (815KB)

[Download File](#)

Hamzah, M., Shaik, M.I. and Sarbon, N.M.

Available Online: 6 JUNE 2021

Hamzah *et al.* studied on the effect of fish protein hydrolysate on the physicochemical properties and oxidative stability of shortfin scad (*Decapterus macrosoma*) emulsion sausage

Microbiological status of some street iftar items collected from chalk bazaar in Dhaka city, Bangladesh



PDF (414KB)

[Download File](#)

Nur, I.T., Talukder, M., Das, T.R., Asaduzzaman, M., Feroz, F. and Munshi, S.K.

Available Online: 6 JUNE 2021

Nur *et al.* evaluated the microbiological status of some street iftar items collected from chalk bazaar in Dhaka city, Bangladesh.

Phytochemical analysis, total phenolic, total flavonoid contents and ferric reducing power of extracts from roots and leaves of *Searsia burchellii*



PDF (376KB)

[Download File](#)

Pillai, M.K., Mpopo, M.S. and Mekbib, S.B.

Available Online: 6 JUNE 2021

The extracts from roots and leaves of *Searsia burchellii* of its phytochemical analysis, total phenolic, total flavonoid contents and ferric reducing power were studied by Pillai *et al.*

Quality characteristics of chicken sausages using a combination of jack bean (*Canavalia ensiformis* L.) and soy protein isolate as a binder



PDF (542KB)
Download File

Canti, M., Murdiati, A., Naruki, S. and Supriyanto

Available Online: 6 JUNE 2021

Canti *et al.* studied the quality characteristics of chicken sausages using a combination of jack bean (*Canavalia ensiformis* L.) and soy protein isolate as a binder.

Anti-salmonellosis agent for foodborne illness from *Mangifera odorata* (kuini) extracts



PDF (578KB)
Download File

Adnan, H., Ismail, N., Hassan, H. and Mat-Ali, M.S.

Available Online: 6 JUNE 2021

Adnan *et al.* studied on the possibility of *Mangifera odorata* (kuini) extracts as an anti-salmonellosis agent for foodborne illness.

Detection of deltamethrin in cabbages using visible shortwave near infrared spectroscopy



PDF (411KB)
Download File

Ishkandar, C.D.M., Naw, N.M., Janius, R., Mazlan, N. and Lin, T.T.

Available Online: 13 JUNE 2021

The detection of deltamethrin in cabbages using visible shortwave near infrared spectroscopy was performed by Ishkandar *et al.*

Chemical and bacteriological risks of shrimp and clams (Gandofly) from Suez Gulf



PDF (315KB)
Download File

Amin A.N., Ahemd, A.M. and Ahmed, O.M.

Available Online: 13 JUNE 2021

The chemical and bacteriological risks of shrimp and clams (Gandofly) from Suez Gulf was evaluated by Amin *et al.*

The effect of combination of sugarcane pressmud compost and potassium fertilizer on vegetative growth of corn in coastal sandy soil



PDF (343KB)
Download File

Budiyanto, G.

Available Online: 13 JUNE 2021

Budiyanto *et al.* studied on the effect of combination of sugarcane pressmud compost and potassium fertilizer on vegetative growth of corn in coastal study soil.

Effect of training program on food safety knowledge, behaviour and practices among female students of Qassim University, KSA



PDF (497KB)
Download File

Asmahan, A.A., Wafa, I.E. and Abd El-Razik, M.M.

Available Online: 13 JUNE 2021

Asmahan *et al.* studied on the effect of a food safety training program on the knowledge, behaviour and practices among the female student of Qassim University, KSA.

Development of biodegradable smart packaging from chitosan, polyvinyl alcohol (PVA) and butterfly pea flower's (*Clitoria ternatea* L.) anthocyanin extract



PDF (589KB)

[Download File](#)

Hidayati, N.A., Wijaya, M.W., Bintoro, V.P., Mulyani, S. and Pratama, Y.

Available Online: 13 JUNE 2021

Hidayati *et al.* developed a biodegradable smart packaging from chitosan, polyvinyl alcohol (PVA) and butterfly pea flower's (*Clitoria ternatea* L.) anthocyanin extract.

Physicochemical and microbial properties of orange-fleshed sweet potato flour produced with sun-drying and sulphiting agent



PDF (351KB)

[Download File](#)

Ogunka, N.P., Ezeama, C.F. and Ukpabi, U.J.

Available Online: 13 JUNE 2021

Ogunka *et al.* evaluated the physicochemical and microbial properties of orange-fleshed sweet potato flour produced by sun-drying and using sulphiting agent.

Morphological characterisation and glycaemic responses of cake developed from carrot and concentrated *Nypa fruticans* sap



PDF (972KB)

[Download File](#)

Tai, Y.Y., Tengku Ismail, T.A. and Wan Rosli, W.I.

Available Online: 13 JUNE 2021

The morphological characteristics and glycaemic response of carrot cake made with concentrated *Nypa fruticans* sap was studied by Tai *et al.*

Effect of different cooking methods on chemical composition, nutritional values, and sensory properties of Jack bean (*Canavalia ensiformis*) tempe



PDF (419KB)

[Download File](#)

Purwandari, F.A., Annisa, E.D.N., Rachmawati, A.T., Puspitasari, D., Wikandari, R., Setyaningsih, W., Ningrum, A. and Sardjono

Available Online: 20 JUNE 2021

The effect of different cooking methods on chemical composition, nutritional values, and sensory properties of Jack bean (*Canavalia ensiformis*) tempe was studied by Purwandari *et al.*

Influence of extraction methods on bioactive compounds from Ngoc

Linh ginseng callus

Lieu, M.D., Nguyen, T.T.L., Nguyen T.H., Dang, T.K.T. and Do, D.

Available Online: 20 JUNE 2021

Lieu *et al.* evaluated the influence of extraction methods on bioactive compounds Ngoc Linh ginseng callus.



PDF (1072KB)

[Download File](#)

Chemical characteristics and organoleptic of gluten free toddler biscuits from mung bean flour, modified cassava flour and flaxseed

Amalia L., Setiarto, R.H.B., Nuraulia G.W. and Mariyani N.

Available Online: 20 JUNE 2021

The chemical characteristics and organoleptic properties of gluten free toddler biscuits made from mung bean flour, modified cassava flour and flaxseed were studied by Amalia *et al.*



PDF (722KB)

[Download File](#)

Effect of extraction conditions (temperature, pH and time) by cellulase on chemical properties of dried oyster mushroom (*Pleurotus sajor-caju*) extract

Nguyen, G.T.N. and Nguyen, T.M.

Available Online: 20 JUNE 2021

Nguyen and Nguyen studied the effect of extraction conditions (temperature, pH and time) by cellulase on the chemical properties of dried oyster mushroom (*Pleurotus sajor-caju*) extract.



PDF (802KB)

[Download File](#)

Effect of different extraction solvents on antioxidant properties of water lily (*Nymphaeaceae antares*) flower petal and stamen

Mohd Zin, Z., Sanuri, A.Z.M., Bashah, N., Ibrahim, K. Yahya, H. and Zainol M.K.

Available Online: 27 JUNE 2021

The effect of different extraction solvents on the antioxidant properties of water lily (*Nymphaeaceae antares*) flower petal and stamen was studied by Mohd Zin *et al.*



PDF (531KB)

[Download File](#)

Residue level and health risk assessment of organophosphorus pesticides in eggplant and cauliflower collected from Dhaka city, Bangladesh

Habib, M., Kaium, A., Khan, M.S.I., Prodhan, M.D.H., Begum, N., Chowdhury, M.T.I. and Islam, M.A.

Available Online: 27 JUNE 2021

Habib *et al.* evaluated the residue level and health risk assessment of organophosphorus pesticides in eggplant and cauliflower collected from Dhaka city, Bangladesh



PDF (658KB)

[Download File](#)

Evaluation of hygiene and food handling attitudes and practices during social isolation due to COVID-19 in Peruvian households: a cross-sectional study



PDF (433KB)
Download File

Cahuana Díaz, E.B., Revoredo Torres, I.G. and Saintila, J.

Available Online: 27 JUNE 2021

Cahuana Diaz *et al.* evaluated the hygiene and food handling attitudes and practices during social isolation due to COVID-19 in Peruvian households

Quantitative *Salmonella enterica* serovar Enteritidis risk assessment from consumption of hard-boiled eggs, half-boiled eggs and raw eggs among Malaysians



PDF (392KB)
Download File

Wong, J.X., Kuan, C.H., Saw, S.H., Chen, S.N., Tan, C.W., Yeo, S.K., Kuan, C.S., Phuah, E.T., Thung, T.Y., Son, R. and New, C.Y.

Available Online: 27 JUNE 2021

Wong *et al.* performed a microbial risk assessment on consumption of hard-boiled, half-boiled and raw eggs contaminated with *Salmonella enterica* serovar Enteritidis among Malaysians.

Antioxidant activity, total phenolic, and aflatoxin contamination in tempeh made from assorted soybeans (*Glycine max* L. Merrill)



PDF (799KB)
Download File

Yudiono, K., Ayu, W.C. and Susilowati, S.

Available Online: 27 JUNE 2021

The antioxidant activity, total phenolic and aflatoxin contamination in tempeh made from assorted soybeans (*Glycine max* L. Merrill) were studied by Yudiono *et al.*

Comparison of conventional frying and microwave frying of beef patty: Effect on oil absorption, texture, physical and chemical properties



PDF (338KB)
Download File

Noor Hidayati, R., Nurul Najihah, I. and Norazatul Hanim, M.R.

Available Online: 27 JUNE 2021

Noor Hidayati *et al.* compared conventional frying and microwave frying of beef patty and studied on the effect on oil absorption, texture, physical and chemical properties.

Proximate composition of the edible part of purple passion fruit and santol and in vitro prebiotic activity of crude polysaccharide extracts



PDF (375KB)
Download File

Wongsariya, K. and Kanchanadumkerng, P.

Available Online: 27 JUNE 2021

Wongsariya and Kanchanadumkerng evaluated the proximate composition of the edible part of purple passion fruit and santol and in vitro prebiotic activity of crude polysaccharide extracts.

Honeybee honey and stingless bee honey quality characteristics and their anti-cancer potential in HeLa cells



PDF (365KB)
[Download File](#)

Imtiazah, S.Z., Zaharah, H., Murni, I., Tufail Ahmad, F. and Yusof, H.M.

Available Online: 27 JUNE 2021

Imtiazah *et al.* studied on the quality characteristics and anti-cancer potential in HeLa cells of honeybee honey and stingless bee honey.

Biochemical content, minerals, and antioxidant activity of fruit jiaosu obtained by natural fermentation



PDF (631KB)
[Download File](#)

Hnin, P.A., Kam, Y.C. and Sit, N.W.

Available Online: 27 JUNE 2021

The biochemical content, minerals, and antioxidant activity of fruit jiaosu obtained by natural fermentation were studied by Hnin *et al.*

Honey-borne infant botulism in tahnik practice: An explanation of the risk and its solutions



PDF (352KB)
[Download File](#)

Harun, M.A.W., Abdullah, A., Mohamad, A.M. and Baharuddin, A.S.

Available Online: 30 JUNE 2021

Harun *et al.* studied on the honey-borne infant botulism in tahnik practice.

Effect of boiling and roasting on physicochemical and antioxidant properties of dark red kidney bean (*Phaseolus vulgaris*)



PDF (424KB)
[Download File](#)

Roy, M., Imran, M.Z.H., Alam, M. and Rahman, M.

Available Online: 30 JUNE 2021

The effect of boiling and roasting on physicochemical and antioxidant properties of dark red kidney bean (*Phaseolus vulgaris*) was studied by Roy *et al.*

Mini Review

Natto and miso: an overview on their preparation, bioactive components and health-promoting effects



PDF (444KB)
[Download File](#)

Chan, E.W.C., Wong, S.K., Kezuka, M., Oshiro, N. and Chan, H.T.

Available Online: 30 JUNE 2021

Chan *et al.* reviewed the preparation, bioactive components and health-promoting effects of natto and miso.

The use of electronic tongue (e-tongue) as a simple and rapid method for honey authentication



PDF (338KB)
[Download File](#)

Lestari, L.A., Triyana, K., Hanifah, A.K. and Wildiana, R.A.

Available Online: 30 JUNE 2021

Lestari *et al.* used the electronic tongue (e-tongue) to authenticate honey.

**Morocco's agricultural system response to the dual shock of the
COVID-19 crisis and drought: learnings and recommendations for the
new normal**

Brouziyne, Y.

Available Online: 30 JUNE 2021

Brouziyne *et al.* reviewed on Morocco's agricultural system response to the dual shock of the
COVID-19 crisis and drought.



PDF (368KB)

Download File

VOL. 5 | ISSUE 2

VOL. 5 | ISSUE 4

[HOME](#)[VIEW ARTICLES](#)[VIEW ISSUES](#)[FOR AUTHORS](#)[FOR REFEREES](#)[ABOUT THE JOURNAL](#)

EDITORIAL BOARD

Chief Editor

Son Radu

Malaysia

Editor

Vivian New Chia Yeung

Malaysia

Associate Editors

Prem Narayan Paudel

Kathmandu University

Nepal

Associate Editors

Borarin Buntong

Royal University of Agriculture

Cambodia

Associate Editors

Yoshitsugu Nakaguchi

Ishikawa Prefectural University

Japan

Nasreldin Elhadi

University of Dammam

Saudi Arabia

Dwaipayan Sikdar

University of Chittagong

Bangladesh

Rungsinee Sothornvit

Kasetsart University,

Thailand

Nikos Tzortzakis

Cyprus University of Technology

Cyprus

Saisamorn Lumyong

Chiang Mai University

Thailand

Ekachai Chukeatirote

Mae Fah Luang University

Thailand

Pin Kar Yong

Forest Research Institute
Malaysia (FRIM)
Malaysia

Chim Chay

University of Agriculture
Cambodia

Le Van Viet Man

Ho Chi Minh City University of
Technology
Viet Nam

Oscar Herrera-Calderon

Universidad Nacional Mayor de
San Marcos
Peru

Hasmadi Mamat

University Malaysia Sabah
Malaysia

Regiane Ribeiro dos Santos

Federal Rural University of Rio de
Janeiro
Brazil

Koh Soo Peng

Malaysian Agricultural Research
and Development Institute
(MARDI)
Malaysia

Dinko Hristov Dinko

Trakia University
Bulgaria

Hanifah Nuryani Lioe

Bogor Agricultural University
Indonesia

Michelle Galindo de Oliveira

Federal University of
Pernambuco
Brazil

Alper Sener

Onzekiz Mart University
Turkey

Angelo Maria Giuffrè

Mediterranea University of
Reggio Calabria
Italy

Javier Garcia-Lomillo

University of Burgos
Spain

Laura Marcela Zárate Polanco

National University of Colombia-
Bogotá
Colombia

Swapan K. Banerjee

HPFB Health Canada,
Canada

Adel Abu Bakr Abd El-Hamid**Shatta**

Suez Canal University
Egypt

Iddya Karunsagar**Saroat Rawdkuen**

Mae Fah Luang University
Thailand

Giovanna Giacalone

Università di Torino
Italy

Lorina Acilo Galvez

Visayas State University
Philippines

Sergio Soto Simental

Universidad Autónoma del
Estado de Hidalgo
Mexico

Fernando Cebola Lidon

Universidade Nova de Lisboa
Portugal

Linda Lim Biaw Leng

Universiti Brunei Darussalam
Brunei Darussalam

J.M. Krishanti Jayarukshi Kurr**Premarathne**

Wayamba University of Sri Lanka
Sri Lanka

Katherine Ann T. Castillo-Isra

University of the Philippines L
Banos
Philippines

Steve Flint

Massey University

Giuseppe Sortino
University of Palermo
Italy

Nitte University
India

New Zealand

COPYRIGHT © 2016 - 2020 RYNNYE LYAN RESOURCES

Effervescent formulation based on variation of nanocapsules matrix type and roselle (*Hibiscus sabdariffa*) nanocapsules percentage

^{1,*}Purbowati, I.S.M., ¹Maksum, A. and ^{2,3}Anggriawan, R.

¹Food Processing Technology Laboratory, Department of Agricultural Technology, Faculty of Agriculture, Universitas Jenderal Soedirman, Dr. Suparno Street, Karangwangkal, Purwokerto 53123, Indonesia.

²Food Business Technology, School of Applied Science, Technology, Engineering and Mathematics (STEM), Prasetya Mulya University, Tangerang 15339, Indonesia

³Head of Research, Community Service and Industry Relation Section, STEM, Prasetya Mulya University, Tangerang 15339, Indonesia

Article history:

Received: 8 October 2020

Received in revised form: 17 November 2020

Accepted: 21 January 2021

Available Online: 10 May 2021

Keywords:

Roselle (*Hibiscus sabdariffa* L),
Effervescent,
Anthocyanin,
Nanocapsule

DOI:

[https://doi.org/10.26656/fr.2017.5\(3\).299](https://doi.org/10.26656/fr.2017.5(3).299)

Abstract

Roselle flowers are usually consumed in the form of fresh steeping. To increase the flexibility of using roselle, effervescent roselle-based products are made which of course more practical, efficient and attractive. Unfortunately, the bioactive compound in roselle extracts is not stable to environmental influences so nanoencapsulation technology needs to be done. The encapsulation matrix type gives a unique effect on each core. The purposes of this study were to determine the effects of variation in the matrix types and percentage of roselle nanocapsules on the physical and chemical effervescent properties. The research design used in this study was a Complete Random Design with 2 factors and 3 replications. The first factor comprised matrix types (A): maltodextrin (A1) and Arabic gum (A2). The second factor was roselle nanocapsule percentages (B): 10% (B1), 15% (B2) and 20% (B3). Observed variables in this study were total phenol, anthocyanin, vitamin C, antioxidant activity and dissolving time. The Arabic gum nano encapsulated matrix type gave a better value of Vitamin C, antioxidant activity and dissolving time than maltodextrin, namely 0.149 ± 0.049 µg/3 g, $34.02 \pm 4.52\%$, 2.93 ± 0.80 mins. The percentage of roselle nanocapsules by 20% gave the highest vitamin C value, namely 0.188 ± 0.032 µg /3 g. The best treatment combination was obtained from the treatment (A2B3) Arabic gum nanocapsules using the third percentage (20%) with the characteristics: total phenol content 4.17 µg /3 g, anthocyanin 0.070 µg /3 g, vitamin C 0.188 µg /3 g when it dissolves in 1.25 mins.

1. Introduction

Roselle has been known as a fiber-producing plant. However, along with the increasing popularity of the slogan back to nature among the public, Rosella's prestige was also raised. The flexibility in using roselle is currently still limited considering the bioactive compounds in plants that have high added value. This is because the roselle raw material used is still in the form of powder and micro-scale extract encapsulation (Selim *et al.*, 2004).

This bioactive content has resulted in roselle being widely used as traditional medicine in Southeast Asia (Tsai *et al.*, 2002), food coloring (Piyarat *et al.*, 2014), antioxidants (Purbowati *et al.*, 2019), and flavoring (Ruangsri *et al.*, 2008). Currently, there are many roselle

-based products such as tea (Olatunji *et al.*, 2014), cupcakes (Abdel-Moemin, 2016), syrup, jam (Arueya and Akomolafe, 2014) and jelly (Kharismawati *et al.*, 2015). In the health and beauty sector, rosella can be used as a raw material for making toothpaste (Mahmud, 2018) and soap. Based on the foregoing, it is necessary to strive for an alternative form of encapsulation at the nanoscale, so that the high added value of bioactive components can be maintained along with the benefits it generates.

Roselle extract which was still in liquid form causes problems in its use in industries, transportation and also relatively short shelf life. Anthocyanin, a bioactive compound in roselle strongly affected by pH, solvent, temperature, oxygen, enzymes, light (Chumsri *et al.*, 2008). To diminish some degradation effects of the

*Corresponding author.

Email: Ikesitoresmi@yahoo.co.id

compound is very common to use encapsulation technique namely microencapsulation. But, according to Mardiah *et al.* (2014), effervescent rosella with formulations using microencapsulation still has a weakness, namely brittle with relatively few bioactive compounds. For these reasons, it was necessary to apply more advanced encapsulation technology that was nanoencapsulation technology on roselle extract. So that its utilization was more optimal.

Nanoencapsulation is a technology for packaging substances in small sizes by utilizing techniques such as nanocomposites, nanoemulsification, and nanostructuring so that functional products can be produced that can release core parts in a controlled manner (Sekhon, 2010). The principle of nanoencapsulation is the mixing of the water phase, the core substance phase and the coating material phase until a stable emulsion is formed, then the process of attaching the coating material to the surface of the core material and the process of reducing the particle size (Dubey, 2009). According to Purbowati *et al.* (2016), nano encapsulated roselle extract had better stability to temperature, time and pH than liquid extract forms. In this research, nanoencapsulation used β -cyclodextrin, capable of trapping 94% of the bioactive compounds therein.

Nano encapsulant matrix selection was a crucial stage in the nanoencapsulation process. Different matrix will provide different protection to the core. Encapsulant must be food grade and GRAS (Generally Recognized as Safe). The compounds can be used as a retaining material. The encapsulant matrix was starch, Arabic gum, methylcellulose, gelatin, whey protein, sugar syrup, β cyclodextrin, disaccharides, pullulan and sodium caseinate (Wandrey *et al.* 2010; Naufalin and Rukmini, 2013; Purbowati *et al.*, 2016). The effect of encapsulant agents is unique, depending on the nature of the protected core material. For this reason, it is necessary to examine the compatibility between the core material and anchoring material. The use of Arabic gum as a coating material can protect, even volatile compounds, from oxidation and evaporation (Kanakdande *et al.*, 2007). Gum arabic has a high viscosity which reaches 38.0 cP, so the wall layer that is formed will better protect the core material because the skin layer (shell) is stronger, so it is able to protect the volatile core material when the drying process goes well (Sugindro *et al.*, 2008). Gum Arabic is also suitable for encapsulation during the drying process because it can protect against oxidation and evaporation and prevent moisture absorption. (Wandrey *et al.*, 2010). Carbohydrates such as starch, maltodextrin are good coatings because they have low

viscosity at high solids and have high solubility properties (Balasubramani *et al.*, 2014)

On the other hand, the development of knowledge and technology in the field of food encourages the development of products with the right formulation to process natural ingredients into a form of preparation that is easily accepted by the consumer. In addition to other quality parameters that must still be met. Thus, it is expected to increase people's interest in consuming.

The effervescent tablet formulation of roselle was developed to make easier for the consumer. This was done to make it more efficient to use. Effervescent has several advantages, namely that it was more easily and quickly absorbed by the body so that the therapeutic effect was more pronounced, optimal compatibility increases body fluid intake and more practical to use (Harrera-Arellano *et al.*, 2007). Roselle has a sour taste. So, it was necessary to determine the exact level of roselle percentage. The combination of roselle percentage and matrix type gives rise to good effervescent characteristics that are acceptable to consumers.

This study aims to determine the effect of variations in nanoencapsulation matrix type and roselle nanocapsule percentage on the physical and chemical effervescent tablet's characteristics.

2. Materials and methods

2.1 Materials

Roselle flower (Bought in Beringharjo Market Yogyakarta), Arabic gum, maltodextrin, artificial sweeteners of aspartame and sorbitol, sodium bicarbonate, citric acid, malic acid, polyethylene glycol, ascorbic acid standard, gallic acid standard, ethanol, maltodextrin, buffer sodium acetate, acetate buffer, reagent folin, distilled water, polyethylene glycol, ethanol (PA), DPPH solution and other chemicals for analysis.

2.2 Roselle extraction method

The roselle extraction method was based on Purbowati *et al.* (2018). Approximately 10 g of the dried roselle petals powder is mixed with 100 mL aquadest. The extraction was done in microwave power extraction 250 W and 5 mins of extraction for each unit combinations. The microwave used for extraction was Electrolux EMM 2007X.

The slurry was radiated in a microwave oven at regular intervals (one-minute radiation and two mins off) to keep the temperature not rising above the boiling

point. Roselle extract was filtered and concentrated with a vacuum evaporator at 70°C, 44 cmHg for 15 mins.

2.2 Roselle extract nanoencapsulation

The roselle extraction method was based on Purbowati (2016). Maltodextrin/Gum Arabic mixed with aquadest with the ratio of the nanoencapsulation matrix (g): water (mL) = 1:5 (w/v). This solution was stirred for 30 mins. Roselle concentrate was added to the solution with the ratio of the solution: extract = 20: 1 (v/v). These solutions were homogenized with OSK 7313 Homogenizer at 40°C for 30 mins until an emulsion is formed. Subsequently, the emulsion particles were reduced into nanoparticles using a 22,000-rpm dispersing machine (Ultra-Turrax) for 30 mins. This nano-sized emulsion was then dried using a spray drier with an inlet temperature of 120°C and a pressure of 4.25 bar.

2.3 Roselle effervescent formulation

The ingredients used were sweetener aspartame and sorbitol, citric acid, malic acid, sodium bicarbonate, polyethylene glycol, and roselle nanocapsule powder. The total base of the material made as much as 3 g with the treatment of the formulation composition is as follows in Table 1.

Table 1. The Ingredients of roselle nanocapsule effervescent

Ingredients	Formula A (%)	Formula B (%)	Formula C (%)
Aspartam	0.06	0.06	0.06
Natrium Bikarbonat	40	40	40
Citric acid	30	30	30
Mallic acid	8.6	8.6	8.6
PEG	0.03	0.03	0.03
Roselle nanocapsule	10	15	20
Sorbitol	11.31	6.31	1.31
Sum	100	100	100

The process of making effervescent drinks was done by mixing all the ingredients well and cast them into a tablet with a tablet casting device with 3 g of the total weight for each. The effervescent tablets were packed using aluminum foil for further analysis.

2.4 Anthocyanin

The total anthocyanin content was determined using following procedure (Fuleki and Francis, 1968): approximately 1 mL of sample solution was added with 1 mL of acidic ethanol (95% ethanol + N HCL (85:15 v/v)). The sample was stored overnight at 40°C. The absorbance was measured using a 535 nm wavelength.

2.5 Total phenol

The total phenol content was determined using the Folin-Ciocalteu method (Chew et al., 2009). About 0.4

mL of sample solution was added with 1.5 mL of Folin-Ciocalteu reagent (10% v/v). After being incubated for 5 mins, it was mixed with 1.5 mL of 7.5% (w/v) of Na₂CO₃ solution. After 90 mins of incubation at room temperature and darkroom, its absorbance was measured using 765 nm. Gallic acid was used as a standard. The result was presented as mg Gallic Acid Equivalent (GAE)/g materials.

2.6 Antioxidant capacity

Capacity antioxidant test using the DPPH method (Blois, 2005). 100 mL of sample was added with 3 mL of methanol. This solution was adding with 1 mL of DPPH and keep in the dark for 15 mins at room temperature. The decrease of DPPH absorbance measure with spectrophotometry using λ 517 nm. As the standard was used gallic acid.

2.7 Experimental design and data analysis

The research design used in this study was a completely randomized design with two factors and three replications. The first factor comprises the type of nano encapsulant matrix (A): maltodextrin (A1) and Arabic gum (A2). The second factor was the percentage of nanocapsule powder (B): 10% (B1) which comprises 15% (B2) and 20% (B3).

Test data were analysed by ANOVA at the level of 5%. If there is a real influence, continue with Duncan's multiple area tests.

3. Results and discussion

From the results of the data processing, the nanoencapsulation matrix type had a significant effect on total phenol, anthocyanin, vitamin C, and dissolved time. While the percentage of roselle nanocapsules had a significant effect on phenol, anthocyanin, vitamin c variables, and had no significant effect on dissolved time. Meanwhile, the interaction between nanoencapsulant matrix type and the percentage of nanocapsules added had a significant effect on total phenol and anthocyanins. And had no significant effect on vitamin C and dissolving time.

3.1 Total phenols

The type of matrix and the percentage of nanocapsules were significantly affected the total phenol in effervescent tablets. Table 2 shows that the combination of Arabic gum with a concentration of 20% roselle nanocapsules gave the highest phenol content of 4.1740 μ g/3 g. Maltodextrin coated with the same percentage had a total phenol value of 2.9075 μ g/3 g. This meant that Arabic gum was better at coating roselle

phenol compounds. Gum Arabic has a high viscosity which reaches 38.0 cP, so the wall layer that is formed will better protect the core material because the skin layer (shell) was stronger, so it can protect the volatile core material when the drying process goes well (Sugindro *et al.*, 2008). This condition as a result of exposure to heat energy received by maltodextrin is greater than Arabic gum, due to the nature of phenolic compounds which are unstable at high temperatures. This was in line with research by Purbowati *et al.* (2016)

Table 2. Effect of combination treatment of encapsulant types and concentrations of rosella nanocapsules on the amount of total phenol and anthocyanin in effervescent

Treatment	Total phenol	Anthocyanin
A1B1	1.96±0.20 ^a	0.012±0.007 ^c
A1B2	2.38±0.11 ^{cd}	0.028±0.006 ^d
A1B3	2.90±0.68 ^{bc}	0.058±0.011 ^b
A2B1	2.06±0.34 ^d	0.007±0.007 ^c
A2B2	3.35±0.21 ^b	0.039±0.008 ^c
A2B3	4.17±0.37 ^a	0.070±0.002 ^a

A1B1 = maltodextrin with concentration 10%, A1B2 = maltodextrin with concentration 15%, A1B3 = maltodextrin with concentration 20%, A2B1 = arabic gum with concentration 10%, A2B2 = arabic gum with concentration 15%, A2B3 = arabic gum with concentration 20%

Values are expressed as mean±standard deviation of four independent data. Values with different superscript in the same column are significantly different in Duncan's test ($P < 0.05$).

3.2 Anthocyanin

The effect of variation in the percentage of Roselle nanocapsules, matrix type, and interaction both of them to anthocyanin was significantly different. Table 2 shows that the increasing nanocapsule percentage will increase the anthocyanin content in effervescent tablets. Nanoparticles formulation is needed to increase the bioavailability of active compounds penetration (Nurkhasanah *et al.*, 2015), the higher the percentage of rosella nanocapsules added, the number of bioactive compounds in it will be even higher. Arabic gum is better at coating rosella bioactive material. This is in line with Khasanah *et al.* (2015). Maltodextrin can form a good matrix network, but its viscosity was lower than Arabic gum and made the drying process last relative short. Gardjito *et al.* (2006) stated that the simpler molecular structure of maltodextrin makes it easy for water to evaporate during the drying process. Anthocyanin, a bioactive compound in roselle strongly affected by pH, solvent, temperature, oxygen, enzymes, light (Chumsri *et al.*, 2008). As a result, maltodextrin compared to gum arabic is not good enough to protect rosella bioactive compounds, namely anthocyanins. The

highest anthocyanin value was in combination with A2B3, which is Arabic gum with 20% of roselle.

3.3 Vitamin C

The different types of the matrix have a significantly different effect on vitamin C content in effervescent. This result can be seen in Table 3. This difference is due to the different chemical structures of maltodextrin and Arabic gum, besides the ability of the coating material to protect the bioactive compounds therein is also different. According to Khasanah *et al.* (2015) maltodextrin did not have a good emulsifying ability. This results in a thinner layer of protection and does not have the ability to properly cover the core material. The matrix must be able to form films, be easily biodegradable, have low viscosity and hygroscopic. The total phenol in rosella 70% is anthocyanin (Purbowati *et al.*, 2016). Anthocyanins are water-soluble dyes and compatible with water-based encapsulants matrices such as maltodextrin and Arabic gum. In addition, Arabic gum molecular weight was greater than maltodextrin (Gardjito *et al.*, 2006). Mean that the solubility of Arabic gum is higher than maltodextrin. This results in the Arabic gum solution being thicker than the maltodextrin solution. So that when drying use a spray dryer using an inlet temperature of 140°C caused the total levels of vitamin C in Arabic gum to be higher. Although spray drying is a fast process, a change in the spray drying temperature can affect the concentration of vitamin C.

Table 3. The effect of variation type nanocapsules matrix on vitamin C, antioxidant activity and time dissolve in effervescent.

Type nanocapsules matrix	Vitamin C (µg/3 g)	Antioxidant activity (%)	Dissolution Time (mins)
Maltodextrin	0.118±0.053 ^b	26.85±8.56 ^b	1.25±0.17 ^b
Arabic gum	0.149±0.049 ^a	34.02±4.52 ^a	2.93±0.80 ^a

Values are expressed as mean±standard deviation of four independent data. Values with different superscript in the same column are significantly different in Duncan's test ($P < 0.05$).

Table 4. The effect of variation nanocapsule concentration on vitamin C in effervescent.

Nanocapsule concentration (%)	Vitamin C (µg/3 g)
10	0.079±0.032 ^c
15	0.135±0.017 ^b
20	0.188±0.032 ^a

Values are expressed as mean±standard deviation of four independent data. Values with different superscript in the same column are significantly different in Duncan's test ($P < 0.05$).

Table 4 shows that the increasing nanocapsule percentage will increase the vitamin C content in

effervescent tablets. The use of the encapsulant matrix served as a wall to coat the active ingredients. The more nano-encapsulation matrix was added, the more vitamin C was trapped.

3.4 Antioxidant activity

The antioxidant properties of the nanocapsules were positively correlated with the content of bioactive compounds therein. Roselle flower contains bioactive compounds, namely phenol 19.45 ± 0.32 mg/g, vitamin C 20.47 ± 0.34 mg/g, anthocyanin 13.51 ± 0.03 mg/g. According to Purbowati *et al.* (2016), the antioxidant activity of rosella flowers was positively correlated with the total content of phenol and vitamin C.

Table 3 shows that the type of matrix had a significant effect on the antioxidant activity of the effervescent tablet. The antioxidant activity of effervescent tablets using Arabic gum is 34.02% higher than effervescent tablets using the maltodextrin matrix which is 26.85%. This is because the content of vitamin C and the total phenols in the nanocapsules made from Arabic gum was greater than nanocapsules that used maltodextrin. Christian and Jackson (2009) stated that the role of antioxidant activity in rosella was not only by anthocyanins as a phenolic group but also the content of vitamin C. The higher levels of anthocyanin in effervescent tablets, the higher antioxidant activity. In this study, effervescent tablets which have formulas with high vitamin C and anthocyanin content will also have a high antioxidant capacity, namely in formulas using Arabic gum with a value of 34.02%.

3.5 Dissolving time

The variation of the matrix significantly affects the effervescent dissolving time. Effervescent tablets are designed to break in contact with liquid such as water or juice, often causing the tablet to dissolve into a solution. The time it takes from the first contact with the water surface until it is completely dissolved is called dissolving time. However, if all the ingredients used are the same, in this research the difference in the effervescent dissolving time is determined by the difference in the matrix used. The nanocapsules used should have high solubility in the commonly used solvent, cold water. The results of the dissolving time test showed in Table 3 that maltodextrin had a lower solubility time of 1.25 mins faster than the solubility time of matrix Arabic gum which was 2.93 mins.

According to Mardiah *et al.* (2014), the addition of maltodextrin fillers in large effervescent tablets greatly affects the hardness value of effervescent tablets made. Fragility is a picture of the bonding strength of the particles forming the tablets. The more fragile the tablet,

the more resistant it is to erosion. As a result, when the maltodextrin's tablets are put into the water the dissolution time is shorter than using Arabic gum fillers.

The solubility of effervescent tablets is influenced by the presence of crushing agents in the form of acidic substances (malic acid and citric acid) and base sources (sodium bicarbonate). Both types of this material, when met with water, will produce CO₂ gas which gives a refreshing effect. In addition to these factors, the amount of coating agent and the method of mixing also affect the dissolution time of the tablet.

4. Conclusion

The Arabic gum gave better value on vitamin C, antioxidant activity and dissolving time than maltodextrin, namely 0.149 ± 0.049 µg/3g, $34.02 \pm 4.52\%$, 2.93 ± 0.80 mins. The percentage of rosella nanocapsules by 20% gave the highest vitamin C value, namely 0.188 ± 0.032 µg/3 g. The best treatment combination was obtained from the treatment using (A2B3) Arabic gum matrix and 20% of roselle nanocapsules with the characteristics: total phenol content 4.17 µg/3 g, anthocyanin 0.070 µg/3 g, vitamin C 0.188 µg/3 g, when it dissolves in 1.25 mins.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgments

The research work was funded by Unsoed's with scheme Competency Improvement Research.

References

- Arueya, G.L. and Akomolafe, B.O. (2014). Stability of microencapsulated anthocyanin of roselle (*Hibiscus sabdariffa*) in native starch and its potential application in jam production. *Journal of Environmental Science, Toxicology and Food Technology*, 8(7), 112-122. <https://doi.org/10.9790/2402-0871112122>
- Abdel Moemin, A.R. (2016). Effect of roselle calyces extract on the chemical and sensory properties of functional cupcakes. *Food Science and Human Wellness*, 5(4), 230-237. <https://doi.org/10.1016/j.fshw.2016.07.003>
- Balasubramani, P., Palaniswamy, P.T., Visvanathan, R.M., Thirupathi, V., Subbayan, A. and Maran, J.P. (2014). Microencapsulation of Garlic Oleoresin using Maltodextrin as Wall Material by Spray Drying Technology. *International Journal of Biological Macromolecules*, 72, 210-217. <https://doi.org/10.1016/j.ijbiomac.2014.05.011>

- doi.org/10.1016/j.ijbiomac.2014.08.011
- Blois, M.S. (2005). Antioxidant determination by the use of stable free radicals. *Nature*, 181, 1191-1200. <https://doi.org/10.1038/1811199a0>
- Chew, Y.L., Goh, J.K. and Lim, Y.Y. (2009). Assessment of *in vitro* antioxidant capacity and polyphenolic composition of selected medicinal herbs from the *Leguminosae* family in Peninsular Malaysia. *Food Chemistry*, 116(1), 13-18. <https://doi.org/10.1016/j.foodchem.2009.01.091>
- Christian, R. and Jackson, C. (2009). Change in total phenols and monomeric anthocyanin composition and antioxidant activity of three variety of sorrel (*Hibiscus sabdariffa*) during maturity. *Journal of Food Composition and Analysis*, 22(7-8), 663-667. <https://doi.org/10.1016/j.jfca.2009.05.007>
- Chumsri, P., Sirichele, A. and Itharat, A. (2008). Studies on the optimum condition for extraction and concentration of roselle (*Hibiscus sabdariffa* Linn) extract. *Songklanakarin Journal of Science and Technology*, 30(Suppl.), 133-139.
- Dubey, R.T. (2009). Microencapsulation Technology and Application. *Journal of Defence Science*, 59(1), 82-95.
- Fuleki, T. and Francis, F.J. (1968). Quantitative methods for anthocyanins, 1. Extraction and determination of total anthocyanin in Cranberries. *Journal of Food Science*, 33(1), 72-77. <https://doi.org/10.1111/j.1365-2621.1968.tb00887.x>
- Gardjito, M., Murdiati, A. and Aini, N. (2006). Mikroenkapsulasi β -karoten buah labu kuning dengan enkapsulan whey dan karbohidrat. *Jurnal Teknologi Pertanian*, 2(1), 13-18. [In Bahasa Indonesia].
- Harrera-Arellanao, A., Miranda-Sanchez, J., Avila-Castro, P., Herrera-Alvarez, S., Jimenez-Ferrer, J.E., Zamilpa, A., Roman-Ramos, R., Ponce-Monter, H. and Tortoriello, J. (2007). Clinical effect produces by standardized herbal medicinal product of *Hibiscus sabdariffa* on patients with hypertension. A randomized, double-blind, lisinopril-controlled clinical trial. *Planta Medica*, 73(1), 6-12. <https://doi.org/10.1055/s-2006-957065>
- Kanakdande, D., Bhosale, R. and Singhal, R.S. (2007). Stability Curcumin Oleoresin Microencapsulated In Different Combination Of Gum Arabic, Maltodextrin, and Modified Starch. *Jurnal Carbohydrate Polymer*, 67(4), 536-541. <https://doi.org/10.1016/j.carbpol.2006.06.023>
- Kharismawati, A.S., Nurhasanah, N. and Widyaningsih, T.D. (2015). Pengaruh minuman fungsional jelly drink kulit buah naga merah dan rosella terhadap stress oksidatif. *Jurnal Pangan dan Agroindustri*, 3 (2), 407-416. [In Bahasa Indonesia].
- Khasanah, L.U., Anandito, B.K., Rachmawaty, T., Utami, R. and Manuhara, G.J. (2015). Pengaruh rasio bahan penyalut maltodekstrin, gum arab dan susu terhadap karakteristik fisik dan kimia mikrokapsul oleoresin daun kayu manis. *Agritech*, 35(4), 414-421. <https://doi.org/10.22146/agritech.9325>
- Machmud, E. (2018). Effectiveness of Roselle effervescent Tablets as traditional medicine plants in preventing growth of *Candida albicans* colonies and *Streptococcus mutans*. *Journal of Contemporary Dental Practice*, 19(8), 925-928. <https://doi.org/10.5005/jp-journals-10024-2358>
- Mardiah., Setiawan, B. and Novidahlia, N. (2014). Minuman fungsional berkarbonasi yang kaya antioksidan berbasis kelopak bunga rosella (*Hibiscus sabdariffa*). *Farmasains*, 2(3), 64-77. [In Bahasa Indonesia].
- Naufalin, R. and Rukmini, H.S. (2013). Nanoencapsulated of natural antioxidant based on kecombrang (*Nicola speciosa* Horan) fruit using cyclodextrin-gelatin as filler ingredient, presented at 13th ASEAN Food Conference 2013. Singapore.
- Nurkhasanah, Yuwono T., Nurani, L.H., Rizki, M.I. and Krisana Kraisintu, K. (2015). The development of chitosan nanoparticles from *Hibiscus sabdariffa* l calyx extract from Indonesia and Thailand. *International Journal of Pharmaceutical Science and Research*, 6(5), 1855-1861.
- Olatunji, S.O. and Juwe, I.S. (2013). Awareness and utilization of Sobo drink (extract of *Hibiscus sabdariffa* roselle calyces) by farm families in Aniocha North Local Government Area of Delta State, Nigeria. *Journal of Agriculture and Social Research*, 13(2), 19-28.
- Piyarat, K., Walaisiri, M. and Pornpen, W. 2014. Comparison of stability of red colorant from natural sources, roselle and lac in micelles. *International Food Research Journal*, 21(1), 325-330.
- Purbowati, I.S.M., Syamsu, K., Warsiki, E. and Rukmini, H.S. (2016). Stabilitas Senyawa Fenolik dalam Ekstrak dan Nanokapsul Kelopak Bunga Rosella pada Berbagai Variasi pH, Suhu dan Waktu. *Agrointek*, 10(1), 31-40. [In Bahasa Indonesia]. <https://doi.org/10.21107/agrointek.v10i1.2023>
- Purbowati, I.S.M., Sujiman and Maksun, A. (2018). Antibacterial activity of roselle (*Hibiscus sabdariffa*) extract phenolics compound produced with varying drying methods and duration. *Jurnal Teknologi Industri*, 28(1), 19-27.

- Purbowati, I.S.M., Sujiman and Maksum, A. (2019). Antioxidant Activity and Isolation Bioactive Compound from Roselle (*Hibiscus sabdariffa*) from Various Methods and Drying Time. *Agrointek*, 13 (1), 1-9. <https://doi.org/10.21107/agrointek.v13i1.4046>
- Ruangsi, P., Chumsri, P., Anschalee, S. and Arunpora, I. (2008). Changes in quality and bioactive properties of concentrated Roselle (*Hibiscus sabdariffa* Linn) extract. *Journal of Agriculture and Food Research*, 1 (2), 62-67.
- Selim, K.A., Khalil, K.E., Abdel-bary, M.S. and Abdel Azeim, N.A. (2005). Extraction, encapsulation and utilization of red pigments from Roselle (*Hibiscus sabdariffa*) as natural food colorant. *Alexandria Journal of Food Science and Technology*, 5(2), 7-20.
- Sekhon, B.S. (2010). Food nanotechnology – An Overview. *Nanotechnology, Science and Applications*, 3, 1–15. <https://doi.org/10.2147/NSA.S8677>
- Sugindro, Mardiyati, E. and Djajadisastra, J. (2008). Pembuatan dan mikroenkapsulasi ekstrak etanol biji jinten hitam pahit (*Nigella sativa* Linn.) *Pharmaceutical Science and Research Journal*, 5(2), 57-66. <https://doi.org/10.7454/psr.v5i2.3419>
- Tsai, P.J., McIntosh, J., Pearce, P., Camden, B. and Jordan, T.B. (2002). Anthocyanin and antioxidant capacity in Roselle (*Hibiscus sabdariffa* L.) extract. *Food Research International*, 35(4), 351-356. [https://doi.org/10.1016/S0963-9969\(01\)00129-6](https://doi.org/10.1016/S0963-9969(01)00129-6)



Make an Impact

Publish Open Access

Make an impact.

tandfonline.com

OPEN

Food Research

COUNTRY

Malaysia



Universities and research
institutions in Malaysia

SUBJECT AREA AND CATEGORY

Agricultural and Biological Sciences
Food Science

PUBLISHER



Gellan Gum for Almond N

For Dairy Products

High Acyl Gellan Gum HAM Type for da
products or milk beverages.

opalgellangum.com

OPEN

PUBLICATION TYPE

Journals

ISSN

25502166

COVERAGE

2017-
2020



Publish Open Access - Regional Studies

Want to find the perfect home for your regional studies research? tandfonline.com

For Dairy Products

High Acyl Gellan Gum HAM Type for dairy products or milk beverages.

opalgellangum.com

OPEN

SCOPE

Food Research welcomes submissions that are relevant to food science and technology, food service management, nutrition, nutraceuticals, food innovation and agriculture food science. Manuscript types accepted are reviews, short reviews, original research articles and short communications that discusses on the current research and scientific views of the following scope: Food science and food chemistry; Food technology, processing, and food engineering; Food safety and quality – microbiological and chemical; Sensory, habits, consumer behaviour/preference; Nutrition and dietetics; Nutraceuticals and functional food/ingredients; Food service management; Food trends, innovation; Post-harvest and agribusiness; Food security; Food packaging.

 Join the conversation about this journal

Submit Your Manuscript With Us

Open

Hindawi

 Quartiles



FIND SIMILAR JOURNALS 



Publish Open Access - Regional Studies

Want to find the perfect home for your regional studies research? tandfonline.com

Submit Your Manuscript With Us

Hindawi

1
International Food Research Journal
MYS

89%
similarity

2
Food Science and Nutrition
GBR

82%
similarity


3
Carpathian Journal of Food Science and Technology
ROU

79%
similarity


4
International Science
EGY

7
s

SJR




Total Documents




Total Cites

Self-Cites




Citations per document



External Cites per Doc

Cites per Doc




% International Collaboration

Citable documents

Non-citable documents

Publish Open Access - Regional Studies

Want to find the perfect home for your regional studies research? tandfonline.com



400

200



← Show this widget in your own website


Just copy the code below and paste within your html code:

`<a href="https://www.scimagojr.com">`

SCImago Graphica

Explore, visually communicate and make sense of data with our **new free tool**.

Get it



Submit Your Manuscript With Us

Op

Hindawi

Metrics based on Scopus® data as of April 2021

T **Tamiur Yazew** 4 weeks ago

Dear,


Thank you for your updated and quality journal.

I have submitted two manuscripts in the Food research journal. The editor of this journal sent me the acceptance letter for the two manuscripts. However, they asked me to pay charge for the manuscripts. I am also working as a reviewer of this journal. I have edited two papers and sent it to the editor. I am currently reviewing a paper and it is ready to send back to the editor of this journal.



Publish Open Access - Regional Studies

Want to find the perfect home for your regional studies research? tandfonline.com



So, please would you help me by considering my issue into consideration!

reply



Melanie Ortiz 4 weeks ago

SCImago Team

Dear Tamiur, thank you very much for your comment. Unfortunately, we cannot help you with your request, we suggest you contact the journal's editorial staff so they could inform you more deeply. Best Regards, SCImago Team

K

Kamal 9 months ago

Hi, Editorial Team members,
I would like to know the topic related to the "hygienic practices along the supply chain of fisheries" is considered or not to review of your Journal. Early response is highly appreciated

reply



Melanie Ortiz 9 months ago

SCImago Team

Dear Kamal,
thank you for contacting us.
We are sorry to tell you that SCImago Journal & Country Rank is not a journal. SJR is a portal with scientometric indicators of journals indexed in Elsevier/Scopus.
Unfortunately, we cannot help you with your request, we suggest you visit the journal's homepage or contact the journal's editorial staff, so they could inform you more deeply.
Best Regards, SCImago Team

Y

yani purbanang 1 year ago

Hi,
This journal is written with Scopus index from 2017-2019, What is the index status in 2021 ??
Thank you for your explanation

reply



Melanie Ortiz 1 year ago

SCImago Team

Dear Yani,
Thank you very much for your comment.
All the metadata have been provided by Scopus /Elsevier in their last update sent to SCImago, including the Coverage's period data. The SJR for 2019 was released on 11



Publish Open Access - Regional Studies

Want to find the perfect home for your regional studies research? tandfonline.com



S **Sigit Susanto** 1 year ago

Whether the supply chain on Runner products could be submitted to this FR Journal?thanks very much

reply



Melanie Ortiz 1 year ago

SCImago Team

Dear Sigit,

Thank you for contacting us. Could you please expand a little bit your comment?

Best Regards, SCImago Team

Leave a comment

Name

Email

(will not be published)



I'm not a robot

reCAPTCHA
[Privacy](#) - [Terms](#)

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.


Developed by:

Powered by:



Publish Open Access - Regional Studies

Want to find the perfect home for your regional studies research? tandfonline.com



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2020. Data Source: Scopus®

EST MODUS IN REBUS
Horatio (Satire 1, 1, 106)



Publish Open Access - Regional Studies

Want to find the perfect home for your regional studies research? tandfonline.com

