

# AGRIVITA

Journal of Agricultural Science

**VOLUME 40 (3) October 2018**



**Agrivita**

Published by Faculty of Agriculture Universitas Brawijaya

**ISSN: 0126-0537 (Printed-Indonesia)**  
**ISSN: 2302-6766 (Printed-English)**  
**E-ISSN: 2477-8516 (Electronic)**

## **AGRIVITA Journal of Agricultural Science**

AGRIVITA Journal of Agricultural Science publishes articles in plant science such as agronomy, horticulture, plant breeding, soil-plant sciences and pest or disease-plant sciences. Published by Faculty of Agriculture Universitas Brawijaya Indonesia in collaboration with Indonesian Agronomy Association (PERAGI). AGRIVITA publishes three times in a year (February, June and October)

AGRIVITA is accredited by Decree of The Directorate General of Higher Education The Ministry of Education and Culture, Republic of Indonesia No: 60/E/KPT/2016.

Indexed in SCImago Journal & Country Rank (SJR), Elsevier Scopus, Emerging Sources Citation Index (ESCI) – Web of Science, Directory of Open Access Journals (DOAJ), ASEAN Citation Index (ACI), EBSCO and Google Scholar.

### **EDITORIAL TEAM**

#### **Editor in Chief**

Prof. Kuswanto

Faculty of Agriculture Universitas Brawijaya, Indonesia (Scopus ID: 57192702058)

#### **Editorial Board**

Prof. Moch. Dawam Maghfoer

Faculty of Agriculture Universitas Brawijaya, Indonesia  
(Scopus ID: 55440224300)

Syahrul Kurniawan, Ph.D

Faculty of Agriculture Universitas Brawijaya, Indonesia  
(Scopus ID: 55876481800)

Dr. Budi Waluyo

Faculty of Agriculture Universitas Brawijaya, Indonesia  
(Scopus ID: 56605006300)

Hagus Tarno, Ph.D

Faculty of Agriculture Universitas Brawijaya, Indonesia  
(Scopus ID: 36163526900)

Dr. Akhmad Rizali

Faculty of Agriculture Universitas Brawijaya, Indonesia  
(Scopus ID: 6507320984)

#### **International Editorial Board**

Prof. Dr. Widiatmaka

(Scopus ID: 56962708400)

Jintana Unartngam, Ph.D

(Scopus ID: 8528493500)

Dr. Dono Wahyuno

(Scopus ID: 650a7969735)

Seca Gandaseca, Ph.D

(Scopus ID: 35071255800)

Prof. Kurniatun Hairiah

(Scopus ID: 6602368888)

Dr. Mohammad Reza Alizadeh

(Scopus ID: 16642292100)

Markus Anda, Ph.D

(Scopus ID: 23024287000)

Prof. Anoop Kumar Srivastava

(Scopus ID: 56688989000)

Dr. Mohammad Ali Shariati

(Scopus ID: 56096567300)

Dr. Mohammad Valipour

(Scopus ID: 55877297968)

Dr. B. Mohan Kumar

(Scopus ID: 55435104500)

Hiran Anjana Ariyawansa, Ph.D

(Scopus ID: 55752594000)

Bogor Agricultural University, Indonesia

Kasertsat University, Thailand

Indonesian Spice and Medicinal Crops Research  
Institute

Universiti Putra Malaysia

Faculty of Agriculture Universitas Brawijaya, Indonesia

Rice Research Institute of Iran (RRII), Islamic Republic  
of Iran

Indonesian Centre for Agricultural Land Resource  
Research and Development

National Research Centre For Citrus, India

Orel State Agrarian University, Russian Federation

Islamic Azad University, Islamic Republic of Iran

Nalanda University, India

National Taiwan University

Prof. Dr. Wani Hadi Utomo  
(Scopus ID: 6507144030)  
Prof. Dr. Eko Handayanto  
(Scopus ID: 6508116842)

Faculty of Agriculture Universitas Brawijaya, Indonesia

Faculty of Agriculture Universitas Brawijaya, Indonesia

**Assistant Editor**

Silvia Santi Wahyuni  
Faculty of Agriculture Universitas Brawijaya, Indonesia  
Aini Nur Laila  
Faculty of Agriculture Universitas Brawijaya, Indonesia  
Endah Apriliani  
Faculty of Agriculture Universitas Brawijaya, Indonesia

**EDITORIAL ADDRESS**

Faculty of Agriculture Universitas Brawijaya  
Jl. Veteran Malang 65145 East Java Indonesia  
Phone/Fax: +62-341-575743  
E-mail : [agrivita@ub.ac.id](mailto:agrivita@ub.ac.id), [agrivitafaperta@yahoo.com](mailto:agrivitafaperta@yahoo.com), [agrivitafaperta@gmail.com](mailto:agrivitafaperta@gmail.com)  
Website: <http://www.agrivita.ub.ac.id>

**Editorial Team would like to thank to peer reviewer who has been invited  
by Agrivita Journal of Agricultural Science Volume 40 number 3, 2018**

- Prof. Dr. Michael Henry Boehme (Scopus ID: 13808322000)  
(Horticulture - Faculty of Life Sciences Humboldt-University of Berlin, Germany)
- Prof. Dr. Meine van Noordwijk (Scopus ID: 7003897764)  
(Agroforestry - World Agroforestry Centre (ICRAF))
- Dr. Robert J. Holmer (Scopus ID: 15829659100)  
(Horticulture - Urban Food International (UFI), Germany)
- Gregory C. Luther, Ph.D (Scopus ID: 8096082400)  
(Entomology - World Vegetable Center, Taiwan)
- Puneet Singh Chauhan, Ph.D (Scopus ID: 12809757600)  
(Microbial Ecology - National Botanical Research Institute, India)
- Dr. Hidenori Tanaka (Scopus ID: 55514958400)  
(Genetics and Plant Breeding - Faculty of Agriculture University of Miyazaki, Japan)
- Munir Ahmad, Ph.D (Scopus ID: 55612573000)  
(Entomology - Faculty of Crop and Food Sciences Arid Agriculture University, Pakistan)
- Dr. Naga Raju Maddela (Scopus ID: 56650913800)  
(Microbiology - Department of Life Science Universidad Estatal Amazónica, Ecuador)
- Shashi Yellareddygar, Ph.D  
(Plant Pathology - Department of Plant Pathology North Dakota State University, United States of America)
- Dr. Morio Tsukada (Scopus ID: 9636907500)  
(Entomology - Graduate School of Bioresources Mie University, Japan)
- Prof. Dr. Sun-Kee Hong (Scopus ID: 34770558100)  
(Landscape Ecology - Institution for Marine & Island Cultures Mokpo National University, South Korea)
- Tetsuhiro Watanabe, Ph.D (Scopus ID: 35397767400)  
(Soil Science - Graduate School of Agriculture Kyoto University, Japan)
- Dr. Endang Arisoelaningsih (Scopus ID: 8236381800)  
(Plant Ecophysiology - Faculty of Mathematics and Natural Sciences Universitas Brawijaya, Indonesia)
- Prof. Yasushi Mitsuda (Scopus ID: 12798575600)  
(Forestry- University of Miyazaki Japan)
- Dr. Tri Joko (Scopus ID: 17345721200)  
(Molecular Biology - Faculty of Agriculture Universitas Gadjah Mada, Indonesia)
- Dr. Ahmad Faizal (Scopus ID: 36727358600)  
(Plant Biotechnology - School of Life Sciences and Technology, Institute of Technology Bandung, Indonesia)
- Luqman Qurata Aini, Ph.D (Scopus ID: 35071053600)  
(Plant Pathology - Faculty of Agriculture Universitas Brawijaya, Indonesia)
- Ahmad Mujib, Ph.D  
(Biotechnology and Plant Breeding - Hamdard University, India)
- Dr. Budi Waluyo (Scopus ID: 56605006300)  
(Genetics and Plant Breeding - Faculty of Agriculture Universitas Brawijaya, Indonesia)
- Nur Wakhid, M.Sc (Scopus ID: 57192961613)  
(Soil Science - Indonesian Swampland Agricultural Research Institute)
- Prof. Moch. Dawam Maghfoer (Scopus ID: 55440224300)  
(Horticulture - Faculty of Agriculture Universitas Brawijaya, Indonesia)
- Dr. Catur Hermanto (Scopus ID: 51863728900)  
(Plant Pathology - Indonesian Vegetables Research Institute)
- Kurniawan Budiarto, Ph.D (Scopus ID: 12243097600)  
(Genetics and Plant Breeding - Indonesian Ornamental Crops Research Institute)
- Hagus Tarno, Ph.D (Scopus ID: 36163526900)  
(Pest and Crop Disease - Faculty of Agriculture Universitas Brawijaya, Indonesia)

Dr. Edy Syahputra (Scopus ID: 55965649800)  
(Plant Toxicology and Physiology - Faculty of Agriculture Universitas Tanjungpura, Indonesia)

Prof. Dr. Kuswanto (Scopus ID: 57192702058)  
(Genetics and Plant Breeding - Faculty of Agriculture Universitas Brawijaya, Indonesia)

Kyaw Min Htut, Ph.D (Scopus ID: 57192314425)  
(Plant Protection - Central Agriculture Research and Training Center, Myanmar)

Dr. Rindang Dwiyani (Scopus ID: 36604233900)  
(Plant Biotechnology - Faculty of Agriculture Universitas Udayana, Indonesia)

Dr. Akhmad Rizali (Scopus ID: 6507320984)  
(Plant Pest and Disease - Faculty of Agriculture Universitas Brawijaya, Indonesia)

Antonius Kasno, M.Si (Scopus ID: 56032982700)  
(Soil Science - Indonesian Soil Research Institute)

Syahrul Kurniawan, Ph.D (Scopus ID: 55876481800)  
(Soil Science - Faculty of Agriculture Universitas Brawijaya, Indonesia)

Dr. Fitri Rachmawati (Scopus ID: 39861797700)  
(Plant Physiology - Indonesian Ornamental Crops Research Institute)

Neneng Laela Nurida, Dr (Scopus ID: 56685751000)  
(Environmental Science – Indonesian Soil Research Institute)

Cahyo Prayogo, Ph.D (Scopus ID: 55935695900)  
(Agricultural and Biological Science - School of Life Sciences The University of Warwick, United Kingdom)

Prof. Jianguo Wang (Scopus ID: 51666293100)  
(Plant Protection – Jiangxi Agricultural University, People's Republic of China)

## TABLE OF CONTENTS

Editorial Team.....	i
Table of Contents.....	v
Allometric Equation for Pinang ( <i>Areca catechu</i> ) Biomass and C Stocks <i>Cahyo Prayogo, Rika Ratna Sari, Degi Harja Asmara, Subekti Rahayu, and Kurniatun Hairiah</i> .....	381
Clonal Fidelity of Micro-propagated <i>Phalaenopsis</i> Plantlets Based on Assessment Using Eighteen <i>Ph-Pto</i> SNAP Marker Loci <i>Erick Raynalta, Juanita Elina, Sudarsono, and Dewi Sukma</i> .....	390
Antifungal Activity of Crude Extracts of <i>Ageratum conyzoides</i> , <i>Cyperus rotundus</i> , and <i>Amaranthus spinosus</i> Against Rust Disease <i>Eriyanto Yusnawan and Alfi Inayati</i> .....	403
Effects of Cytokinin and Auxin on <i>In Vitro</i> Organ Development and Plumbagin Content of <i>Drosera peltata</i> Thunb. <i>Thanakorn Wongsap, Phithak Inthima, Maliwan Nakkuntod, Duangporn Premjet, and Anupan Kongbangkerd</i> .....	415
Physico-Chemical, Viability Evaluations and Efficacy Assessment of <i>Bacillus subtilis</i> Against Soft Rot Disease in <i>Phalaenopsis</i> <i>Wakiah Nuryani, Hanudin, Evi Silvia Yusuf, and Kurniawan Budiarto</i> .....	425
Developments of Rice Cell Suspension Culture and A Novel Strategy for Screening New Resistant Lines to Rice Blight Disease Caused by <i>Xanthomonas oryzae</i> pv. <i>oryzae</i> <i>Restu Rizkyta Kusuma, Liliek Sulistyowati, Chiu-Chsiung Cheng, and Yi-Hsien Lin</i> .....	434
Shoot and Root Growth in Common Bean ( <i>Phaseolus vulgaris</i> L.) Exposed to Gradual Drought Stress <i>Laily Ilman Widuri, Benyamin Lakitan, Erizal Sodikin, Mery Hasmeda, Mei Meihana, Kartika Kartika, and Erna Siaga</i> .....	442
Utilization of Liquid Smoke to Suppress Blood Diseases on Bananas and Its Effects on the Plant Growth <i>Imas Aisyah, Meity Suradji Sinaga, Abdjad Asih Nawangsih, Giyanto, and Gustan Pari</i> .....	453
Response of Temperate, Subtropical and Tropical Soybean Genotypes to Type-B Overflow Tidal Swamp of Indonesia <i>Danner Sagala, Munif Ghulamahdi, Trikoesoemaningtyas, Iskandar Lubis, Tatsuhiko Shiraiwa, and Koki Homma</i> .....	461
The Strategy of Organic Pest Control in Ecuador: Capture Effectiveness of Fruit Fly ( <i>Anastrepha</i> ) Species in Orange Tree Regulated by Volatile Alimentary Attractants <i>Enny Ganchozo-Mendoza, Julio Muñoz-Rengifo, Ronald Villamar-Torres, Jhon Molina-Villamar, Luz Cecilia Garcia Cruzaty, José Andrés Intriago-Quintana, Jefferson Herrera-Torres, Malena Martinez-Chevez, Edison Segura-Chavez, and Seyed Mehdi Jazayeri</i> .....	472
Compatibility Test of Four <i>Trichoderma</i> spp. Isolates on Several Synthetic Pesticides <i>Loekas Soesanto, Endang Mugiastuti, Ruth Feti Rahayuniati, Abdul Manan, and Ratna Stia Dewi</i> .....	481
Study on Trap Barrier System Towards Rodent Population and Rice Production in Tidal-Area of South Sumatera Indonesia <i>Yulia Pujiastuti, Kristian B. Sitompul, Suparman, Hastin WS Weni, Siti Herlinda, and Buyung A. Hadi</i> .....	490
Storage Temperature of Botanical Insecticide Mixture Formulations and Its Activity Againsts <i>Crociodolomia pavonana</i> (F.) (Lepidoptera: Crambidae) <i>Eka Candra Lina, Nia Yulianti, Gustria Ernisa, Arneti, and Novri Nelly</i> .....	498
The Conservation of Osingnese Traditional Home Garden Agroforestry in Banyuwangi, East Java, Indonesia <i>Luchman Hakim, Nila Ratih Pamungkas, Karuniawan Puji Wicaksono, and Soemarno</i> .....	506
Harvesting of Residual Soil Phosphorus on Intensive Shallot Farming in Brebes, Indonesia <i>Muliana, Arief Hartono, Syaiful Anwar, Anas Dinurohman Susila, and Supiandi Sabiham</i> .....	515
The Use of Fly Ash in Peat Soil on the Growth and Yield of Rice <i>Nelvia Nelvia</i> .....	527
Radiation Use Efficiency on the Different Varieties and the Number of Seedlings of Rice ( <i>Oryza sativa</i> L.) <i>Agus Suryanto, Moch. Dawam Maghfoer, and Tietyk Kartinaty</i> .....	536
Preliminary Study of WOFOST Crop Simulation in Its Prospect for Soybean ( <i>Glycine max</i> L.) Optimum Harvest Time and Yield Gap Analysis in East Java <i>F. R. Abadi, I. K. Tastra and B. S. Koentjoro</i> .....	544
Wheat ( <i>Triticum aestivum</i> ) Adaptation to Lombok Island Indonesia <i>Akhmad Zubaidi, Mansur Ma'shum, Gurjeet Gill, and Glenn K. McDonald</i> .....	556
Corrigendum .....	567
Instruction to Authors .....	App. 1
Partners .....	App. 2

[HOME](#) | [ABOUT](#) | [LOGIN](#) | [REGISTER](#) | [SEARCH](#) | [CURRENT](#) | [ARCHIVES](#) | [ARTICLE IN PRESS](#) | [ANNOUNCEMENTS](#)
[Home > Vol 44, No 3 \(2022\)](#)

## AGRIVITA, Journal of Agricultural Science

**AGRIVITA** *Journal of Agricultural Science* is a peer-reviewed, scientific journal published by Faculty of Agriculture Universitas Brawijaya Indonesia in collaboration with **Indonesian Agronomy Association** (PERAGI). The aims of the journal are to publish and disseminate high quality, original research papers and article review in plant science i.e. agronomy, horticulture, plant breeding, soil sciences, plant protection and other pertinent field related to plant production. AGRIVITA is published three times per year.

The Journal has been indexed in SCOPUS, Scimago Journal Ranks (SJR), Emerging Source Citation Index (ESCI-Web of Science), EBSCO, ProQuest, Google Scholar and others international indexing.

**AGRIVITA** is accredited first grade (Sinta 1/S1) for five years (2018-2023) based on Decree No: 30/E/KPT/2018 by Ministry of Research, Technology and Higher Education (Ristek Dikti), The Republic of Indonesia.

We accept submission from all over the world. All submitted articles shall never been published elsewhere, original and not under consideration for other publication.

### Announcements

#### News: AGRIVITA October 2022 Edition

AGRIVITA October 2022 Edition

Posted: 2022-11-21

[More...](#)

#### News: AGRIVITA February 2022

**AGRIVITA** *Journal of agricultural science* Vol 44 no 1 February 2022 Edition is available online on 1st February 2022.

Posted: 2022-01-31

#### News: DOI for 2012-2015 Edition

For citation purpose of 2012-2015 edition, we inform you to use DOI that stated in the document (not in website page).

If you have question please feel free to email us at [agrivita@ub.ac.id](mailto:agrivita@ub.ac.id)

Posted: 2019-07-08

[More...](#)[More Announcements...](#)

### Vol 44, No 3 (2022)

### Table of Contents

#### Articles

 **Mycelial Growth and Basidiocarp Production of *Pleurotus* Species in Substrates Supplemented with Fermented Plant Juice**  
Leilidyn Y. Zurbano, Diana A. Antones, Carla Mae C. Almoradie

 PDF

391-404

### About Agrivita

- Aim and Scope
- Editorial Team
- Publication Ethics
- Visitor Statistics
- Reviewer Acknowledgment

### Issues

- Current Issue
- Back Issues
- Article in Press
- Accepted Papers

### Information For Author



- Author Guidelines
- Template (docx)
- Template (pdf)

### Indexed By



SJR Rank







 Effect of Pre-Harvest Foliar Calcium and Silicon Fertilization on Pineapple Quality and Fruit Collapse Incidence <i>Diego Mauricio Cano-Reinoso, Loekas Soesanto, Kharisun Kharisun, Condro Wibowo</i>	 PDF 405-418
 Resistance Level and Enzyme Activity of <i>Spodoptera litura</i> F. to Chlorpyrifos and Their Sensitivity to the Oil Formulation of <i>Azadirachta indica</i> Juss. and <i>Cymbopogon nardus</i> (L.) Rendl. <i>R. Arif Malik Ramadhan, Neneng Sri Widayani, Danar Dono, Yusup Hidayat, Safri Ishmayana</i>	 PDF 419-430
 Chemical Properties and Micromorphology of Biochars Resulted from Pyrolysis of Agricultural Waste at Different Temperature <i>Nur Indah Mansyur, Eko Hanudin, Benito Heru Purwanto, Sri Nuryani Hidayah Utami</i>	 PDF 431-446
 Nitrogen Sources Take Roles on Different Growth Balance of Red Beet ( <i>Beta vulgaris</i> ) <i>Mochammad Roviq, Ellis Nihayati, Sitawati Sitawati, Soemarno Soemarno</i>	 PDF 447-458
 Assessment of Mulch Material Effect on Surface Runoff, Soil Loss, and Water Quality in an Agricultural Region <i>Riyanto Haribowo, Runi Asmaranto, L. Tri Wijaya Nata Kusuma, Berlian Gari Amrina</i>	 PDF 459-469
 Phytochemical Analysis of Leaves and Cherries of Coffee and Sensory Evaluation of Tea Products of Robusta Coffee in Songkhla, Thailand <i>Yudithia Maxiselly, Pisamai Anusornwanit, Adirek Rugkong, Rawee Chiarawipa</i>	 PDF 470-478
 Contribution of Agricultural Landscape Composition on Shaping the Interaction Between Pests and Natural Enemies in Cacao Agroforestry <i>Akhmad Rizali, Toto Himawan, Novita Yuniasari, Nely Yulianti, Muhamad Ari Bachtiar, Emha Dwi Rifqi Rafid</i>	 PDF 479-489
 The Effect of Nutrition and Planting Media on the Productivity and Quality of Baby Kai-Lan ( <i>Brassica oleracea</i> var. <i>alboglabra</i> ) Cultivated Using Nutrient Film Technique System <i>Azmi Alvian Gabriel, Muhammad Hadziq Shafri</i>	 PDF 490-499
 Acoustic Playback Stimulus Experiment to Study Mating Behavioral Responses of <i>Bactrocera cucurbitae</i> Coquillett (Diptera: Tephritidae) <i>Ayu Purnamasari, Susilo Hadi, Suputa Suputa</i>	 PDF 500-512
 Maize Media Enhance the Conidia Production of Entomopathogenic Fungi <i>Lecanicillium lecanii</i> also Its Effective to Control the Weevil <i>Cylas formicarius</i> (Fabricius) (Coleoptera: Brentidae) <i>Lutfi Afifah, Aulia Corry Aena, Nurcahyo Widyodaru Saputro, Anik Kurniati, Rosalia Maryana, Ani Lestari, Slamet Abadi, Ultach Enri</i>	 PDF 513-525
 The Efficiency of Aboriginal Entomopathogenic Nematodes from Semi-Arid Zone Against Tenebrionidae Larvae with Comparison to Commercial Bio-Insecticides <i>Dauren Kaliaskar, Aigerim Shibaeva, Nariman Zhappar, Valentin Shaikhutdinov, Laura Asherbekova, Sayakhat Bekbulatov, Almagul Kalyaskarova</i>	 PDF 526-536
 Spatial Analysis of Soil Available Potassium and Plant Brix Content for Site Specific Nutrient Management in Sugarcane <i>Yagus Wijayanto, Moch. Reza Wahyu Abdilah, Ika Purnamasari, Tri Wahyu Saputra</i>	 PDF 537-548
 Organics Acids from Cocoa Pod Waste Inoculated by Basidiomycota Fungi to Enhance the Performance of Shallots <i>Iradhatullah Rahim, Harsani Harsani, Hakzah Hakzah, Selis Meriem, Elkheir Hassaballah Abdallah Ahamed</i>	 PDF 549-558
 Characterization and Potential of Plant Growth-Promoting Rhizobacteria (PGPR) Isolates Capacity Correlating with Their Hydrocarbon Biodegradation Capability <i>Pujawati Suryatmana, Mieke Rochimi Setiawati, Diyan Herdiantoro, Betty Natalie Fitriatin, Nadia Nuraniya Kamaluddin</i>	 PDF 559-574
 Incompatibility Selected Dwarf Rootstock and Scion of Citrus sp. regard to Abiotic Stress Tolerant <i>Norry Eka Palupi, Moch. Dawam Maghfoer, Nunun Barunawati, Didik Hariyono</i>	 PDF 575-585

Short Communication

 Characteristics of Virus Symptoms in Chili Plants ( <i>Capsicum frutescens</i> ) Based on RGB Image Analysis <i>Asmar Hasan, Widodo Widodo, Kikin Hamzah Mutaqin, Muhammad Taufik, Sri Hendrastuti Hidayat</i>	 PDF 586-594
---	--

Review

 Pests and Diseases Management of Konjac ( <i>Amorphophallus muelleri</i> Blume) <i>Siti Hardiyanti, Supriadi Supriadi, Sri Rahayuningsih, Titiek Yulianti</i>	 PDF 595-603
 Synergy of Innovation between Hybrid Corn Seed Production and Seed Companies: A Review <i>Bahtiar Bahtiar, Darmawan Salman, Muhammad Arsyad, Muhammad Azrai</i>	 PDF 604-615

Editorial

 Front Matter	 PDF
--	---



User

Username

Password

☐ Remember me

Login

Notifications

- [View](#)
- [Subscribe](#)

Visitors

ID	120694	US	12017
IN	11378	PH	10877
MY	5603	TH	4408
VN	3551	IR	3535
CN	2959	NG	2175
Newest:	AP You:	ID	
Today:		83	
Month:		4532	
Total:		211711	
Supercounters.com			

Keywords

agri bisnis Agroforestry Biological control Botanical insecticides Chrysanthemum Correlation Endophytic fungi Lettuce Morphology Nitrogen Organic fertilizer Oryza sativa PGPR Rice Selection Soil amendment Soybean diversity drought rice soil fertility soybean

Journal Content

Search

Search Scope

All

▼

Search

Browse

- [By Issue](#)
- [By Author](#)
- [By Title](#)
- [Other Journals](#)

Information

- [For Readers](#)
- [For Authors](#)
- [For Librarians](#)





**AGRIVITA**

Journal of Agricultural Science  
Universitas Brawijaya Online Journal - © 2016  
Powered by [Open Journal System 2.4.7.1](#)



This work is licensed under a  
[Creative Commons Attribution-NonCommercial 4.0 International License](#)

**Agrivita**

**Q3**

Agronomy and  
Crop Science

best quartile

**SJR 2021**

**0.27**



powered by scimagojr.com



# AGRIVITA

## Journal of Agricultural Science

### Compatibility Test of Four *Trichoderma* spp. Isolates on Several Synthetic Pesticides

Loekas Soesanto<sup>1\*)</sup>, Endang Mugiastuti<sup>1)</sup>, Ruth Feti Rahayuniati<sup>1)</sup>, Abdul Manan<sup>1)</sup>, and Ratna Stia Dewi<sup>2)</sup>

<sup>1)</sup> Faculty of Agriculture, Jenderal Soedirman University, Purwokerto, Central Java, Indonesia

<sup>2)</sup> Faculty of Biology, Jenderal Soedirman University, Purwokerto, Central Java, Indonesia

#### ARTICLE INFO

##### Keywords:

Compatibility  
Synthetic pesticides  
*Trichoderma* spp

##### Article History:

Received: October 28, 2016

Accepted: August 28, 2018

\*) Corresponding author:

E-mail: lukassus26@gmail.com

#### ABSTRACT

This research aimed to study the compatibility of some *Trichoderma* spp. isolates on some synthetic chemical pesticides carried out at the Laboratory of Plant Protection, Faculty of Agriculture, Jenderal Soedirman University from April up to July 2014. *Trichoderma* isolates were derived from rhizosphere exploration on ginger, banana, pineapple and shallot. The synthetic pesticides used were mancozeb and propineb (fungicides), oxytetracycline and streptomycin sulfate (agrimycin, bactericides), carbofuran (nematicide), and deltamethrin and prephenophos (insecticides: synthetic pyrethroids and chiral organophosphates, respectively). The compatibility test used food poisoning method in a completely randomized design with three replicates. Variables observed were discolouration, sporulation, colony diameter, conidia density, and fungal growth at pesticides treatment. The data were analyzed by F test at 5 % significant level and continued by Duncan Multiple Range Test (DMRT) when there was a significant difference. The result of the research showed that the most significant decreasing of *Trichoderma* spp. was found on mancozeb for shallot, ginger, and banana isolates, and propineb for pineapple isolate, respectively, 89.4, 97.7, 93.3, and 95.2 %. This result was in line with colour, sporulation, and inhibition level observation.

#### INTRODUCTION

*Trichoderma* spp. are soil-borne fungus to control biologically of plant pathogens especially resulated by soil-borne pathogens given much attention and play an important role in integrated diseases management (IDM). Genus *Trichoderma* is the most characteristic beneficial isolated soil fungi because of its competence to guard plants and maintain pathogen community under various soil circumstances. This antagonist has been either everywhere studied or marketable as biopesticides, biofertilizers, and soil amendments (Harman, 2006; Howell, 2003). *Trichoderma* spp. also provides abundant biologically active suspension, including enzymes to degrade pathogen cell wall and more secondary metabolites (Vinale et al., 2008).

*Trichoderma* spp. have been isolated and

screened from several rhizospheres, that was ginger (Soesanto et al., 2005), shallot (Santoso, Soesanto, & Haryanto, 2007), banana (Haryono, Prihatiningsih, Wardhana, & Soesanto, 2009), and pineapple (L. Soesanto collection) rhizospheres. These isolates have been tested to suppress some soil-borne plant pathogens. *Trichoderma* spp. was used to manage *Fusarium oxysporum* (Probowo, Prihatiningsih, & Soesanto, 2006; Santoso, Soesanto, & Haryanto, 2007; Soesanto et al., 2005; Wardhana, Soesanto, & Utami, 2009), *Phytophthora* spp. (Karthikeyan, Kumar, & Kumar, 2003), *Sclerotium rolfsii* (Elad, Barak, & Chet, 1984; Henis, Adams, Lewis, & Papavizas, 1983; Jegathambigai, Wilson Wijeratnam, & Wijesundera, 2010), *Rhizoctonia solani* (Erper et al., 2013), and *Botrytis cinerea* (Barakat, Abada, Abou-Zeid, & El-

ISSN: 0126-0537 Accredited by DIKTI Decree No: 60/E/KPT/2016

**Cite this as:** Soesanto, L., Mugiastuti, E., Rahayuniati, R. F., Manan, A., & Dewi, R. S. (2018). Compatibility test of four *Trichoderma* spp. isolates on several synthetic pesticides. *AGRIVITA Journal of Agricultural Science*, 40(3), 481-489. <http://doi.org/10.17503/agrivita.v40i3.1126>

Gammal, 2014; Zimand, Elad, & Chet, 1996). Study of inhibition mechanisms on *Trichoderma* spp. have been done, such as on competition ability (Bailey et al., 2008; Howell, 2003), antibiosis (Bailey et al., 2008; Benítez, Rincón, Limón, & Codón, 2004), mycoparasitism (Bailey et al., 2008; Reithner, Ibarra-Laclette, Mach, & Herrera-Estrella, 2011), enzyme activities (Benhamou & Chet, 1997; Haggag & Abo-Sedera, 2005), plant growth promoting fungi (Benítez, Rincón, Limón, & Codón, 2004), and organic waste decomposer (Haggag & Abo-Sedera, 2005; Soyong & Quyet, 2013).

Synthetic pesticides are highly used in almost all agricultural systems especially IDM system to prevent and to control plant diseases. Pesticides are easily found in the market and practically applied in the field so that the establishment of pesticides in Indonesia market is highly increasing (Direktorat Pupuk dan Pestisida, 2016). The pesticides are tremendously and commonly utilized to improve plant productivity by combating plant pests and diseases (Aktar, Sengupta, & Chowdhury, 2009; Cooper & Dobson, 2007). The other benefits of pesticides are for protection of crop losses, control of vector disease, and quality of food. Besides their advantages, pesticides also have numerous disadvantages such as adverse impact to human health, to food, on environment, on water and soil contamination, and to non-target organisms (Aktar, Sengupta, & Chowdhury, 2009; Chowdhury, Pradhan, Saha, & Sanyal, 2008; Sengupta, Aktar, Alam, & Chowdhury, 2010).

Integrated disease management combines biological, physical, cultural, and chemical component strategies applied to all plants and crops and many plant pathogens including fungi, bacteria, viruses, and nematodes. The pathogens can cause extremely harmful to crops from small to significant losses primarily in the tropical developing country. Application of single IDM component to the plant pathogens resulted in an insignificant result. The application of IDM has been proved to be more effective in the way of holistic manner than the use of single IDM component strategy (El Khoury & Makkouk, 2010). As a component of IDM, the antagonist *Trichoderma* spp. should be compatible with other IDM components in the agricultural system. The compatibility of *Trichoderma* spp. with the other components is needed to improve the performance of the plant disease management, so that the disease could be prevented and controlled,

and the agricultural product could be saved and increased (Paret, Dufault, Momol, Marois, & Olson, 2015). The role of synthetic chemical pesticides in the agricultural system is still needed in the disease management, and the compatibility of both components need to be studied. The compatibility of *Trichoderma* spp. with the other components is required.

*Trichoderma* spp. has wide variability in controlling plant pathogens included its compatibility with chemical pesticides. This variability is caused by many influencing factors (Gómez, Chet, & Herrera-Estrella, 1997), especially the genetics factor. Exploration of *Trichoderma* spp. indicated some potential isolates that could control plant pathogens (Haryono, Prihatiningsih, Wardhana, & Soesanto, 2009; Santoso, Soesanto, & Haryanto, 2007; Soesanto et al., 2005). However, the compatibility of *Trichoderma* spp. isolates with general pesticides are not known yet. Therefore, this research was carried out to determine the compatibility of selected *Trichoderma* spp. isolates with commonly applied synthetic chemical pesticides including fungicides, insecticides, bactericide, and nematocides.

## MATERIALS AND METHODS

The research was conducted at the Laboratory of Plant Protection, Agricultural Faculty, Jenderal Soedirman University, Purwokerto, from April to July 2014. Four *Trichoderma* spp. isolates and six pesticides used arranged split-plot design with three replicates.

### *Trichoderma* spp. Isolates Preparation

*Trichoderma* spp. were prepared by plating on PDA (Tuite, 1969), then incubated for 3 days at room temperature ( $26 \pm 1^\circ\text{C}$ ) daylight. The *Trichoderma* isolates were derived from rhizosphere exploration on ginger (Soesanto et al., 2005), banana (Haryono, Prihatiningsih, Wardhana, & Soesanto, 2009), pineapple (L. Soesanto collection), and shallot (Santoso, Soesanto, & Haryanto, 2007). The isolates were identified as *T. harzianum* Rifai for ginger, banana, and shallot isolates, while the pineapple one had not been identified.

### Synthetic Pesticides Preparation

The synthetic pesticides were mancozeb and propineb (fungicides), oxytetracycline and streptomycin sulfate (agrimycin, bactericides), carbofuran (nematicide), and deltamethrin and prephenophos (insecticides: synthetic pyrethroids

and chiral organophosphates, respectively). The choice of the pesticides was based on the frequency of their usage in Indonesia. All pesticides were prepared by mixing with sterile water according to label recommended.

### Compatibility Test *In Vitro*

The compatibility test was conducted with food poisoning method (Khan & Shahzad, 2007) in completely randomized design with three replicates, by adding 1 droplet of the pesticides in Petri dish mixed with PDA just before plating. The Petri dish was homogenized and after solid, each *Trichoderma* spp. isolates were inoculated 5 mm discs of a seven-day-old culture of *Trichoderma* spp. isolates. After that, they were punched by sterilized cork borer and put a single disc in each Petri dish containing PDA and pesticide with the help of inoculating needle under an aseptic condition or PDA without pesticides as a control. Then, they were incubated at room temperature ( $26 \pm 1^\circ\text{C}$ ) for 5 days or at least the growth of control reached the edge of Petri dish. Each treatment was repeated four times.

### Observation and Measurements

Variables observed were discolouration, sporulation, colony diameter, conidia density, fungal growth at pesticides treatment compared to control with the formula.

$$I = 100(C-T)/C$$

Where: I = inhibition percentage; C = *Trichoderma* spp. colony diameter at control; T = *Trichoderma* spp. colony diameter at pesticides treatment (Gowdar, Babu, Nargund, & Krishnappa, 2006). The mycelial dry weight of the antagonist was measured based on Lilly & Barnett (1951) and Sutton & Starzyk (1972).

### Data Analysis

The data were analyzed by F test at 5 % significant level and continued by Duncan Multiple Range Test (DMRT) when there was a significant difference.

## RESULTS AND DISCUSSION

### Growth Inhibition

Some pesticides significantly affected colony discolouration (Table 1). The colony of all *Trichoderma* spp. isolates had discolouration from green to white after growing on PDA supplemented by mancozeb, propineb, and preferenophos, while on

PDA supplemented by carbofuran, oxytetracycline and streptomycine sulphate (agrimycin), and deltamethrin, the colony was still green in colour as the control. Moreover, the spotted isolate of *Trichoderma* spp. which change colour to white, i.e., in the treatment of mancozeb, propineb, and preferenophos, is considered low, whereas green colony isolates produce more conidium. (Table 1 and Table 2).

**Table 1.** Discolouration of treated all *Trichoderma* spp. colony after three days incubation

Treatments	Colour
Control	Green
Carbofuran	Green
Mancozeb	White
Propineb	White
Oxytetracycline and streptomycine sulphate	Green
Deltamethrin	Green
Preferenophos	White

At Fig. 1, some pesticides resulted in higher growth inhibition of all *Trichoderma* spp., especially mancozeb, propineb, and preferenophos, although among isolates of *Trichoderma* spp. showed indifferent growth inhibition at similar pesticide. This result is in line with colony discolouration (Table 1). The highest growth inhibition was found at preferenophos in all *Trichoderma* spp. isolates by a range of 74.10 - 80.38 %, while mancozeb and propineb inhibited the growth as a range of 29.81 - 43.18 % and 27.27 - 33.21 %, respectively. The lowest growth inhibition was showed by bactericide (oxytetracycline and streptomycin sulphate or agrimycin), i.e., in a range of 0.38 - 4.55 %. Based on kinds of *Trichoderma* spp. isolate, the ginger isolate was the most sensitive isolate on preferenophos compared to other isolates, while different growth inhibition between preferenophos and mancozeb and propineb as high as 47.4 and 65.7 %, respectively.

Antagonists against plant pathogenic fungi have been applied to control plant pathogens, and 90 % of the applications have been done with various fungus *Trichoderma* strains (Benítez, Rincón, Limón, & Codón, 2004). The influence of fungicides toward the advantageous action of microbes is significant to understand due to its value of danger analogous by synthetic fungicide applied in agriculture. Maximized crop productiveness and economic profit will be

achieved with the utilise of the products to suppress the pathogens well, but sustaining beneficial organisms (Yang, Hamel, Vujanovic, & Gan, 2011). Research of the antagonist *Trichoderma* spp. on pesticides have escorted to a better comprehension of compatibility instrument to gain better-integrated application in an agricultural system, especially in IDM. Modes of action from fungicide have never been successfully demonstrated, and the auxiliary effects of the synthetic fungicides are not sufficiently

perceived. Hence, the synthetic fungicide used may have contradictive impacts which are obscure to forecast (Lo, 2010). If the formulations made by the recommended dose of insecticides with the bio-control agent *Trichoderma* spp. and used for the management of various plant pests, they will show a promising effect than the chemicals alone. It costs effectively and also environment-friendly (Mahfut, Joko, & Daryono, 2016; Singh, Srivastava, Shrivastava, & Singh, 2012).

**Table 2.** The effect of synthetic pesticides on colony diameter, conidia density, and mycelial dry weight of *Trichoderma* spp. isolates

Treatments	Colony diameter (mm)	Conidia density ( $\times 10^6$ conidia ml <sup>-1</sup> )	Mycelial dry weight (mg)
F cal.T	159.57 **	64.31 **	17.76 **
F tab 5 %	2.78	2.78	2.78
Trichoderma shallot	41.3 b	93.7 a	0.0203 b
Trichoderma ginger	65.3 a	86.4 b	0.0193 b
Trichoderma banana	64.3 a	60.1 d	0.0194 b
Trichoderma pineapple	64.9 a	80.9 c	0.0261 a
F cal.F	305.87 **	450.06 **	73.67 **
F tab 5 %	2.27	2.27	2.27
Control	80.0 a	123.0 b	0.0305 a
Carbofuran	72.6 b	102.0 d	0.0268 b
Mancozeb	48.8 d	14.0 f	0.0118 c
Propineb	56.2 c	14.8 f	0.0138 c
Agrimicyn	78.4 a	134.3 a	0.0258 b
Deltamethrin	58.4 c	112.5 c	0.0298 a
Prefenofos	18.3 e	61.5 e	0.0108 c
F cal. TXF	5.28 **	36.12 **	19.68 **
F tab 5%	1.80	1.80	1.80
Shallot, Control	55.3 ef	160.0 b	0.031 c-e
Shallot, Carbofuran	51.0 fg	139.0 c	0.026 d-g
Shallot, Mancozeb	32.0 h	17.0 l	0.015 j-n
Shallot, Propineb	37.7 h	32.0 k	0.009 n-p
Shallot, Agrimicyn	54.3 ef	147.0 bc	0.029 c-f
Shallot, Deltamethrin	44.7 g	110.0 e	0.025 e-h
Shallot, Prefenofos	14.3 j	51.0 j	0.007 op
Ginger, Control	88.3 a	133.0 cd	0.020 g-j
Ginger, Carbofuran	83.3 abc	106.0 ef	0.016 j-m
Ginger, Mancozeb	51.0 fg	3.0 l	0.009 n-p
Ginger, Propineb	64.0 d	12.0 l	0.009 n-p
Ginger, Agrimicyn	88.7 a	160.0 b	0.019 h-k
Ginger, Deltamethrin	64.3 d	174.0 a	0.055 a
Ginger, Prefenofos	17.3 ij	17.0 l	0.007 op
Banana, Control	88.0 a	75.0 hi	0.032 cd
Banana, Carbofuran	80.0 bc	67.0 i	0.032 cd
Banana, Mancozeb	50.0 fg	5.0 l	0.012 l-p
Banana, Propineb	64.0 d	9.0 l	0.013 k-o
Banana, Agrimicyn	84.0 ab	92.0 fg	0.023 f-i
Banana, Deltamethrin	61.3 de	84.0 gh	0.018 i-l
Banana, Prefenofos	22.7 i	89.0 gh	0.006 p
Pineapple, Control	88.3 a	124.0 d	0.039 b
Pineapple, Carbofuran	76.0 c	96.0 efg	0.033 c
Pineapple, Mancozeb	62.0 de	31.0 k	0.011 m-p
Pineapple, Propineb	59.0 de	6.0 l	0.024 f-i
Pineapple, Agrimicyn	86.7 ab	138.0 cd	0.032 cd
Pineapple, Deltamethrin	63.3 d	82.0 gh	0.021 g-j
Pineapple, Prefenofos	18.7 ij	89.0 gh	0.023 f-i

Remarks: Numbers accompanied by the same letter at the same column are not significantly different at DMRT  $\alpha$  %5






Fig. 1. Growth inhibition of *Trichoderma* spp. isolates due to pesticides

The growth and development of *Trichoderma* spp. isolates were influenced by given pesticides (Table 2). All isolates were affected by the pesticides in colony diameter, conidia density, and mycelial dry weight. Colony diameter of *Trichoderma* spp. shallot isolate was smaller than other isolates and in line with its mycelial dry weight. However, the highest conidia density was found in the isolate ( $93.7 \times 10^6$  conidia  $\text{ml}^{-1}$ ) and differ significantly with other isolates ( $p < 0.01$ ). *Trichoderma* spp. pineapple isolate gave the highest colony diameter, and this was in line with its mycelial dry weight.

Based on the results, almost all chemical synthetic pesticides indicated a negative effect on colony diameter of all *Trichoderma* spp. isolates, which resulted in decreasing mycelial dry weight and conidia density. Prefenphos could decrease the highest colony diameter but not in conidia density and mycelial dry weight as high as 77.1, 50.0, and 64.6 %, respectively. The highest decreasing conidia density of *Trichoderma* spp. isolates were found in mancozeb and propineb, i.e. 88.6 and 88.0 %, respectively; while bactericide (oxytetracycline and streptomycine sulfate or agrimycin) seems to stimulate increasing conidia density only 8.4 %.

Deltamethrin is a synthetic pyrethroid insecticide widely used on fruits and vegetables to control the household, industrial, and veterinary pests. Deltamethrin is highly toxic to some aquatic organisms, such as Nile tilapia (*Oreochromis niloticus* L.) fingerlings (Yildirim et al., 2006), freshwater mussel (Köprücü & Seker, 2008), and

*Daphnia magna* (Day & Maguire, 1990). Oda & El-Maddawy (2012) point out that deltamethrin has danger effects for male reproductive systems and the protective effect of vitamin E and selenium combination on deltamethrin induced the deleterious effects of deltamethrin on male potency. However, deltamethrin has less effect on conidia density of *Trichoderma* spp. isolates.

Mancozeb and propineb inhibit conidia density of all *Trichoderma* spp. isolates wherein the highest inhibition happened. This result is contrast to Bagwan's (2010) research which stated *Trichoderma* applied for seed coating or irrigation would be harmonious with several synthetic or non-synthetic chemical, such as synthetic fungicides (thiram, copper oxychloride, mancozeb); pesticides; herbicides; and botanical pesticides (neem oil, neem leaves extract, wild sorghum leaves extract, neem cake, castor cake and mustard cake extracts) for the IDM of soil-borne pathogens on peanut. Another activity of mancozeb fungicide affecting metabolism of target cells, can also influence bacteria included in both cycle of soil C and N (Černohlávková, Jarkovský, & Hofman, 2009; Cycoń, Piotrowska-Seget, & Kozdrój, 2010). These fungicides performance are widely applied in agronomic system because of the wide spectrum activity of plant pathogen control, but the synthetic fungicides may have side effects on other microbes due to their various sites impacts of biochemistry (Dwimartina, Arwiyanto, & Joko, 2017; Yang, Hamel, Vujanovic, & Gan, 2011).



Table 2 showed significant interaction between *Trichoderma* spp. isolates and pesticides for all parameters. Based on the isolates, ginger, banana, and pineapple isolates showed the highest colony diameter grown on media supplemented by carbofuran and agrimycin, and in line with mycelial dry weight and conidia density. The lowest inhibition of colony diameter and mycelial dry weight of these isolates was indicated on agrimycin treatment compared to other pesticides, but on ginger isolate, this bactericide could stimulate colony diameter growth though only 0.5 %. The lowest colony diameter of all *Trichoderma* spp. Isolates were found at prefenophos, i.e., 74.1, 80.4, 74.2, and 78.8 %, respectively.

The combination between fungicide tolerant biological control agents and reduced levels of fungicide IDM strategies would increase the level of plant pathogens suppression identical to that obtained by the synthetic fungicides with full dosage (Monte, 2001). The effect of integration of *Trichoderma* with fungicides was reported by Sharma, Singh, & Sugha (1992) in controlling *Sclerotinia sclerotiorum*. When biocontrol agents *Trichoderma harzianum* and *Aspergillus niger* were incorporated with two synthetic fungicides, Foltaf 80W (Captafol 80 %) and Blue Copper-50, for protection from pigeon-pea wilt. The combination of biological agents and the fungicides was more suppressed the disease effectively than only the fungicides were used (Bhatnagar, 1995). Biopesticides can degrade more quickly than synthetic chemical pesticides and can supplement the synthetic pesticides used in integrated pest management (IPM) programs, which offer potentially higher crop yield and can reduce the use of conventional pesticides (Thakore, 2006). *Trichoderma* spp. are a soil microorganism. The soil microbial populations can influence plant growth and production in agricultural systems so that understanding the effects of synthetic fungicides on soil microbes can be important manner (Joko et al., 2012).

There were some isolates resulted in the highest conidia density and mycelial dry weight but gave the lowest colony diameter, such as shallot isolate on deltamethrin, ginger isolates on deltamethrin, and banana isolates on deltamethrin and prefenophos. In all isolates, all pesticides decreased conidia density compared to control except ginger, banana, and pineapple isolates showed increasing the density because of

agrimycin, i.e., 16.9, 18.5, and 10.2 %, respectively, and deltamethrin and prefenophos could increase banana isolate conidia density, i.e., 10.7 and 15.7 %, respectively. The greatest decrease was found on mancozeb for shallot, ginger, and banana isolates, and propineb for pineapple isolate respectively 89.4, 97.7, 93.3, and 95.2 %. This result was in line with colour, sporulation, and inhibition level observation (Table 1 and Fig. 1).

## CONCLUSION

Mancozeb for shallot, ginger, and banana isolates, and propineb for pineapple isolate decreased the growth of *Trichoderma* spp., respectively, 89.4, 97.7, 93.3, and 95.2 %. This result was in line with colour, sporulation, and inhibition level observation.

## ACKNOWLEDGEMENT

This research was a part of the main study funded by the Competency Fund Batch I from Directorate of Higher Education, Indonesian Minister of Education and Culture; for that, I deeply thank for the financial support.

## REFERENCES

- Aktar, M. W., Sengupta, D., & Chowdhury, A. (2009). Impact of pesticides use in agriculture: Their benefits and hazards. *Interdisciplinary Toxicology*, 2(1), 1–12. <http://doi.org/10.2478/v10102-009-0001-7>
- Bagwan, N. B. (2010). Evaluation of *Trichoderma* compatibility with fungicides, pesticides, organic cakes and botanicals for integrated management of soil borne diseases of soybean [*Glycine max* (L.) Merrill]. *International Journal of Plant Protection*, 3(2), 206–209. Retrieved from <https://pdfs.semanticscholar.org/5d06/16b3822e866d7d80fd520b3176781785ae95.pdf>
- Bailey, B. A., Bae, H., Strem, M. D., Crozier, J., Thomas, S. E., Samuels, G. J., ... Holmes, K. A. (2008). Antibiosis, mycoparasitism, and colonization success for endophytic *Trichoderma* isolates with biological control potential in *Theobroma cacao*. *Biological Control*, 46(1), 24–35. <http://doi.org/10.1016/j.biocontrol.2008.01.003>
- Barakat, F. M., Abada, K. A., Abou-Zeid, N. M., & El-Gammal, Y. H. E. (2014). Effect of volatile and non-volatile compounds of *Trichoderma* spp. on *Botrytis fabae* the causative agent of faba bean chocolate spot. *American Journal of Life Sciences*, 2(6–2), 11. <http://doi.org/10.11648/j>

Loekas Soesanto *et al.*: *Compatibility Test of Trichoderma spp. on Synthetic Pesticides* .....

ajls.s.2014020602.12

- Benhamou, N., & Chet, I. (1997). Cellular and molecular mechanisms involved in the interaction between *Trichoderma harzianum* and *Pythium ultimum*. *Applied and Environmental Microbiology*, 63(5), 2095–2099. Retrieved from <https://aem.asm.org/content/63/5/2095/article-info>
- Benítez, T., Rincón, A. M., Limón, M. C., & Codón, A. C. (2004). Biocontrol mechanisms of *Trichoderma* strains. *International Microbiology*, 7, 249–260. Retrieved from <http://www.im.microbios.org/0704/0704249.pdf>
- Bhatnagar, H. (1995). Integrated use of biocontrol agents with fungicides to control wilt incidence in pigeon-pea. *World Journal of Microbiology & Biotechnology*, 11(5), 564–566. <http://doi.org/10.1007/BF00286374>
- Černohlávková, J., Jarkovský, J., & Hofman, J. (2009). Effects of fungicides mancozeb and dinocap on carbon and nitrogen mineralization in soils. *Ecotoxicology and Environmental Safety*, 72(1), 80–85. <http://doi.org/10.1016/j.ecoenv.2008.07.001>
- Chowdhury, A., Pradhan, S., Saha, M., & Sanyal, N. (2008). Impact of pesticides on soil microbiological parameters and possible bioremediation strategies. *Indian Journal of Microbiology*, 48(1), 114–127. <http://doi.org/10.1007/s12088-008-0011-8>
- Cooper, J., & Dobson, H. (2007). The benefits of pesticides to mankind and the environment. *Crop Protection*, 26(9), 1337–1348. <http://doi.org/10.1016/j.cropro.2007.03.022>
- Cycoń, M., Piotrowska-Seget, Z., & Kozdrój, J. (2010). Responses of indigenous microorganisms to a fungicidal mixture of mancozeb and dimethomorph added to sandy soils. *International Biodeterioration and Biodegradation*, 64(4), 316–323. <http://doi.org/10.1016/j.ibiod.2010.03.006>
- Day, K. E., & Maguire, R. J. (1990). Acute toxicity of isomers of the pyrethroid insecticide deltamethrin and its major degradation products to *Daphnia magna*. *Environmental Toxicology and Chemistry*, 9, 1297–1300. <http://doi.org/10.1002/etc.5620091009>
- Direktorat Pupuk dan Pestisida. (2016). *Pestisida pertanian dan kehutanan tahun 2016* [Pesticides of agriculture and forestry 2016]. Jakarta: Ditjen PSP. Retrieved from [http://psp.pertanian.go.id/assets/file/2016/Pestisida\\_Pertanian\\_dan\\_Kehutanan\\_Tahun\\_2016.pdf](http://psp.pertanian.go.id/assets/file/2016/Pestisida_Pertanian_dan_Kehutanan_Tahun_2016.pdf)
- Dwimartina, F., Arwiyanto, T., & Joko, T. (2017). Potential of endophytic and rhizobacteria as an effective biocontrol for *Ralstonia syzygii* subsp. *syzygii*. *Asian Journal of Plant Pathology*, 11, 191–198. <http://doi.org/10.3923/ajppaj.2017.191.198>
- El Khoury, W., & Makkouk, K. (2010). Integrated plant disease management in developing countries. *Journal of Plant Pathology*, 92(4, supplement), 35–42. Retrieved from <https://www.researchgate.net/publication/268380542/download>
- Elad, Y., Barak, R., & Chet, I. (1984). Parasitism of sclerotia of *Sclerotium rolfsii* by *Trichoderma harzianum*. *Soil Biology and Biochemistry*, 16(4), 381–386. [http://doi.org/10.1016/0038-0717\(84\)90037-3](http://doi.org/10.1016/0038-0717(84)90037-3)
- Erper, I., Turkkan, M., Atanasova, L., Druzhinina, I. S., Karaca, G. H., & Cebi-Kilicoglu, M. (2013). Integrated assessment of the mycoparasitic and phytostimulating properties of *Trichoderma* strains against *Rhizoctonia solani*. *Bulgarian Journal of Agricultural Science*, 19(4), 742–748. Retrieved from [https://www.researchgate.net/profile/Guersel\\_Karaca/publication/257233286\\_Integrated\\_assessment\\_of\\_the\\_mycoparasitic\\_and\\_phytostimulating\\_properties\\_of\\_Trichoderma\\_strains\\_against\\_Rhizoctonia\\_solani/links/00463537479445d716000000/Integrated-assessment-of-the-mycoparasitic-and-phytostimulating-properties-of-Trichoderma-strains-against-Rhizoctonia-solani.pdf](https://www.researchgate.net/profile/Guersel_Karaca/publication/257233286_Integrated_assessment_of_the_mycoparasitic_and_phytostimulating_properties_of_Trichoderma_strains_against_Rhizoctonia_solani/links/00463537479445d716000000/Integrated-assessment-of-the-mycoparasitic-and-phytostimulating-properties-of-Trichoderma-strains-against-Rhizoctonia-solani.pdf)
- Gómez, I., Chet, I., & Herrera-Estrella, A. (1997). Genetic diversity and vegetative compatibility among *Trichoderma harzianum* isolates. *Molecular and General Genetics*, 256(2), 127–135. <http://doi.org/10.1007/s004380050554>
- Gowdar, S. B., Babu, H. N. R., Nargund, V. B., & Krishnappa, M. (2006). Compatibility of fungicides with *Trichoderma harzianum*. *Agricultural Science Digest*, 26(4), 279–281. Retrieved from [https://www.researchgate.net/file.PostFileLoader.html?id=50fa787ee5438f3a2600007e&assetKey=AS%3A271751534055425%4014418020\\_22498](https://www.researchgate.net/file.PostFileLoader.html?id=50fa787ee5438f3a2600007e&assetKey=AS%3A271751534055425%4014418020_22498)
- Haggag, W. M., & Abo-Sedera, S. A. (2005). Characteristics of three *Trichoderma* species in peanut haulms compost involved in biocontrol

Loekas Soesanto *et al.*: Compatibility Test of *Trichoderma* spp. on Synthetic Pesticides .....

- of cumin wilt disease. *International Journal of Agriculture and Biology*, 7(2), 222–229. Retrieved from [http://www.fspublishers.org/published\\_papers/94350\\_.pdf](http://www.fspublishers.org/published_papers/94350_.pdf)
- Harman, G. E. (2006). Overview of mechanisms and uses of *Trichoderma* spp. *Phytopathology*, 96(2), 190–194. <http://doi.org/10.1094/PHYTO-96-0190>
- Haryono, J., Prihatiningsih, N., Wardhana, R. A., & Soesanto, L. (2009). Pengaruh pemasteuran tanah tunggal atau digabung agensia hayati terhadap pengelolaan penyakit busuk hati di pembibitan pisang [The effect of soil pasteurization alone or in combination with biological agents on heart rot disease management of banana seed]. *Akta Agrosia*, 12(1), 21–28. Retrieved from [http://repository.unib.ac.id/214/1/joko\\_akta\\_Vol12No.1.pdf](http://repository.unib.ac.id/214/1/joko_akta_Vol12No.1.pdf)
- Henis, Y., Adams, P. B., Lewis, J. A., & Papavizas, G. C. (1983). Penetration of sclerotia of *Sclerotium rolfsii* by *Trichoderma* spp. *Phytopathology*, 73, 1043–1046. Retrieved from [https://www.apsnet.org/publications/phytopathology/backissues/Documents/1983Articles/Phyto73n07\\_1043.PDF](https://www.apsnet.org/publications/phytopathology/backissues/Documents/1983Articles/Phyto73n07_1043.PDF)
- Howell, C. R. (2003). Mechanisms employed by *Trichoderma* species in the biological control of plant diseases: The history and evolution of current concepts. *Plant Disease*, 87(1), 4–10. <http://doi.org/10.1094/PDIS.2003.87.1.4>
- Jegathambigai, V., Wilson Wijeratnam, R. S., & Wijesundera, R. L. C. (2010). Effect of *Trichoderma* sp. on *Sclerotium rolfsii*, the causative agent of collar rot on *Zamioculcas zamiifolia* and an on farm method to mass produce *Trichoderma* species. *Plant Pathology Journal*, 9, 47–55. <http://doi.org/10.3923/ppj.2010.47.55>
- Joko, T., Koentjoro, M. P., Somowiyarjo, S., Rohman, M. S., Liana, A., & Ogawa, N. (2012). Response of rhizobacterial communities in watermelon to infection with cucumber green mottle mosaic virus as revealed by cultivation-dependent RISA. *Archives of Phytopathology and Plant Protection*, 45(15), 1810–1818. <http://doi.org/10.1080/03235408.2012.707526>
- Karthikeyan, A., Kumar, S., & Kumar, S. (2003). *Trichoderma viride*: A mycoparasite for the control of *Phytophthora cinnamomi*. *Indian Forester*, 129(5), 631–634. Retrieved from [https://www.researchgate.net/publication/307546182\\_Trichoderma\\_viride\\_A\\_mycoparasite\\_for\\_the\\_control\\_of\\_Phytophthora\\_cinnamomi](https://www.researchgate.net/publication/307546182_Trichoderma_viride_A_mycoparasite_for_the_control_of_Phytophthora_cinnamomi)
- Khan, M. O., & Shahzad, S. (2007). Screening of *Trichoderma* species for tolerance to fungicides. *Pakistan Journal of Botany*, 39(3), 945–951. Retrieved from [http://www.pakbs.org/pjbot/PDFs/39\(3\)/PJB39\(3\)945.pdf](http://www.pakbs.org/pjbot/PDFs/39(3)/PJB39(3)945.pdf)
- Köprücü, K., & Seker, E. (2008). Acute toxicity of deltamethrin for freshwater mussel, *Unio elongatulus eucirrus bourguignat*. *Bulletin of Environmental Contamination and Toxicology*, 80(1), 1–4. <http://doi.org/10.1007/s00128-007-9254-z>
- Lilly, V. G., & Barnett, H. L. (1951). *Physiology of the fungi*. New York, Toronto, London: McGraw-Hill Book Company, Inc. Retrieved from <https://www.cabdirect.org/cabdirect/abstract/19521603587>
- Lo, C.-C. (2010). Effect of pesticides on soil microbial community. *Journal of Environmental Science and Health Part. B*, 45(5), 348–359. <http://doi.org/10.1080/10934520903467873>
- Mahfut, Joko, T., & Daryono, B. S. (2016). Molecular characterization of odontoglossum ringspot virus (ORSV) in Java and Bali, Indonesia. *Asian Journal of Plant Pathology*, 10(1–2), 9–14. <http://doi.org/10.3923/ajppaj.2016.9.14>
- Monte, E. (2001). Understanding *Trichoderma*: between biotechnology and microbial ecology. *International Microbiology*, 4(1), 1–4. <http://doi.org/10.1007/s101230100001>
- Oda, S. S., & El-Maddawy, Z. K. (2012). Protective effect of vitamin E and selenium combination on deltamethrin-induced reproductive toxicity in male rats. *Experimental and Toxicologic Pathology*, 64(7–8), 813–819. <http://doi.org/10.1016/j.etp.2011.03.001>
- Paret, M., Dufault, N., Momol, T., Marois, J., & Olson, S. (2015). *Integrated disease management for vegetable crops in Florida*. Florida: University of Florida. Retrieved from <http://edis.ifas.ufl.edu/pdf/PP/PP11100.pdf>
- Probowo, A., Prihatiningsih, N., & Soesanto, L. (2006). Potensi *Trichoderma harzianum* dalam mengendalikan sembilan isolat *Fusarium oxysporum* Schlecht. f. sp. *zingiberi* Trujillo pada kencur [Potency of *Trichoderma harzianum* in

Loekas Soesanto et al.: Compatibility Test of *Trichoderma* spp. on Synthetic Pesticides .....

- controlling nine isolates of *Fusarium oxysporum* Schlecht. f. sp. *zingiberi* Trujilo on galanga]. *Jurnal Ilmu-Ilmu Pertanian Indonesia*, 8(2), 76–84. Retrieved from <http://repository.unib.ac.id/47/1/76JIPi-2006.pdf>
- Reithner, B., Ibarra-Laclette, E., Mach, R. L., & Herrera-Estrella, A. (2011). Identification of mycoparasitism-related genes in *Trichoderma atroviride*. *Applied and Environmental Microbiology*, 77(13), 4361–4370. <http://doi.org/10.1128/AEM.00129-11>
- Santos, S. E., Soesanto, L., & Haryanto, T. A. D. (2007). Penekanan hayati penyakit moler pada bawang merah dengan *Trichoderma harzianum*, *Trichoderma koningii*, dan *Pseudomonas fluorescens* P60 [Biological suppression of moler disease on shallot by *Trichoderma harzianum*, *Trichoderma koningii*, and *Pseudomonas fluorescens* P60]. *Jurnal Hama Dan Penyakit Tumbuhan Tropika*, 7(1), 53–61. Retrieved from <http://journal.unila.ac.id/index.php/jhtrop/article/view/275>
- Sengupta, D., Aktar, M. W., Alam, S., & Chowdhury, A. (2010). Impact assessment and decontamination of pesticides from meat under different culinary processes. *Environmental Monitoring and Assessment*, 169(1–4), 37–43. <http://doi.org/10.1007/s10661-009-1148-6>
- Sharma, B. K., Singh, B. M., & Sugha, S. K. (1992). Integrated effect of biological and chemical control on sclerotial viability of *Sclerotinia sclerotiorum* (Lib.) de Bary. *Journal of Biological Control*, 6(1), 29–34. Retrieved from <http://www.informaticsjournals.com/index.php/jbc/article/viewFile/15197/12978>
- Singh, V. P., Srivastava, S., Shrivastava, S. K., & Singh, H. B. (2012). Compatibility of different insecticides with *Trichoderma harzianum* under *in vitro* condition. *Plant Pathology Journal*, 11(2), 73–76. <http://doi.org/10.3923/ppj.2012.73.76>
- Soesanto, L., Sudarmono, Prihatiningsih, N., Manan, A., Iriani, E., & Pramono, J. (2005). Potensi agensi hayati dan nabati dalam mengendalikan penyakit busuk rimpang jahe [Potency of biological and botanical agents in controlling ginger rhizome rot]. *Jurnal Hama Dan Penyakit Tumbuhan Tropika*, 5(1), 50–57. Retrieved from <http://jhtropika.fp.unila.ac.id/index.php/jhtropika/article/view/159>
- Soytong, K., & Quyet, N. T. (2013). Production of organic compost from mushroom producing substances waste and tested for Kangkong organic cultivation. *International Journal of Agricultural Technology*, 9(1), 115–123. Retrieved from <https://www.cabdirect.org/cabdirect/abstract/20133059295>
- Sutton, L. M., & Starzyk, M. J. (1972). Procedure and analysis of a useful method in determining mycelial dry weights from agar plates. *Applied Microbiology*, 24(6), 1011–1012. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC380718/>
- Thakore, Y. (2006). The biopesticide market for global agricultural use. *Industrial Biotechnology*, 2(3), 194–208. <http://doi.org/10.1089/ind.2006.2.194>
- Vinale, F., Sivasithamparam, K., Ghisalberti, E. L., Marra, R., Woo, S. L., & Lorito, M. (2008). *Trichoderma*-plant-pathogen interactions. *Soil Biology and Biochemistry*, 40(1), 1–10. <http://doi.org/10.1016/j.soilbio.2007.07.002>
- Wardhana, D. W., Soesanto, L., & Utami, D. S. (2009). Penekanan hayati penyakit layu *Fusarium* pada subang gladiol [Biological suppression of Fusarial Wilt on gladiolus corms]. *Jurnal Hortikultura*, 19(2), 199–206. Retrieved from <http://ejurnal.litbang.pertanian.go.id/index.php/jhort/article/view/802>
- Yang, C., Hamel, C., Vujanovic, V., & Gan, Y. (2011). Fungicide: Modes of action and possible impact on nontarget microorganisms. *ISRN Ecology*, 2011, 1–8. <http://doi.org/10.5402/2011/130289>
- Yildirim, M. Z., Benli, A. Ç. K., Selvi, M., Özkul, A., Erkoç, F., & Koçak, O. (2006). Acute toxicity, behavioral changes, and histopathological effects of deltamethrin on tissues (gills, liver, brain, spleen, kidney, muscle, skin) of Nile tilapia (*Oreochromis niloticus* L.) fingerlings. *Environmental Toxicology*, 21(6), 614–620. <http://doi.org/10.1002/tox.20225>
- Zimand, G., Elad, Y., & Chet, I. (1996). Effect of *Trichoderma harzianum* on *Botrytis cinerea* pathogenicity. *Phytopathology*, 86, 1255–1260. <http://doi.org/10.1094/Phyto-86-1255>