

FOOD RESEARCH

OPEN ACCESS





FOOD RESEARCH

Volume 6, Issue 6

December 2022

Mini Review

Factors affecting the properties of rice flour: a review

Ronie, M.E. and Hasmadi, M.

Available Online: 16 OCTOBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).531](https://doi.org/10.26656/fr.2017.6(6).531)

The factors affecting the properties of rice flour were reviewed by Ronie and Hasmadi.

Full Papers

Effects of pickling treatments and species on the properties of processed mushrooms (*Pleurotus* species)

Tanimola, A.R., Abai, O.P., Ibitoye, W.O. and Otegbayo, B.O.

Available Online: 4 NOVEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).508](https://doi.org/10.26656/fr.2017.6(6).508)

Tanimola *et al.* studied the effects of pickling treatments and species on the properties of processed mushrooms (*Pleurotus* species).

Formulation of sprouted cowpea yogurt rich in antioxidant, as functional drink for diabetics

Winarsi, H., Erminawati, E. and Ramadhan, G.R.

Available Online: 4 NOVEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).750](https://doi.org/10.26656/fr.2017.6(6).750)

The formulation of sprouted cowpea yogurt rich in antioxidant, as functional drink for diabetics was studied by Winarsi *et al.*

Effect of nutraceutical from mulberry on anthropometric measures in overweight and obese adults: a randomized, double-blinded, and placebo-controlled trial

Mangkalard, P., Limwikrant, W., Kanchanadumkerng, P., Pongpirul, W.A. and Lomarat, P.

Available Online: 4 NOVEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).617](https://doi.org/10.26656/fr.2017.6(6).617)

The effect of nutraceutical from mulberry on anthropometric measures in overweight and obese adults was studied by Mangkalard *et al.*

Development of γ -aminobutyric acid (GABA)-rich yoghurt using *Tetragenococcus halophilus* strain KBC isolated from a commercial soy sauce moromi

Azizan, N.A.Z., Lim T.C., Raseetha, S. and Wan-Mohtar, W.A.A.Q.I.

Available Online: 4 NOVEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).676](https://doi.org/10.26656/fr.2017.6(6).676)

Azizan *et al.* developed a γ -aminobutyric acid (GABA)-rich yoghurt using *Tetragenococcus halophilus* strain KBC isolated from commercial soy sauce moromi.

Comparative analysis of nutritional composition and droplet size of coconut milk due to dilution and emulsification

Abdullah, Z., Taip, F.S. and Mustapa Kamal, S.M.

Available Online: 11 NOVEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).763](https://doi.org/10.26656/fr.2017.6(6).763)

Abdullah *et al.* performed a comparative analysis of the nutritional composition and droplet size of coconut milk due to dilution and emulsification.

Detection and impact of heat treatments on tetracycline residues in chicken meats sold in Amman city, Jordan

Al Aboudi, A.R., Almashhadany, D.A. and Al Hijazin, M.

Available Online: 11 NOVEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).718](https://doi.org/10.26656/fr.2017.6(6).718)

Al Aboudi *et al.* detected tetracycline residues and studied the impact of heat treatments on tetracycline residues in chicken meats sold in Amman city, Jordan.

Effect of pH and temperature on antioxidant enzymes activities in *Morinda citrifolia* L. (Mengkudu) leaves extract

Mohd Zin, Z., Azman, S.N.S., Yahya, F., Hasmadi, M. and Zainol, M.K.

Available Online: 11 NOVEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).642](https://doi.org/10.26656/fr.2017.6(6).642)

The effect of pH and temperature on antioxidant enzymes activities in *Morinda citrifolia* L. (Mengkudu) leaves extract was studied by Mohd Zin *et al.*

Hepatoprotective effect of *Carica pubescens* juice and rutin through improving the Adenosine Monophosphate-activated Protein Kinase (AMPK) pathway in type 2 diabetes Mellitus Wistar Rats

Purbasari, E., Rachmawati, S.N., Anjani, G., Widyastiti, N.S. and Muniroh, M.

Available Online: 11 NOVEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).648](https://doi.org/10.26656/fr.2017.6(6).648)

Purbasari *et al.* evaluated the hepatoprotective effect of *Carica pubescens* juice and rutin through improving the Adenosine Monophosphate-activated Protein Kinase (AMPK) pathway in type 2 diabetes Mellitus Wistar Rats.

Effects of roasted defatted sesame seed (*Sesamum indicum*) flour addition on the nutritional quality, dietary fibre, and glycaemic index of malted amaranth grain (*Amaranthus hybridus*) flour

Ogundele, O.M., Ojedokun, F.O., Fasogbon, B.M. and Bamidele, O.P.

Available Online: 19 NOVEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).698](https://doi.org/10.26656/fr.2017.6(6).698)

The effects of roasted defatted sesame seed (*Sesamum indicum*) flour addition on the nutritional quality, dietary fibre, and glycaemic index of malted amaranth grain (*Amaranthus hybridus*) flour were studied by Ogundele *et al.*

Optimization of vitamin c extraction of roselle petals (*Hibiscus sabdariffa*) assisted with microwaves using response surface methodology

Purbowati, I.S.M., Maksum, A, Ritonga, A.M. and Kurniawan, R.E.K.

Available Online: 19 NOVEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).639](https://doi.org/10.26656/fr.2017.6(6).639)

Purbowati *et al.* optimized the extraction of roselle petals (*Hibiscus sabdariffa*) assisted with microwaves using response surface methodology.

The effect of high protein milk supplementation on satiety in normal weight subjects

Muliadi, R.D., Kartawidjajaputra, F. and Antono, L.

Available Online: 19 NOVEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).805](https://doi.org/10.26656/fr.2017.6(6).805)

Muliadi *et al.* evaluated the effects of high protein milk supplementation on satiety in normal weight subjects.

Physicochemical properties, sensory acceptance and glycaemic index of processed stingless bee honey and processed honeybee honey

Najeeb, I.M.A., Norhayati M.K., Zaharah, H., Mohd Isa, N.S., Nur Nadrah, M.R. and Yusof, H.M.

Available Online: 28 NOVEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).745](https://doi.org/10.26656/fr.2017.6(6).745)

The physicochemical properties, sensory acceptance and glycaemic index of processed stingless bee honey and processed honeybee honey were studied by Najeeb *et al.*

A qualitative study on fishery export refusals due to food safety concerns: identification of product handling, corrective actions, risk factors, and risk mitigation

Indrotristanto, N., Andarwulan, N., Fardiaz, D. and Dewanti-Hariyadi, R.

Available Online: 7 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).781](https://doi.org/10.26656/fr.2017.6(6).781)

Indrotristanto *et al.* performed a qualitative study on fishery export refusal from the product handling to risk mitigation.

High levels of bacterial and antimicrobial drug residues contamination in chicken eggs for human consumption in Morogoro municipality, Tanzania

Sineque, C.I.I.I.M., Machado, A.Da C., Macuamule, C.J. and Ngowi, H.A.

Available Online: 28 NOVEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).725](https://doi.org/10.26656/fr.2017.6(6).725)

Sineque *et al.* evaluated the high levels of bacterial and antimicrobial drug residues contamination in chicken eggs for human consumption in Morogoro municipality, Tanzania.

Consumer attitude toward Halal food in the case of the United Kingdom: the role of product attributes and marketing stimuli

Moghaddam, H.K., Khan, N., Tan, B.C. and Khan, S.

Available Online: 28 NOVEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).861](https://doi.org/10.26656/fr.2017.6(6).861)

Moghaddam *et al.* studied the consumer attitude toward Halal food in United Kingdom.

Appealing to the health-conscious: low glycaemic food purchase behaviour

Tan, B.Z., Lau, T.C., Tan, B.C. and Lo, Y.T.

Available Online: 7 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).890](https://doi.org/10.26656/fr.2017.6(6).890)

Tan *et al.* studied on the purchase behaviour of health-conscious consumers on the low glycaemic food.

Changing consumption behaviour towards immune-boosting food amidst COVID-19 outbreak: tracing the future trajectory

Rizwana., Padmalini, S., Ahalya, N., Sanjay, N., Narmada and Nayanika, B.

Available Online: 7 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).798](https://doi.org/10.26656/fr.2017.6(6).798)

Rizwana *et al.* evaluated the changing consumption behaviour towards immune-boosting food.

Improvement the texture and prebiotic properties of steam bun quality using cultivated banana (Musa ABB cv. Kluai 'Namwa') flour

Oupathumpanont, O. and Wisalsakkul, S.

Available Online: 7 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).800](https://doi.org/10.26656/fr.2017.6(6).800)

The improvement of texture and prebiotic properties of steam bun quality using cultivated banana (Musa ABB cv. Kluai 'Namwa' flour were studied by Oupathumpanont and Wisalsakkul.

Drying edible jellyfish (*Lobonema smithii*) using a parabolic greenhouse solar dryer

Maisont, S., Samutsri, W., Sansomboon, A. and Limsuwan, P.

Available Online: 18 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).883](https://doi.org/10.26656/fr.2017.6(6).883)

Maisont *et al.* studied on the drying edible jellyfish (*Lobonema smithii*) using a parabolic greenhouse solar dryer.

Effect of additional carrageenan concentration on the characteristics of wet noodle based on mangrove fruit flour variation

Sumartini and Putri, W.R.

Available Online: 18 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).709](https://doi.org/10.26656/fr.2017.6(6).709)

The effect of additional carrageenan concentration on the characteristics of wet noodle based on mangrove fruit flour variation was studied by Sumartini and Putri.

Household level determinants of food security in the City of Tshwane, South Africa

Mazenda, A., Manzi, P., Mushayanyama, T. and Ngarava, S.

Available Online: 18 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).915](https://doi.org/10.26656/fr.2017.6(6).915)

Mazenda *et al.* evaluated the household level determinants of food security in the City of Tshwane, South Africa.

Antioxidant potential of *Moringa stenopetala* leaf extract on laagered beer stored at room temperature

Gedefaw M.Y., Engeda D.A. and Tilku D.M.

Available Online: 18 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).550](https://doi.org/10.26656/fr.2017.6(6).550)

Gedefaw *et al.* evaluated the antioxidant potential of *Moringa stenopetala* leaf extract on laagered beer stored at room temperature.

Green tea extract improved minced mutton quality during chilled storage

Alnori, H.M., Saeed, O.A., Alnoori, M.A., Leo, T.K. and Sani, U.M.

Available Online: 21 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).672](https://doi.org/10.26656/fr.2017.6(6).672)

Alnori *et al.* evaluated the quality of minced mutton with green text extract during chilled storage.

The effect of concentration calcium chloride and immersion time on changes in colour and texture of fresh-cut mangosteen

Suriati, L.

Available Online: 21 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).752](https://doi.org/10.26656/fr.2017.6(6).752)

The effect of concentration calcium chloride and immersion time on changes in colour and texture of fresh-cut mangosteen was studied by Suriati.

Substitution of rice flour with banana flour for developing Khanom Thong Att products

Songpranam, P., Chowthanakij, R. and Noppakun, M.

Available Online: 21 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).655](https://doi.org/10.26656/fr.2017.6(6).655)

Songpranam *et al.* developed Khanom Thong Att products with substitution of rice flour with banana flour.

Consumer acceptance of fresh-cut peppers and tomatoes and their enhancement by edible coatings

Cloete, L., Hosany, H., Rungasamy, I., Ramful-Baboolall, D., Ramasawmy, B. and Neetoo, H.

Available Online: 21 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).661](https://doi.org/10.26656/fr.2017.6(6).661)

Cloete *et al.* evaluated consumer's acceptance of fresh-cut peppers and tomatoes enhanced with edible coatings.

Quality attributes of Black-Chin Tilapia (*Sarotherodon melanotheron*) surimis treated with different cryoprotectants

Perdio, A.C., Valerio, R.B., Consigna, M.J.S., Llave, D.S., Gonzales, A.J.V. and Turtur, M.

Available Online: 23 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).610](https://doi.org/10.26656/fr.2017.6(6).610)

The quality attributes of Black-Chin Tilapia (*Sarotherodon melanotheron*) surimis treated with different cryoprotectants were evaluated by Perdio *et al.*

The effect of tempeh flour on LDL (low-density lipoprotein)-level lowering in *Rattus norvegicus* fed a high-fat diet

Kusuma, H.S., Febrianti, D., Rosidi, A. and Ulvie, Y.N.S.

Available Online: 23 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).780](https://doi.org/10.26656/fr.2017.6(6).780)

Kusuma *et al.* studied of the effect of tempeh flour on LDL-level lowering in *Rattus norvegicus* fed with a high-fat diet.

Conceptual paper: Malaysian youth's perception and consumption behaviour of roasted chicken products towards a better quality of life

Enthira, P.T., Yusnita, H., Azizulyadi, Y., Asma', A. and Siti Nur'afifah, J.

Available Online: 23 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).774](https://doi.org/10.26656/fr.2017.6(6).774)

Enthira *et al.* proposed to study Malaysian youth's perception and consumption behaviour of roasted chicken products towards a better quality of life.

Effects of cooking methods on physicochemical properties, antioxidant properties and sensory acceptability of purple sweet potato (*Ipomoea batatas*)

Cheong, J.J., Ahmad, F. and Tengku Rozaina, T.M.

Available Online: 23 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).1008](https://doi.org/10.26656/fr.2017.6(6).1008)

Cheong *et al.* evaluated the effects of cooking method on physicochemical properties, antioxidant properties and sensory acceptability of purple sweet potato (*Ipomoea batatas*).

Determination of total phenolic and flavonoid content of Ghontoghe (*Kleinhovia hospita* L.) leaves: in vitro antioxidant study

Nuralifah, Yamin, Halidi, D.R., Arba, M. and Zubaydah, W.O.S.

Available Online: 27 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).440](https://doi.org/10.26656/fr.2017.6(6).440)

Nuralifah *et al.* determined the total phenolic and flavonoid content of Ghontoghe (*Kleinhovia hospita* L.) leaves.

Malaysian Muslim consumers' awareness, confidence, and purchase behaviour on halal meat and its products after the meat cartel scandal

Maifiah, M.H.M., Ahmad, A.N., Azam, M.S.E., Norazmi, A.R.M. and Nawawi, K.A.

Available Online: 27 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).714](https://doi.org/10.26656/fr.2017.6(6).714)

Maifiah *et al.* studied on the awareness, confidence and purchase behaviour of Malaysian Muslim consumers' on halal meat and its products after the meat cartel scandal.

Comparative study of physicochemical, nutritional and functional properties of whole and defatted legume flours

Jamalullail, N.A., Chan, Y.L., Tang, T.K., Tan, C.P., Mat Dian, N.L.H., Cheong, L.Z. and Lai, O.M.

Available Online: 27 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).659](https://doi.org/10.26656/fr.2017.6(6).659)

Jamalullail *et al.* performed a comparative study to investigate the physicochemical, nutritional and functional properties of whole and defatted legume flours.

Incorporation of *Neolamarckia cadamba* leaf extract into chitosan coating for maintaining the quality of white shrimp (*Litopenaeus vannamei*)

Mubarak, A., Zulkipli, N.N., Norasmadi, I.N.S., Suhaimi, N.I., Wan Abdullah, W.Z. and Karim, N.U.

Available Online: 27 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).734](https://doi.org/10.26656/fr.2017.6(6).734)

The incorporation of *Neolamarckia cadamba* leaf extract into chitosan coating to maintain the quality of white shrimp (*Litopenaeus vannamei*) was evaluated by Mubarak *et al.*

Storage stability of microencapsulated anthocyanin extracted from Kokum (*Garcinia indica*) rind

Ranveer, R.C., Rathod, N.B., Nandane A.S., Sahoo A.K. and Ganorkar P.M.

Available Online: 31 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).870](https://doi.org/10.26656/fr.2017.6(6).870)

The storage stability of microencapsulated anthocyanin extracted from Kokum (*Garcinia indica*) rind was studied by Ranveer *et al.*

Knowledge and attitude of selected non-Muslim college students towards Halal-certified food products

Barrion, A.S.A., Orillo, A.T.O., Nguyen-Orca, M.F.R. and Tandang, N.A.

Available Online: 31 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).869](https://doi.org/10.26656/fr.2017.6(6).869)

Barrion *et al.* evaluated the knowledge and attitude of selected non-Muslim college students towards Halal-certified food products.

Spent brewer yeast status in halal food industry: An analysis from Shariah and science perspectives

Ramli, S.N.H., Jamaludin, M.A., Abdullah Sani, M.S. and Nordin, N.F.H.

Available Online: 31 DECEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).794](https://doi.org/10.26656/fr.2017.6(6).794)

Ramli *et al.* performed an analysis on spent brewer yeast status in halal food industry from Shariah and science perspectives.

Mini Review

Anti-noncommunicable diseases attributes of pineapple (*Ananas comosus*): a concise review

Kumalawati, D.A., Nurkolis, F., Wewengkang, D.S., Permatasari, H.K., Rotinsulu, H., Taslim, N.A., Joseph, V. and Samtiya, M.

Available Online: 19 NOVEMBER 2022 | [https://doi.org/10.26656/fr.2017.6\(6\).887](https://doi.org/10.26656/fr.2017.6(6).887)

The review of anti-noncommunicable disease attributes of pineapple (*Ananas comosus*) was performed by Kumalawati *et al.*

VOL. 6 | ISSUE 5

VOL. 6 | SUPPLEMENTARY ISSUE 1

[HOME](#)[VIEW ARTICLES](#)[VIEW ISSUES](#)[FOR AUTHORS](#)[FOR REFEREES](#)[ABOUT THE JOURNAL](#)

EDITORIAL BOARD

Chief Editor

Son Radu

Malaysia

Editor

Vivian New Chia Yeung

Malaysia

Associate Editors

Prem Narayan Paudel

Kathmandu University

Nepal

Associate Editors

Borarin Buntong

Royal University of Agriculture

Cambodia

Associate Editors

Yoshitsugu Nakaguchi

Ishikawa Prefectural University

Japan

Nasreldin Elhadi

University of Dammam

Saudi Arabia

Dwaipayan Sikdar

University of Chittagong

Bangladesh

Rungsinee Sothornvit

Kasetsart University,

Thailand

Nikos Tzortzakis

Cyprus University of Technology

Cyprus

Saisamorn Lumyong

Chiang Mai University

Thailand

Ekachai Chukeatirote

Mae Fah Luang University

Thailand

Optimization of vitamin c extraction of roselle petals (*Hibiscus sabdariffa*) assisted with microwaves using response surface methodology

^{1,*}Purbowati, I.S.M., ¹Maksum, A., ²Ritonga, A.M. and ³Kurniawan, R.E.K.

¹Food Processing Technology Laboratory, Department of Agriculture Technology Faculty of Agriculture, Universitas Jenderal Soedirman, Dr Suparno Street, Karangwangkal, Purwokerto 53123, Indonesia

²Thermal system engineering and renewable energy Laboratory, Department of Agriculture Technology Faculty of Agriculture, Universitas Jenderal Soedirman, Dr Suparno Street, Karangwangkal, Purwokerto 53123, Indonesia

³Soil Science and Land Resource Planning Laboratory, Department of Agriculture Technology Faculty of Agriculture, Universitas Jenderal Soedirman, Dr Suparno Street, Karangwangkal, Purwokerto 53123, Indonesia

Article history:

Received: 17 September 2021

Received in revised form: 29 October 2021

Accepted: 25 February 2022

Available Online: 19 November 2022

Keywords:

Roselle (*Hibiscus sabdariffa*), vitamin C, Extraction, Response Surface Methodology

DOI:

[https://doi.org/10.26656/fr.2017.6\(6\).639](https://doi.org/10.26656/fr.2017.6(6).639)

Abstract

Vitamin C contained in roselle petals acts as an antioxidant that provides health benefits. In order to make it easier to use, vitamin C needs to be extracted out of the roselle petal cells in the form of extracted raw roselle juice. The commonly used conventional method has disadvantages that can damage heat-sensitive compounds such as vitamin C. An alternative extraction method used was microwave-assisted extraction. The success of this method in extracting vitamin C was influenced by a few factors, such as the ratio of the roselle petals-water, microwave power, and extraction time. The purpose of this study was to determine the optimum conditions for microwave-assisted extraction of vitamin C in roselle extract and to determine the physical and chemical characteristics of roselle extract produced in the optimum conditions. This study utilized response surface methodology (RSM) in a central composite design (CCD). The optimized factors included the roselle petals-water ratio (5 - 10%), microwave power (150 - 350 W), and extraction time (3 - 7 mins). The parameter response tested in this study was vitamin C. While the extract produced under optimum conditions was also characterized by the total anthocyanin content, total phenols, antioxidant activity, pH value and colour (L, a* and b*). The results showed that the recommended optimum conditions were a 5% roselle petals-water ratio, 250 W of microwave power, and 6.09 mins of extraction time. The verification results show that this optimum condition produced vitamin C higher than the conventional method. The characteristics of roselle extract produced by microwave-assisted extraction at optimum conditions were vitamin C at 9.973 ± 0.508 mg/g, total anthocyanins at 1.771 ± 0.124 mg/100 g, phenol at 30.848 ± 1.899 mg/g, IC_{50} at 6.699 ± 0.627 ppm, pH at 2.133 ± 0.057 , colour with an L value of 31.033 ± 0.404 , a* value of 1.066 ± 0.155 , and b* value of 18.100 ± 0.458 .

1. Introduction

Ascorbic acid (Vitamin C) is one of the most important active substances which humans should consume in sufficient amounts together with food. Recent epidemiological studies have associated the natural antioxidant properties of vitamin C with possible health benefits in the prevention of the development of coronary heart disease and anticancer activities (Bai *et al.*, 2015). Recently, vitamin C has been widely recommended as an immune-boosting supplement during

the COVID-19 pandemic.

In line with increasing public knowledge and awareness of health, natural antioxidants are increasingly in demand. One plant that has the potential to be used is roselle (*Hibiscus sabdariffa* L.). Roselle does not just taste good but it also contains large amounts of vitamin C (Purbowati *et al.*, 2020). According to Babajide *et al.* (2004), roselle petals contain several vitamins and minerals such as vitamin C, calcium, niacin, riboflavin, and flavonoids which play a role in the body's cell

*Corresponding author.

Email: ikesitoresmi@yahoo.co.id



Make an Impact

Publish Open Access

Make an impact.

tandfonline.com

OPEN

Food Research

COUNTRY

Malaysia



Universities and research
institutions in Malaysia

SUBJECT AREA AND CATEGORY

Agricultural and Biological Sciences
Food Science

PUBLISHER



Gellan Gum for Almond N

For Dairy Products

High Acyl Gellan Gum HAM Type for da
products or milk beverages.

opalgellangum.com

OPEN

PUBLICATION TYPE

Journals

ISSN

25502166

COVERAGE

2017-
2020



Publish Open Access - Regional Studies

Want to find the perfect home for your regional studies research? tandfonline.com

For Dairy Products

High Acyl Gellan Gum HAM Type for dairy products or milk beverages.

opalgellangum.com

OPEN

SCOPE

Food Research welcomes submissions that are relevant to food science and technology, food service management, nutrition, nutraceuticals, food innovation and agriculture food science. Manuscript types accepted are reviews, short reviews, original research articles and short communications that discusses on the current research and scientific views of the following scope: Food science and food chemistry; Food technology, processing, and food engineering; Food safety and quality – microbiological and chemical; Sensory, habits, consumer behaviour/preference; Nutrition and dietetics; Nutraceuticals and functional food/ingredients; Food service management; Food trends, innovation; Post-harvest and agribusiness; Food security; Food packaging.

 Join the conversation about this journal

Submit Your Manuscript With Us

Open

Hindawi

 Quartiles



FIND SIMILAR JOURNALS 



Publish Open Access - Regional Studies

Want to find the perfect home for your regional studies research? tandfonline.com

Submit Your Manuscript With Us

Hindawi

1
International Food Research Journal
MYS

89%
similarity

2
Food Science and Nutrition
GBR

82%
similarity


3
Carpathian Journal of Food Science and Technology
ROU

79%
similarity


4
International Science
EGY

7
s

SJR




Total Documents




Total Cites

Self-Cites




Citations per document



External Cites per Doc

Cites per Doc




% International Collaboration

Citable documents

Non-citable documents

Publish Open Access - Regional Studies

Want to find the perfect home for your regional studies research? tandfonline.com



400

200



← Show this widget in your own website


Just copy the code below and paste within your html code:

`<a href="https://www.scimagojr.com">`

SCImago Graphica

Explore, visually communicate and make sense of data with our **new free tool**.

Get it



Submit Your Manuscript With Us

Op

Hindawi

Metrics based on Scopus® data as of April 2021

T **Tamiur Yazew** 4 weeks ago

Dear,


Thank you for your updated and quality journal.

I have submitted two manuscripts in the Food research journal. The editor of this journal sent me the acceptance letter for the two manuscripts. However, they asked me to pay charge for the manuscripts. I am also working as a reviewer of this journal. I have edited two papers and sent them to the editor. I am currently reviewing a paper and it is ready to send back to the editor of this journal.



Publish Open Access - Regional Studies

Want to find the perfect home for your regional studies research? tandfonline.com



So, please would you help me by considering my issue into consideration!

reply



Melanie Ortiz 4 weeks ago

SCImago Team

Dear Tamiur, thank you very much for your comment. Unfortunately, we cannot help you with your request, we suggest you contact the journal's editorial staff so they could inform you more deeply. Best Regards, SCImago Team

K

Kamal 9 months ago

Hi, Editorial Team members,
I would like to know the topic related to the "hygienic practices along the supply chain of fisheries" is considered or not to review of your Journal. Early response is highly appreciated

reply



Melanie Ortiz 9 months ago

SCImago Team

Dear Kamal,
thank you for contacting us.
We are sorry to tell you that SCImago Journal & Country Rank is not a journal. SJR is a portal with scientometric indicators of journals indexed in Elsevier/Scopus.
Unfortunately, we cannot help you with your request, we suggest you visit the journal's homepage or contact the journal's editorial staff, so they could inform you more deeply.
Best Regards, SCImago Team

Y

yani purbanang 1 year ago

Hi,
This journal is written with Scopus index from 2017-2019, What is the index status in 2021 ??
Thank you for your explanation

reply



Melanie Ortiz 1 year ago


SCImago Team

Dear Yani,
Thank you very much for your comment.
All the metadata have been provided by Scopus /Elsevier in their last update sent to SCImago, including the Coverage's period data. The SJR for 2019 was released on 11



Publish Open Access - Regional Studies

Want to find the perfect home for your regional studies research? tandfonline.com



S **Sigit Susanto** 1 year ago

Whether the supply chain on Runner products could be submitted to this FR Journal?thanks very much

reply



Melanie Ortiz 1 year ago

SCImago Team

Dear Sigit,

Thank you for contacting us. Could you please expand a little bit your comment?

Best Regards, SCImago Team

Leave a comment

Name

Email

(will not be published)



I'm not a robot

reCAPTCHA
[Privacy](#) - [Terms](#)

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.


Developed by:

Powered by:



Publish Open Access - Regional Studies

Want to find the perfect home for your regional studies research? tandfonline.com



Follow us on @ScimagoJR


Scimago Lab, Copyright 2007-2020. Data Source: Scopus®

EST MODUS IN REBUS
Horatio (Satire 1, 1, 106)



Publish Open Access - Regional Studies

Want to find the perfect home for your regional studies research? tandfonline.com



rejuvenation process. The content of vitamin C in roselle petals per 100 grams is 260-280 mg. According to Purbowati *et al.* (2016), vitamin C, total phenol, anthocyanin, and yield of microwave-assisted extraction respectively were 10.74 ± 0.14 mg/g, 23.77 ± 0.25 mg/g, 14.80 ± 0.08 mg/g, and $22.09 \pm 3.3\%$. Maksum and Purbowati (2017), reported that the extraction of roselle using the MAE method with an extraction time of 7 mins, a roselle weight of 10 g, and a power of 400 W produced 74.8 mg/100 mg of Vitamin C. The optimized extraction parameters were determined at a soaking time of 48 mins, the solid-solvent ratio of 44% and the extraction temperature of 55°C produced 7.8 mg/100 g L-ascorbic acid (Peter *et al.*, 2014).

It is known that vitamin C inhibits oxidation processes and can be used as an antioxidant. Purbowati *et al.* (2015) reported that the antioxidant activity in rosella flowers is not only caused by phenol compounds but is also determined by the amount of vitamin C. Therefore, technological processes used for extraction have an impact on the amount of vitamin C. Most of it was lost during the extraction process. Extracting vitamin C using conventional extraction methods was time-consuming and less efficient, and according to Hapsari *et al.* (2021), extraction of roselle by maceration method with sample-solvent ratio 1:12.5 took an extraction time of 72 hrs, while the methanol solvent yielded 27.3% of the extract, total phenolic compounds of 29.2 mg GAE/g, total flavonoid compounds of 36.7 mg QE/g, and DPPH scavenging activity of 78%. Whereas Kurtulbaş *et al.* (2020) reported that extraction by the MAE method with power 550 W, time 180 s, DES in 50% water solvent produced total phenolic compounds of 31.90 mg GAE/g, total anthocyanin compounds of 3.00 mg Cy-3-GE/g and DPPH scavenging activity of 95.89%. An alternative extraction method that can be done is the Microwave-Assisted Extraction (MAE) method.

MAE had several advantages, such as lower extraction temperature and shorter extraction time compared to the conventional one, thereby reducing the thermolabile compounds (Wei *et al.*, 2009). Water is a suitable solvent for MAE because water has a dielectric coefficient of 78.3 throughout it can absorb extraction microwaves well (Markom, 2007).

Solvent volume, material weight, extraction time, and microwave power will interact to influence the yield and activity of bioactive compounds produced in the microwave-assisted extraction process (Mandal *et al.*, 2007). The solvent-raw material ratio affects the ability of the solvent to dissolve the compound from the material until the solution reaches its saturation point. Microwave power and extraction time affect the amount

of heat energy received by vitamin C. These three factors need to be known to optimize extraction conditions during microwave-assisted extraction. Aulia (2018), reported that the extraction of soursop leaves in total phenol and antioxidant response obtained the optimum point from the extraction time of 9 mins 84 s with a solvent ratio of 25.19 with the raw material in the form of dried soursop leaf powder as much as 25 g.

Based on the description above, the objectives to be achieved in this study were to determine the optimum conditions for microwave-assisted extraction to the response of vitamin C content in roselle extract, as well as to know the physical and chemical characteristics of roselle extract produced in optimum conditions.

2. Materials and methods

2.1 Equipment

The tools used to make roselle powder include cabinet dryers, grinders, 80 mesh sieves, and microwaves (Electrolux EMM 2007X) for extraction. While the tools for analysis include a UV-Vis spectrophotometer, centrifuge, colour reader, rotary evaporator, water bath, vortex, pH meter, Erlenmeyer flask, measuring flask, measuring cup, beaker glass, test tube, drop pipette, measuring pipette, analytical balance, paper, filter, funnel, plastic cup, metal spatula, aluminium foil.

2.2 Material

The materials used were roselle flowers, while the materials for analysis included NaCl, NaOH, HCl, Na_2CO_3 , 95% ethanol, methanol, DPPH, Folin-Ciocalteu, starch, iodine, and distilled water.

2.3 Experimental design

This study used the Response Surface Method using the Central Composite Design (CCD). The factors observed were the roselle petals-water ratio (5, 10, 15% w/v) (Chumsri, *et al.*, 2008), microwave power (150, 200, and 250 W) (Purbowati *et al.*, 2015), and extraction time (3, 5, and 7 mins) (Purbowati *et al.*, 2015).

2.4 Preparation of roselle powder

The preparation was conducted according to the method of Purbowati *et al.* (2019). The roselle petals were sorted and then dried in a cabinet dryer at a temperature of 50°C for 6 hrs, after drying they were ground into powder form. It was then sieved with an 80 mesh sieve.

2.5 Optimization

Starting from the determination of the independent variable to determine the maximum-minimum value

range which was then inputted into the Design-Expert program using the Central Composite Design. Twenty treatment combinations obtained were analysed for vitamin C. The program will determine the optimization based on the input response measurement data, thus providing recommendations for some optimal combinations according to the program. The most optimal response was chosen with the highest desirability value.

2.6 Extraction

Approximately 10 g of the dried roselle petals were ground for 1 min using a grinder and sieved in a 60 mesh sieve. The extraction was done in 100 mL distilled water, microwave power extraction 250 W for 5 mins. The microwave used for extraction was Electrolux EMM 2007X. The slurry was radiated in a microwave oven at regular intervals (one-min radiation and two mins off) to ensure the temperature did not rise above the boiling point (Purbowati *et al.*, 2016). Roselle extract was filtered and concentrated with a vacuum evaporator at 70°C, 44 cmHg and blowing with N₂ to ensure the solvent evaporate.

2.7 Vitamin C analysis

The optimum conditions of the extraction process are determined based on the vitamin C response as measured by (Jacob, 1999). Standardization of sodium thiosulfate solution, this standardization was done by pipetting 5.00 mL of KIO₃ solution + 10 mL of 10 % KI solution + 10 mL of dilute H₂SO₄ into a conical flask. The liberated I₂ was titrated against NS₂S₂O₃ solution from a burette until a faint yellow colour was reached. Then 10 drops of the starch solution were added and the titration was continued until the blue-black colour of the starch-triiodide complex just disappeared. The titration was repeated until concordant results were obtained. The normality of the sodium thiosulfate solution was calculated. For the titration of the standard solution of ascorbic acid, 5.00 mL of the sample solution was pipetted and the steps for the standardization of sodium thiosulfate solution were repeated. The concentration of the ascorbic acid was calculated from the relation, Milliequivalent of sodium iodate-milliequivalent of sodium thiosulfate = milliequivalent of ascorbic acid. Titration of samples, 5.00 mL of the treated sample was pipetted into a conical flask and the steps were repeated as with standard ascorbic acid. The data resulting from these measurements were processed by the Design Expert to determine the optimum conditions.

2.8 Data analysis

The results of vitamin C data were analysed using RSM. The optimization design was carried out using

CCD (Central Composite Design) and analysed using the Response Surface Methodology. The equation with a single expected response (Y) was:

$$Y = \beta_0 + \beta_{ixi} + \beta_{iixi^2} + \beta_{ijx_i x_j} + \epsilon$$

Where 0 was the value of the expected response at the centre of the design being developed, namely (0, 0, 0). i, ii, and ij were linear, quadratic, and cross-product regression model constants, respectively.

2.9 Verification

The optimum conditions were obtained by running the program of Central Composite Design with the highest response to vitamin C. After the optimum condition was chosen, the verification experiments were conducted 3 times and each replication will be followed by physical and chemical characterization, colour (Soewarno, 1990), total anthocyanins (Fuleki and Francis, 1968), total phenol (Bray and Thorpe, 1954), IC₅₀ (Sheikh *et al.*, 2009 with modification), pH (Suwetja, 2007). The verification results obtained were then compared with the response variable values predicted by RSM.

The results of the verification were carried out also by comparing the characteristics of the roselle extract between microwave-assisted extraction and extraction using conventional heating. This conventional one was carried out in the optimum conditions for microwave-assisted extraction on a water bath device.

3. Results and discussion

3.1 Optimization

The determination of the model to predict the relationship between treatment and response was based on the data in Table 1, which is the basis for determining model selection. The data from the optimization were tested in the Analysis of Variance (ANOVA) to find a suitable equation model that relates the solvent-solvent ratio, microwave power, and extraction time with the vitamin C content as the response in this study. ANOVA results from the extraction optimization process in these studies can be seen in Table 2.

Based on the results of the ANOVA test, the optimization results of roselle extraction with vitamin C response were obtained with a significant quadratic model because it had a p-value of <0.0001. These showed that the regression model can represent the independent variables in influencing vitamin C. Supported by the insignificant Lack of Fit value because it had a value of 0.5040 (p-value > 0.05) which means that there was a fit of the model with the response data of vitamin C. The model used can be described by the

Table 1. Central Composite Design (CCD) of research factor

Run	Factor 1 A: roselle petals-water ratio (%)	Factor 2 B: power microwave (Watt)	Factor 3 C:time extraction (mins)	Response Vitamin C (mg/g)
1	10.00	250.00	5.00	7.04
2	15.00	150.00	7.00	4.106667
3	10.00	250.00	5.00	7.92
4	5.00	150.00	3.00	7.04
5	5.00	350.00	7.00	8.8
6	15.00	350.00	3.00	4.106667
7	15.00	150.00	3.00	4.693333
8	10.00	250.00	5.00	7.92
9	15.00	350.00	7.00	3.52
10	5.00	350.00	3.00	7.04
11	5.00	150.00	7.00	8.8
12	10.00	250.00	5.00	7.92
13	10.00	250.00	1.64	5.28
14	10.00	100.00	5.00	6.16
15	10.00	250.00	5.00	7.92
16	1.59	250.00	5.00	11.06918
17	10.00	250.00	8.36	6.16
18	18.49	250.00	5.00	5.235262
19	10.00	250.00	5.00	7.92
20	10.00	418.18	5.00	4.4

Table 2. Analysis of variance

Source	Sum of Squares	df	Mean Square	F-Value	p-value Prob > F	
Block	0.24	2	0.12			
Model	71.95	9	7.99	58.63	< 0.0001	significant
A-roselle petals-water ratio	46.00	1	46.00	337.39	< 0.0001	
B-microwave power	1.25	1	1.25	9.17	0.0163	
C-extraction time	1.07	1	1.07	7.86	0.023	
AB	0.17	1	0.17	1.26	0.2938	
AC	2.75	1	2.75	20.19	0.002	
BC	0.00	1	0.00	0.00	1.0000	
A ²	0.07	1	0.07	0.51	0.4965	
B ²	12.89	1	12.89	94.55	< 0.0001	
C ²	9.00	1	9.00	66.01	< 0.0001	
Residual	1.09	8	0.14			
Lack of Fit	0.70	5	0.14	1.09	0.504	not significant
Pure Error	0.39	3	0.13			
Cor Total	73.28	19				

equation $Y = 7.77 - 1.84A - 0.30B + 0.28C - 0.15AB - 0.59AC + 0.000BC + 0.069A^2 - 0.95B^2 - 0.79C^2$.

3.2 Effect of roselle petals-water ratio, microwave power and extraction time to the total of vitamin C in crude roselle extract

Based on Figure 1, it was known that microwave power has a significant effect on the total vitamin c of roselle crude extract with a significance value of 0.0163 less than 0.05. According to Kusuma and Mahfud (2015), the microwave will break the cell walls of the material. An increase in microwave power can cause damage to the cell wall more quickly and the solvent will

easily diffuse into the material to dissolve the compound. The higher the extraction temperature, the wider the cell pores are, making it easier for the solvent to enter cell-matrix and extract the compound (Wuryantoro and Susanto, 2014). However, too high a temperature causes a compound with a certain degree of resistance to temperature to be decomposed into its components. According to Pakaya (2014) this vitamin C was easily damaged by the oxidation process, especially when exposed to heat. The results of previous studies by Kadakal *et al.* (2018), ascorbic acid is very unstable at 95°C with a k value of 15×10^{-3} .




Figure 1. Surface response from the optimization between treatment and responses

Figure 1 also show that the roselle petal-water ratio has a significant effect on the total vitamin C of roselle crude extract with a significance value of 0.0001. The roselle petals-water ratio affected the amount of surface area the solvent could contact. According to Gertenbach (2002), the greater the roselle petals-water ratio would increase the concentration of dissolved solutes on the surface of the particles, thus increasing the concentration gradient inside and on the surface of the solid particles.

When the extraction time increases, the solvent has more opportunity to come into contact with the material, making the solvent diffused more into the pores of the material to dissolve vitamin C. The extracted material components would continue to increase until the solution became saturated. After passing the saturation point of the solution, there would be no increase in the extraction yield with the addition of solvent (Siregar, 2005). Figure 1 shows that extraction time also has a significant effect on the total vitamin C of roselle crude extract with a significance value of 0.0230. Winata (2015) agreed the longer the extraction time, the greater the amount of material extracted because the contact time between the

material and the solvent is getting longer, and the results would increase to the saturation point of the solution.

The next step after getting the equation in modelling is to determine the level of importance of each optimization variable. This was done to get the desired response (desirability) in the setting of criteria, as in Table 3. Based on the Design Expert v.10 program/application of the modelling used for optimization and the criteria for the level of importance, the program provides the optimum combination of conditions described by the response surface (Response Surface). Referring to the 3D contour in Figure 1, the optimum point was indicated by a red dot at the top of the saddle. The optimum conditions were obtained at a roselle petals -water ratio of 5%, microwave power of 250 W, and an extraction time of 6.09 mins with a desirability value of 0.920. The desirability value indicates that the optimum conditions for roselle extraction in accordance with the determination of the criteria of interest for optimization are 92.0%, which means that the optimum value was suitable because the desirability value was close to 1.

3.3 Verification

Table 3. Solution

No	Roselle petals-water ratio	Microwave power	Extraction time	vitamin C	StdErr (vitamin C)	Desirability	
1	5	250	6.099	9.916	0.178	0.92	Selected

Based on the results of verification at optimum conditions, it was known that the actual data falls within the Prediction Interval (PI) range. It was shown that the optimum condition was obtained by a combination of a roselle petals-water ratio of 5%, microwave power of 250 W, and an extraction time of 6.09 mins which will produce 9.91 mg/g of vitamin C. A previous study by Purbowati (2014), extracted roselle using ethanol as a solvent, and obtained optimum conditions at the extraction time of 4.91 mins, 78.36% ethanol and 250 W microwave power with vitamin C of 10.74 ± 0.14 mg/g. The amount of vitamin C extracted (Wang *et al.* 2010) performed on mulberry leaf extraction with MAE, resulted at 255 W for 5 mins with a yield of 8.37 mg/g.

Table 4 shows that extraction using MAE had higher characteristics than conventional extraction. During MAE, heat transfer occurs from the material to the solvent in the MAE and was distributed volumetrically throughout the irradiated sample. In this phenomenon, the heat was created in the microwave absorbent agent and transferred to the extraction medium. The similarity of the two transport phenomena, which are heat and mass transfer, makes the MAE extraction rate higher than conventional solvent extraction (Taghvaei *et al.*, 2014). This could prove that microwave-assisted extraction could provide the advantage of faster mass transfer, leading to a shorter extraction time to obtain optimum results. According to Amir *et al.* (2016), the energy in the extraction process using the MAE method goes directly to the specific target sample reducing the heat that was lost to the environment. Also, the bioactive compounds in the material were easier to dissolve because the heating process takes place in the closed system.

Vitamin C produced from the extraction of the MAE method was higher than that of the conventional method, which was 9.973 mg/g. This is in line with the higher

Table 4. Characteristics of roselle petals extract under optimum conditions with MAE and conventional

Characteristics	MAE	Conventional
Vitamin C (mg/g)	9.973 ± 0.508	8.213 ± 0.508
Total anthocyanin (mg/g)	1.771 ± 0.124	0.901 ± 0.184
Total Phenol (mg/ 100 g)	30.848 ± 1.899	10.705 ± 0.490
IC ₅₀ (ppm)	6.699 ± 0.627	10.170 ± 1.128
pH	2.133 ± 0.057	2.466 ± 0.057
Colour		
L	31.033 ± 0.404	27.533 ± 0.568
a*	1.066 ± 0.155	0.667 ± 0.115
b*	18.100 ± 0.458	16.866 ± 0.665

total anthocyanin content and lower pH in the MAE method. According to Harris and Karmas, (1989), the stability of ascorbic acid (vitamin C) will increase with decreasing pH value. In addition, Wong *et al.* (2002) also agreed that in acidic conditions or if the pH approaches pH 1, more anthocyanin pigments are in the form of flavour cations which are red and stable.

4. Conclusion

This research recommends that the optimum conditions for vitamin C extraction in roselle were obtained at a roselle petal-water ratio of 5%, microwave power of 250 W, and an extraction time of 6.09 mins. This optimum condition is expected to extract 9.91 mg/g of vitamin C. The verification results show that this optimum condition produces vitamin C higher than the conventional method. The characteristic of roselle extract produced by microwave-assisted extraction at optimum conditions, having vitamin C of 9.973 ± 0.508 mg/g, total anthocyanins of 1.771 ± 0.124 mg/100 g, phenol of 30.848 ± 1.899 mg/g, IC₅₀ of 6.699 ± 0.627 ppm, pH of 2.133 ± 0.057 , colour with L* value of 31.033 ± 0.404 , a* value of 1.066 ± 0.155 , b* value of 18.100 ± 0.458 .

Acknowledgements

This research was funded through the Jenderal Soedirman University Basic Research Scheme in 2021.

References

- Amir, A., Wiraningtyas, A. and Annafi, N. (2016). Perbandingan metode ekstraksi natrium alginat: metode konvensional dan Microwave Assisted Extraction (MAE). *Chempublish Journal*, 1(2), 7–13. [In Bahasa Indonesia].
- Aulia, L.P. (2018). Optimasi proses ekstraksi daun sirih (*Annona muricata* L) metode MAE (Microwave Assisted Extraction) dengan respon aktivitas antioksidan dan total fenol. *Jurnal Agroindustri Halal*, 4(1), 9–87. [In Bahasa Indonesia]. <https://doi.org/10.30997/jah.v4i1.1142>
- Babajide, J.M., Bodunde, J.G. and Salami, A.A. (2004). Quality and sensory evaluation of processed calyces of six varieties of roselle (*Hibiscus sabdariffa*). *Nigerian Journal of Horticultural Science*, 9(1), 3388. <https://doi.org/10.4314/njhs.v9i1.3388>
- Bai, X.Y., Qu, X., Jiang, X., Xu, Z., Yang, Y., Su, Q., Wang, M. and Wu, H. (2015). Association between

- dietary vitamin C intake and risk of prostate cancer: A meta-analysis involving 103,658 subjects. *Journal of Cancer*, 6(9), 913–921. <https://doi.org/10.7150/jca.12162>
- Bray, H.G. and Thorpe, W.V. (1954). Analysis of phenolic compounds of interest in metabolism. *Methods of Biochemical Analysis*, 1, 27–52. <https://doi.org/10.1002/9780470110171.ch2>
- Fuleki, T. and Francis, F.J. (1968). Quantitative methods for anthocyanins. 1. extraction and determination of total anthocyanin in cranberries. *Journal of Food Science*, 33(1), 72–77. <https://doi.org/10.1111/j.1365-2621.1968.tb00887.x>
- Gertenbach, D. (2002). Solid-Liquid Extraction Technologies for Manufacturing Nutraceutical. USA: CRC Press. <https://doi.org/10.1201/9781420012873.ch11>
- Hapsari, B.W., Manikhanda and Setyaningsih, W. (2021). Methodologies in the analysis of phenolic compounds in roselle (*Hibiscus sabdariffa* L.): composition, biological activity, and beneficial effects on human health. *Horticulturae*, 7(35), 1–41. <https://doi.org/10.3390/horticulturae7020035>
- Harris, R.S. and Karmas, E. (1989). Evaluasi Gizi Pada Pengolahan Bahan Pangan. Bandung, Indonesia: ITB. [In Bahasa Indonesia].
- Jacob, R. (1999). Vitamin C. 9th ed. Baltimore, USA: Williams and Wilkins.
- Kadalk, Ç., Duman, T. and Ekin, R. (2018). Thermal degradation kinetics of ascorbic acid, thiamine and riboflavin in rosehip (*Rosa canina* L) nectar. *Food Science and Technology*, 38(4), 667–673. <https://doi.org/10.1590/1678-457x.11417>
- Kurtulbaş, E., Pekel, A.G., Bilgin, M., Makris, D.P. and Şahin, S. (2020). Citric acid-based deep eutectic solvent for the anthocyanin recovery from *Hibiscus sabdariffa* through microwave-assisted extraction. *Biomass Conversion and Biorefinery*, 12, 351–360. <https://doi.org/10.1007/s13399-020-00606-3>
- Kusuma, H.S. and Mahfud, M. (2015). Box-Behnken design for investigation of microwave-assisted extraction of patchouli oil. *AIP Conference Proceedings*, 1699, 050014. <https://doi.org/10.1063/1.4938350>
- Maksum, A. and Purbowati, I.S.M. (2017). Optimasi ekstraksi senyawa fenolik dari kelopak bunga rosella (*Hibiscus sabdariffa*) berbantu gelombang mikro. *Agrin*, 21(2), 91–104. <https://doi.org/10.20884/1.agrin.2017.21.2.368>
- Mandal, V., Mohan, Y. and Hemalatha, S. (2007). Microwave assisted extraction – an innovative and promising extraction tool for medicinal plant research. *Pharmacognosy Reviews*, 1(1), 7–18.
- Pakaya, D. (2014). Peranan vitamin C pada kulit. *Jurnal Ilmiah Kedokteran*, 1(3), 45–54.
- Peter, E., Mashoto, K.O., Rumisha, S.F., Malebo, H.M., Shija, A. and Oriyo, N. (2014). Iron and ascorbic acid content in *Hibiscus sabdariffa* calyces in Tanzania: modeling and optimization of extraction conditions. *International Journal of Food Science and Nutrition Engineering*, 4(2), 27–35.
- Purbowati, I.S.M., Syamsu, K., Warsiki, E. and Rukmini, H. (2015). Evaluasi toksisitas, aktivitas antibakteri dan antioksidan komponen bioaktif roselle dengan variasi jenis pelarut. *Jurnal Teknologi Industri*, 25(2), 182–189. [In Bahasa Indonesia].
- Purbowati, I.S.M. (2014). Nanoenkapsulasi ekstrak kelopak bunga rosella (*Hibiscus sabdariffa*) hasil optimasi ekstraksi berbantu gelombang mikro sebagai bahan antibakteri dan antioksidan. Indonesia: IPB, Dissertation. [In Bahasa Indonesia].
- Purbowati, I.S.M., Karseno and Maksum, A. (2020). Acidity level control formulation of roselle tea functional drinks based on variations in the addition of flower petals and sugar type. *Food Research*, 4(3), 772–779. [https://doi.org/10.26656/fr.2017.4\(3\).343](https://doi.org/10.26656/fr.2017.4(3).343)
- Purbowati, I.S.M., Syamsu, K., Warsiki, E. and Sri, H. (2016). Stabilitas senyawa fenolik dalam ekstrak dan nanokapsul kelopak bunga rosella pada berbagai variasi pH, suhu. *Agrointek*, 10(1), 31–40. [In Bahasa Indonesia]. <https://doi.org/10.21107/agrointek.v10i1.2023>
- Sheikh, T.Z.B., Yong, C.L. and Lian, M.S. (2009). *In vitro* antioxidant activity of the hexane and methanolic extract of *Sargassum baccularia* and *Cladophora patentiramea*. *Journal of Applied Science*, 9(13), 2490–2493. <https://doi.org/10.3923/jas.2009.2490.2493>
- Siregar, S. (2005). Instalasi Pengolahan Air Limbah. Yogyakarta, Indonesia: Kanisius. [In Bahasa Indonesia].
- Soewarno, T.S. (1990). Dasar-dasar Pengawasan dan Standarisasi Mutu Pangan. Bogor, Indonesia: Pusat Antar Universitas Pangan dan Gizi. [In Bahasa Indonesia].
- Suwetja, I.K. (2007). Biokimia Hasil Perikanan. 3rd ed. Fakultas Perikanan dan Ilmu Kelautan. Manado, Indonesia: Universitas Sam Ratulangi. [In Bahasa Indonesia].
- Taghvaei, M., Jafari, S.M., Assadpoor, E., Nowrouzieh, S. and Alishah, O. (2014). Optimization of microwave-assisted extraction of cottonseed oil and evaluation of its oxidative stability and physicochemical properties. *Food Chemistry*, 160,

- 90–97. <https://doi.org/10.1016/j.foodchem.2014.03.064>
- Wang, Y.L., Xi, G.S., Zheng, Y.C. and Miao, F.S. (2010). Microwave-assisted extraction of flavonoids from Chinese herb *Radix puerariae* (Ge Gen). *Journal of Medicinal Plants Research*, 4(4), 304–308.
- Wei, L., Tao, L. and Keji, T. (2009). Flavonoids from mulberry leaves by microwave-assisted extract and anti-fatigue activity. *African Journal of Agricultural Research*, 4, 898–902.
- Winata, E.W. (2015). Extraction of Anthocyanin Mulberry (*Morus alba* L.) with Ultrasonic Bath (Study of Extraction Time and Solid: liquid Ratio). *Jurnal Pangan dan Agroindustri*, 3, 773–783.
- Wong, P., Yusof, S., Ghazali, H.M. and Che Man, Y.B. (2002). Physicochemical characteristics of roselle (*Hibiscus sabdariffa* L.). *Nutrition and Food Science*, 32(2), 68–73. <https://doi.org/10.1108/00346650210416994>
- Wuryantoro, H. and Susanto, W.H. (2014). Penyusunan standar operating procedures industri rumah tangga pangan pemanis alami instan sari stevia (*Stevia rebaudiana*). *Jurnal Pangan dan Agroindustri*, 2, 76–87. [In Bahasa Indonesia].

Pin Kar Yong

Forest Research Institute
Malaysia (FRIM)
Malaysia

Chim Chay

University of Agriculture
Cambodia

Le Van Viet Man

Ho Chi Minh City University of
Technology
Viet Nam

Oscar Herrera-Calderon

Universidad Nacional Mayor de
San Marcos
Peru

Hasmadi Mamat

University Malaysia Sabah
Malaysia

Regiane Ribeiro dos Santos

Federal Rural University of Rio de
Janeiro
Brazil

Koh Soo Peng

Malaysian Agricultural Research
and Development Institute
(MARDI)
Malaysia

Dinko Hristov Dinko

Trakia University
Bulgaria

Hanifah Nuryani Lioe

Bogor Agricultural University
Indonesia

Michelle Galindo de Oliveira

Federal University of
Pernambuco
Brazil

Alper Sener

Onzekiz Mart University
Turkey

Angelo Maria Giuffrè

Mediterranea University of
Reggio Calabria
Italy

Javier Garcia-Lomillo

University of Burgos
Spain

Laura Marcela Zárate Polanco

National University of Colombia-
Bogotá
Colombia

Swapan K. Banerjee

HPFB Health Canada,
Canada

Adel Abu Bakr Abd El-Hamid**Shatta**

Suez Canal University
Egypt

Iddya Karunsagar**Saroat Rawdkuen**

Mae Fah Luang University
Thailand

Giovanna Giacalone

Università di Torino
Italy

Lorina Acilo Galvez

Visayas State University
Philippines

Sergio Soto Simental

Universidad Autónoma del
Estado de Hidalgo
Mexico

Fernando Cebola Lidon

Universidade Nova de Lisboa
Portugal

Linda Lim Biaw Leng

Universiti Brunei Darussalam
Brunei Darussalam

J.M. Krishanti Jayarukshi Kurr**Premarathne**

Wayamba University of Sri Lanka
Sri Lanka

Katherine Ann T. Castillo-Isra

University of the Philippines L
Banos
Philippines

Steve Flint

Massey University

Giuseppe Sortino
University of Palermo
Italy

Nitte University
India

New Zealand

COPYRIGHT © 2016 - 2020 RYNNYE LYAN RESOURCES