

ISSN: 1412-033X
E-ISSN: 2085-4722

BIODIVERSITAS

Journal of Biological Diversity

Volume 23 - Number 11 - November 2022



BIODIVERSITAS

Journal of Biological Diversity
Volume 23 – Number 11 – November 2022

ISSN/E-ISSN:

1412-033X (printed edition), 2085-4722 (electronic)

EDITORIAL BOARD:

Abdel Fattah N.A. Rabou (Palestine), **Agnieszka B. Najda** (Poland), **Ajay Kumar Gautam** (India), **Alan J. Lymbery** (Australia), **Annisa** (Indonesia), **Bambang H. Saharjo** (Indonesia), **Dalane H. Nunes** (Brazil), **Darlina Md. Naim** (Malaysia), **Ghulam Hassan Dar** (India), **Hassan Pourbabaie** (Iran), **Joko R. Witono** (Indonesia), **Kartika Dewi** (Indonesia), **Katsuhiko Kondo** (Japan), **Kusumadewi Sri Yulita** (Indonesia), **Livia Wanntorp** (Sweden), **M. Jayakara Bhandary** (India), **Mahdi Reyahi-Khoram** (Iran), **Mahendra K. Rai** (India), **Mahesh K. Adhikari** (Nepal), **Maria Panitsa** (Greece), **Mochamad A. Soendjoto** (Indonesia), **Mohamed M.M. Najim** (Srilanka), **Mohib Shah** (Pakistan), **Nurhasanah** (Indonesia), **Praptiwi** (Indonesia), **Rasool B. Tareen** (Pakistan), **Sayed Aliakbar Hedayati** (Iran), **Sayed Mehdi Talebi** (Iran), **Shahabuddin** (Indonesia), **Shahir Shamsir** (Malaysia), **Shri Kant Tripathi** (India), **Subhash C. Santra** (India), **Sugeng Budiharta** (Indonesia), **Sugiyarto** (Indonesia), **Taufiq Purna Nugraha** (Indonesia), **Yosep S. Mau** (Indonesia)

EDITOR-IN-CHIEF:

S u t a r n o

EDITORIAL MEMBERS:

English Editors: **Graham Eagleton** (grahameagleton@gmail.com), **Suranto** (surantouns@gmail.com); Technical Editor: **Solichatun** (solichatun_s@yahoo.com), **Artini Pangastuti** (pangastuti_tutut@yahoo.co.id); Distribution & Marketing: **Rita Rakhmawati** (oktia@yahoo.com); Webmaster: **Ari Pitoyo** (aripitoyo@yahoo.com)

MANAGING EDITORS:

Ahmad Dwi Setyawan (unsjournals@gmail.com)

PUBLISHER:

The Society for Indonesian Biodiversity

CO-PUBLISHER:

Department of Biology, Faculty of Mathematics and Natural Sciences, Sebelas Maret University, Surakarta

ADDRESS:

Jl. Ir. Sutami 36A Surakarta 57126. Tel. +62-271-7994097, Tel. & Fax.: +62-271-663375, email: editors@smujo.id

ONLINE:

biodiversitas.mipa.uns.ac.id; smujo.id/biodiv

.....



Society for Indonesia
Biodiversity



Sebelas Maret University
Surakarta

保護網站安全


使用 GoDaddy 的 SSL 憑證，保護相關客戶資料。

GoDaddy.com

Biodiversitas

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
Indonesia Universities and research institutions in Indonesia Media Ranking in Indonesia	Agricultural and Biological Sciences Animal Science and Zoology Plant Science Biochemistry, Genetics and Molecular Biology Molecular Biology	Biology department, Sebelas Maret University Surakarta Universitas Negeri Sebelas Maret in Scimago Institutions Rankings	19
PUBLICATION TYPE	ISSN	COVERAGE	INFORMATION
Journals	1412033X, 20854722	2014-2021	Homepage How to publish in this journal grahameagleton@gmail.com
SCOPE			
"Biodiversitas, Journal of Biological Diversity" or Biodiversitas encourages submission of manuscripts dealing with all biodiversity aspects of plants, animals and microbes at the level of gene, species, and ecosystem.			



 Join the conversation about this journal







Quartiles

FIND SIMILAR JOURNALS ?

1 Diversity CHE 41% similarity	2 PeerJ USA 39% similarity	3 Biologia (Poland) DEU 37% similarity	4 Tropical Life Research MYS 3 s
--	--	--	--






← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç"
```

SCImago Graphica

Explore, visually communicate and make sense of data with our [new data visualization tool](#).



Metrics based on Scopus® data as of April 2022

H **Hasrianti Hasrianti** 2 months ago

60 articles in each month is a large number. Is it true that this journal meets the standards of scientific publication so that it is worthy of being included in Scopus?


reply



Melanie Ortiz 2 months ago

SCImago Team

Dear Hasrianti,
thank you for your comment.
Our data source is Scopus, SCImago doesn't participate in the journal's selection. Please contact Scopus Support regarding this matter here:
https://service.elsevier.com/app/answers/detail/a_id/14883/kw/scimago/supporthub/scopus/
Best Regards, SCImago Team



W **Widi Sunaryo** 9 months ago

Good Journal, fast response, good quality, good reviewing process, divers in diversity.....

reply

W **Wahyu** 2 months ago

60 article in every month is same with predatory journal



Melanie Ortiz 9 months ago

SCImago Team

Dear Widi, thanks for your participation! Best Regards, SCImago Team

B **BOURAADA KHALID** 1 year ago

I want to submit my manuscript in this journal

reply



Melanie Ortiz 1 year ago

SCImago Team

Dear Bouraada, thank you very much for your comment, we suggest you look for the author's instructions/submission guidelines in the journal's website. Best Regards, SCImago Team

L **Lyes** 2 years ago

I am very pleased with the responsiveness of the journal's editor stuff. Without a doubt, this journal has the potential to become a great journal

reply




Melanie Ortiz 2 years ago

SCImago Team

Dear Lyes, thanks for your participation! Best Regards, SCImago Team

E **Eko** 2 years ago

How long i wait to get email accepted jurnal?



**Melanie Ortiz** 2 years ago

Dear Sir/Madam,
thank you for contacting us.
Unfortunately, we cannot help you with your request, we suggest you visit the journal's homepage or contact the journal's editorial staff , so they could inform you more deeply.
Best Regards, SCImago Team

N **Niken Dharmayanti** 2 years ago

The biodiversity journal has a good response, is published periodically with a frequency every month, is communicative and the response is fast, move on to a higher ranking, thank you

reply

**Melanie Ortiz** 2 years ago

SCImago Team

Dear Niken, thanks for your participation! Best Regards, SCImago Team

Z **zainal abidin** 2 years ago

This journal management is good, fast response, and communicative.

reply

**Melanie Ortiz** 2 years ago

SCImago Team

Dear Zainal, thanks for your participation! Best Regards, SCImago Team


R **Rubiyo** 2 years ago

Thank yuo SCImago Team

reply

K **KETUT SUADA** 3 years ago

Dear Editors
May I know the reason/s of Why my article title "The potential of various indigenous Trichoderma spp. to suppress Plasmodiophora brassicae the pathogen of clubroot disease on cabbage" DOI: 10.13057/biodiv/d180418, in BIODIVERSITAS VOL 18/4 OCT 2017, PAGES:1424-1429, was justified as "SHORT COMMUNICATION", WHILE THE DATA IN THE ARTICLE WAS COMPLETE INCLUDING TO DIVERSITY AND EVEN ITS EFFECT TO THE TRICHODERMA IN PLANT (CABBAGE). CAN YOU



reply**Melanie Ortiz** 3 years ago

SCImago Team

Dear Ketut,
thank you for contacting us.
We are sorry to tell you that SCImago Journal & Country Rank is not a journal. SJR is a portal with scientometric indicators of journals indexed in Elsevier/Scopus.
Unfortunately, we cannot help you with your request, we suggest you contact the journal's editorial staff , so they could inform you more deeply.
Best Regards, SCImago Team

F **Fitra Syawal Harahap** 3 years ago

I want to submit my manuscript in this journal

reply

I**Istiyanto Samidjan** 3 years ago

I want to submit my manuscript in this journal
my regards

Istiyanto Samidjan

**Melanie Ortiz** 3 years ago

SCImago Team

Dear Istiyanto, thank you very much for your comment, we suggest you look for author's instructions/submission guidelines in the journal's website. Best Regards,
SCImago Team

**Melanie Ortiz** 3 years ago


SCImago Team

Dear Fitra, thank you very much for your comment, we suggest you to look for author's instructions/submission guidelines in the journal's website. Best Regards, SCImago Team

J**Joko Prasetyo** 3 years ago

I want to submit my manuscript in this journal

reply





Melanie Ortiz 3 years ago

Dear Joko, thank you very much for your comment, we suggest you to look for author's instructions/submission guidelines in the journal's website. Best Regards, SCImago Team

S **salamiah** 3 years ago

I see the subject area

reply

A **Afik** 3 years ago

Bagus kali jurnal nya

reply



Melanie Ortiz 3 years ago

SCImago Team

Dear user, thanks for your participation! Best Regards, SCImago Team

F **Feron** 3 years ago

Wow bagus banget..samangat berkarya untuk Indonesia yang lebih maju

reply



Melanie Ortiz 3 years ago

SCImago Team

Dear user, thanks for your participation! Best Regards, SCImago Team




muhamad iksan 4 years ago

i want submit my journal thanks

reply

A **ahmad** 4 years ago

Kindly submit your paper here, <https://smujo.id/biodiv/about/submissions>



Leave a comment

Name

Email

(will not be published)

☐ I'm not a robot

reCAPTCHA
[Privacy](#) - [Terms](#)

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:




Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2022. Data Source: Scopus®

EST MODUS IN REBUS

Horatio (Satire 1,1,106)

[Cookie settings](#)[Cookie policy](#)

Isolation and molecular identification of halotolerant diazotrophic bacteria from the Northern Coastal of Pemalang, Central Java, Indonesia

PURWANTO*, EKA OKTAVIANI, NI WAYAN ANIK LEANA

Department of Agrotechnology, Faculty of Agriculture, Universitas Jenderal Soedirman. Jl. Dr. Suparno KP 125 Purwokerto 53122, Central Java, Indonesia. Tel./Fax.: +62-81392645838, *email: purwanto0401@unsoed.ac.id

Manuscript received: 4 October 2022. Revision accepted: 6 November 2022.

Abstract. Purwanto, Oktaviani E, Leana NWA. 2022. Isolation and molecular identification of halotolerant diazotrophic bacteria from the Northern Coastal of Pemalang, Central Java, Indonesia. *Biodiversitas* 23: 5814-5821. Plant Growth Promoting Bacteria can fix nitrogen, a very important macronutrient for plant growth. However, the application of urea as a source of this macronutrient has a negative impact on the environment. Therefore, developing biofertilizers using N-fixing bacteria is an environmentally friendly technology. Therefore, this research aimed to isolate and analyze the diversity of N₂-fixing bacteria from saline rice fields. The soil samples were taken from the rhizosphere of rice plants in Nyamplung Sari Village, Petarukan Sub-district, Pemalang District. Meanwhile, 9 (nine) isolates of nitrogen-fixing bacteria have been isolated, which can fix N₂. The isolates can bind to N₂, but only a few can produce IAA. The nitrogen-fixing ability of diazotrophic bacteria ranged from 17.85 ppm to 29.05 ppm. Isolate Jn3 has the highest ability to fix nitrogen, reaching 29.05 ppm, while Jn3, J, J12, J5, Kn1, and A3 can produce IAA with values of 2.00, 2.69, 2.22, 1.76, 3.47, and 1.79 ppm, respectively. Based on the 16S rRNA analysis and phylogeny construction, the isolated bacteria were identified in 3 (three) clusters. The first cluster was identified as *Pseudomonas stutzeri* and *Acinetobacter junii* (Kn1 and Jn3), while the second was identified as *Bacillus cereus*, *Bacillus tropicus*, *Bacillus altitudinis*, and *Bacillus subtilis* (Jn, Jn1, A3, and K3 isolates). The third cluster was identified as *Bacillus pumilus*, *Acinetobacter baumannii*, and *Acinetobacter schindleri* (J12, J5, and J). In addition, our study reported the findings of *Acinetobacter baumannii* and *Acinetobacter schindleri* species that can fix nitrogen.

Keywords: Bacteria, diazotroph, halotolerant, N₂, saline

INTRODUCTION

National rice production is faced with the problem of increasingly narrowing productive fields. The rate of land conversion every five years shows a relatively high number, reaching 650,000 ha for various purposes such as housing, infrastructure, and industrial area development (Firmansyah et al. 2021). Furthermore, the north coast of Java is a national rice barn but cannot be separated from the shrinking of productive land for other purposes. Zuhri (2018) reported that the rate of land use change on the northern coast of Central Java reaches 0.31% per year. Efforts to maintain national rice production can be made by expanding the area. Since fertility is very low, the alternative is to use marginal land with the potential for rice cultivation. Land with saline is one potential for developing new paddy fields. The total area of saline land in Indonesia reaches 440,300 ha, with the criteria of slightly and totally saline reaching 304,000 ha and 140,300 ha (Karolinoerita and Yusuf 2020).

The North Coast of Pemalang is another of Central Java's rice production centers. However, its location causes seawater intrusion into the rice fields, causing saline conditions in the paddy fields. Salinity occurs when the concentration of soluble salts is high in the soil, and the electrical conductivity is above 4 dS m⁻¹ or 40 mM NaCl (Abbas et al. 2019; Karolinoerita and Yusuf 2020).

According to Karolinoerita and Yusuf (2020), using irrigation water with high salt content can cause accumulation in the root areas of plants and negatively influence growth. The negative effect of salinity stress affects plants, especially in decreasing the rate of photosynthesis and accumulation of plant biomass (Yan et al. 2013). Suryaman et al. (2021) reported that salinity decreases photosynthetic activity, both stomata and non-stomata factors, and decreases chlorophyll content, leaf area, plant height, and biomass of mung bean plants. Soils with high salt content cause ion toxicity, nutritional deficiencies, oxidative and osmotic stress, limiting water uptake, disrupting nutrient balance, and affecting plant nutrient absorption (Abbas et al. 2019). The high salt content in the form of Na, Ca, and Mg as chloride or sulfate causes plants to be deficient in nutrients N, P, Mn, Fe, and osmotic stress (Shrivastava and Kumar 2015; Subowo 2015; Sulasih and Widawati 2016).

The biological technology approach can be carried out using tolerant varieties and beneficial microbes or Plant Growth Promotion Rhizobacteria (PGPR). Furthermore, the utilization can stimulate plant growth and increase the availability of nutrients. PGPR interacts with plants through the colonization of root areas and stimulates growth through direct and indirect effects (Shrivastava and Kumar 2015). The direct and indirect effects improve plant growth and increase tolerance to saline stress (Abbas et al.

2019). Agustiyani et al. (2021) reported the isolation of N-fixing bacteria from various ecosystems with the genus *Sinomonas* strain and *Arthrobacter*. Furthermore, Sultana et al. (2021) stated that the bacteria *A. denitrificans*, *B. aryabhatai*, and *Pseudomonas* spp. were tolerant to salinity and could produce siderophores.

It is imperative to study the diversity of nitrogen-fixing bacteria in saline paddy fields on the Northern Coast of Pemalang. Using saline-tolerant nitrogen-fixing bacteria is significant for developing biofertilizers to support plant growth and yield in saline soils. Tirry et al. (2021) reported applying saline-tolerant bacteria such as *Pseudomonas putida*, *Alcaligenes* sp., *Klebsiella* sp., and *Pseudomonas cedrina* could reduce the negative impact of stress caused by NaCl, increasing chlorophyll content compared to uninoculated plants. Therefore, this study aims to isolate and analyze the diversity of N₂-fixing bacteria from saline rice fields.

This research is essential in providing information about nitrogen-fixing bacteria in coastal ecosystems. The nitrogen-fixing bacteria found can be an important source of inoculants in supporting the plants' growth and development in coastal areas, both for food and conservation needs. Based on the previous research, finding bacteria that can fix free nitrogen in the Northern Coastal Area, Pemalang District, Central Java Province, Indonesia, was possible.

MATERIALS AND METHODS

Sampling area

Soil samples were taken from Nyamplung Sari Village, Petarukan Sub-district, and Pemalang District, Central Java Province, Indonesia rice fields about 500 meters from the north coast of the Java Sea. Furthermore, the sampling location has a pH and EC above 7 and 4 mS/cm. The samples were obtained from the roots of rice plants in the maximum vegetative phase. The soil from the rice roots was excavated and taken with the plant roots, put into a plastic bag, placed in a cooler box at a temperature of 4°C, and brought to the laboratory for further isolation and characterization.

Isolation andp

The sum of 10 g of soil sample was mixed with 90 mL of sterile distilled water, and graded dilution was conducted by taking 1 mL mixed with 9 mL of sterile distilled water to obtain a dilution of 10⁻². The graded dilutions were carried out up to 10⁻⁶, and a total of 0.1 mL of the last two dilutions were inoculated on Nfb medium (the composition of Nfb medium: malic acid 5.0 g/L; K₂HPO₄ 0.5 g/L; MgSO₄·7H₂O 0.2 g/L; NaCl 0.1 g/L; CaCl₂·2H₂O 0.02 g/L; KOH 4.5 g/L; NH₄Cl 1 g/L; FeEDTA 4 mL (solution 16.4 g/L); micronutrient solution 2 mL (CuSO₄·5H₂O 0.04 g/L; ZnSO₄·7H₂O 0.12 g/L; H₃BO₃ 1.40 g/L; Na₂MoO₄·2H₂O 1.0 g/L; MnSO₄·H₂O 1.175 g/L); vitamin solution 1 mL (biotin 100 mg/L; pyridoxal-HCl 200 mg/L); pH to 6.5, without the addition of bromothymol blue). Each medium was incubated at room temperature for 2-4 x 24

hours. Meanwhile, the isolates that grew and differed were purified and stored in a slanted medium.

IAA production capability test

The ability to produce IAA was tested using the Salkowski reagent (Agustiyani et al. 2021). About fifty milliliters of NB medium (for 1L, beef extract 0.3g, peptone 0.5g, sodium chloride 0.5g, and distilled water 1L) containing 0.1% DL-Tryptophan was inoculated with 500 mL of bacterial culture aged 24 hours with a population density above 10⁷. Subsequently, the bacterial culture was shaken at 180 rpm for 48 hours in the dark and centrifuged at 10,000 rpm for ten minutes. About one milliliter of supernatant was added to the test tube with 4 mL of Salkowski reagent and incubated for 30 minutes until a pink color appeared. The absorbance level was measured using a spectrophotometer at 535 nm, and IAA production was calculated by a standard curve regression equation expressed in ppm.

N₂ fixation assay

N₂ fixing ability was tested on 9 isolates using the Kjeldahl method (Oedjijono et al. 2014). Each isolate was grown in a semisolid Nfb medium and incubated for 20 days at room temperature. The bacterial culture was put into a Kjeldahl flask containing a mixture of salts in a ratio of 40:2.5:1.5 of K₂SO₄, CuSO₄, and metallic selenium, and 3 mL of concentrated sulfuric acid was added. The flask was heated in a fume hood in the Digester at 420°C for 20 minutes and was cooled by adding distilled water to reach a final volume of 50 mL. A total of 20 mL of the digestion sample was poured into a distillation tube and inserted into the apparatus. A 250 mL Erlenmeyer flask containing 20 mL of 4% boric acid and 6 drops of Conway reagent (1,000 mg methyl red, 150 mg bromocresol green, 200 mL 96% ethanol) was placed below the condenser of the distillation apparatus, and the outlet of the condenser was under the solution. Distillation was performed using a semi-automatic distillation unit. Meanwhile, 30 mL of 40% NaOH and 100 mL of distilled water were automatically poured through the apparatus in the distiller. The solution containing NH₃, boric acid, and mixed indicator was titrated with 0.05 N HCl using an auto titrator. The following formula calculated the N₂ value fixed by each isolate:

$$N_2 \text{ (ppm)} = \frac{\text{sample titer} - \text{blank titer} \times \text{normality of HCl} \times 14 \times 1000000}{\text{sample weight (g)} \times 1000}$$

Pathogenicity test

A pathogenicity test was performed on tobacco plants (Zuraidah 2013), and bacterial isolates were propagated using NB media. They were incubated for 24 hours in a shaker at 120 rpm until the population reached above 10⁷ cfu/mL. Furthermore, 1 mL of inoculum was injected using a sterile syringe without a needle on the back of a healthy tobacco leaf. Sterile distilled water was used as a negative control, and tobacco leaves were labeled according to the injected isolate. Plant response was observed within 24-48 hours to observe the level of necrosis on tobacco leaves.

Salinity tolerance test

A tolerance test for salinity was carried out using NB media with varying concentrations of NaCl (Shultana et al. 2020) at 3, 5, and 7%. Furthermore, 20 mL of bacterial culture was inoculated on NB media with NaCl concentrations of 3%, 5%, and 7% and incubated for 24 hours in a shaker at 120 rpm. Finally, the population was measured using a 10-fold dilution and total plate count (TPC) technique.

Molecular identification

Isolation of bacterial genomic DNA followed the protocol of the TIANamp Bacteria kit (Tiangen). First, the results were visualized with 1% agarose electrophoresis gel, followed by visualization under UV light. Meanwhile, absorbance measurements were performed at a wavelength of 260nm with a UV-Vis (NanoDrop) spectrophotometer to determine the concentration of genomic DNA.

DNA concentration and purity were determined according to the protocol (Green and Sambrook 2012) using an Ultra Violet (UV) spectrophotometer at 260 nm. The amount of radiation absorbed is directly proportional to the concentration of DNA, i.e., $A = 1$ is proportional to 50 μ g of double-stranded DNA per mL. Therefore, a comparison of absorbance at wavelengths of 260 and 280 nm indicates the purity of DNA. The 16S rDNA sequence amplification reaction was conducted using universal primers 27F (AGAGTTTGATCCTGGCTCAG) and 1429R (GGTACCTTGTACGACTT). The amplification reaction was carried out with the following reaction composition in Table 1.

The 16S sequence amplification reaction followed the pre-denaturation protocol at 95°C for 4 minutes, then by 35 repeated cycles consisting of denaturation at 95°C for 40 seconds, annealing at 54°C for 30 seconds, and extension (elongation) at 72°C for 1 minute, followed by the final extension stage at 72°C for 10 minutes. Furthermore, the amplification results were visualized by agarose gel electrophoresis with a concentration of 2% and observed with a UV transilluminator (UV light). The 16S bands are located at the base length of about 1,500 bp, and the PCR products were sent to 1st Base (Singapore) for the rRNA sequencing determination. The sequence was determined using a two-way reaction with DNA primers forward 27F and reverse 1429R. Furthermore, 16S rRNA sequences from bidirectional sequencing reactions (with forward and

reverse primers) were assembled using BioEdit 7.2 software (Hall 1999). Bacterial species were identified based on nucleotide sequences using BLASTn (Basic Local Alignment Search Tool) analysis at <https://blast.ncbi.nlm.nih.gov/Blast.cgi>.

Phylogenetic tree construction

The obtained sequences were analyzed using Molecular Evolutionary Genetic Analysis-11 (MEGA-11) software downloaded at <http://www.megasoftware.net> (Felsenstein 1985; Tamura et al. 2021). The assembly of 16S rRNA sequences was carried out using BioEdit software (Hall 1999). Furthermore, the nucleotide sequence of each isolate was aligned (multiple sequence alignment was performed) with the MUSCLE program on MEGA 11 software. The phylogenetic tree construction was performed using the Maximum-Likelihood statistical method and was tested with 1,000 times bootstrap replication. The nucleotide substitution model follows the Tamura and Nei (1993) model, and this analysis involved 9 isolated bacterial nucleotide sequences downloaded from the National Center for Biotechnology Information (NCBI) website, as well as 1 (one) outgroup sequence from the 18S rRNA genetic material of maize (*Zea mays*). There were 1441 positions for the 16S rRNA gene used as the final dataset.

RESULTS AND DISCUSSION

Nitrogen fixation capacity and IAA production

The exploration of diazotrophic bacteria from saline fields on the north coast of Pemalang obtained 9 isolates with the highest nitrogen-fixing ability. The nitrogen-fixing ability ranged from 17.85 ppm to 29.05 ppm (Table 2), and Jn3 has the highest ability to fix nitrogen, reaching 29.05 ppm. The results showed that all isolates from saline rice fields could fix nitrogen, and the fixation ability varies greatly due to environmental conditions.

Diazotrophic bacteria were isolated from the rhizosphere of rice plants, and the plant root environment was very dominant by diazotrophs. Purwanto et al. (2017) reported that the presence of nitrogen-fixing bacteria is more dominant in the plant rhizosphere of the total microflora and a flooded environment strongly supports the fixation activity.

Table 1. The reaction composition used for the 16S rRNA sequence amplification reaction

Composition	Volume	Description
DNA template	>50 ng	The volume is adjusted according to the concentration of each genomic DNA isolate
Primer F (IDT)	1 μ L	Concentration 10 μ M
Primer R (IDT)	1 μ L	Concentration 10 μ M
MasterMix PCR (Thermo Scientific)	12.5 μ L	Half of the total reaction
Nuclease-free water (Thermo Scientific)	Adjusted	Added up to a total volume of 25 μ L
Total Reaction	25 μ L	

Nitrogen-fixing bacteria play a vital role in maintaining nitrogen availability in the soil. The contribution of the bacteria to nitrogen availability varies greatly depending on several factors, including environmental variability, management, cropping patterns, and genotype differences (Aasfar et al. 2021). The variations reflect the diversity of microorganisms resistant or tolerant to saline environmental conditions. Saline conditions are negatively correlated with the abundance and diversity of diazotrophic bacteria (Barua et al. 2011).

Limited isolates of diazotrophic bacteria can produce phytohormones, especially indole acetic acid (IAA). Table 2 showed that the ability to produce IAA was only on isolates Jn3, J, J12, J5, Kn1, and A3 at 2.00, 2.69, 2.22, 1.76, 3.47, and 1.79 ppm, respectively. The highest IAA production capability by isolate Kn1 was 3.47 ppm, and diazotrophic bacteria that can produce this compound are potential Plant Growth Promotion Rhizobacteria. Oedjiono et al. (2014) stated that the ability of rhizobacteria to produce phytohormones is one of the principal mechanisms of candidates for PGPR. Diazotrophic bacteria capable of producing IAA can convert L-tryptophan as the primary precursor in synthesizing the compound. Goswami et al. (2014) noted that IAA is a product of L-tryptophan in the metabolism of microorganisms; hence some PGPRs can produce the compound depending on the availability of L-tryptophan in the media. The level of L-tryptophan in the medium was 0.1%, but some isolates could not synthesize IAA, especially Jn1, Jn, and K3. The ability to produce IAA by microorganisms varies widely between species. Meanwhile, culture conditions and substrate strongly influence the inability to produce IAA (Yousef 2018). Jabborova et al. (2022) reported that *B. endophyticus* IGPEB 33 could produce PGPR by producing proteases, amylase, catalase, and IAA.

Salinity tolerance

The results showed that all diazotrophs could live in media with high salt concentrations of up to 7%. Almost all

isolates experienced a population decline at this concentration, but Jn1, J12, Kn1, and A3 remain high, reaching 1.13×10^8 , 7.10×10^8 , 6.00×10^8 , and 4.30×10^8 cfu/mL (Figure 1). The lowest population was at 7% salt concentration, namely, isolate Jn1 at 3.80×10^5 cfu/mL.

The high salt content decreased the diazotrophic bacteria population in all isolates. Salinity is one of the stress factors that affect the decrease in bacterial diversity and controls the abundance, composition, and function due to dissolved Na^+ ions greater than 0.15 M. In addition, it causes hyperosmotic conditions and diffusion of water out of the microbial cells (Moradi et al. 2011). Microorganisms adapt to saline conditions by maintaining internal salt levels at the same environmental conditions, or microorganisms actively secrete NaCl and produce compatible organic osmoregulatory such as betaine (Moradi et al. 2011). Based on the experimental results, 9 bacterial isolates can live in high salt concentrations and be categorized as halotolerant. According to (Moradi et al. 2011), microorganisms, based on their optimal growth, can be categorized into non, slight, moderate, and extreme halophiles, which grow best on media with a salt content below 0.2 M (1%), 0.2-0.5 M (1-3%), 0.5-2.5 M (3-15%), and 2.5-5.2 M (15-32%), respectively.

Table 2. Nitrogen fixation capacity and IAA production of diazotrophic bacteria from The Northern Coastal of Pemalang

Isolate	Fixation of N_2 (ppm)	IAA (ppm)
Jn3	29.05	2.00
Jn1	23.10	-
J	22.75	2.69
J12	20.30	2.22
J5	19.95	1.76
Kn1	18.90	3.47
A3	18.55	1.79
Jn	17.85	-
K3	17.85	-




Figure 1. The population of diazotrophic bacteria in different salt concentration (data was transformed in Log (x))

Pathogenicity test

Diazotrophic bacteria pathogenicity test was conducted on tobacco plant leaves and observed 48 hours after injection. The experimental results reported that all isolates did not show necrosis symptoms (Figure 2). Therefore, the isolates of diazotrophic bacteria are not pathogenic to tobacco plants and do not cause collapse or tissue death symptoms.

Analysis of 16S rRNA

Visualization of the 16S rRNA sequence from 9 (nine) bacteria isolates can be seen in Figure 3 with PCR reaction products along ± 1500 bp. The results of species identification by BLASTn analysis are shown in Table 3. The phylogenetic tree was generated from the analysis using alignment (MUSCLE) and the Maximum Likelihood statistical method (Figure 4).

The bacteria identification analysis of 16S rRNA sequences showed that 5 isolates were categorized into the genus *Bacillus*, 3 from the genus *Acinetobacter*, and 1 from *Pseudomonas*. Research on the isolation and identification of nitrogen-fixing bacteria from the rhizosphere of plant roots has been carried out in the root rhizosphere of rice and mangrove plants on saline land in coastal areas.

The isolated bacterial species clustering based on the phylogenetic tree consisted of 3 clusters. The first cluster consisted of Kn1 and Jn3 isolates, identified as

Pseudomonas stutzeri and *Acinetobacter junii*. The second was Jn, Jn1, A3, and K3 isolates, identified as *Bacillus cereus*, *Bacillus tropicus*, *Bacillus altitudinis*, and *Bacillus subtilis*, respectively. Meanwhile, the third cluster, separated from the reference nucleotide sequences, isolates J12, J5, and J, identified as *Bacillus pumilus*, *Acinetobacter baumannii*, and *Acinetobacter schindleri*, respectively.

The bacterium *Bacillus pumilus* dominated coastal marine habitats and had enzyme activity of amylase, protease, lipase, oxidase, and catalase, showing a positive response to the Voges-Proskauer test (Parvathi et al. 2009). In addition, Faizah et al. (2017) reported the ability to inhibit the growth of the plant pathogen *Xanthomonas campestris* by producing an inhibition zone of 1.84 mm in diameter. However, there was no report of each species isolated among the various *Bacillus* species.

The results indicate that in cluster 1, isolate Kn1 was identified as *Pseudomonas stutzeri* isolated from various ecosystems. That is reinforced by the statement of Lalucat et al. (2006) the bacterium is gram-negative with a wide ecosystem. Yan et al. (2008) reported that the *Pseudomonas stutzeri* colonizes the roots of rice and wheat plants. Based on genomic analysis, it has several genes regulating osmotolerance. Furthermore, Li et al. (2017) reported that the bacterial species were isolated from the rhizosphere of sugarcane plants.




Figure 2. Symptoms of Diazotrophic Bacteria injection on tobacco leaves. Note: 1 = J5; 2 = J12; 3 = Jn3; 4 = J; 5 = K3; 6 = Kn1; 7 = A3; 8 = Jn; 9 = Jn1




Figure 3. Visualization of the 16S rRNA sequence (1 = J5; 2 = J12; 3 = Jn3; 4 = J; 5 = K3; 6 = Kn1; 7 = A3; 8 = Jn; 9 = Jn1; M = 1Kbp DNA Ladder)




Figure 4. The original tree generated from the MEGA11 software

Table 3. Identification of isolated bacterial species based on 16S rRNA sequences

No. of well	Code of isolates	Description	Scientific name	Percentage identification	Accession
1	J5	<i>Acinetobacter baumannii</i> strain MS14413 chromosome, complete genome	<i>Acinetobacter baumannii</i>	99.57%	CP054302.1
2	J12	<i>Bacillus pumilus</i> strain MGB1027 16S ribosomal RNA gene, partial sequence	<i>Bacillus pumilus</i>	98.69%	MH260924.1
3	Jn3	<i>Acinetobacter junii</i> strain WR4 16S ribosomal RNA gene, partial sequence	<i>Acinetobacter junii</i>	98.67%	KT886456.1
4	J	<i>Acinetobacter schindleri</i> strain HZE30-1 chromosome, complete genome	<i>Acinetobacter schindleri</i>	98.57%	CP044483.1
5	K3	<i>Bacillus subtilis</i> strain WR10-qm 16S ribosomal RNA gene, partial sequence	<i>Bacillus subtilis</i>	100%	OM320447.1
6	Kn1	<i>Pseudomonas stutzeri</i> strain SGAir0442 chromosome, complete genome	<i>Pseudomonas stutzeri</i>	100%	CP025149.2
7	A3	<i>Bacillus altitudinis</i> strain GUCC 1037 16S ribosomal RNA gene, partial sequence	<i>Bacillus altitudinis</i>	100%	OM319557.1
8	Jn	<i>Bacillus cereus</i> group sp. strain HD1.3 16S ribosomal RNA gene, partial sequence	<i>Bacillus cereus</i>	100%	OM368635.1
9	Jn1	<i>Bacillus tropicus</i> strain CU96 16S ribosomal RNA gene, partial sequence	<i>Bacillus tropicus</i>	100%	OM108144.1

The first cluster also identified isolate Jn3 as *Acinetobacter junii* bacteria. In the third cluster, isolates J5 and J were identified as *Acinetobacter baumannii* and *Acinetobacter schindleri*. The results of previous studies reported by Sachdev et al. (2010) have successfully isolated the bacterium *Acinetobacter* sp. from wheat roots. The most dominant strains were *Ac. calcoaceticus*, *Ac. baumannii*, *Ac. lwoffii*, *Ac. baylyi* and *Ac. calcoaceticus*. Rokhbakhsh-Zamin et al. (2011) successfully conducted isolation from the rhizosphere of *Pennisetum glaucum*. The genus *Acinetobacter* sp. as PGPR has been reported by Padmavathi et al. (2016), where strains of *Acinetobacter junii* have been identified from tomato plants and dual inoculation with mycorrhizae.

Bacillus sp. in this study is based on the phylogenetic tree. The second cluster consists of isolates J12, K3, A3, Jn, and Jn1, identified as *Bacillus pumilus* strain MGB1027, *Bacillus subtilis* strain WR10-qm, *Bacillus altitudinis* strain GUCC1037, *Bacillus cereus* strain HD1.3, and *Bacillus tropicus* strain CU96. *Bacillus* sp. is adaptive to marginal environmental conditions and is dominant in various environments, up to 95% of the population of gram-negative bacteria (Miljakovic et al. 2020). According to Kumar et al. (2014), it has a higher tolerance to abiotic stress than *Pseudomonas* sp. That is due to the ability to form endospore, a dormant resistant form that can survive in unfavorable environmental conditions. Ibarra-Villarreal et al. (2021) explained that *Bacillus* sp. can survive in abiotic stress through structural adaptation and has metabolite characteristics such as the production of multi-layered cell wall structures, antibiotic secretion, signal peptide molecules, extracellular enzymes, and formation of cell walls. Meanwhile, several strains of *Bacillus* sp. that can fix nitrogen have been previously reported. Antunes et al. (2017) successfully isolated bacteria from the genus *Bacillus* sp. originating from the rhizosphere of sugarcane plants with a direct and positive influence on plants through their ability to fix nitrogen, solubilize P, and produce various types of hormones. Kumar et al. (2021) successfully isolated the *Bacillus pumilus* strain from saline soil with attributes such as PGPR to produce IAA, PSB, ACC deaminase, ammonia, HCN, siderophore, and EPS production. Ibarra-Villarreal et al. (2021) obtained 10 saline-tolerant *Bacillus* sp. strains from wheat plants' rhizosphere, namely, *B. subtilis*, *B. megaterium*, *B. cereus*, *B. amyloliquefaciens*, *B. pumilus*, *B. licheniformis*, *B. endophyticus*, *B. atrophaeus*, *B. safensis*, and *B. mojavensis*. The highest abundance was dominated by *B. subtilis* and *B. megaterium*.

In conclusion, nine isolates of nitrogen-fixing bacteria have been isolated, which can fix N₂. The entire isolates could fix N₂, but few could produce IAA. The nitrogen-fixing ability of diazotrophic bacteria ranged from 17.85 ppm to 29.05 ppm. Furthermore, isolate Jn3 has the highest ability to fix nitrogen, reaching 29.05 ppm. The ability to produce IAA was only on Jn3, J, J12, J5, Kn1, and A3, with values of 2.00, 2.69, 2.22, 1.76, 3.47, and 1.79 ppm, respectively. Based on the 16S rRNA analysis and phylogeny construction, the isolated bacteria were identified in 3 clusters. The first cluster was identified as

Pseudomonas stutzeri and *Acinetobacter junii* (Kn1 and Jn3), while the second was identified as *Bacillus cereus*, *Bacillus tropicus*, *Bacillus altitudinis*, and *Bacillus subtilis* (Jn, Jn1, A3, and K3 isolates). Furthermore, the third cluster was identified as *Bacillus pumilus*, *Acinetobacter baumannii*, and *Acinetobacter schindleri* (J12, J5, and J).

ACKNOWLEDGEMENTS

The author is grateful to LPPM Universitas Jenderal Soedirman (UNSOED), Indonesia, which has provided research funding through the UNSOED basic research scheme in 2021. In addition, the author thanks Dwi Ayu Lutfiana, Retna Susanti, and Fenti Chakimatul Isnaeni, who helped conduct the research.

REFERENCES

- Aasfar A, Bargaz A, Yaakoubi K, Hilali A. 2021. Nitrogen fixing Azotobacter species as potential soil biological enhancers for crop nutrition and yield stability. *Front Microbiol* 12: 628379. DOI: 10.3389/fmicb.2021.628379.
- Abbas R, Rasul S, Aslam K, Baber M, Shahid M, Mubeen F, Naqqash T. 2019. Halotolerant PGPR: A hope for cultivation of saline soils. *J King Saud Univ Sci* 31 (4): 1195-1201. DOI: 10.1016/j.jksus.2019.02.019.
- Agustiyani DWI, Dewi TK, Laili NUR, Nditasari A, Antonius S. 2021. Exploring biofertilizer potential of plant growth-promoting rhizobacteria candidates from different plant ecosystems. *Biodiversitas* 22 (5): 2691-2698. DOI: 10.13057/biodiv/d220529.
- Antunes JEL, Lyra MCCP, Ollero FJ, Freitas ADS, Oliveira LMS, Araújo ASF, Figueiredo MVB. 2017. Diversity of plant growth-promoting bacteria associated with sugarcane. *Genet Mol Res* 16 (2): gmr16029662. DOI: 10.4238/gmr16029662.
- Barua S, Tripathi S, Chakraborty A. 2011. Studies on non-symbiotic diazotrophic bacterial populations of coastal arable saline soils of India. *Indian J Microbiol* 51 (3): 369-376. DOI: 10.1007/s12088-011-0082-9.
- Faizah LN, Budiharjo A, Kusdiyantini E. 2017. Optimasi pertumbuhan dan potensi antagonistik *Bacillus pumilus* terhadap patogen *Xanthomonas campestris* serta identifikasi molekuler gen penyandi PKS dan NRPS. *Jurnal Akademika Biologi* 6 (1): 38-48. [Indonesian]
- Felsenstein J. 1985. Confidence Limits on Phylogenies: An Approach Using the Bootstrap. *Author (s): Joseph Felsenstein Stable. Evolution* 39 (4): 783-791. DOI: 10.2307/2408678.
- Firmansyah F, Yusuf M, Argarini O, Perencanaan D, Sipil FT, Kebumian P. 2021. Strategi pengendalian alih fungsi lahan sawah di Provinsi Jawa Timur. *Jurnal Penataan Ruang* 16 (1): 47-53. DOI: 10.12962/j2716179X.v16i1.8726. [Indonesian]
- Goswami D, Pithwa S, Dhandhukia P, Janki N. 2014. Delineating *Kocuria turfanensis* 2M4 as a credible PGPR: a novel IAA-producing bacteria isolated from saline desert. *J. Plant Interact* 9 (1): 566-576. DOI: 10.1080/17429145.2013.871650.
- Green MR, Sambrook J. 2012. *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, New York.
- Hall T. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows. *Nucleic Acid Symp* 41: 95-98. DOI: 10.14601/Phytopathol_Mediterr-14998u1.29.
- Ibarra-Villarreal AL, Gandara-Ledezma A, Godoy-Flores AD, Herrera-Septulveda A, Alondra Diaz-Rodriguez M, Fannie Isela Parra-Cota, FI, Santos-Villalobos SD. 2021. Salt-tolerant *Bacillus* species as a promising strategy to mitigate the salinity stress in wheat (*Triticum turgidum* subsp. durum). *J Arid Environ* 186: 104399. DOI: 10.1016/j.jaridenv.2020.104399.
- Jaborova D, Davranov K, Jabbarov Z, Bhowmik SN, Ercisli S, Danish S, Singh S, Desouky SE, Elazazy AM, Nasif O, Datta R. 2022. Dual inoculation of plant growth-promoting *Bacillus endophyticus* and *Fumelliformis mosseae* improves plant growth and soil properties in

- ginger. ACS Omega 7: 34779-34788. DOI: 10.1021/acscomega.2c02353.
- Karolinoerita V, Yusuf WA. 2020. Salinisasi lahan dan permasalahannya di Indonesia. Jurnal Sumberdaya Lahan 4 (2): 91-99. DOI: 10.21082/jsdl.v14n2.2020.91-99. [Indonesian]
- Kumar A, Kumar A, Pratush A. 2014. Molecular diversity and functional variability of environmental isolates of *Bacillus* species. SpringerPlus 3: 312. DOI: 10.1186/2193-1801-3-312.
- Kumar A, Maurya BR, Raghuwanshi R. 2021. The microbial consortium of indigenous rhizobacteria improving plant health, yield and nutrient content in wheat (*Triticum aestivum*). J Plant Nutr 44 (13): 1942-1956. DOI: 10.1080/01904167.2021.1884706.
- Lalucat J, Bennasar A, Bosch R, Garc   E. 2006. Biology of *Pseudomonas* stutzeri. Microbiol Mol Biol Rev 70 (2): 510-547. DOI: 10.1128/MMBR.00047-05.
- Li H, Singh RK, Singh P, Song Q, Xing Y. 2017. Genetic diversity of nitrogen-fixing and plant growth promoting *Pseudomonas* species isolated from sugarcane rhizosphere. Front Microbiol 8: 1-20. DOI: 10.3389/fmicb.2017.01268.
- Miljakovic D, Marinkovi   J, Bale  evic-Tubi  c S. 2020. The Significance of *Bacillus* spp. in disease suppression and growth promotion of field and vegetable crops. Microorganisms 8: 1037. DOI: 10.3390/microorganisms8071037.
- Moradi A, Tahmourespour A, Hoodaji M, Khorsandi F. 2011. Effect of salinity on free living-diazotroph and total bacterial populations of two saline soils. Afr J Microbiol Res 5 (2): 144-148. DOI: 10.5897/AJMR10.838.
- Oedjiono Soetarto ES, Moeljopawiro S, Djatmiko HA. 2014. Promising plant growth promoting rhizobacteria of *Azospirillum* spp. isolated from iron sand soils, Purworejo Coast, Central Java, Indonesia. Adv Appl Sci Res 5 (3): 302-308.
- Padmavathi T, Dikshit R, Seshagiri S. 2016. Effect of *Rhizophagus* spp. and plant growth-promoting *Acinetobacter junii* on *Solanum lycopersicum* and *Capsicum annum*. Braz J Bot 38: 273-280. DOI: 10.1007/s40415-015-0144-z.
- Parvathi A, Krishna K, Jose J, Joseph N, Nair S. 2009. Biochemical and molecular characterization of *Bacillus pumilus* isolated from coastal environment in Cochin, India. Braz J Microbiol 40: 269-275. DOI: 10.1590/S1517-838220090002000012.
- Purwanto, Yuwariah Y, Sumadi, Simarmat T. 2017. Nitrogenase activity and IAA production of indigenous diazotroph and its effect on rice seedling growth. Agrivita J Agric Sci 39 (81): 31-37. DOI: 10.17503/AGRIVITA.V39I1.653.
- Rokhbakhsh-Zamin F, Sachdev D, Kazemi-pour N, Engineer A, Pardesi KR, Zinjarde S, Dhakephalkar PK, Chopade BA. 2011. Characterization of plant-growth-promoting traits of *Acinetobacter* species isolated from rhizosphere of *Pennisetum glaucum*. J Microbiol Biotechnol 21 (6): 556-566. DOI: 10.4014/jmb.1012.12006.
- Sachdev D, Nema P, Dhakephalkar P. 2010. Assessment of 16S rRNA gene-based phylogenetic diversity and promising plant growth-promoting traits of *Acinetobacter* community from the rhizosphere of wheat. Microbiol Res 165 (8): 627-638. DOI: 10.1016/j.micres.2009.12.002.
- Shrivastava P, Kumar R. 2015. Soil salinity: A serious environmental issue and plant growth promoting bacteria as one of the tools for its alleviation. Saudi J Biol Sci 22 (2): 123-131. DOI: 10.1016/j.sjbs.2014.12.001.
- Shultana R, Tan A, Zuan K, Yusop MR, Saud H. M. 2020. Characterization of salt-tolerant plant growth-promoting rhizobacteria and the effect on growth and yield of saline-affected rice. Plos One 15 (9): e0238537. DOI: 10.1371/journal.pone.0238537.
- Subowo YB. 2015. Penambahan pupuk hayati jamur sebagai pendukung pertumbuhan tanaman padi (*Oryza sativa*) pada tanah salin. Pros Sem Nas Masy Biodiv Indon 1 (1): 150-154. DOI: 10.13057/psnmbi/m010126. [Indonesian]
- Sulasih, Widawati S. 2016. Pengaruh salinitas dan inokulan bakteri terhadap pertumbuhan tanaman terung (*Solanum melongena* L.). Berita Biologi 15 (1): 17-25. DOI: 10.14203/BERITABIOLOGI.V15I1.2854.
- Sultana S, Alam S, Manjurul M. 2021. Screening of siderophore-producing salt-tolerant rhizobacteria suitable for supporting plant growth in saline soils with iron limitation. J Agric Food Res 4 (April): 100150. DOI: 10.1016/j.jafr.2021.100150.
- Suryaman M, Sunarya Y, Istarimila I, Fudholi A. 2021. Biocatalysis and agricultural biotechnology effect of salinity stress on the growth and yield of mungbean (*Vigna radiata* (L.) R Wilczek) treated with mangosteen pericarp extract. Biocatalysis Agric Biotechnol 36: 102132. DOI: 10.1016/j.bcab.2021.102132.
- Tamura K, Nei M. 1993. Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. Mol Biol Evol 10 (3): 512-526. DOI: 10.1093/oxfordjournals.molbev.a040023.
- Tamura K, Stecher G, Kumar S. 2021. MEGA11: Molecular Evolutionary Genetics Analysis Version 11. Mol Biol Evol 38 (7): 3022-3027. DOI: 10.1093/molbev/msab120.
- Tirry N, Kouchou A, Laghmari G, Lemjereb M, Hnadi H, Amrani K, Bahafid W, Ghachtouli NEI. 2021. Biocatalysis and agricultural biotechnology improved salinity tolerance of *Medicago sativa* and soil enzyme activities by PGPR. Biocatal Agric Biotechnol 31: 101914. DOI: 10.1016/j.bcab.2021.101914.
- Yan K, Shao H, Shao C, Chen X. 2013. Physiological adaptive mechanisms of plants grown in saline soil and implications for sustainable saline agriculture in coastal zone. Acta Physiol Plant 35 (10): 2867-2878. DOI: 10.1007/s11738-013-1325-7.
- Yan Y, Yang J, Dou Y, Chen M, Ping S, Peng J, Lu W, Zhang W, Yao Z, Li H, Liu W, He S, Geng L, Zhang X, Yang F, Yu H, Zhan Y, Li D, Lin Z, Jin Q. 2008. Nitrogen fixation island and rhizosphere competence traits in the genome of root-associated *Pseudomonas* stutzeri A1501. PNAS 105 (21): 7564-7569. DOI: 10.1073/pnas.0801093105.
- Yousef NMH. 2018. Capability of Plant Growth-Promoting Rhizobacteria (PGPR) for producing indole acetic acid (IAA) under extreme conditions. Eur J Biol Res 8 (4): 174-182. DOI: 10.5281/zenodo.1412796.
- Zuhri M. 2018. Alih fungsi lahan pertanian di Pantura Jawa Tengah (studi kasus Kabupaten Brebes). Jurnal Litbang Provinsi Jawa Tengah 6 (1): 119-130. DOI: 10.36762/litbangjateng.v16i1.756. [Indonesian]
- Zuraidah. 2013. Pengujian beberapa bakteri penghambat pertumbuhan *Xanthomonas oryzae* pv. *oryzae* pada tanaman padi. Jurnal Ilmiah Pendidikan Biologi Biologi Edukasi 5 (1): 18-24. [Indonesian]