

6. Formulation of low-fat cheese analogue from sweet corn extract using papain and lime extract as coagulant

Jurnal internasional bereputasi

Nama jurnal: Food Research

Vol 4 no 4, Agustus 2020

1. Cover
2. Editor
3. Daftar isi
4. Artikel
5. Indexing (Q3, SJR 0,218)
6. Bukti koresponden





FOOD RESEARCH

Volume 4, Issue 4

August 2020

Review

Monoacylglycerols and diacylglycerols for fat-based food products: a review

Subroto, E.

Available Online: 21 JANUARY 2020

Subroto reviewed the monoacylglycerols and diacylglycerols for fat-based food products.



PDF (378KB)

[Download File](#)

Mini Review

Emerging of Bio-nano composite gelatin-based film as bio-degradable food packaging: a review

Tuan Zainazor, T.C., Fisal, A., Goh, E.G., Che Sulaiman, N.F. and Sarbon, N.M.

Available Online: 19 FEBRUARY 2020

Tuan Zainazor *et al.* reviewed the emerging of bio-nano composite gelatine-based film as bio-degradable food packaging.



PDF (432KB)

[Download File](#)

Full Paper

The quality of nata de coco from sawarna and mapanget coconut varieties to the time of storing coconut water



PDF (685KB)

[Download File](#)

Santosa, B., Wignyanto, W., Hidayat, N. and Sucipto, S.

Available Online: 23 DECEMBER 2019

Santosa *et al.* studied the quality of nata de coco from sawarna and mapanget coconut varieties produced from stored coconut water.

Storage stability of freeze-dried *Lactobacillus plantarum* S20 starter culture as affected by various formulations of drying medium, and its fermentation characteristics in mung bean (*Vigna radiata* L.) slurry

Sulabo, A.S.L., Villasanta, M.E.L., Hermo, K.G., Lascano, R.A., Collado, L.S. and Babaran, G.M.O.

Available Online: 28 FEBRUARY 2020

The storage stability of freeze-dried *Lactobacillus plantarum* S20 starter culture as affected by various formulations of drying medium, and its fermentation characteristics in mung bean (*Vigna radiata* L.) slurry was studied by Sulabo *et al.*



PDF (585KB)

[Download File](#)

Regression analysis for determination of antioxidant activity of coconut sap under various heating temperature and concentration of lysine addition

Sulistyo, S.B. and Haryanti, P.

Available Online: 20 FEBRUARY 2020

Sulistyo and Haryanti studied the regression analysis to determine the antioxidant activity of coconut sap under various heating temperature and concentration of lysine addition.



PDF (518KB)

[Download File](#)

Prevalence of *Bacillus cereus* s.l. in ultra-high temperature chocolate milk from selected milk manufacturers in Malaysia

Ubong, A., New, C.Y., Chai, L.C., Loo, Y.Y., Nor Khaizura, M.A.R., Kayali, A.Y. and Son, R.

Available Online: 8 MARCH 2020

The prevalence of *Bacillus cereus* s.l. in ultra-high temperature chocolate milk from selected milk manufacturers in Malaysia was studied by Ubong *et al.*



PDF (461KB)

[Download File](#)

Physicochemical and functional properties of cassava flour grown in different locations in Sabah, Malaysia

Hasmadi, M., Harlina, L., Jau-Shya, L., Mansoor, A.H., Jahurul, M.H.A. and Zainol, M.K.

Available Online: 9 MARCH 2020



PDF (367KB)

[Download File](#)

The physicochemical and functional properties of cassava flour grown in different locations in Sabah, Malaysia was studied by Hasmadi *et al.*

The effect of processed *Tempeh Gembus* to triglycerides levels and insulin resistance status in women with obesity



PDF (458KB)
Download File

Nadia, F.S., Wati, D.A., Isnawati, M., Sulchan, M. and Afifah, D.N.

Available Online: 5 MARCH 2020

Nadia *et al.* studied the effect of consumption of *Tempeh Gembus* on the triglyceride levels and insulin resistance status in women with obesity.

Purification engineering of mono-diacylglycerol using creaming demulsification technique in 20 kg-scale



PDF (874KB)
Download File

Mursalin, M., Sahrial, S. and Yernisa, Y.

Available Online: 12 MARCH 2020

Mursalin *et al.* studied the engineering of purification of mono-diacylglycerol using creaming demulsification technique in 20 kg-scale.

Cytotoxic effect of *Spirulina platensis* extract and *Ulva compressa* Linn. on cancer cell lines



PDF (538KB)
Download File

Maryati, M., Saifudin, A., Wahyuni, S., Rahmawati, J., Arrum, A., Priyunita, O., Aulia, A., Putra, F.P.H., As'hari, Y., Rasyidah, U.M., Fadhilah, A., Muflihah, C.H. and Da'I, M

Available Online: 14 MARCH 2020

The cytotoxic effect of *Spirulina platensis* extract and *Ulva compressa* Linn. on cancer cell lines was studied by Maryati *et al.*

Formulation, characterization and stability of o/w nanoemulsion containing rice bran oil prepared by emulsion phase inversion



PDF (427KB)
Download File

Sari, Y. P., Raharjo, S., Santoso, U. and Supriyadi

Available Online: 9 MARCH 2020

Sari *et al.* studied on the formulation, characterization and stability of o/w nanoemulsion containing the rice bran oil prepared by emulsion phase inversion.

Sensorial and physicochemical characteristics of herbal noodle enriched with *Centella asiatica*



PDF (425KB)
Download File

Fadzil, N.F., Abu Bakar, M.F., Yusop, M.H.M. and Abu Bakar, F.I.

Available Online: 26 FEBRUARY 2020

Fadzil *et al.* studied the sensorial and physicochemical characteristics of herbal noodle enriched with *Centella asiatica*.

Effect of *Lactobacillus plantarum* and *Leuconostoc mesenteroides* starter cultures in lower salt concentration fermentation on the sauerkraut quality



PDF (482KB)
Download File

Zubaidah, E., Susanti, I., Yuwono, S.S., Rahayu, A.P., Srinta, I. and Blanc, P.J.

Available Online: 10 MARCH 2020

Zubaidah *et al.* studied the effect of *Lactobacillus plantarum* and *Leuconostoc mesenteroides* starter cultures on the quality of low salt concentration sauerkraut.

Modeling of food safety knowledge, attitude, and behavior characteristics



PDF (415KB)
Download File

Ellinda-Patra, M.W., Dewanti-Hariyadi, R. and Nurtama, B.

Available Online: 20 MARCH 2020

Ellinda-Patra *et al.* modelled the food safety knowledge, attitude and behaviour characteristics.

In vitro antioxidant properties and digestibility of chicken feather protein hydrolysates



PDF (814KB)
Download File

Oluba, O.M., Akpor, O.B., Alabi, O.O., Shoyombo, A.J., Adeyonu, A.G. and Adebisi, F.D.

Available Online: 10 FEBRUARY 2020

Oluba *et al.* studied the in vitro antioxidant properties and digestibility of chicken feather protein hydrolysates

Improving plant layout and processing flow of jelly mushroom drink to reduce microbial contamination



PDF (809KB)
Download File

Changchai, S., Sumrit, T., Chunlakan, P. and Inriri, P.

Available Online: 23 MARCH 2020

Changchai *et al.* studied on improving the plant layout and processing flow of jelly mushroom drink to reduce microbial contamination

Effect of chemical agents, metallic salts on the stability of α -amylase, protease and comparative analyses of enzyme activity of selected salad vegetables



PDF (359KB)
Download File

Rahman, M.M., Shahjadee, U.F., Khanam, F., Rupa, A.Z. and Azad, M.A.K.

Available Online: 9 MARCH 2020

The effect of chemical agents, metallic salts on the stability of α -amylase, protease and comparative analyses of enzyme activity of selected salad vegetables was studied by Rahman *et al.*

Formulation of low-fat cheese analogue from sweet corn extract using papain and lime extract as coagulant



PDF (825KB)

[Download File](#)

Aini N., Sustriawan, B., Prihananto, V., Sumarmono, J., Ramadan R.N. and Romadhon, D.

Available Online: 24 MARCH 2020

Aini *et al.* studied the formulation of low-fat cheese analogue from sweet corn extract using papain and lime extract as coagulant.

The study of angiotensin converting enzyme isolation from crossbred rabbit lung and enzyme storage capacity in frozen conditions



PDF (445KB)

[Download File](#)

Nguyen, V.M., Tran, T.T. and Vo, H.N.

Available Online: 26 MARCH 2020

Nguyen *et al.* studied the angiotensin converting enzyme isolated from crossbred rabbit lungs and the enzyme storage capacity in frozen conditions.

Texture, microstructure and *in-vitro* starch digestibility of waxy rice cooked with hydrocolloid



PDF (941KB)

[Download File](#)

Srikaeo, K. and Paphonyanyong, W.

Available Online: 26 MARCH 2020

The texture, microstructure and *in-vitro* starch digestibility of waxy rice cooked with hydrocolloid was studied by Srikaeo and Paphonyanyong.

Effect of the different encapsulation methods on the physicochemical and biological properties of *Clitoria ternatea* flowers microencapsulated in gelatine



PDF (833KB)

[Download File](#)

Liew, S.Y., Mohd Zin, Z., Mohd Maidin, N.M., Mamat, H. and Zainol, M.K.

Available Online: 26 MARCH 2020

Liew *et al.* evaluated the effect of different encapsulation methods on the physicochemical and biological properties of *Clitoria ternatea* flowers microencapsulated in gelatine.

Quality and sensory profile evaluation of gluten-free sapodilla-wild almond seed bar with stevia as partial sugar substitution



PDF (432KB)

[Download File](#)

Akesowan, A., Choonhahirun, A. and Jariyawaranugoon, U.

Available Online: 27 MARCH 2020

The quality and sensory profile evaluation of gluten-free sapodilla-wild almond seed bar with stevia as partial sugar substitution was studied by Akesowan *et al.*

Phytochemical screening, phenolic acid profiling and antioxidant activity analysis of peels from selected mango (*Mangifera* spp.) genotypes in the Philippines



PDF (1080KB)

[Download File](#)

Ocampo, E.T.M., Libron, J.A.M.A., Guevarra, M.L.D. and Mateo, J.M.C.

Available Online: 30 MARCH 2020

The phytochemical screening, phenolic acid profiling and antioxidant activity analysis of peels from selected mango (*Mangifera* spp.) genotypes in the Philippines were evaluated by Ocampo *et al.*

Phytochemical composition and essential mineral profile, antioxidant and antimicrobial potential of unutilized parts of jackfruit



PDF (402KB)

[Download File](#)

Adan, A.A., Ojwang, R.A., Muge, E.K., Mwanza, B.K. and Nyaboga, E.N.

Available Online: 1 APRIL 2020

The phytochemical composition and essential mineral profile, antioxidant and antimicrobial potential of unutilized parts of jackfruit were studied by Adan *et al.*

Separation and analysis of *Monascus* yellow pigment produced on durian seed substrate



PDF (569KB)

[Download File](#)

Srianta, I., Nugerahani, I. and Ristiarini, S.

Available Online: 27 MARCH 2020

Srianta *et al.* separated and analysed *Monascus* yellow pigment produced on durian seed substrate.

Effect of air drying on quality characteristics and mass transfer kinetics of osmotically dehydrated sea buckthorn by stevia



PDF (589KB)

[Download File](#)

Lentzou, D., Templalexis, Ch. and Xanthopoulos, G.

Available Online: 27 MARCH 2020

The effect of air drying on quality characteristics and mass transfer kinetics of osmotically dehydrated sea buckthorn by stevia was studied by Lentzou *et al.*

Consumption of young barley leaf extract increases fecal short-chain fatty acid levels: a before-after clinical trial



PDF (359KB)

[Download File](#)

Unno, T., Tanaka, H. and Kohara, A.

Available Online: 1 APRIL 2020

Unno *et al.* studied the effects of consumption of young barley leaf extract on the levels on short-chain fatty acid in feces.

Determination of simple carbohydrates in beer using SPME, on fiber derivatization and gas chromatography mass spectrometry



PDF (636KB)
Download File

D'Auria, M., Emanuele, L., Racioppi, R. and Stefanizzi, N.

Available Online: 1 APRIL 2020

D'Auria *et al.* determined the simple carbohydrates in beer using SPME on fiber derivatization and gas chromatography mass spectrometry.

The effects of different pre-treatments on the physicochemical composition and sensory acceptability of 'Kacang Koro' energy bar



PDF (389KB)
Download File

Wan Mohamad Din, W.N.I., Mohd Zin, Z., Abdullah, M.A.A. and Zainol, M.K.

Available Online: 5 APRIL 2020

The effects of different pre-treatments on the physicochemical composition and sensory evaluation of 'Kacang Koro' energy bar were studied by Wan Mohamad Din *et al.*

Effects of various grated sago sizes, steeping periods and trunk sections on sago starch recovery (*Metroxylon* spp.).



PDF (676KB)
Download File

Wan Mohd Fariz, W.A., Rosnah, S., Mohd Zuhir, M.N., Azman, S. and Noramina, H.

Available Online: 30 MARCH 2020

Wan Mohd Fariz *et al.* studied the effects of various grated sago sizes, steeping periods and trunk sections on sago starch recovery (*Metroxylon* spp.).

Physicochemical, antioxidant and sensory characteristics of cookies supplemented with different levels of spent coffee ground extract



PDF (376KB)
Download File

Azuan, A.A., Mohd Zin, Z., Mamat, H., Rusli, N.D. and Zainol, M.K.

Available Online: 8 APRIL 2020

The physicochemical, antioxidant and sensory characteristics of cookies supplemented with different levels of spent coffee ground extract was studied by Azuan *et al.*

The viability of probiotic *Lactobacillus acidophilus* IFO 13951 and *Bifidobacterium longum* ATCC 15707 in gummy candies decreased during 4 weeks of storage



PDF (343KB)
Download File

Lestari, L.A., Kusuma, W.I., Nurhayati, F., Kusuma, R.J. and Erwanto, Y.

Available Online: 3 APRIL 2020

Lestari *et al.* studied the viability of probiotic *Lactobacillus acidophilus* IFO 13951 and *Bifidobacterium longum* ATCC 15707 in gummy candies during 4 weeks of storage.

Development of indigenous *Streptococcus salivarius* TUC 1253-fermented soy oral strip using Box-Behnken design for oral health



PDF (709KB)

[Download File](#)

How, Y.H., Ewe, J.A., Song, K.P., Kuan, C.S., Kuan, C.H. and Yeo, S.K.

Available Online: 9 APRIL 2020

How *et al.* developed indigenous *Streptococcus salivarius* TUC 1253-fermented soy oral strip using the Box-Behnken design for oral health.

Effect of cooking methods on nutritional composition and antioxidant properties of lotus (*Nelumbo nucifera*) rhizome



PDF (449KB)

[Download File](#)

Chen, S.Y. and Tengku Rozaina, T.M.

Available Online: 11 APRIL 2020

The effect of cooking methods on nutritional composition and antioxidant properties of lotus (*Nelumbo nucifera*) rhizome was studied by Chen and Tengku Rozaina.

Physical and chemical characteristics of goldenberry (*Physalis peruviana*) grown in Lam Dong province, Vietnam



PDF (528KB)

[Download File](#)

Thuy, N.M., Phuong, N.P, Suong, C.T.D. and Tai, N.V.

Available Online: 12 APRIL 2020

The physical and chemical characteristics of goldenberry (*Physalis peruviana*) grown in Lam Dong province, Vietnam were studied by Thuy *et al.*

Microstructure and physicochemical characteristics of modified taro starch after annealing, autoclaving-cooling and heat moisture treatment



PDF (474KB)

[Download File](#)

Setiarto, R.H.B., Kusumaningrum, H.D., Jenie, B.S.L., Khusniati, T., Widhyastuti, N. and Ramadhani, I.

Available Online: 15 APRIL 2020

The microstructure and physicochemical characteristics of modified taro starch after annealing, autoclaving-cooling and heat moisture treatment were studied by Setiarto *et al.*

Detection of enterotoxin gene (*sea*) and biofilm formation ability among multi-drug resistant *Staphylococcus aureus* isolated from



PDF (451KB)

[Download File](#)

shawarma sandwich sold at selected kiosks in Klang Valley, Malaysia

Salahaldin Fathalla M., Mahyudin, N.A., Mohamad Ghazali, F., Rukayadi, Y. and Jaafar, S.N.

Available Online: 16 APRIL 2020

Salahaldin Fathalla *et al.* investigated the prevalence of enterotoxin gene (*sea*) and biofilm formation ability multi-drug resistant *Staphylococcus aureus* from shawarma sandwich sold at selected kiosks in Klang Valley, Malaysia

Physicochemical and sensory properties of deep fried battered squid containing Brownstripe red snapper (*Lutjanus vitta*) protein hydrolysate

Chew, R.M., Mohd Zin, Z., Ahmad, A., Mohtar, N.F., Rusli, N.D. and Zainol, M.K.

Available Online: 17 APRIL 2020

Chew *et al.* studied the physicochemical and sensory properties of deep fried battered squid containing Brownstripe red snapper (*Lutjanus vitta*) protein hydrolysate.



PDF (714KB)
Download File

Accelerated shelf-life test (ASLT) of batuan [*Garcinia binucao* (Blanco) Choisy] fruit powder

Ancheta, A.K.G., Yaptenco, K.F., Mopera, L.E., Bainto, L.C., Lizardo, R.C.M. and Dizon, E.I.

Available Online: 17 APRIL 2020

Ancheta *et al.* perform the accelerated shelf-life test (ASLT) to study the shelf-life of batuan [*Garcinia binucao* (Blanco) Choisy] fruit powder.



PDF (498KB)
Download File

Effect of starch modification on physico-chemical, functional and structural characterization of cassava starch (*Manihot esculenta* Crantz)

Dolas, K.A., Ranveer, R.C., Tapre, A.R., Nandane, A.S. and Sahoo, A.K.

Available Online: 14 APRIL 2020

Dolas *et al.* studied the effects of starch modification on physico-chemical, functional and structural characterization of cassava starch (*Manihot esculenta* Crantz)



PDF (462KB)
Download File

Ethnic differences on cardiovascular disease risk and quality of life in Selangor

Lee, H.S., Aman, A., Kamaruddin, K.S., Ali, A. and Yusof, H.M.

Available Online: 19 APRIL 2020

Lee *et al.* studied the ethnic differences on cardiovascular disease risk and quality of life in Selangor.



PDF (627KB)
Download File

Formulation and quality evaluation of instant mango drink powder

Akther, S., Alim, M.A., Badsha, M.R., Matin, A., Ahmad, M. and Hoque, S.M.

Available Online: 8 APRIL 2020

Akther *et al.* formulated instant mango drink powder and evaluated the quality of the developed product.



PDF (389KB)

[Download File](#)

Microbiological quality analysis of different types of popular dried food items

Ahmed, T., Sharmin, A., Paul, A., Dipu, M.R. and Islam, T.

Available Online: 3 APRIL 2020

The microbiological quality of different types of popular dried food items were analysed by Ahmed *et al.*



PDF (386KB)

[Download File](#)

Ramification of pH in pectinase-assisted extraction on the antioxidant capacity of Arabica spent coffee ground (SCG) extract

Shukri, N.A., Mohd Zin, Z., MohdMaidin, N., Mamat, H. and Zainol, M.K.

Available Online: 22 APRIL 2020

Shukri *et al.* studied the ramification of pH in pectinase-assisted extraction on the antioxidant capacity of Arabica spent coffee ground (SCG) extract.



PDF (639KB)

[Download File](#)

Isolation of pathogenic microorganisms and determination of their antibiotic resistance patterns collected from different bakery products of Dhaka city

Das, K.K., Sarkar, A. and Hossain, A.

Available Online: 22 FEBRUARY 2020

Das *et al.* isolated pathogenic microorganisms from different bakery products of Dhaka city and determined their antibiotic resistance patterns.



PDF (338KB)

[Download File](#)

Optimization of vacuum infiltration before blanching of black cherry tomatoes (*Solanum lycopersicum* cv. OG) using response surface methodology

Ha, H.T.N. and Thuy, N.M.

Available Online: 24 APRIL 2020

Ha and Thuy optimized the vacuum infiltration before blanching of black cherry tomatoes (*Solanum lycopersicum* cv. OG) using response surface methodology.



PDF (895KB)

[Download File](#)

Physicochemical and emulsifying properties of pre-treated octenyl succinic anhydride (OSA) sago starch in simple emulsion system



PDF (547KB)
Download File

Saw, S.S., Shariffa, N.Y., Ruri, A.S. and Uthumporn, U.

Available Online: 24 APRIL 2020

Saw *et al.* evaluated the physicochemical and emulsifying properties of pre-treated octenyl succinic anhydride (OSA) sago starch in simple emulsion system.

Development of composite biscuits made from amadumbe (*Colocasia esculenta* L. Schott) and soybean (*Glycine max* L. Merrill) and investigation of their nutritional and organoleptic properties



PDF (565KB)
Download File

Mokhele, T.M., Amonsou, E.O. and Tabit, F.T.

Available Online: 14 APRIL 2020

Mokhele *et al.* developed composite biscuits made from amadumbe (*Colocasia esculenta* L. Schott) and soybean (*Glycine max* L. Merrill) and investigated their nutritional and organoleptic properties.

Consumers' willingness to pay for functional rice: a survey from Indonesia



PDF (661KB)
Download File

Annur, N.D., Nugrohoningtyas, B.S.H., Rodríguez Doderó, M.C. and Setyaningsih, W.

Available Online: 26 APRIL 2020

Annur *et al.* surveyed the consumers' willingness to pay for functional rice in Indonesia.

Microbiological analysis of different categories of food items in Dhaka city, Bangladesh



PDF (335KB)
Download File

Uddin, M.A., Jabin, T., Siam, E.A., Afreen, M.T., Uddin, S.S. and Sayeda, U.

Available Online: 25 APRIL 2020

Uddin *et al.* performed a microbiological analysis on the different food items in Dhaka city, Bangladesh.

Clitoria ternatea anthocyanin extract suppresses inflammation in carrageenan-induced rat paw edema via down-regulating genes of phosphoinositide 3-kinase signaling pathway



PDF (376KB)
Download File

Yanti, Sabella, D., Gunawan, A.W. and Lay, B.W.

Available Online: 8 APRIL 2020

Yanti *et al.* studied the effects of *Clitoria ternatea* anthocyanin extract in suppressing the inflammation in carrageenan-induced rat paw edema via down-regulating genes of phosphoinositide 3-kinase signaling pathway.

Effect of different molecular weight on the antioxidant activity and physicochemical properties of golden apple snail (*Ampullariidae*) protein hydrolysates

Saallah, S., Ishak, N.H. and Sarbon, N.M.

Available Online: 17 FEBRUARY 2020

The effect of different molecular weight on the antioxidant activity and physicochemical properties of golden apple snail (*Ampullariidae*) protein hydrolysates were studied by Saallah *et al.*



PDF (431KB)

[Download File](#)

Optimization of processing parameters for vacuum fried oyster mushroom (*Pleurotus ostreatus* (Jacquin) P. kummer)

Hilapad, M.R., Esguerra, E.B. and Castillo - Israel, K.A.T.

Available Online: 28 APRIL 2020

Hilapad *et al.* optimized the processing parameters for vacuum fried oyster mushrooms (*Pleurotus ostreatus* (Jacquin) P. kummer)



PDF (1.67MB)

[Download File](#)

Articles 1 - 53

VOL. 4 | ISSUE 3

VOL. 4 | ISSUE 5

[HOME](#)[VIEW ARTICLES](#)[VIEW ISSUES](#)[FOR AUTHORS](#)[FOR REFEREES](#)[ABOUT THE JOURNAL](#)

EDITORIAL BOARD

Chief Editor

Son Radu

Malaysia

Editor

Vivian New Chia Yeung

Malaysia

Associate Editors

Prem Narayan Paudel

Kathmandu University

Nepal

Associate Editors

Borarin Buntong

Royal University of Agriculture

Cambodia

Associate Editors

Yoshitsugu Nakaguchi

Ishikawa Prefectural University

Japan

Nasreldin Elhadi

University of Dammam

Saudi Arabia

Dwaipayan Sikdar

University of Chittagong

Bangladesh

Rungsinee Sothornvit

Kasetsart University,

Thailand

Nikos Tzortzakis

Cyprus University of Technology

Cyprus

Saisamorn Lumyong

Chiang Mai University

Thailand

Ekachai Chukeatirote

Mae Fah Luang University

Thailand

Pin Kar Yong

Forest Research Institute
Malaysia (FRIM)
Malaysia

Chim Chay

University of Agriculture
Cambodia

Le Van Viet Man

Ho Chi Minh City University of
Technology
Viet Nam

Oscar Herrera-Calderon

Universidad Nacional Mayor de
San Marcos
Peru

Hasmadi Mamat

University Malaysia Sabah
Malaysia

Regiane Ribeiro dos Santos

Federal Rural University of Rio de
Janeiro
Brazil

Koh Soo Peng

Malaysian Agricultural Research
and Development Institute
(MARDI)
Malaysia

Dinko Hristov Dinko

Trakia University
Bulgaria

Hanifah Nuryani Lioe

Bogor Agricultural University
Indonesia

Michelle Galindo de Oliveira

Federal University of
Pernambuco
Brazil

Alper Sener

Onzekiz Mart University
Turkey

Angelo Maria Giuffrè

Mediterranea University of
Reggio Calabria
Italy

Javier Garcia-Lomillo

University of Burgos
Spain

Laura Marcela Zárate Polanco

National University of Colombia-
Bogotá
Colombia

Swapan K. Banerjee

HPFB Health Canada,
Canada

Adel Abu Bakr Abd El-Hamid**Shatta**

Suez Canal University
Egypt

Iddya Karunsagar**Saroat Rawdkuen**

Mae Fah Luang University
Thailand

Giovanna Giacalone

Università di Torino
Italy

Lorina Acilo Galvez

Visayas State University
Philippines

Sergio Soto Simental

Universidad Autónoma del
Estado de Hidalgo
Mexico

Fernando Cebola Lidon

Universidade Nova de Lisboa
Portugal

Linda Lim Biaw Leng

Universiti Brunei Darussalam
Brunei Darussalam

J.M. Krishanti Jayarukshi Kurr**Premarathne**

Wayamba University of Sri Lanka
Sri Lanka

Katherine Ann T. Castillo-Isra

University of the Philippines L
Banos
Philippines

Steve Flint

Massey University

Giuseppe Sortino
University of Palermo
Italy

Nitte University
India

New Zealand

COPYRIGHT © 2016 - 2020 RYNNYE LYAN RESOURCES

Formulation of low-fat cheese analogue from sweet corn extract using papain and lime extract as coagulant

^{1*}Aini N., ¹Sustriawan, B., ¹Prihananto, V., ²Sumarmono, J., ¹Ramadan R.N. and ¹Romadhon, D.

¹Department of Food Science and Technology, Jenderal Soedirman University, Jalan Dr. Soeparno, Purwokerto, 53123, Central Java, Indonesia

²Department of Animal Husbandry, Jenderal Soedirman University, Jalan Dr. Soeparno, Purwokerto, 53123, Central Java, Indonesia

Article history:

Received: 24 November 2019

Received in revised form: 13 January 2020

Accepted: 15 January 2020

Available Online: 24 March 2020

Keywords:

Cheese analogues,

Corn extract,

Low fat,

Carotenoid

DOI:

[https://doi.org/10.26656/fr.2017.4\(4\).395](https://doi.org/10.26656/fr.2017.4(4).395)

Abstract

Cheese is not only created using cow's milk and can also be made from a mixture of vegetable extracts, including corn extract. Cheese from corn extract has the advantages of low-fat and high-carotene. Notably, papain can be used as a coagulant in the production of cheese analogue, while maltodextrin functions to increase volume and total solids for greater yield. The objectives of the present study was 1) to optimize the formula composition between lime extract, papain, and maltodextrin to create a cheese analogue from sweet corn extract with high yield and protein as well as good sensory properties, 2) to study the physicochemical and sensory characteristics of the cheese analogue using the optimal formula, and 3) to compare analog cheese from corn milk to cow's milk cheese. The experimental design involved response surface methodology with three factors (lime extract, papain, and maltodextrin). The results of the study produced the optimal cheese analogue formula from corn extract with the addition of lime extract (2.283%), papain (0.022%), and maltodextrin (15%). The characteristics of this cheese analogue include a yield of 20.3%; pH of 5.4; 14°Brix soluble solids; water content of 65.3%; protein content of 13.5%; total-carotene of 544.4 ppm and of fat content 4.6%. The cheese analogue has sensory characteristics of soft texture, the ability to spread evenly, the typical color of cheese (i.e. yellowish-white), and was preferred by panelists. Cheese analogue has protein content of 7.1%, fat content of 4.55%, total carotene of 544.4 mg/g, cholesterol 0.02 mg/g; while commercial cheese from cow's milk has protein content 6.3%, fat content 24.53%, total carotene 5.32 mg/g and cholesterol 0.19 mg/g. Thus, sweet corn can potentially be used as a raw material for producing low-fat cheese analogues.

1. Introduction

Cheese is not only used in bakery products but also used in various other types of culinary items such as for appetizers, entrees, desserts, and various snacks. Grated, sliced, and liquid cheeses have been developed for various purposes in the modern food industry. A trend of increasing cheese popularity has led to an increased demand for cheese. In order to satisfy the increasing demand for cheese, products are developed from ingredients other than cow's milk. This product is called cheese analogue or imitation cheese. Cheese analogue is a type of cheese product, with fat, milk protein, or both being replaced in part or entirely by non-milk components, especially from vegetable ingredients (e.g. apricots, sunflower seeds, or other plants) (Cunha *et al.*,

2010; O'Riordan *et al.*, 2011; Mohamed and Shalaby, 2016). The fat originating from milk is replaced by vegetable oil or fat in cheese analogue production, resulting in saturated fatty acid levels being very low, thus reducing the risk of cardiovascular disease among consumers (Liu *et al.*, 2008; Tuntragul *et al.*, 2010). Sweet corn extract can be processed into yogurt (Aini, Prihananto, Wijonarko, Astuti *et al.*, 2017); therefore, it is assumed that it can also be processed into cheese analogue. The production of cheese analogue from corn extract is not commonly performed; as such, there is room for development in cheese analogue production using corn extract. According to Aini, Prihananto, Wijonarko, Astuti *et al.* (2017), corn extract has low fat content and high beta carotene. The cheese analogue

*Corresponding author.

Email: nur.aini@unsoed.ac.id

from corn extract produced the expected low in fat and have high levels of beta carotene as the benefits of its products.

The technique of making cheese through direct acidification can produce soft white cheese that can be consumed without going through the ripening process (González *et al.*, 2018). This technique can produce soft cheese, high meltability, and good stretchability (i.e. forming fiber when stretched, making it suitable for use in making pizza and as cheese spread). However, the exploration of direct acidification techniques remains very limited. In this technique, the acidification stage is typically performed through the addition of organic acids, such as vinegar, lactic acid, or citric acid (Aini, Prihananto, Sustriawan *et al.*, 2019). Research on direct acidification techniques remains very limited. High acidity fruit extracts can also coagulate milk in the production of soft cheese (e.g. lime and pineapple) (Sumarmono and Suhartati, 2016; Aini, Sustriawan, Prihananto *et al.*, 2019).

The cheese production process typically involves using the enzyme rennet as a coagulant (Hashim *et al.*, 2011). Rennet enzymes used as coagulants of cheese are very expensive, and their availability is limited; as such, an alternative to the rennet enzyme is required. According to (Mahajan and Chaudhari, 2014), in order to reduce dependence on rennet enzymes, vegetable extract can be used as a substitute. An ingredient that can be used as a substitute for rennet enzyme is papain (Aini, Prihananto, Sustriawan *et al.*, 2019). The use of papain in cheese production is expected to replace the rennet enzyme in order to reduce production cost.

Moreover, the addition of fillers such as maltodextrin can increase the yield of cheese. According to (Moser *et al.*, 2017), maltodextrin functions to increase the volume and raise the total solids of the material to increase yield. Thus, the addition of maltodextrin as a filler in cheese analogue can increase its yield (Aini, Prihananto, Sustriawan *et al.*, 2019).

The objectives of this study were: 1) to optimize the formula composition between lime extract, papain, and maltodextrin to create a cheese analogue from sweet corn extract with high yield and protein as well as good sensory properties and 2) to study the physicochemical and sensory characteristics of the cheese analogue using the optimal formula.

2. Materials and methods

2.1 Materials

The main ingredients used in this study were sweet corn and lime obtained from the local market in

Purwokerto, Indonesia; fresh cow's milk (moisture content of 86%, protein content of 3%, and fat content of 3.2%) from the Experimental Farm, Jenderal Soedirman University, Purwokerto, Indonesia; commercial papain (Paya); and *Lactobacillus casei* (in the form of liquid) from the Indonesian Institute of Sciences, Cibinong, Indonesia. The main tools used were blenders (Miyako), analytical balance (AND), an oven (Mettler), presses, a spectrophotometer (Shimano UV-VIS 1800), a refractometer (Atago), and a pH meter.

2.2 Formulation and characterization of the initial products

This research was performed in three stages, namely: formulation and characterization of the initial products, determine the optimal formula, and optimization stage. We followed the method of Aini, Prihananto, Sustriawan *et al.* (2019) with modification. The first stage aimed to determine the lower and upper limits of the concentration of the independent variables. Determination of the lower limits was based on trial and error, while the upper limits were determined through the organoleptic test. The experimental design of this study involved RSM with three independent variables (the concentrations of lime extract, papain, and maltodextrin). Each factor had a lower limit (1%, 0.018%, and 5%) and an upper limit (3%, 0.022%, and 15%). The determination of variations in formula combinations using the RSM method was assisted by Design Expert 11 software. Based on the results of Design Expert 11 calculations for three independent variables, 20 variations of formula were obtained. In this stage, each response was tested using quadratic model equations. Based on the test results of variance (ANOVA) when the p-value less than 0.05 indicates a real effect (significant), which means that the model represents the independent variable in influencing the response.

2.3 Determine the optimal formula

In the second stage, the optimization process was performed to determine the formula with the optimal response. In this section, it is determined criteria that include variables and each response that influences. The optimized components included the research factors (concentrations of lime extract, papain, and maltodextrin) and variables observed in stage I (moisture content, yield, soluble protein, soluble solids, pH, and fat). At this stage, the expected criteria for each variable are set by determining the level of importance and the target values. Yield, soluble solids and protein content have a higher importance level (5) than pH, water content and fat content (importance level 3).

Target values can be minimized, maximized, or in

range, depending on the desired value. The components of maltodextrin, papain, and lime extract were optimized with a target range so that the values at the lower and upper limit concentrations have the same opportunity to be selected in determining the optimum formula. Moisture content is optimized in ranged because the moisture content of cheese analogues ranged between 63.5 and 68.3%, which is within the range of moisture content for fresh cheese (55-80%) (Lee *et al.*, 2015). Expected texture in spreadable cheese analogue is semi-solid. Moisture content that is too low can produce a hard texture, while moisture content that is too high cause the texture of the cheese to be runny and difficult to solidify when applied. The yield and protein content were optimized for a maximum target range. The fat content of cheese analogue was optimized for the minimum targets because it was expected to contain less fat. Also, pH was optimized for the target range so that the produced cheese analogue would be less sour, but remain within the isoelectric point range to obtain the maximum yield.

2.4 Optimization stage and verification

From the previous stage, 10 optimum formulas are obtained, but only three optimum formulas that have the highest desirability values are taken for this stage. Optimization was performed in order to achieve the optimal cheese analogue formula (Bezerra *et al.*, 2008). At the optimization stage using Response Surface Methodology also obtained predictive value for each variable.

The three optimum formulas were verified in this stage. The verification phase is done by comparing the results of analysis in the laboratory with the predicted value of the response generated from the response surface methodology method. The three optimum formulas were also analyzed for sensory properties using ranking test. The tests were performed by twenty trained panelists using a ranking test based on sensory properties. Panelists are asked to rank products based on sensory qualities (color, aroma, taste, ease smeared and preferences). Products with the best sensory quality are ranked first. Rank test results were analyzed using ANOVA and when significant difference followed by the least significant difference test level 5%.

Three formulas obtained were then analyzed for effectiveness index to get the best product. Determining the best optimum formula is based on the chemical and sensory performed using the index effectiveness (DeGarmo *et al.*, 2003). The best products are then analyzed levels of protein, fat content, total carotene, and cholesterol. The same analysis was also conducted on spreadable cheese from cow's milk. The two products

were compared using LSD test.

2.5 Cheese analogue production

Cheese analogue was prepared based on previous research by Aini *et al.* (Aini, Prihananto, Sustriawan *et al.*, 2019) with slight modifications. Corn extract is made from corn and water in a ratio of 1: 2 which is mashed using a blender for 3 minutes, then filtered. Corn extract then mixed with cow's milk (ratio 1: 1) pasteurized at 72°C for 30 minutes, after which it was cooled to 30°C. The next step is the addition of 1 mL starter (in liquid form), 2 grams of CaCl₂ and 30 grams of sugar per 500 mL of corn extract and cow's milk. Subsequently, the mixture was incubated at 30°C for 2 hours. The next step is the addition of lime extract, papain and maltodextrin and stir until homogeneous, then allowed to stand 30 minutes. The mixture of materials that have undergone coagulation is inserted into the filter cloth and pressed. Pressing is done at 30°C until all whey and curd are separated. The next step is the addition of salt (1%) and sugar (6%) and then do the blanching at 40°C for 15 s. Cheese analogue from corn extract is ready to be analyzed.

2.6 Analysis of cheese analogue

The cheese analogue was analyzed for yield (Aini, Sustriawan, Prihananto *et al.*, 2019), pH using pH meter (Obloh and Imafidon, 2018), soluble solids (Dhanraj *et al.*, 2017), moisture content (AOAC, 2005), protein content (AOAC, 2005), and fat content (AOAC, 2005). The three best products were analyzed for sensory properties using the ranking method. The best products are then analyzed levels of protein, fat content, total carotene using AOAC 970.64 (Hoffmann *et al.*, 2017) and cholesterol

The yield is the ratio between the cheese analogues and the ingredients. The determination of actual yield requires measurement of the weight of production inputs and outputs (Badem and Ucar, 2016). The actual yield may then be calculated using the following equation:

$$\text{Yield} = \frac{\text{the weight of cheese}}{\text{the weight of ingredients}} \times 100\%$$

Moisture content was analyzed using the AOAC method (AOAC, 2005) as follows: 1-2 g of sample is weighed and placed in a bowl that has been dried. Then, the sample and bowl were dried in a 105°C temperature oven until a constant weight was obtained. Moisture content is the difference between the weight of the starting material and the weight of the final material after drying.

The pH was assessed out using pH meter (Obloh and Imafidon, 2018). The pH meter was standardized using a

buffer solution with a pH of 4.0 and 7.0. pH readings were taken after 10 mins stabilization. Then, a 10% w/v suspension of the sample was prepared using distilled water. The mixture was mixed vigorously by manual shaking, and pH was measured using a functional pH meter (Extech instruments, model DO700, China).

Determination of soluble solids is carried out using a refractometer (Cunha *et al.*, 2010). The measurement method is as follows: the sample is stirred until homogeneous, then filtered through filter paper. The filtrate is collected, then taken and dropped on the refractometer prism. Read scale on the tool, and then converted to the value of the refractive index of soluble solids.

Soluble protein was analyzed using the method of Lowry-Folin as follows: 0.1-1 g of sample was dissolved in 100 mL of distilled water. Then, 1 mL of the sample was taken and placed in a test tube. Then, 5.5 mL of copper sulphate was added and left for 10-15 mins at room temperature. Thereafter, 0.5 mL of Folin Ciocalteu was added to each of the test tubes, which were immediately beaten quickly and evenly. The solution was left for 30 mins until a blue color was formed. The intensity of blue was measured by its absorbance using a spectrophotometer at a wavelength of 600 nm. The calculation of soluble protein levels was determined using the standard bovine serum albumin curve.

Fat content measurements were performed using a soxhlet extractor as follows: 2-5 g of finely ground sample wrapped in filter paper was inserted into a soxhlet extraction tube. Then, a fat dish and extraction tube are installed on the distillation apparatus. The soxhlet extractor that was filled with solvents was then drained with cool water and the appliance was turned on. Extraction was performed for 4-5 hrs. Then, the solvent was separated from the fat, while the fat-filled dish was dried in an oven at 100-105°C for 30 mins. The residual weight in the fat dish is expressed as fat weight.

Total carotene was analyzed using AOAC 970.64 (Hoffmann *et al.*, 2017). Samples of 2.5 g and 15 mL of a mixture of solution (hexane: acetone: ethanol: toluene in a ratio of 10: 7: 6: 7) were mixed using vortex for 1 min. After that, add 1 mL of potassium hydroxide in 10% methanol, and stir for 1 min. The next stage is heating the water bath at 56°C for 20 min. After that it is allowed to stand at room temperature for 1 hr, then add 15 mL of hexane and 29 mL of 10% sodium sulfate solution. After 1 hr, absorbance was taken at 450 nm. Standard curves are made using beta carotene (Sigma-Aldrich) dissolved in hexane.

Cholesterol was analyzed using Beyer and Jensen method (Osman and Chin, 2006). The sample was saponified using 2% alcoholic KOH for an hour. After that, the unsaponified fraction was extracted with 10 mL hexane twice. The extracts were collected and washed with 5 mL distilled water. The next, extract was heated in water bath at 45°C until dry. The dried extracts diluted in 3 mL glacial acetic acid and added with 2 mL of FeCl₃. The suspension is read on a spectrophotometer at 565 nm. The standard solution is prepared by making a solution of cholesterol at concentrations of 0, 0.03, 0.15, 0.30 and 0.60 mg/mL cholesterol than standard cholesterol (Sigma Chemical).

3. Results and discussion

3.1 Characterization of the initial product

Cheese analogue yields ranged from 17.1 to 22.9%. This yield is greater than the yield of cheese from cow's milk (Sumarmono and Suhartati, 2016), which is 7.75 – 10.29%. Based on quadratic models, the yield was a significant effect ($p < 0.05$). Notably, increasing papain and maltodextrin concentrations increased the yield of cheese analogues (Figure 1). Papain is a protease enzyme that can break down casein in milk. According to Fernández-Lucas *et al.* (2017), papain interferes with kappa-casein to form para-kappa-casein. Papain cuts the peptide bond between phenyl and methionine in the kappa-casein, damages its structure, and produces para-kappa-casein—which has a hydrophobic portion. Then, when pH approaches the isoelectric point, casein joins and agglomerates. This casein can be joined due to the interaction of the hydrophobic portions of the kinetic kappa. The presence of calcium in milk aids the coagulation process as a bridge between micelles.




Figure 1. Cheese analogue yield was influenced by papain and maltodextrin concentration

The addition of maltodextrin increases cheese analogue yield because, according to (Jittanit *et al.*, 2010), maltodextrin serves as an enhancer and filler in

order to increase yield. This result is similar to that of other studies (Dhanraj *et al.*, 2017) highlighting that maltodextrin addition increases the yield of mozzarella cheese analogue. Cheese analogue has previously been produced from corn extract, and the addition of will increased its yield (Aini, Prihananto, Sustriawan *et al.*, 2019).

According to Hjalmarsson (2015), the most important factor affecting yield is the composition of milk—particularly the concentrations of fat and caseins. Casein forms the continuous paracasein network and occludes the fat and moisture phases. Moisture contributes to cheese yield directly by weight and indirectly by containing soluble solids. The contribution from fat results in moisture content increasing with increased fat content. On its own, fat has low water-holding capacity, though it impedes syneresis when occluded in the casein network. Fat globules physically limit the aggregation of paracasein, resulting in less matrix contraction and reduced moisture expulsion (Al-Baarri *et al.*, 2018).

According to Ogunlade *et al.* (2019), pH also has an influence on yield. This can be explained through the effect that pH has on the casein matrix. At pH values above 5.1, the casein matrix is in intact form, whereas at pH below 4.8, casein will be hydrolyzed. A disintegrated, looser matrix can influence the yield and the texture of cheese analogue because of high water levels can maintain texture. This cheese analogue has a pH of 4.8 to 6.0; as such, pH does not significantly affect yield in this case.

Increasing the concentration of lime extract decreases the pH of cheese analogue; however, the addition of papain and maltodextrin does not affect the pH value. The lowest pH value observed in this study for cheese analogue was 4.8, while the highest value was 6.3 (Figure 2).




Figure 2. The pH of cheese analogue was influenced by the concentration of papain and lime extract

The effect of the lime extract on pH was lime contains citric and ascorbic acid. The citric acid content in lime has an acidic pH range of 2.48-2.5 (Chaudhari and Pathan, 2015). According to Grassi *et al.* (2014), if milk—which has a pH range of 6.5-6.8—and the lime extract is added in increasing concentrations, it will reduce the pH of the milk to near the isoelectric point. Decreasing the pH of milk causes the release of calcium ions from calcium caseinate due to the availability of increasing H ions, which can break down calcium phosphate compounds. The rupture of calcium phosphate compounds causes the stability of casein to falter so that coagulation occurs.

Maltodextrin has no effect on pH. According to Farimin and Nordin (2009), maltodextrin originates from oligosaccharide compounds that contain a large hydroxyl (OH) group capable of neutralizing acidic properties. If the pH value is neutral, the process of milk casein agglomeration will be inhibited because protein agglomeration can only occur under acidic conditions. However, in this study, pH values tended to be acidic; therefore, the casein protein could still agglomerate.

Soluble solids consist of sugars, pigments, organic acids, and proteins. The soluble solid range of cheese analogue was 11-18°Brix. The addition of maltodextrin was in line with an increase in soluble solids (Figure 3). Maltodextrin includes fillers that have a high total soluble solid value. According to Jittanit *et al.* (2010), the properties possessed by maltodextrin are rapidly dispersed and have high solubility that causes high levels of soluble solids. Therefore, an increase in the concentration of maltodextrin will increase soluble solids of cheese. This is congruent with results from Farimin and Nordin (2009) in the drying of purple sweet potato flour, which exhibited an increase in soluble solids due to maltodextrin being able to disperse in water.

Increased papain concentration increases moisture content, as shown in Figure 4. The moisture content of cheese analogue ranged from 63.5 to 68.3%. According to Codex 275-2011, cream cheese has a high moisture content (over 67%). Papain cuts the peptide bonds between phenyl and methionine in the kappa-casein, thereby damaging the structure and producing para-kappa casein, which has a hydrophobic portion (Farimin and Nordin, 2009). This hydrophobic portion is insoluble in water; therefore, as the concentration of papain increases, water content should be lower due to the peptide bonds being increasingly cut. However, the work of the papain enzyme also depends on the substrate (casein). Therefore, if casein content is low, a small amount of cut peptide bonds causes the water content to be high. Water content can also be influenced by other




Figure 3. Soluble solids in cheese analogue from corn extract was influenced by the papain and maltodextrin concentration

factors, namely pasteurization. According to Rizzolo and Cortellino (2017), the pasteurization process results in a decrease in moisture content during the process, as the water content in milk will evaporate.

Cheese analogues exhibited protein content ranging from 7.05 to 18.18%. Notably, increasing the concentration of lime extract increased the soluble protein content (Figure 5). Adding lime extract to protein caused cheese casein to agglomerate. When lime is added, the pH of the milk is reduced, and the casein coagulation process occurs. According to Shah *et al.* (2010), casein is easy to settle at the isoelectric point (pH 4.6-5.0). The addition of acid results in the addition of H^+ ions, which will neutralize proteins and achieve isoelectric pH, thus making the proteins hydrophobic (Ah *et al.*, 2015). At the isoelectric point, water is separated and protein will bind between its own contents, forming a fold and resulting in coagulation.

Protein solubility will increase if excessive acid treatment is performed, since according to Pang *et al.* (2016)—positive ions in acids that cause proteins (initially neutral or zero) to be positively charged. The




Figure 5. The protein content of cheese analogue was influenced by lime extract and maltodextrin concentration




Figure 4. The moisture content of cheese analogue was influenced by papain and maltodextrin concentration

further that the acidity of the protein solution is from the isoelectric point, the more soluble it will be. The observed protein level (22.9%) of cheese analogue produced by regulating the acidity of lime juice was lower than that of cheese produced from cow's milk (Razig and Babiker, 2009). This is because cow milk has a higher protein content of 16.5% (Colmenero and Broderick, 2006) compared to sweet corn (5.9%) (Almodares *et al.*, 2009). The greater the addition of maltodextrin, the more soluble protein will present. This is due to maltodextrin having good consistency and stability, as it can bind water-soluble proteins even in small amounts (Akhtar and Dickinson, 2007).

The lipid content of cheese analogue was 2.34 to 5.95%. Figure 6 shows that increasing the extract of lime will increase the fat content of cheese analogue. According to Seth and Bajwa (2015), the addition of acid during cheese production will increase fat-binding capacity. This is because the addition of acid can lead to the coagulation of the proteins that form fat globules. The protein coagulation process occurs in the presence of acids, resulting in an increased fat-binding capacity as




Figure 6. The fat content of cheese analogue was influenced by concentrations of lime extract and papain

more acid is added.

Furthermore, fat content tends to increase with the addition of papain (Figure 6). According to Fernández-Lucas *et al.* (2017), papain exhibits proteolytic activity during the milk coagulation process, which causes the phospholipid-protein layer to be damaged. As a result, fat globules will be trapped at the point of protein clumping and eventually unite with the curd.

According to Thorning *et al.* (2016), the fat content found in cheese is heavily dependent on the type of milk and other ingredients used as raw material for cheese production. The low-fat content of cheese analogue in this study is due to the basic ingredients of cheese that do use cow's milk but rather corn extract. According to Aini, Prihananto, Wijonarko, Astuti *et al.* (2017), sweet corn has a fat content of 1%, while milk fat content is 3.4%. This causes the fat content of corn-based cheese analogue to be relatively low. According to Mohamed *et al.* (2017), substituting milk fat with vegetable fat changes the nutritional profile of cheese analogue, making it lower in cholesterol and the composition of saturated and unsaturated fatty acids.

The cheese analogue in this study included low-fat cheese, which has lower fat content compared to full-fat cheese (Wang *et al.*, 2016). Based on wet weight, full-fat cheese in fresh form has a fat content of 24.5%. Based on the fat content of cheese analogue, it can be concluded that fat content is generally in accordance with the USDA (2001) suggestion (below 5%).

This cheese analogue has 472-1146 ppm of beta-carotene. Moreover, levels of maltodextrin, lime, and papain do not affect the levels of beta-carotene in cheese analogue. This level of beta-carotene is comparable to Kareish cheese, which is fortified with 1 to 1.5% of *Spirulina platensis*, which has beta-carotene levels of 600-1020 ppm (Mohamed and Darwish, 2017).

3.2 Formula optimization and verification of the optimum formula

Based on the optimization phase, three formulae were obtained that had the highest desirability value and according to the objectives of this study. Cheese analogue formula I was made with the addition of 15% maltodextrin, 2.283% lime extract, and 0.022% papain. Formula II was produced from sweet corn with the addition of 15% maltodextrin, 2.178% lime extract, and 0.022% papain. Formula III made with the addition of 15% maltodextrin, 2.39% lime extract, and 0.022% papain.

The testing accuracy results for these three formulae are presented in Table 1. Based on the verification results of Formula I, II and III, all responses were deemed appropriate and within the prediction interval range. The results of sensory testing are presented in Table 2. The cheese with a minimal yellow color was the best choice among panelists. Panelists chose formula II as the best color, since it has a yellowish-white color. According to Ben-Amira *et al.* (2017), cheese made from cow's milk without coloring will produce cheese that is yellowish-white. Yellow comes from fat-soluble carotene pigments. However, the fat content of cheese analogue is low, thus the fat-soluble carotene pigments levels are also low. Therefore, white is the more dominant color among the cheese analogues.

Products with the best aroma are those possessing the maximum or strongest distinctive cheese aroma. Based on the analysis of various rankings, results from tests of aroma were not significantly different among the three optimum formulae. The distinctive aroma of cheese originates from the fatty acids found in milk (Villalobos-Chaparro *et al.*, 2018). According to Everard *et al.* (2007), a change in dairy products such as cheese is caused by the fermentation of lactose, citrate, and other organic compounds into various acids, esters, alcohols, flavor-forming compounds, and volatile aromas. However, the use of lime ingredients can affect the distinctive aroma of spreadable cheese analogue.

Table 1. Verification results for the optimal formula

Response	Formula I			Formula II			Formula III		
	Actual data	Predictive value	Verification	Actual data	Predictive value	Verification	Actual data	Predictive value	Verification
Moisture Content (%)	65.692	66.33	Appropriate	65.329	66.388	Appropriate	62.083	64.001	Appropriate
Yield (%)	21.178	21.757	Appropriate	20.314	21.745	Appropriate	20.456	21.778	Appropriate
Soluble solids (°Brix)	14	16	Appropriate	14	16	Appropriate	14	16	Appropriate
Soluble protein (% db)	11.223	13.439	Appropriate	13.516	13.255	Appropriate	14.078	13.593	Appropriate
pH	5.4	5.4	Appropriate	5.4	5.5	Appropriate	5.3	5.4	Appropriate
Fat (% db)	4.344	3.718	Appropriate	4.553	3.6527	Appropriate	5.428	3.795	Appropriate

Table 2. Test rank results for the three best cheese analogue formulae

Formula	Attribute ^a				
	Color	Aroma	Taste	Spreadability	Preference
I	2.75 ^b	1.85 ^a	2.15 ^a	2.2 ^a	2.3 ^a
II	1 ^a	1.75 ^a	1.65 ^a	1.35 ^a	1.2 ^a
III	2.25 ^b	2.58 ^a	2.2 ^a	2.45 ^a	2.5 ^b

Values in the same column followed by different letters indicate a significant difference in the level of 5% LSD test.

According to Chaudhari and Pathan (2015), lime has a strong aroma and distinctive flavor. In optimum formula II, the concentration of lime is the lowest; thus, the panelists assessed the distinctive cheese aroma as stronger compared to the other optimum formulae.

The products with the best taste are those with the least or lowest sour taste. Based on the analyses on cheese analogue taste, the formulae were not significantly different. However, formula II was ranked as having the best taste. The lime extract contains a distinctive and strong sour taste. In formula II, the concentration of lime is lowest; therefore, the typical taste of lime in this cheese analogue is lower. The effect of the lime extract is not only sour taste, but it also causes a slightly bitter taste. According to Ben-Amira *et al.* (2017), the bitter taste in oranges is influenced by the presence of two compounds, namely flavonoids and limonoids. The main component of flavonoids is naringin, which gives the impression of an astringent to bitter taste. Therefore, the panelists tended to rank optimum formula II as having the best taste.

Products with the best spreadability performance are those possessing the best ease of spreading when applied. Based on the analysis of various ranking tests on the applicable performance of cheese, no significant differences were observed. However, the panelists tended to assess optimum formula II as having a softer texture, making it easier to spread than the other two formulae.

Based on the complete ranking test, panelists judged optimum formula II as the preferred spreadable cheese analogue. This preference was based on various sensory attributes, for which most panelists chose optimum formula II as being optimal. Determination of the optimal formula based on chemical and sensory properties was performed using the effectiveness index. Based on this index, the total value of each formula is presented in Table 3. With the highest total value of 1.504, optimum formula II was chosen as the best formula. Formula II has the lowest concentration of lime extract, an extract that results in a distinctive taste that can reduce consumer acceptance of cheese analogue. Therefore, the selection of formula II is appropriate since the taste of lime in regular cheese is low, while other physicochemical and sensory properties are as expected.

Thus, sweet corn extract has the potential to be developed as a raw material for the production of cheese analogue.

Table 3. Total values for optimum formula

Characteristics	Formula I	Formula II	Formula III
Chemical properties	0.63	0.504	0.406
Sensory properties	0.271	1	0.037
Total	0.901	1.504	0.443

The advantage of cheese analogue from sweet corn is lower fat and cholesterol content compared to cow's milk cheese, as shown in Table 4. Another advantage of analog cheese from corn is that it has higher carotene (544.4 mg/g) than commercial cheese (5.32 mg/g). The protein content of products has the same protein content, which is 7.1 and 6.3 %. These products can be designed to fulfill special dietary needs through changes in the formulation (e.g. lactose-free, low calorie, low saturated fat, and low cholesterol) (Chavan and Jana, 2007).

Table 4. Characteristics of selected spreadable cheese analogue and s commercial spreadable cheese (from cow's milk)

	Cheese analogues	Commercial cheese
Protein (%)	7.1	6.3
Fat (%)	4.55 ^b	24.53 ^a
Total carotene (mg/g)	544.4 ^a	5.32 ^b
Cholesterol (mg/g)	0.02 ^b	0.19 ^a

Values in the same column followed by different letters indicate a significant difference in the level of 5% LSD test.

4. Conclusion

The optimal formula for cheese analogue used 2.178% lime extract, 0.022% papain, and 15% maltodextrin, and provided the highest-ranked responses for chemical and sensory properties. This cheese analogue has yields of 21.178%, pH of 5.4, 14% of total soluble solids, moisture content of 65.3%, protein content of 7.1%, total carotene of 544.4 mg/g, fat content 4.55%, and cholesterol 0.02 mg/g. This cheese analogue has minimal yellow sensory properties, a distinctive strong cheese aroma, a minimal sour taste, while it was also the easiest analogue to apply due to its soft texture. Thus, sweet corn has the potential for use as a raw material for producing low-fat cheese.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgments

The research work was funded by Jenderal Soedirman University through *Riset Unggulan Terapan* [Grant No 267/UN23.14/PN/2019]. We also give acknowledgment to Ministry of Research, Technology and Higher Education for the funding in the *Program Bantuan Seminar Luar Negeri* 2019.

References

- Ah, J., Rahul, S., Kd, A. and Dhanraj, P. (2015). Influence of rennet casein levels on the chemical, baking and sensory quality of mozzarella cheese analogue. *Journal of Dairy, Veterinary and Animal Research*, 2(3), 39–45. <https://doi.org/10.15406/jdvar.2015.02.00039>
- Aini, N., Prihananto, V., Sustriawan, B., Romadhon, D. and Ramadhan, R.N. (2019). The formulation of cheese analogue from sweet corn extract. *International Journal of Food Science*, 2019, 1–8. <https://doi.org/10.1155/2019/8624835>
- Aini, N., Sustriawan, B., Prihananto, V. and Heryanti, T. (2019). Characteristics of cheese analogue from corn extract added by papain and pineapple extract. *IOP Conference Series: Earth and Environmental Science*, 255, 012016. <https://doi.org/10.1088/1755-1315/255/1/012016>
- Aini, N., Prihananto, V., Wijonarko, G., Astuti, Y., Maulina, M.R. and Muthmainah, M. (2017). Quality deterioration and shelf life estimation of corn yogurt was packaged by glass bottle. *Advanced Science Letters*, 23(6), 5796–5798. <https://doi.org/10.1166/asl.2017.8835>
- Akhtar, M. and Dickinson, E. (2007). Whey protein–maltodextrin conjugates as emulsifying agents: An alternative to gum arabic. *Food Hydrocolloids*, 21(4), 607–616. <https://doi.org/10.1016/j.foodhyd.2005.07.014>
- Al-Baarri, A.N., Legowo, A.M., Arum, S.K. and Hayakawa, S. (2018). Extending shelf life of Indonesian soft milk cheese (dangke) by lactoperoxidase system and lysozyme. *International Journal of Food Science*, 2018, 1–7. <https://doi.org/10.1155/2018/4305395>
- Almodares, A., Jafarinia, M. and Hadi, M.R. (2009). The effects of nitrogen fertilizer on chemical compositions in corn and sweet sorghum. *American-Eurasian Journal of Agricultural and Environmental Science*, 6(4), 441–446.
- AOAC. (2005). Official methods of analysis of the Association of Official Agricultural Chemists International. USA: AOAC.
- Badem, A. and Ucar, G. (2016). Cheese analogues: review article. *Journal of Food and Dairy Technology*, 4(3), 44–48.
- Ben-Amira, A., Besbes, S., Attia, H. and Blecker, C. (2017). Milk-clotting properties of plant rennets and their enzymatic, rheological, and sensory role in cheese making: A review. *International Journal of Food Properties*, 20(Suppl. 1), 1–18. <https://doi.org/10.1080/10942912.2017.1289959>
- Bezerra, M.A., Santelli, R.E., Oliveira, E.P., Villar, L.S. and Escalera, L.A. (2008). Response surface methodology (RSM) as a tool for optimization in analytical chemistry. *Talanta*, 76(5), 965–977. <https://doi.org/10.1016/J.TALANTA.2008.05.019>
- Chaudhari, S.N. and Pathan, A.H. (2015). Study of development of pome-lime nectar fortified with sweet lime peel powder. *International Journal of Science and Research*, 4(10), 2013–2016.
- Chavan, R.S. and Jana, A. (2007). Cheese substitutes: an alternative to natural cheese: a review. *International Journal of Food Science, Technology and Nutrition*, 2(2), 25–39.
- Colmenero, J.J.O. and Broderick, G.A. (2006). Effect of dietary crude protein concentration on milk production and nitrogen utilization in lactating dairy cows. *Journal of Dairy Science*, 89(5), 1704–1712. [https://doi.org/10.3168/jds.S0022-0302\(06\)72238-X](https://doi.org/10.3168/jds.S0022-0302(06)72238-X)
- Cunha, C.R., Dias, A.I. and Viotto, W.H. (2010). Microstructure, texture, colour and sensory evaluation of a spreadable processed cheese analogue made with vegetable fat. *Food Research International*, 43(3), 723–729. <https://doi.org/10.1016/j.foodres.2009.11.009>
- DeGarmo, E.P., Black, J.T. and Kohser, R.A. (2003). *Materials and Processes in Manufacturing*. 9th ed. New York, USA: John Wiley and Sons Inc.
- Dhanraj, P., Jana, A., Modha, H. and Aparnathi, K.D. (2017). Influence of using a blend of rennet casein and whey protein concentrate as protein source on the quality of Mozzarella cheese analogue. *Journal of Food Science and Technology*, 54(3), 822–831. <https://doi.org/10.1007/s13197-017-2528-5>
- Everard, C.D., O'Donnell, C.P., O'Callaghan, D.J., Sheehan, E.M., Delahunty, C.M., O'Kennedy, B.T. and Howard, V. (2007). Prediction of sensory textural properties from rheological analysis for process cheeses varying in emulsifying salt, protein and moisture contents. *Journal of the Science of Food and Agriculture*, 87(4), 641–650. <https://doi.org/10.1007/s13197-017-2528-5>

doi.org/10.1002/jsfa.2756

- Farimin, A.O.A. and Nordin, E. (2009). Physical properties of powdered roselle-pineapple juice - Effects of maltodextrin presented at the National Conference on Postgraduate Research (NCON-PGR) 2009. Pahang: Universiti Malaysia Pahang.
- Fernández-Lucas, J., Castañeda, D. and Hormigo, D. (2017). New trends for a classical enzyme: Papain, a biotechnological success story in the food industry. *Trends in Food Science and Technology*, 68, 91–101. <https://doi.org/10.1016/j.tifs.2017.08.017>
- González, M.L., Sánchez, H.C., Franco, F.M.J., Güemes, V.N. and Soto, S.S. (2018). Physical, chemical and texture characteristics of aro cheese. *Food Research*, 2(1), 61–67. [https://doi.org/10.26656/fr.2017.2\(1\).211](https://doi.org/10.26656/fr.2017.2(1).211)
- Grassi, S., Alamprese, C., Bono, V., Casiraghi, E. and Amigo, J.M. (2014). Modelling milk lactic acid fermentation using multivariate curve resolution-alternating least squares (MCR-ALS). *Food and Bioprocess Technology*, 7(6), 1819–1829. <https://doi.org/10.1007/s11947-013-1189-2>
- Hashim, M.M., Dong, M., Iqbal, M.F., Li, W. and Chen, X. (2011). Ginger protease used as coagulant enhances the proteolysis and sensory quality of Peshawari cheese compared to calf rennet. *Dairy Science and Technology*, 91(4), 431–440. <https://doi.org/10.1007/s13594-011-0021-x>
- Hjalmarsson, M. (2015). The effect of vegetable fat on cheese yield and cheese properties. Uppsala: Swedish University of Agriculture, MSc. Thesis.
- Hoffmann, J.F., Zandoná, G.P., dos Santos, P.S., Dallmann, C.M., Madruga, F.B., Rombaldi, C.V. and Chaves, F.C. (2017). Stability of bioactive compounds in butiá (*Butia odorata*) fruit pulp and nectar. *Food Chemistry*, 237, 638–644. <https://doi.org/10.1016/j.foodchem.2017.05.154>
- Jittanit, W., Niti-Att, S. and Techanuntachai, O. (2010). Study of spray drying of pineapple juice using maltodextrin as an adjunct. *Chiang Mai Journal of Science*, 37(3), 498–506.
- Lee, S.K., Klostermeyer, H. and Anema, S.G. (2015). Effect of fat and protein-in-water concentrations on the properties of model processed cheese. *International Dairy Journal*, 50, 15–23. <https://doi.org/10.1016/j.idairyj.2015.06.001>
- Liu, H., Xu, X.M. and Guo, S.D. (2008). Comparison of full-fat and low-fat cheese analogues with or without pectin gel through microstructure, texture, rheology, thermal and sensory analysis. *International Journal of Food Science and Technology*, 43(9), 1581–1592. <https://doi.org/10.1111/j.1365-2621.2007.01616.x>
- Mahajan, R.T. and Chaudhari, G.M. (2014). Plant latex as vegetable source for milk clotting enzymes and their use in cheese preparation. *International Journal of Advanced Research*, 2(5), 1173–1181.
- Mohamed, A. and Darwish, I. (2017). Physicochemical properties, bioactive compounds and antioxidant activity of Kareish Cheese fortified with spirulina platensis. *World Journal of Dairy and Food Sciences*, 12(2), 71–78. <https://doi.org/10.5829/idosi.wjdfs.2017.71.78>
- Mohamed, A.G., Abo-El-Khair, B.E. and Shalaby, S. (2017). Quality of novel healthy processed cheese analogue enhanced with marine microalgae *Chlorella vulgaris* biomass. *World Applied Sciences Journal*, 23(7), 914–925. <https://doi.org/10.5829/idosi.wasj.2013.23.07.13122>
- Mohamed, A.G. and Shalaby, S.M. (2016). Texture, chemical properties and sensory evaluation of a spreadable processed cheese analogue made with apricot pulp (*Prunus armeniaca* L.). *International Journal of Dairy Science*, 11(2), 61–68. <https://doi.org/10.3923/ijds.2016.61.68>
- Moser, P., Telis, V.R.N., de Andrade Neves, N., García-Romero, E., Gómez-Alonso, S. and Hermosín-Gutiérrez, I. (2017). Storage stability of phenolic compounds in powdered BRS Violeta grape juice microencapsulated with protein and maltodextrin blends. *Food Chemistry*, 214, 308–318. <https://doi.org/10.1016/j.foodchem.2016.07.081>
- Oboh, F.O. and Imafidon, J. (2018). Antioxidant and sensory properties of new beverage formulations composed of palm sugar, *Aframomum melegueta*, and citric acid. *Beverages*, 4(3), 59–72. <https://doi.org/10.3390/beverages4030059>
- Ogunlade, A.O., Oyetayo, V.O. and Ojokoh, A.O. (2019). Effect of different biocoagulants on the microbial quality and mineral composition of west African cheese produced from sheep milk. *Food Research*, 3(3), 272–279. [https://doi.org/10.26656/fr.2017.3\(3\).238](https://doi.org/10.26656/fr.2017.3(3).238)
- O’Riordan, E.D., Duggan, E., O’Sullivan, M. and Noronha, N. (2011). Production of Analogue Cheeses. In Tamime, A.Y. (Ed.) *Processed Cheese and Analogues*, p. 219–244. United Kingdom: Wiley Online Library. <https://doi.org/10.1002/9781444341850.ch9>
- Osman, H. and Chin, Y.K. (2006). Comparative sensitivities of cholesterol analysis using GC, HPLC and spectrophotometric methods. *The Malaysian Journal of Analytical Sciences*, 10(2), 205–210. https://doi.org/http://www.ukm.my/mjas/v10_n2/29_osman.pdf

- Pang, Z., Deeth, H., Prakash, S. and Bansal, N. (2016). Development of rheological and sensory properties of combinations of milk proteins and gelling polysaccharides as potential gelatin replacements in the manufacture of stirred acid milk gels and yogurt. *Journal of Food Engineering*, 169, 27-37. <https://doi.org/10.1016/j.jfoodeng.2015.08.007>
- Razig, K.A.A. and Babiker, N.A. (2009). Chemical and microbiological properties of Sudanese white soft cheese made by direct acidification technique. *Pakistan Journal of Nutrition*, 8(8), 1138–1143. <https://doi.org/10.3923/pjn.2009.1138.1143>
- Rizzolo, A. and Cortellino, G. (2017). Ricotta cheese whey-fruit-based beverages: pasteurization effects on antioxidant composition and color. *Beverages*, 3 (1), 15. <https://doi.org/10.3390/beverages3010015>
- Seth, K. and Bajwa, U. (2015). Effect of acidulants on the recovery of milk constituents and quality of Mozzarella processed cheese. *Journal of Food Science and Technology*, 52(3), 1561–1569. <https://doi.org/10.1007/s13197-013-1176-7>
- Shah, R., Jana, A.H., Aparnathi, K.D. and Prajapati, P.S. (2010). Process standardization for rennet casein based Mozzarella cheese analogue. *Journal of Food Science and Technology*, 47(5), 574–578. <https://doi.org/10.1007/s13197-010-0104-3>
- Sumarmono, J. and Suhartati, F.M. (2016). Yield dan komposisi keju lunak (soft cheese) yang dibuat dengan teknik direct acidification menggunakan ekstrak buah lokal. *Jurnal Aplikasi Teknologi Pangan*, 1(3), 65–68. [In Bahasa Indonesia].
- Thorning, T.K., Raben, A., Tholstrup, T., Soedaman-Muthu, S.S., Given, I. and Astrup, A. (2016). Milk and dairy products: Good or bad for human health? An assessment of the totality of scientific evidence. *Food and Nutrition Research*, 60(1), 32527. <https://doi.org/10.3402/fnr.v60.32527>
- Tuntragul, S., Surapat, S. and Hongsprabhas, P. (2010). Influence of rice bran oil and rice flours on physicochemical properties of a mozzarella cheese analog. *Kasetsart Journal (Natural Science)*, 44, 924–934.
- USDA. (2001). Specification for Cottage Cheese and Dry Curd Cottage Cheese. Agriculture marketing Service USDA. USA: USDA.
- Villalobos-Chaparro, S., Salas-Muñoz, E., Gutiérrez-Méndez, N. and Nevárez-Moorillón, G.V. (2018). Sensory profile of Chihuahua cheese manufactured from raw milk. *International Journal of Food Science*, 2018, 1–7. <https://doi.org/10.1155/2018/8494105>
- Wang, F., Tong, Q., Luo, J., Xu, Y. and Ren, F. (2016). Effect of carrageenan on physicochemical and functional properties of low fat Colby Cheese. *Journal of Food Science*, 81(8), E1949–E1955. <https://doi.org/10.1111/1750-3841.13369>



Enhance your knowledge

Gain a holistic overview of property analysis with a flexible online learning experience.

MIT SA+P

Open

Food Research

COUNTRY

Malaysia



Universities and
research institutions in
Malaysia

SUBJECT AREA AND CATEGORY

Agricultural and
Biological Sciences
Food Science

PUBLISHER

H-INDEX

7

PUBLICATION TYPE

Journals

ISSN

25502166

COVERAGE

2017-2020

INFORMATION

[Homepage](#)

[How to publish in this
journal](#)

[foodresearch.my@outlook.
com](mailto:foodresearch.my@outlook.com)

SCOPE

Food Research welcomes submissions that are relevant to food science and technology, food service management, nutrition, nutraceuticals, food innovation and agriculture food science. Manuscript types accepted are reviews, short reviews, original research articles and short communications that discusses on the current research and scientific views of the following scope: Food science and food chemistry; Food technology, food processing, and food engineering; Food safety and quality – microbiological and chemical; Sensory, habits, consumer behaviour/practice and preference; Nutrition and dietetics; Nutraceuticals and functional food/ingredients; Food service management; Food trends, innovation and business; Post-harvest and agribusiness; Food security; Food packaging.



Proses Pasang Loker Jadi Mudah

Cukup WA ke tim JobStreet, kami siap bantu pasang iklan loker perusahaan Anda.
JobStreet Indonesia

Submit Your Manuscript With Us

Accepting Research Focused on Wireless Power Transf

Hindawi

Submit Your Manuscript With Us

Accepting Research Focused on Wireless Power Transf

Hindawi

Submit Your Manu:


Accepting Research Focused or

Hindawi

FIND SIMILAR JOURNALS ?

1 International Food Research Journal MYS	2 Food Science and Nutrition GBR	3 Carpathian Journal of Food Science and Technology ROU	4 Internationa Science EGY
--	--	--	-------------------------------------





free tool.

Get it



Submit Your Manuscript With Us

Accepting Research Focused on Wireless Power Transfer. Publish

Hindawi

Metrics based on Scopus® data as of April 2021

T **Tamiur Yazew** 2 months ago

Dear,

Thank you for your updated and quality journal.

I have submitted two manuscripts in the Food research journal. The editor of this journal sent me the acceptance letter for the two manuscripts. However, they asked me to pay charge for the manuscripts. I am also working as a reviewer of this journal. I have edited two papers and sent it to the editor. I am currently reviewing a paper and it is ready to send back to the editor of this journal.

But I am from poor country, Ethiopia and I am unable to pay it. The condition in Ethiopia may also not allow me due to lack of accessibility of this 150 USD.

So, please would you help me by considering my issue into consideration!

reply



Melanie Ortiz 2 months ago

SCImago Team

Dear Tamiur, thank you very much for your comment. Unfortunately, we cannot help you with your request, we suggest you contact the journal's editorial staff so they could inform you more deeply. Best Regards, SCImago Team

K **Kamal** 10 months ago

Hi, Editorial Team members,

I would like to know the topic related to the "hygienic practices along the supply chain of fisheries"

Proses Pasang Loker Jadi Mudah

Cukup WA ke tim JobStreet, kami siap bantu pasang iklan loker perusahaan Anda.
JobStreet Indonesia



KORESPONDENSI



Gmail



foodresearch.my@outlook.com



58

FR-2019-395 Inbox x



Food Research <foodresearch.my@outlook.com>

to me

Dear Ain,

This message is to acknowledge the receipt of the above manuscript that you submitted with manuscript ID number in any correspondence with the Food Research Editorial C

Your paper will be reviewed by three or more reviewers assigned by the Food R and editor's comments will be then made available via email attached file. You ca

If your manuscript is accepted for publication, Food Research editorial office will c

Thank you very much for submitting your manuscript to Food Research.

Sincerely,

Professor Dr. Son Radu

Chief Editor

Email: foodresearch.my@outlook.com

From: nur.aini 1 <nur.aini@unsoed.ac.id>

Sent: Saturday, 23 November, 2019 10:34 PM

To: foodresearch.my@outlook.com <foodresearch.my@outlook.com>



Gmail



foodresearch.my@outlook.com



58

FR-2019-395 Inbox x



Food Research <foodresearch.my@outlook.com>

to me

Dear Aini,

Manuscript FR-2019-395 entitled " Formulation of low-fat cheese analogue form sw comments of the reviewer(s) are included in the attached file.

The reviewer(s) have recommended publication, but also suggest some revisior revised manuscript is prepared, please send it back to me for further processing.

Because we are trying to facilitate timely publication of manuscripts submitted to revision by this date, please let us know.

Once again, thank you for submitting your manuscript to Food Research and I loc

Sincerely,

Professor Dr. Son Radu

foodresearch.my@outlook.com

Chief Editor, Food Research

3 Attachments





Gmail



foodresearch.my@outlook.com



58

FR-2019-395 - Decision on your manuscript Inbox x



Food Research <foodresearch.my@outlook.com>

to me

Dear Dr. Nur Aini,

It is a pleasure to accept your manuscript for publication in Food Research journal. Please

Please note that all accepted manuscripts are subjected to Article Processing Charges (APC) within five (5) working days. Once we have received the form, your article will be transferred to the journal.

Thank you for your fine contribution. We look forward to your continued contributions to the journal.

Sincerely,

Dr. Vivian New

Editor

Food Research

From: nur.aini 1 <nur.aini@unsoed.ac.id>

Sent: Monday, 13 January, 2020 1:40 PM

To: foodresearch.my@outlook.com <foodresearch.my@outlook.com>

Subject: Revision of article FR-2019-395

Dear Prof Son Radu,

Thank you very much for the feedback to my article.

Hereby submit a revision of my article for Food Research

There are 3 files I sent, i.e.

1. Articles that have been revised



Gmail



foodresearch.my@outlook.com



58

MANUSCRIPT SUBMISSION



nur.aini 1 <nur.aini@unsoed.ac.id>

to foodresearch.my

Dear Prof Son Radu,

I hereby submit an article to be published in Food Research.

There are 3 files,

1. Manuscript
2. Cover letter
3. Manuscript submission form

We really hope and grateful that this article can be a reference for similar articles as they

Sincerely

Dr. Nur Aini

Department of Food Technology,

Jenderal Soedirman University

Purwokerto, Indonesia

3 Attachments

