



Open Agriculture

ISSN: 2391-9531

EDITORIAL OVERVIEW LATESTISSUE ISSUES RANKING SUBMIT

Editorial

Editors-in-Chief Hans R. Herren, Millenium Institute, USA Vijay Singh, Texas A and M University, USA

Managing Editor Oskar Szczepaniak, De Gruyter oskar.szczepaniak@degruyter.com

Editorial Advisory Board

Filiberto Altobelli, Council for Agricultural Research and Analysis of Agricultural Economics (CRA), Italy

Fred Asiegbu, University of Helsinki, Finland

Hassan Azaizeh, Institute of Applied Research, Galilee Society, Israel

Zohar Ben-Asher, EuCRF European Centre for Research & Financing, Israel

Furio Brighendi, University of Parma, Italy

Mladen Brnčić, University of Zagreb, Croatia

Katerina Demnerova, Institute of Chemical Technology, Czech Republic

Santa Olga Cacciola, University of Catania, Italy

Gary Dykes, Monash University, Australia

Hernâni Gerós, Universidade do Minho, Portugal

Martin Greimel, Austrian Ministry for Agriculture, Forestry, Environment and Water Management, Austria

Zoran Herceg, University of Zagreb, Zagreb, Croatia

Julie Ingram, University of Gloucestershire, UK

Anet Režek Jambrak, University of Zagreb, Zagreb, Croatia

Manuel Jamilena, University of Almería, Spain

Anders H. Karlsson, University of Copenhagen, Denmark

Roy Kennedy, Warwickshire Colleges, UK

Ferando Lidon, New University of Lisbon, Portugal

Ornella Mikuš, University of Zagreb Faculty of Agriculture, Croatia

Piergiuseppe Morone, Unitelma-Sapienza, University of Rome, Italy

Peter Morgan, Aquamor Pvt Ltd., Zimbabwe

Theodoor Niewold, Katholieke Universiteit Leuven, Belgium

Norman Olson, University of Wisconsin-Madison, USA

Ari Pappinen, University of Eastern Finland, Finland

Paola Pittia, University of Teramo, Italy

Richard E. Plant, University of California, Davis, USA

Saskia van Ruth, Wageningen University and Research Centre, Netherlands

Andrea Vannini, Università degli Studi della Tuscia, Italy

Rimantas Venskutonis, Kaunas University of Technology, Lithuania

Stefan Vidal, Universitat Gottingen, Germany

Wim de Vries, Wageningen University and Research Centre, Netherlands

Xinmin Zhan, National University of Ireland, Ireland

Editors

Agro-ecology

Clifford S. Gold, Independent consultant, USA

Mehmood Ali Noor, Chinese Academy of Agricultural Sciences, China

Sergio de los Santos Villalobos, Instituto Tecnológico de Sonora, Mexico

Ulrich Schmutz, Coventry University, United Kingdom

Christoph Steiner, University of Kassel, Germany

Debjani Sihi, University of Maryland, Center for Environmental Science, USA

Srdjan Šeremešić, University of Novi Sad, Serbia

Priit Tammeorg, University of Helsinki, Finland

Sebastian Wolfrum, Technische Universität München, Germany

Agro-forestry

Mirza Barjees Baig, King Saud University, Kingdom of Saudi Arabia (agricultural education, rural development, agricultural information and communication)

Rodolfo Alves Barbosa, University of Viçosa, Brazil

Bohdan Lojka, Czech University of Life Sciences Prague, Czech Republic

Mohammad Valipour, Payame Noor University, Iran

Pedro Gonçalves Vaz, CEABN - Centre of Applied Ecology "Prof. Baeta Neves", Portugal

Sebastian Wolfrum, Technische Universität München, Germany

Soil eciance

Agustín Alesso, National University of the Littoral, Argentina

Elena Aydin, Slovak University of Agriculture, Slovakia (soil physics)

Biswanath Dari, University of Florida, USA

Rahul Datta, Mendel University In Brno, Czech Republic

Fabián Fernández-Luqueño, Cinvestav Saltillo, Mexico

Shahid Hussain, Bahauddin Zakariya University, Pakistan

Charanjit Singh Kahlon, Independent consultant, USA (soil biology)

Noémi Kappel, Corvinus University of Budapest, Hungary (soil physics)

Mario Lira, Universidade Federal Rural de Pernambco, Brazil

Sergio de los Santos Villalobos, Instituto Tecnológico de Sonora, Mexico

Riccardo Scalenghe, Università degli Studi di Palermo, Italy

Andres Seijo, Universidade do Porto Centro Interdisciplinar de Investigação Marinha e Ambiental, Portugal

Parmodh Sharma, New Mexico State University, USA

Debjani Sihi, University of Maryland, Center for Environmental Science, USA (soil biology)

Christoph Steiner, University of Kassel, Germany (soil biology, biochar)

Priit Tammeorg, University of Helsinki, Finland (soil chemistry, soil physics, biochar)

Mohammad Valipour, Payame Noor University, Iran

Sebastian Wolfrum, Technische Universität München, Germany (soil biology, organic matter)

Plant science

Bartosz Adamczyk, Natural Resources Institute Finland (Luke), Finland

Duska Delic, University of Banjaluka, Republic of Srpska (plant pathology)

Mahendra Dia, North Carolina State University, NC, USA (plant breeding, plant biotechnology, plant genetics)

Clovis Douanla-Meli, Julius Kühn-Institut, Germany (plant pathology, plant ecology)

Baoguo Du, University of Freiburg, Germany (plant physiology)

Malleshwari Gelli, SIBS, DuPont-Pioneer, USA

Clifford S. Gold, Independent consultant, USA (plant ecology, plant pathology)

Dr. Thomas Graham, University of Guelph, Guelph, Canada

Alireza Haghighi Hasanalideh, CWARice, Iran

Vinod Jakkula, Bayer U.S - Crop Science, USA

Charanjit Singh Kahlon, Independent consultant, USA (plant physiology, plant ecology, plant breeding; agronomy)

Hristofor Kirchev, Agricultural University, Bulgaria

Gunars Lacis, Latvia State Institute of Fruit-Growing, Latvia (plant breeding)

Mehmood Ali Noor, Chinese Academy of Agricultural Sciences, China

Mehdi Rahimi, Graduate University of Advanced Technology, Iran

Sergio de los Santos Villalobos, Instituto Tecnológicode Sonora, Mexico

Daljit Singh, Bayer - Crop Science, Monsanto Company, USA

Konstantinos Spanos, Forest Research Institute, Greece (plant pathology, plant breeding)

Khalid Ul Rehman Hakeem, King Abdulaziz University, Kingdom of Saudi Arabia

Ramesh Vetukuri, Swedish University of Agricultural Sciences, Sweden

Horticulture

Pinaki Acharyya, Central Institute for Arid Horticulture, India (olericulture, pomology, postharvest physiology, floriculture)

Cédric Camps, Agroscope, Switzerland (pomology, postharvest physiology)

Mário Cunha, University of Porto, Portugal (viticulture)

Boris Duralija, University of Zagreb, Croatia (pomology)

Ebrahim Hadavi, Islamic Azad University, Iran

Lorentz Jäntschi, Universitatea Tehnica din Cluj-Napoca, Romania

Noémi Kappel, Corvinus University of Budapest, Hungary (olericulture)

Sandeep Rana, Bayer U.S. - Crop Science, USA

Christoph Steiner, University of Kassel, Germany

Forestry

Alina M. Balu, Universidad de Cordoba, Spain (forest biomass)

Baoguo Du, University of Freiburg, Germany (silviculture)

Carol S.K. Lin, City University of Hong Kong, Hong Kong (forest biomass)

Konstantinos Spanos, Forest Research Institute, Greece (silviculture, arboriculture, conservation and management, biomass, forest genetics and forest tree breeding, forest biodiversity)

Wood technology

Alina M. Balu, Universidad de Cordoba, Spain (wood properties, wood products)

Ramon Gonzalez, J. Rettenmaier und Sohne USA, Cambridge, USA (wood, pulp and paper products)

Jaya Sundaram, United States Department of Agricultural, USA

Animal Science

Michael Boateng, Kwame Nkrumah University of Science and Technology (KNUST), Ghana

Tomislav Mikuš, University of Zagreb, Croatia

Jana Pickova, Swedish University of Agricultural Sciences, Sweden

Arya Sobhakumari, University of California Davis, USA

Gunnar Sundstøl Eriksen, Norwegian Veterinary Institute, Norway

Entomology

Clifford S. Gold, Independent consultant, USA

Aquaculture

Sunil Kadri, Aquaculture Innovation, United Kingdom

Sven Würtz, Leibniz-Institute of Freshwater Ecology and Inland Fisheries, Germany

Jana Pickova, Swedish University of Agricultural Sciences, Sweden

Hydrology, Irrigation

Babak Mohammadi, Lund University, Sweden

Prahlad Jat, University of North Carolina, USA

Elena Kondrlova, Slovak University of Agriculture, Slovakia

Sergio Salazar-Galán, Universidad Nacional de Colombia, Colombia

Ulrich Schmutz, Coventry University, United Kingdom

Biswanath Dari, University of Florida, USA

Babak Vaheddoost, Bursa Technical University, Turkey

Mohammad Valipour, Payame Noor University, Iran

Omid Aminoroayaie Yamini, K.N. Toosi University of Technology, Iran

Food science

Maria Manuela Abreu da Silva, ESEAG, University Lusofona, Portugal

Asif Ahmad, Institute of Food and Nutritional Sciences, PMAS-Arid Agriculture University Rawalpindi, Pakistan

Felix Aladedunye, Max Rubner-Institut, Germany

Mehraj Ahmad, University Sains Malaysia, Malaysia (nutrition)

Ali Asghar, University of Agriculture, Faisalabad, Pakistan

Alina M. Balu, Universidad de Cordoba, Spain

Mustafa Bayram, Gaziantep University, Turkey

Michael Boateng, Kwame Nkrumah University of Science and Technology (KNUST), Ghana

Julieta Correa-Betanzo, University of Guelph, Canada (Food Chemistry, Nutrition)

Cédric Camps, Agroscope, Switzerland (food engineering)

Rui Cruz, Universidade do Algarve, Portugal

Jose A. Egea, Center for Edaphology and Applied Biology of Segura (CEBAS-CSIC), Spain

Andronoiu Doina Georgeta, Dunarea de Jos University of Galati, Romania (food engineering, food technology, food texture, nutrition)

Ahmed I. Gomaa, Laval University, Canada

Ramon Gonzalez, J. Rettenmaier und Sohne USA, Cambridge, USA (food engineering, food technology)

Galani Yamdeu Joseph Hubert, University of Leeds, UK

Annie King, UC Davis, USA

Diganta Kalita, Colorado State University, USA (food technology, food chemistry, nutrition)

<u>Przemysław Kowalczewski</u>, Poznan University of Life Sciences, Poland

Carol S.K. Lin, City University of Hong Kong, Hong Kong (food engineering)

Tomislav Mikuš, University of Zagreb, Croatia

Pranabendu Mitra, University of Wisconsin Stout, USA

Pankaj B Pathare, Newcastle University, UK (food engineering, food preservation)

Jana Pickova, Swedish University of Agricultural Sciences, Sweden

Subash Shrestha, ConAgra Foods, USA (food microbiology)

Ajaypal Singh, United States Department of Agriculture, USA (Food engineering, Food chemistry, Food technology, Food preservation)

Jaya Sundaram, United States Department of Agricultural, USA

Gunnar Sundstøl Eriksen, Norwegian Veterinary Institute, Norway (food microbiology)

Agricultural economics

Mirza Barjees Baig, King Saud University, Kingdom of Saudi Arabia (agricultural education, rural development, agricultural information and communication)

Antonino Galati, University of Palermo, Italy

Antonio Lopolito, University of Foggia, Italy

Donato Morea, University of Cagliari, Italy

Piergiuseppe Morone, Unitelma-Sapienza, University of Rome, Italy (bio-based economics)

Arkadiusz Piwowar, Wroclaw University of Economics and Business, Poland

Ulrich Schmutz, Coventry University, United Kingdom

Roberta Sisto, University of Foggia, Italy

Ariel Soto-Caro, Universidad de Concepción - Campus Chillán, Chile

Agricultural engineering

Mário Cunha, University of Porto, Portugal (environmental engineering, biological systems engineering)

Rui S. Oliveira, University of Coimbra, Portugal (environmental engineering, biological systems engineering)

George S. Skouteris, Loughborough University, UK (environmental engineering, biological systems engineering)

Carol S.K. Lin, City University of Hong Kong, Hong Kong (biological systems engineering)

Cédric Camps, Agroscope, Switzerland (biological systems engineering)

Prahlad Jat, University of North Carolina, USA

Pankaj B Pathare, Newcastle University, UK (postharvest technology)

Pablo Martín Ramos, Universidad de Zaragoza, Spain

Gurdeep Singh, The Climate Corporation, USA

Christoph Steiner, University of Kassel, Germany

Mohammad Valipour, Payame Noor University, Iran

Jose A. Egea, Center for Edaphology and Applied Biology of Segura (CEBAS-CSIC); Campus Universitario de Espinardo, Spain

CHIHAIE-DASEU AGIICUITUIE

Mário Cunha, University of Porto, Portugal (environmental engineering, biological systems engineering)

Cliford S. Gold, Independent consultant, USA

Debjani Sihi, University of Maryland, Center for Environmental Science, USA

Christoph Steiner, University of Kassel, Germany

Mohammad Valipour, Payame Noor University, Iran

Martin Wiesmeier, Bavarian State Research Center for Agriculture, Germany

Amelioration

Biswanath Dari, University of Florida, USA

Elena Aydin, Slovak University of Agriculture, Slovakia

Christoph Steiner, University of Kassel, Germany

Mohammad Valipour, Payame Noor University, Iran

Sebastian Wolfrum, Technische Universität München, Germany (soil improvement, erosion control)

Tropical Agriculture

Pinaki Acharyya, Central Institute for Arid Horticulture, India

Lorenzo Barbanti, University of Bologna, Italy

Vittorio Farina, University of Palermo, Italy

Clifford S. Gold, Independent consultant, USA

Sergio de los Santos Villalobos, Instituto Tecnológico de Sonora, Mexico

Christoph Steiner, University of Kassel, Germany

Debjani Sihi, University of Maryland, Center for Environmental Science, USA

Mohammad Valipour, Payame Noor University, Iran

Microbiology

Sergio de los Santos Villalobos, Instituto Tecnológico de Sonora, Mexico

Debjani Sihi, University of Maryland, Center for Environmental Science, USA

Jian Wu, Virginia Polytechnic Institute and State University, USA

Sustainable developement

Felix Arion, AgroTransilvania Cluster: Dezmir, Romania

Joseph Awuni, University for Development Studies, Ghana

Mario D'Amico, Università di Catania, Italy

Giuseppe Di Vita, University of Turin, Italy

Sartaj Bhat, Gifu University, Japan

Prahlad Jat, University of North Carolina, USA

Piotr Prus, University of Science and Technology, Poland

Mohammad Valipour, Payame Noor University, Iran

Sebastian Wolfrum, Technische Universität München, Germany (farm management systems, GIS)

Agronomy

Karkanis Anestis, University of Thessaly, Greece

Mirza Barjees Baig, King Saud University, Kingdomof Saudi Arabia (agricultural education, rural development, agricultural information and communication)

Moritz Von Cossel, University of Hohenheim: Stuttgart, Germany

 $\underline{\text{M\'{a}rio Cunha}}, \text{University of Porto, Portugal (environmental engineering, biological systems engineering)}$

Vinod Jakkula, Bayer U.S - Crop Science, USA

Mirza Hasanuzzaman, Sher-e-Bangla Agricultural University, Bangladesh

Mehdi Rahimi, Graduate University of Advanced Technology, Iran

Domenico Ronga, Centro Ricerche Produzioni Animali SpA, Italy

Parmodh Sharma, New Mexico State University, USA

Daljit Singh, Bayer - Crop Science, Monsanto Company, USA

Agnieszka Synowiec, University of Agriculture in Krakow, Poland

 $\underline{Mohammad\ Valipour}, Payame\ Noor\ University, Iran$

Plant biology

<u>Dr. Anket Sharma</u>, Guru Nanak Dev University, India

Biomass, Biofuel

Muhammad Irfan, University of Sargodha, Pakistan

Dahunsi Olatunde, Landmark University, Nigeria

Rural development

Fotios Chatzitheodoridis, University of Western Macedonia, Greece

Vitor Martinho, Instituto Politécnico de Viseu - Escola Superior Agrária, Portugal

Juan Rodríguez-Cohard, Universidad de Jaén, Spain

Kate Stephen, Scotland's Rural College, UK

Sri Krishna Sudheer Patoju, Tata Institute of Social Sciences, India

Agricultural information and communication

Festus Victor Bekun, Gelisim Üniversitesi, Turkey

Christian Franco-Crespo, Universidad Técnica de Ambato, Ecuador

Giuseppina Migliore, University of Palermo, Italy

Language Editors

Dorice Ago, Independent Consultant, UK

Graham Brodie, The University of Melbourne, Australia

Emmanuel G. Escobar, University of Sheffield, UK

Galani Yamdeu Joseph Hubert, University of Leeds, UK

Jasmin Hufschmid, The University of Melbourne, Australia

Renuka Kholkute, India

Neil R. McEwan, School of Pharmacy and Life Sciences, Robert Gordon University, Scotland, UK

Publisher

DE GRUYTER Poland

Bogumiła Zuga 32A Str.


01-811 Warsaw, Poland

T: +48 22 701 50 15

Editorial Contact

openagriculture@degruyter.com

Terindeks Scimagojr.com (Q2)



SCOPE


Open Agriculture is an open access journal that publishes original articles reflecting the latest achievements on agro-ecology, soil science, plant science, horticulture, forestry, wood technology, zootechnics and veterinary medicine, entomology, aquaculture, hydrology, food science, agricultural economics, agricultural engineering, climate-based agriculture, amelioration, social sciences in agriculture, smart farming technologies, farm management.

 $\ensuremath{\bigcirc}$ Join the conversation about this journal





Terindeks Scopus


В	C	D	E	F	G
Source Title (Medline-sourced journals are indicated in Green)	Print-ISSN	E-ISSN	Active or Inactive	Coverage	Titles discontinued by Scopus due to quality issues
	- 1	•			
Open Access Journal of Medicinal and Aromatic Plants	09747877		Inactive	2010-2019	
Open Access Journal of Urology	11791551		Inactive	2010-2011	
Open Access Macedonian Journal of Medical Sciences		18579655	Inactive	2014-2022	Discontinued by Scopu
Open Access Rheumatology: Research and Reviews	1179156X		Active	2009-ongoing	
Open Access Surgery		11787082	Inactive	2016-2017	
Open Agriculture		23919531	Active	2016-ongoing	
Open Agriculture Journal		18743315	Active	2012-ongoing	
Open AIDS Journal		18746136	Active	2009-ongoing	
Open Allergy Journal		18748384	Inactive	2010-2015	
Open Anesthesia Journal	25896458		Active	2019-ongoing	
Open Anesthesiology Journal		18743218	Inactive	2010-2018	
Open Archaeology		23006560	Active	2015-ongoing	
Open Astronomy		25436376	Active	2016-ongoing, 2003	
Open Atmospheric Science Journal		18742823	Inactive	2011-2015	
Open Automation and Control Systems Journal		18744443	Inactive	2011-2015, 2009	
Open Biochemistry Journal		1874091X	Active	2011-ongoing, 2007	
Open Bioinformatics Journal		18750362	Active	2017-ongoing, 2010-2015	
Open Biology		20462441	Active	2011-ongoing	
Open Biomarkers Journal		18753183	Active	2018-ongoing, 2015, 2009-2013	
Open Biomedical Engineering Journal		18741207	Active	2009-ongoing	
Open Biotechnology Journal		18740707	Active	2009-ongoing	
Open Bone Journal		18765254	Inactive	2011-2015	
Open Breast Cancer Journal		18768172	Inactive	2011-2015	
Open Cancer Journal		18740790	Inactive	2009-2015	
Open Cardiovascular Medicine Journal		18741924	Active	2009-ongoing	
Open Catalysis Journal		1876214X	Inactive	2009-2014	
Open Ceramics		26665395	Active	2020-ongoing	
Open Chemical Engineering Journal		18741231	Active	2009-ongoing	
Open Chemistry	23915420		Active	2015-ongoing, 2003	
Onen Civil Engineering Journal		18741495		2010-ongoing	
Scopus Sources October 2022 Accepted titles Feb. 2	2023 / Disco	ntinued title:	s Feb. 2023	Serial Conf. Proc. with profile All (Con / 4




3 Open Access April 6, 2022 Soil test-based phosphorus fertilizer recommendation for malting barley production on Nitisols Kassu Tadesse, Birhan Abdulkadir, Wubengeda Admasu, Dawit Habte, Almaz Admasu, Amare Tadesse, Anbessie Debebe Page range: 171-180 More ▼ Cite this Download PDF Open Access April 7, 2022 Effects of domestication and temperature on the growth and survival of the giant freshwater prawn (Macrobrachium rosenbergii) postlarvae Jolene Tay, Adriana Suhanizen, Malik Aziz, Noorlela Yassin, Takaomi Arai Page range: 181-190 More ▼ Cite this Download PDF 3 Open Access April 7, 2022 Influence of irrigation regime on gas exchange, growth, and oil quality of field grown, Texas (USA) olive trees Staci Parks, Thayne Montague Page range: 191-206 More ▼ Cite this Download PDF 8 Open Access April 13, 2022 Present status and prospects of value addition industry for agricultural produce-A review Alaa Al Hinai, Hemantha Jayasuriya, Pankaj B. Pathare, Talal Al Shukaili Page range: 207-216 More ♥ Cite this Download PDF 3 Open Access April 18, 2022 Competitiveness and impact of government policy on chili in Indonesia Saptana Saptana, Ening Ariningsih, Ashari Ashari, Endro Gunawan, Atika Dyah Perwita, Syahrul Ganda Sukmaya, Handewi Purwati Saliem, Helena Juliani Purba, Kurnia Suci Indraningsih, Atika Dian Pitaloka, Nur Qomariah Hayati Page range: 226-237 More ▼ Cite this Download PDF 3 Open Access April 18, 2022 Growth of Rucola on Mars soil simulant under the influence of pig slurry and earthworms Gerrit Willem Wieger Wamelink, Line Schug, Joep Yolanda Frissel, Ingrid Lubbers Page range: 238-248 More ▼ Cite this Download PDF 8 Open Access April 19, 2022 Effect of potassium fertilizer application in teff yield and nutrient uptake on Vertisols in the central highlands of Ethiopia Evasu Elias, Fanosie Mekonen, Gizachew Kebede Biratu, Wassie Hail Page range: 257-266 More ▼ Cite this Download PDF 3 Open Access April 27, 2022 Dissection of social interaction and community engagement of smallholder oil palm in reducing conflict using soft system methodology Herdis Herdiansvah Page range: 267-283 More ▼ Cite this Download PDF












a Open Access October 14, 2022 Do poor farmers have entrepreneurship skill, intention, and competence? Lessons from transmigration program in rural Gorontalo Province, Indonesia Amelia Murtisari, Irham Irham, Jangkung Handoyo Mulyo, Lestari Rahayu Waluyati Page range: 794-807 More * Cite this Download PDF a Open Access October 22, 2022 Communication networks used by smallholder livestock farmers during disease outbreaks: Case study in the Free State, South Africa Christopher Ugochukwu Nwafor, Ifeoma Chinyelu Nwafor Page range: 808-819 More * Cite this Download PDF 3 Open Access October 22, 2022 Sustainability of Arabica coffee business in West Java, Indonesia: A multidimensional scaling approach Eddy S. Yusuf, Ening Ariningsih, Ashari, Endro Gunawan, Helena J. Purba, Sri H. Suhartini, Herlina Tarigan, Syahyuti, Juni Hestina, Yonas H. Saputra, Suci Wulandari, Nyak Ilham, Mewa Ariani Page range: 820-836 More ▼ Cite this Download PDF 3 Open Access October 26, 2022 Farmers' perspectives on the adoption of smart farming technology to support food farming in Aceh Province, Indonesia Agussabti Agussabti, Rahmaddiansyah Rahmaddiansyah, Ahmad Humam Hamid, Zakaria Zakaria, Agus Arip Munawar, Basri Abu Baka Page range: 857-870 3 Open Access October 27, 2022 Rice yield grown in different fertilizer combination and planting methods: Case study in Buru Island, Indonesia Reginawanti Hindersah, Agusthinus Marthin Kalay, Abraham Talahaturuson Page range: 871-881 More ♥ Cite this Download PDF 3 Open Access November 7, 2022 Paclobutrazol and benzylaminopurine improve potato yield grown under high temperatures in lowland and medium land Svariful Mubarok, Anne Nuraini, Sumadi Sumadi, Jajang Sauman Hamdar Page range: 882-888 More ▼ Cite this Download PDF a Open Access November 16, 2022 Agricultural sciences publication activity in Russia and the impact of the national project "Science." A bibliometric analysis Page range: 889-898 More ▼ Cite this Download PDF a Open Access November 30, 2022 Storage conditions and postharvest practices lead to aflatoxin contamination in maize in two counties (Makueni and Baringo) in Kenya Hannah Mugure Kamano, Michael Wandayi Okoth, Wambui-Kogi Makau, Patrick Kuloba, Nduhiu Gitahi Page range: 910-919 More ♥ Cite this Download PDF



Short Communication





Special Issue on Agriculture, Climate Change, Information Technology, Food and Animal (ACIFAS 2020)




Special Issue of International Web Conference on Food Choice and Eating Motivation

3 Open Access October 27, 2022

Can ingredients and information interventions affect the hedonic level and (emo-sensory) perceptions of the milk chocolate and cocoa drink's consumers?

Dimas Rahadian Aji Muhammad, Nanik Maya Marettama, Gusti Fauza, Dian Rachmawanti Affandi

Page range: 847-856



10.1515_opag-2022-0080+Preceived Atribute

by 10.1515_opag-2022-0080+preceived Atribute Eni Sumarni

Submission date: 27-Feb-2023 02:46PM (UTC+0700)

Submission ID: 2024138178

File name: 10.1515_opag-2022-0080_Preceived_Atribute.pdf (389.06K)

Word count: 5771

Character count: 32104

DE GRUYTER Open Agriculture 2022; 7: 217–225

Research Article

Poppy Arsil*, Yeong Sheng Tey, Mark Brindal, Ardiansyah, Eni Sumarni, Masrukhi

Perceived attributes driving the adoption of system of rice intensification: The Indonesian farmers' view

https://doi.org/10.1515/opag-2022-0080 received July 19, 2020; accepted March 11, 2022

Abstract: This article argues that the System of Rice Intensification (SRI) future promotion should be based on the potential users' good understanding of sustainable agriculture. A qualitative approach was used to examine the perceptions of SRI attributes among Indonesian rice farmers, which is built upon the developing theory of diffusion of innovation. Through focus group discussions in three Indonesian provinces, compatibility, complexity, and relative advantage were identified as essential factors for SRI adoption. SRI was seen as incompatible with current farming practices, labour capacity, budget, and time available for additional labour inputs. SRI was seen as relatively complicated in terms of compost processing and application as well as mechanised agricultural technologies. As a result of the economic surplus provided by SRI rice, organised farmers may be able to obtain a higher price for SRI rice than non-organised farmers. Environmental and agronomic benefits were thought to have a long-term payoff. Such results demonstrate the subjective evaluation of SRI by farmers, which is important to its implementation.

Keywords: compatibility, complexity, diffusion of innovation, Rogers' theory, relative advantage

Ardiansyah, Eni Sumarni, Masrukhi: Agricultural Technology Department, Universitas Jenderal Soedirman, Purwokerto 53122, Indonesia

1 Introduction

One innovation aimed at increasing rice productivity is the System of Rice Intensification (SRI). This innovation emphasises sustainability principles in managing local plants, soil, water, and nutrients and their incorporation into farmers' current practices (where they deem compatible). SRI, as promoted to rice farmers in our study areas, is a set of principles [1]. Core principles of the SRI are (1) younger seedling, (2) one seedling planted at one clump, (3) wide square planting (more than $20 \text{ cm} \times 20 \text{ cm}$), and (4) intermittent irrigation [2]. SRI is a fluid technological package [1]. It needs to be adjusted according to local nuances. Even though the sudden conversion from standard practice to fully organic is not recommended [3], organic fertilisers are still advised to reduce synthetic fertilisers and improve the soil structure and quality [4]. Different kinds of biological control further differentiate SRI from conventional weed and pest management [1,4]. Through the SRI principles, rice plants are reported more resistant to pests and pathogens as their leaves are bolder, larger, and stronger than those planted using conventional systems [5]. When properly followed and implemented, chemical inputs, water, and seed are used efficiently [2]. Because the system diminishes external inputs, SRI principles have positive impacts on resource and environmental conservation [1]. As such, SRI offers a means to realise the goals of sustainable agriculture. Importantly, too, given its flexibility and robustness, SRI principles are applicable to small-scale farmers.

Despite offering great potential, the adoption rate of the SRI generally remains low, especially among Indonesian rice farmers [6–9]. However, little research effort has been made to understand why adoption rates remain low. This issue has, to the best of our knowledge, only been investigated by Takahashi [8]. Like many farmers' adoption studies (i.e. [10]), their investigation focused on relating the heterogeneity of producer, farm, institutional, and intervention variables to adoptive decisions.

^{*} Corresponding author: Poppy Arsil, Agricultural Technology
Department, Universitas Jenderal Soedirman, Purwokerto 53122,
Jonesia, e-mail: poppy.arsil@unsoed.ac.id
Yeong Sheng Tey: Laboratory of Agricultural and Food Policy
Studies, Institute of Tropical Agricultural and Food Policy Security,
Universiti Putra Malaysia, Selangor Darul Ehsan 43400, Malaysia
Mark Brindal: Department of Agricultural Sciences, School of
Agriculture, Food and Wine, University of Adelaide, Urrbrae SA 5064,
Australia

Open Access. © 2022 Poppy Arsil *et al.*, published by De Gruyter. © This work is licensed under the Creative Commons Attribution 4.0 International License.

Others investigate the effect of intermittent irrigation on rice yield [11]. Such approaches are criticised for providing a few practical insights, which might help extension agents understand how to encourage greater adoption [12,13]. Farmers' opinion and ideas are based on their experience over the years when practising farming. Therefore, their knowledge can be used as an explanation of the variation of input and output quantities in rice production that existed and sometimes cannot be explained statistically. The objective of SRI adoption set by policymakers is to increase farmer incomes and food security as well as improved environmental quality. Policymakers and change agents should understand SRI from the subjective view of farmers as potential users. Thus, policymakers can achieve the goal of adoption of the SRI method. Focus group discussion (FGD) is one way to obtain farmers' insight from different SRI exposure in a natural discussion guided by a moderator.

SRI was introduced in Indonesia during 1997. Local farmers were introduced to the SRI methods through self-help training in plant ecology lessons in 2002 in Tasikmalaya. As the pioneer, this area serves as a key learning resource to farmers or group of farmers, extension agents, and researchers. In contrast to the history of SRI in the Tasikmalaya regency, Purbalingga and Tabanan regencies followed the SRI program initiated by the Indonesian Government. The program was introduced to local farmers in the mid of 2000s. However, the adoption rate, for example in Purbalingga, was considered low, which was less than half of the local rice farmers who participated in the SRI programme initiated by the local Government.

In Rogers' seminal synthesis on the adoption of innovations, innovation attributes are identified as having a profound impact on farmers' adaptive decisions [14]. Agricultural studies have also demonstrated the importance of the perceived attributes of innovation in relation to its adoption (i.e. [15]). However, as posited in ref. [12], the current explanatory power of past empirical studies is less than adequate. Their work suggests that a lot of important information on perceived innovation attributes is likely to remain unaccounted when using restrictive empirical methods. To address this research gap that we have identified, this study examines the perceptions of SRI attributes held by current and potential Indonesian users. We propose to do this using the exploratory framework as prepositioned by Rogers [14]. In this framework, innovation attributes are theorised to influence farm decision-making in relation to its adoption. This study utilises a qualitative approach to understand adoption, focusing on the role of SRI attributes. The study aims to investigate farmers' views of perceived attributes driving the adoption of SRI through the use of a focus group in a frame of Rogers' theory of the adoption of innovation. A key benefit of this approach is that it does not involve judgement from the researchers. It values what individual farmers believe the attributes are and evaluates which attributes are acceptable from their point of view [16].

In Rogers' theory of diffusion of innovation [14], there are five common attributes of innovations: relative advantage, complexity, compatibility, observability, and trialability. Individuals are likely to vary in their perceptions according to a matrix based on their congruence stretching across attributes. The term "relative advantage" refers to the extent to which new ideas, behaviours, and objects are viewed as more innovative and superior to the innovations they are replacing [14]. It is commonly evidenced through financial costs and/or gains. Sustainable innovations that generate a net financial advantage, both perceived and actual, are more likely to be adopted [17]. Additional relative advantages include timesaving, reduction of discomfort, social prestige, and immediacy of the benefits from the innovation.

Compatibility is defined as the degree to which potential adopters perceived the innovation consistent with their existing values and past experiences [14]. This is traditionally interpreted in terms of the compatibility with an existing system, with little modification [18]. Sustainable innovations that are believed to be necessary and applicable are more likely to be adopted [19]. Similar inclination is also likely to crystallise when a sustainable innovation is aligned with the value of a social system [20]. Complexity is described as the degree to which potential users perceived the innovation as relatively difficult to comprehend and use ([14], p. 15). Complex innovations typically involve a new learning curve before initiating them into practice. They are less likely to be adopted [21]. Trialability is characterised by "the degree to which an innovation may be experimented with on a limited basis" ([14], p. 16). Trial on a small scale allows users to experiment with the integration of innovations within an existing system and to learn relative advantages and handle the complexity of innovations prior to their full implementation. Trialability thereby reduces the risk associated with and increases the likelihood of adoption [22]. Observability is the degree to which others can see the results of an

innovation [14]. Visibility is split into practice observability and benefit observability. Being able to see the actual implementation and the associated benefits strengthen the inclination towards adoption [23]. According to Rogers [14], the perception of innovation attributes affects an individual's action. Favourable ones are more likely to induce farmers to adopt and continue using the innovation.

2 Methods

Guided by Rogers' conceptual framework [14], FGDs were conducted in the Tasikmalaya, Purbalingga, and Tabanan regencies. Tasikmalaya regency is known as the pioneer of SRI implementation and becomes the central learning of SRI farming practices. Purbalingga and Tabanan regencies followed through a government assistance program. Tasikmalaya and Purbalingga are regencies on the island of Java, the most populated island in Indonesia, whereas the Tabanan regency is located on the Bali island. Our target participants were rice farmers who have heard of the SRI. A total of 40 key informants participated in the FGD held. They were selected according to the recommendations made by local extension agents and the leader of farmer organisations. The participants were selected based on their knowledge and experience and their roles in the community. Two subgroup FGDs consisted of five to nine participants for each regency to give all participants enough time to share. The first subgroup consisted of farmers who received the SRI program's government assistance, whereas the other did not receive the assistance program for data cross-checking. Each subgroup involved SRI adopters and dis-adopters. Careful consideration was given to the definition of "SRI adopters." Figure 1 shows how the research was conducted to cross-check the data.

Some researchers have argued that farmers who applied at least one core practice of SRI can be classified as adopters [6,24]. That same definition is applied in local standards and, in turn, is used in this study. Dis-adopters are rice farmers who are aware of, have applied, but then have discontinued using SRI principles. A skilled moderator led the FGDs using a semi-structured interview. The moderator would stop the FGDs if no more comprehensive new information is identified.

As the FGDs were conducted in Indonesia, the collected information was transcribed and translated into English to achieve a standard understanding across the individual researchers involved in this study. Two researchers identified the common keywords, the subsequent themes (attributes), and patterns, which the FGD participants valued following Rogers' theory of the adoption of innovation.

3 Results and discussion

The characteristics of the participants of the FGDs are presented in Table 1. On average, the participants are approximately 50 years old, and most of them completed senior high school education. Two-thirds of the participants




Figure 1: Research model.

Table 1: Respondents' characteristics of FGDs

Characteristics	Tasikmalaya (n = 12) Mean	Purbalingga (n = 18) Mean	Tabanan (n = 10) Mean	Overall (n = 40) Mean
Age (years old)	48.4	49.1	54.1	50.33
Male (%)	83	78	100	87
Education level				
Primary school (%)	16	_	10	8.7
Junior high school (%)	_	17	20	12.3
Senior high school (%)	42	50	60	50.7
University (%)	42	33	10	27.7
SRI adopters (%)	58.3	55.6	60	58
SRI dis-adopters (%)	41.7	44.4	40	42
Duration applying the SRI (year)	10	2.8	3.8	5.53
Land area planted using the SRI (ha)	6.146	3.740	3.681	4.52
Fertiliser and pest control methods				
Fully organic inputs (%)	16.7	_	20	12,3
Mixed organic and chemical inputs $(\%)$	83.3	100	80	87.7

were adopters of the SRI. The pioneers in Tasikmalaya regency started using the SRI 10 years ago on 6.15 hectares of rice farm. Their counterparts in Purbalingga and Tabanan have had less than 5 years' experience with the innovation and applied it on smaller scales. It was observed that only a fraction of the adopters had entirely used organic inputs. The majority employed a mix of organic and chemical inputs.

FGD participants frequently mentioned common keywords such as rice yields, production costs, price of the SRI rice, the use of organic fertiliser, labour capacity, land preparation, and water irrigation control. The keywords were categorised into three significant attributes of SRI, according to Rogers' theory of the adoption of innovation. They are compatibility, complexity, and relative advantages. These findings are substantially congruent with Rogers' general framework [14]. Each of these significant attributes is interrelated with the others. Little mention was made of perceived trialability by implementing a trial on a small scale of paddy field to experiment with the innovation. The FGD participants did not mention perceived observability such as success story. These two less significant attributes are similarly identified by Tornatzky and Klein [25] and Rogers [14]. Table 2 shows the dimensions driving the adoption and disadoption of SRI.

3.1 Compatibility

Despite several years of experience, previous rice farming experience and local culture still constrained their SRI

practises. This remains, therefore, a major concern. Application of the SRI principles requires the modification of existing farming practices. Adopters from the Purbalingga regency remained uncomfortable with planting seeds in a tray, shallow planting, and the land preparation methods, all of which are promoted under SRI. Similar concerns were also expressed by adopters from the Tabanan regency, specifically concerning the discipline required to grow seed for less than 10 days before transplanting and then transplant the seed at a rate of one seed per hole.

"Although SRI techniques are well-intended to boost seedling survival rate and vegetation, it is difficult to change the mindset of farmworkers who are used to conventional methods." Dis-adopter from Tabanan, two years of SRI experience.

Resistance to change in technical aspect was commonly mentioned as partial reasoning contributing to dis-adoption. Those who chose to observe from the sidelines remained sceptical, preferring to wait to see how SRI techniques would affect farm productivity. The same observation is shared by Sato et al. [9] and Handono [26]. Other researchers also reported that some farmers are lazy to learn and adopt new technology [27].

The implementation of sustainable weed and pest management is labour intensive [28]. Participants typically observed rapid growth of weeds and snails as a result of the SRI because it involves cultivating rice with wide spacing between plants, the use of organic fertilisers, and less water. Snail invasion was said to be particularly prevalent soon after the transplanting cycle.

Table 2: Determinant attributes driving the adoption and dis-adoption of SRI following Rogers' theory of the adoption of innovation

Dimensions of innovation	Benefits	Obstacles
Compatibility	_	Need a modification of existing farming practices involving planting seeds technique, land preparation, and water control
		Persistent to change in the technical view of rice agriculture
Complexity	-	Difficult to adjust the appropriate organic fertiliser according to spatial farming requirement
		2. Farmers face a shortage of raw material for making organic fertiliser.
Relative advantages	 Farmers believe that the SRI method yields higher productivity than conventional practices if farmers follow the SRI principles 	 Extra labour is needed for planting, weeding, applying fertiliser, and water control
	2. The production cost of seed might be reduced due to less seed used	2. Scarce farm labour against high industrial demand
	 Price is higher than traditional farming of rice if the farmer cooperatives have bargaining power at the markets 	3. Price seems the same as conventional rice farming if the farmers or farmer cooperatives have no bargaining power at the markets
	4. SRI increases natural pest and disease control in	
	the paddy field 5. Soil condition improved	
	6. Water usage decreased	
Immediacy of results	_	A significant time gap between the time of adoption and returns of the application of SRI methods

3.2 Complexity

As SRI principles encourage organic fertiliser use, adopters have to grapple with a degree of complexity in judging the farm's nutrient needs and applying the appropriate organic fertiliser according to their farm's spatial requirements. The knowledge of integrated nutrient management is crucial for sustaining high yield of SRI [29]. In our study areas, composts are promoted as suitable organic fertilisers that restore organic matter and enhance soil properties. Subsidised by local Government, such organic fertiliser is sold at affordable prices. Given its affordability, high local demand is often unsatisfied because of supply shortages. Consequently, participants in the SRI program are often forced to do the extra task of making their own compost. Although most of them received training in both composting and application techniques, production and application processes in respect to composts are not always straightforward. Therefore, it is recommended that there is an unmet need to educate participants to become competent to troubleshoot composting problems (e.g. anaerobic fermentation or incorrect N:P:K balance) and determine the resultant quality of compost. This is in line with

ref. 27 that increasing training access is useful to prepare farmers to practise SRI.

3.3 Relative advantages

Yield: Nearly all participants of the FGDs agreed that the SRI promises high productivity. Notwithstanding this, it must be noted that, in fact, productivity levels varied between enterprises.

However, the participants believed that the productivity of the SRI depends mainly on the adherence to its principles: use 5 tons/ha of compost, plant a single seed-ling in each hole with wide spacing between clumps, apply local microorganisms and at least four times, and conduct four periods of weeding each season. Liquid fermentation contains local microorganism derived from base material such as cow rumen or rabbit urine provided a useful decomposition tool for making nutrient available to plant. Then, the organism will reproduce with natural ingredients containing carbohydrates, proteins, vitamins, and minerals. Indeed, following these practices, adopters from Purbalingga and Tasikmalaya regencies produced approximately 7–8 ton/ha of milled

rice in comparison to the yield of conventional methods. Similar findings are also recorded by [30], who reported that Bangladesh farmers who implement water-saving technology (WST) of rice agriculture recorded higher productivity than farmers who were practising the conventional irrigation method. The farmers' income also increased by 24.6% when using the WST method.

SRI's superior yield was said to be a key factor contributing to the inclination to adopt and continue using the SRI principles.

"Most SRI farmers in my area do not strictly follow the recommendations. We applied 3 ton/ha of compost, sole cropping systems, one time of local microorganisms, and 2–3 times of weeding in a season. We only produced about 7–8 ton/ha of milled rice on average. Nevertheless, the yield is still considered high compared to the 5 ton/ha of milled rice produced through conventional farming in the local area. Although we do not entirely follow the SRI, we are more directed to organic farming. Collectively, nearly 70% of land in Manonjaya sub-district is planted using organic methods, and the certified organic farmland is about 37 hectares."

Adopter from Tasikmalaya, 10 years of SRI experience.

Production costs: Among participants of the FGDs, who have had experience with SRI techniques, there were mixed opinions with regard to production costs of SRI. Such findings were related to variations in the cost of seed, labour input, and organic fertilisers. Rice cultivated under the SRI generally uses less seed than conventional systems. Adopters from the Tabanan regency indicated that the cost of seeds was reduced as much as 65% and this had led them to save more money.

As previously mentioned, extra labour hours were necessarily allocated for planting, weeding, fertilisation, and irrigation activities. The greater demand for human input further squeezed the already scarce farm labour, which has increasingly shifted to other industries. The availability of farm labour thus becomes a critical issue in some local contexts. For example, access to labour has been identified as a key factor determining the continuity of SRI in the Jeneponto district of the South Sulawesi province [8] and Madagascar [27]. Farmers who have an opportunity to more labour resources have increased their apability to adopt SRI [27].

As demonstrated above, any calculation of production costs is not straightforward. The application of SRI techniques could save nearly 20% production cost [31] and decrease production cost with the benefit—cost ratio of 1.49 [30]. However, FGD participants relied on their subjective evaluation rather than an objective one when weighing the cost and benefit for their decision-making.

"We support the government program aiming at achieving 10 ton/ha of milled rice. Through SRI techniques, I used to produce around 8 ton/ha of milled rice, but its production cost was high. In opposite, the *Jajar Legowo* technique, a planting rice method with the pattern of multiple rows of rice plant interspersed with an empty row, is simpler in terms of planting and crop maintenance. Although I achieved slightly lower yield (7–7.7 ton/ha) using the traditional method, the associated cost was significantly lower."

Dis-adopter from the Tabanan regency, 6 years of SRI experience.

Price of SRI rice: The price of SRI rice was suggested to be the most important factor driving farmer decisions in relation to adoption and dis-adoption. Rice produced using the SRI principles, especially organic rice, is considered to be of higher quality and to have health benefits. Consequently, it should follow that SRI farmers should reap higher prices. However, two divisions were noted among our focus group participants. Organic SRI rice in the Purbalingga regency typically commands a price premium of about 50% above the standard market price. At the time of study, the SRI rice was sold for around 12,000-13,000 IDR/kg, and undifferentiated rice was priced at approximately 8,000 IDR/kg. This pricing outcome was due to the collective bargaining power of group action. Local organic farmers were engaged and organised through the Pamorbangga Farmer Association. The farmer association worked as a marketing agent, distributing local organic rice to Jakarta - the capital city of Indonesia - and selling directly to consumers. Without going through any middlemen, the farmer association recorded a higher profit margin and returned greater profits to its members.

"Price of the organic rice that sold through the Pamorbangga Farmer Association is lucrative. However, this farmer association only covers a sub-district, and there are many organic farmers out there. We hope the local Government will help and support us to extend the outreach of the farmer association." Adopter from Purbalingga, 5 years of SRI experience.

In contrast, participants from the Tasikmalaya and Tabanan regencies had significantly less bargaining power on an individual basis. The price achieved for rice grown using SRI principles was only marginally higher than the price of conventional rice. Such pricing was already generally lower (8,000–9,000 IDR/kg) than the returns in the Purbalingga regency. This occurred because SRI rice was sold directly to a farmers' group (gapoktan) in the Tasikmalaya regency. With little premium gained for the extra effort involved, participants were demotivated and expressed an intention to quit the SRI.

"Healthy rice is what we called for rice produced using SRI methods. However, its demand is still low. That leads to the low prices of SRI rice."

Dis-adopter from Tabanan, 1 year of SRI experience.

Agronomic benefits: Through the integrated pest management that is promoted under the SRI, participants of our FGDs believed that they are likely to strike a natural balance in which pests and diseases are well controlled. Anecdotal evidence was provided by an adopter from Purbalingga that his rice plants cultivated using SRI techniques were more pest and disease resistant. The use of SRI is also believed to improve environmental quality such as improved soil aeration. Plants were also more resistant to diseases and pests [30]. Adopters of SRI observed that their soil conditions differed from non-users of SRI techniques.

Water-saving derived from the irrigation management that is promoted under the SRI was another significant impact that is valued by participants of the FGDs. Under controlled environment, water usage of SRI can be reduced up to 86% [4,30]. This translates into a significant improvement in water productivity [30]. Such benefit was said to be particularly critical during dry seasons.

Immediacy of results: Participants in our FGDs emphasised that they cannot afford to wait for long periods before they benefit from adopting SRI. Adopters have invested significant effort to learn and master SRI techniques to produce more satisfactory yields. In other words, there is a significant time gap between learning SRI methods and optimising returns from its use. Such lead times have led dis-adopters to believe that the SRI does not promise lucrative benefits in the short-term. This concern was particularly highlighted by participants who worked on leased lands under time-limited tenancies across all three study areas. As a result, such farmers had the minimal motivation to invest in the SRI.

Arsil et al. [32] who conducted a study regarding perceived importance and performance of SRI attributes between adopters and dis-adopters reported that "profit," "risk," and "effort" are three critical attributes for rice farmers. The performance of those attributes was reported below the average. In this study, profit is related to rice yield, production cost, and price. As other business-like attributes, the benefit is an important attribute by farmers. Therefore, the promotion of SRI should involve any effort related to increasing the SRI price. The risk was identified as a second essential attribute for both adopters and disadopters. Crop failure due to the complexity of mastering

the SRI technique such as organic fertiliser application, farm nutrient needs according to their spatial requirement, land preparation, and planting the seed in trays are identified as barriers to adopt SRI for farmers. Applying the SRI technique is sometimes thought to be a waste of time and effort for farmers.

This study highlights that it is a subjective evaluation of SRI attributes that drives its adoption. While learning through experience, adopters remain objective. For example, SRI users expect their organic rice to command higher prices both in view of higher production costs and to achieve acceptable profit levels. Failure to meet their objectives is likely to result in discontinuation of SRI. The adoption process varied slightly between farmers and regions during the application of fertiliser and pest control method. Bali and Tasikmalaya farmers seem to use more organic fertiliser and pest control during the adoption of SRI. The heterogeneity of adoption might be affected by psychological, behavioural, economic, and technological factors [33].

4 Conclusions and policy implications

Through FGDs, a number of perceived attributes have been identified. Compatibility, complexity, and relative advantage appeared to be the key attributes driving the adoption of SRI. As their motivation for the adoption of SRI is centred on the relative advantages of economic returns, future promotions of SRI should highlight and inform potential users of the degree to which SRI is more profitable than competing rice farming systems in both the short- and long-run. Having convinced them to use and stay in the program seems likely to overcome farmer difficulty in the perception of SRI's non-economic relative advantages in the long-term.

Funding information: This study was partially supported by Universitas Jenderal Soedirman under the scheme of the competency research grant (contract number 2448/UN 23.14/PN/2018).

Author contributions: PA: conceptualisation, formal analysis, funding acquisition, writing – original draft, and writing – review and editing; YST and MB: conceptualisation and writing – original draft; A: formal analysis and

writing – review and editing; ES and M: project administration and formal analysis.

Conflict of interest: The authors state no conflict of interest.

Data availability statement: The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

References

- Stoop WA, Uphoff N, Kassam A. A review of agricultural research issues raised by the System of Rice Intensification (SRI) from Madagascar: opportunities for improving farming systems for resource-poor farmers. Agric Syst. 2002;71(3):249-74.
- [2] Styger E, Aboubacrine G, Attaher MA, Uphoff N. The system of rice intensification as a sustainable agricultural innovation: introducing, adapting and scaling up a system of rice intensification practices in the Timbuktu region of Mali. Int J Agric Sustain. 2011;9(1):67-75.
- [3] Ardiansyah ArifC, Hardanto A, Mustofa A, Nishida K. Performance of SRI Rice Growth on Soil Accustomed to conventional cultivation methods. Agric Eng Int CIGR J. 2020;22:11–8.
- [4] Kassam A, Friedrich T, Derpsch R, Lahmar R, Mrabet R, Basch G, et al. Conservation agriculture in the dry Mediterranean climate. F Crop Res. 2012;132:7–17.
- [5] Thakur AK, Uphoff N, Antony E. An assessment of physiological effects of System of Rice Intensification (SRI) practices compared with recommended rice cultivation practices in India. Exp Agric. 2010;46(1):77–98.
- [6] Berkhout E, Glover D. The evolution of the System of Rice Intensification as a socio-technical phenomenon: A report to the Bill & Melinda Gates Foundation [Internet]. 2011. http://papers. ssrn.com/sol3/papers.cfm?abstract_id=1922760.
- [7] Natawidjaja R, Djuwendah E, Mukti G. Socio-economic impact assessment of SRI paddy cultivation for farmers and society in Tasikmalaya regency. Local government report. Bandung (Indonesia); Universitas Padjadjaran and Departement of Agriculture, Tasikmalaya regency; 2008.
- [8] Takahashi K. The roles of risk and ambiguity in the adoption of the system of rice intensification (SRI): evidence from Indonesia. Food Secur. 2013:5:513–24.
- [9] Sato S, Yamaji E, Kuroda T. Strategies and engineering adaptions to disseminate SRI methods in large-scale irrigation systems in Eastern Indonesia. Paddy Water Env. 2011;9(1):79–88.
- [10] Abebaw D, Haile MG. The impact of cooperatives on agricultural technology adoption: Empirical evidence from Ethiopia. Food Policy. 2013;38:82-91.
- [11] Nugroho BDA, Toriyama K, Kobayashi K, Arif C, Yokoyama S, Mizoguchi M. Effect of intermittent irrigation following the system of rice intensification (SRI) on rice yield in a farmer's paddy fields in Indonesia. Paddy Water Environ. 2018;16(4):715–23.

- [12] Knowler D, Bradshaw B. Farmers' adoption of conservation agriculture: A review and synthesis of recent research. Food Policy. 2007;32(1):25–48.
- [13] Prokopy LS, Floress K, Klotthor-Weinkauf D, Baumgart-Getz A. Determinants of agricultural best management practice adoption: Evidence from the literature. J Soil Water Conserv. 2008;63:300-11.
- [14] Rogers E. Diffusion of Innovations. New York: Free Press; 2003.
- [15] Tey YS, Li E, Bruwer J, Abdullah AM, Brindal M, Radam A, et al. The relative importance of factors influencing the adoption of sustainable agricultural practices: a factor approach for Malaysian vegetable farmers. Sustain Sci. 2014;9(1):17–29.
- [16] Reimer AP, Weinkauf DK, Prokopy LS. The influence of perceptions of practice characteristics: An examination of agricultural best management practice adoption in two Indiana watersheds. J Rural Stud. 2012;28(1):118-28.
- [17] Barr N, Cary J. Greening a Brown Land: An Australian Search for Sustainable Land Use. Melbourne: Macmillan; 1992.
- [18] Gamon JA. Educational delivery methods to encourage adoption of sustainable agricultural practices. J Agric Educ. 1994;35:38–42.
- [19] Gillespie JM, Kim S-A, Paudel KP. Why don't producers adopt best management practices? An analysis of the beef cattle industry. Agric Econ. 2007;36:89–102.
- [20] Webb T. Understanding behavior: Social and economic influences on land practice change, Australia. A Paper Presented at the Workshop on Land Management Practices Information Priorities, Classification and Mapping Towards an Agreed National Approach, Kamberra Winery, Canberra 2004 May 11–12. Australia: Bureau of Rural Sciences; 2004. p. 1–7.
- [21] Vanclay F, Lawrence G. Agricultural extension as social welfare. Rural Soc. 1995;5(1):20-33.
- [22] Pannell DJ, Marshall GR, Barr N, Curtis A, Vanclay F, Wilkinson RL. Understanding and promoting adoption of conservation practices by rural landholders. Aust J Exp Agric. 2006;46(11):1407–24.
- [23] Guerin LJ, Guerin TF. Constraints to the adoption of innovations in agricultural research and environmental management: a review. Aust J Exp Agric. 1994;34:549–71.
- [24] Takahashi K, Barrett CB. The System of Rice Intensification and its impacts on Household Income and Child Schooling: evidence from Rural Indonesia. Am J Agric Econ. 2014;96(1):269–89.
- [25] Tornatzky LG, Klein KJ. Innovation characteristics and innovation adoption-implementation: A meta-analysis of findings. IEEE Trans Eng Manag. 1982;EM-29(1):28-45.
- [26] Handono SY. Problems and obstacles of SRI (System of Rice Intensification) method in the practice. Habitat. 2013;24:10-9.
- [27] Whitman K, Ratsimbazafy J, Stevens N. The use of System of Rice Intensification (SRI) near Maromizaha Protected Area, Madagascar. Madag Conserv Dev. 2020;15(1):1–8.
- [28] Barrett CB, Moser CM, McHugh OV, Barison J. Better technology, better plots, or better farmers? Identifying changes in productivity and risk among Malagasy Rice Farmers. Am J Agric Econ. 2004;86(4):869–88.
- [29] Gairhe J, Thapa S. System of Rice Intensification: boon or burden? I Agric Env. 2020:21:216–28.
- [30] Uddin MT, Dhar AR. Assessing the impact of water-saving technologies on Boro rice farming in Bangladesh: economic and environmental perspective. Irrig Sci. 2020;38(2):199–212.

- [31] Sato S. A review of on-farm evaluations of system of rice intensification methods in Eastern Indonesia. Cab Rev Perspect Agric Vet Sci Nutr Nat Resour. 2007;2:2.
- [32] Arsil P, Tey YS, Brindal M, Ardiansyah A, Sahirman S. Perceived importance and performance of System of Rice Intensification
- (SRI) between adopters and dis-adopters: insights from Indonesia. J Soc Sci Res. 2018;2018(Special Issue 6):14-21.
- [33] Chavas J-P, Nauges C. Uncertainty, learning, and technology adoption in agriculture. Appl Econ Perspect Policy. 2020;42(1):42-53.

10.1515_opag-2022-0080+Preceived Atribute

ORIGINALITY REPORT

9% SIMILARITY INDEX

9%
INTERNET SOURCES

0%
PUBLICATIONS

U% STUDENT PAPERS

PRIMARY SOURCES

1

doaj.org
Internet Source

5%

2

www.researchgate.net

Internet Source

4%

Exclude quotes

On

Exclude matches

< 2%

Exclude bibliography