

## 5. Formulation and characterization of emergency food based on instan corn flour supplemented by instan tempeh (or soybean) flour

Jurnal internasional bereputasi

Nama jurnal: International Food Research Journal

Vol 25 no 1, Februari 2018

1. Cover
2. Editor
3. Daftar isi
4. Artikel
5. Indexing (Q3, SJR 0,312)
6. Bukti koresponden
7. Turnitin



# INTERNATIONAL FOOD RESEARCH JOURNAL

Home Editorial Board Author's Guide Code of Ethics List of Issues Contact Us FOOD Website

## OPEN ACCESS



Now Indexed in SCOPUS, EBSCO, CHEMICAL ABSTRACT, ProQuest, CABI, AGRICOLA & WORLD ISLAMIC SCIENCE CITATION, MYCITE CITATION Report

SUBMIT YOUR MANUSCRIPTS - <http://mc.manuscriptcentral.com/upm-ifrj>

**ISSN (Online): 2231 7546**

## RECENTLY PUBLISHED ISSUES

- [2018 - Volume 25 Issue 1](#) **NEW!**
- [2017 - Volume 24 Issue 6](#)
- [2017 - Volume 24 Issue 5](#)
- [2017 - Volume 24 Issue 4](#)
- [2017 - Volume 24 Issue 3](#)
- [2017 - Volume 24 Issue 2](#)
- [2017 - Volume 24 Issue 1](#)

### [SEE ALL ISSUES](#)



**[SCImago Journal and Country Rank](#)**  
**[MyJurnal - Malaysian Citation Centre](#)**

The International Food Research Journal (IFRJ) publishes papers in English, six (6) issues a year with the coverage of:

- Food Science and Technology
- Nutrition and Dietetics
- Agriculture, multidisciplinary
- Chemistry, multidisciplinary

The scope of the Journal includes:

- Food Science, Food Technology and Food Biotechnology
- Product Development and Sensory Evaluation
- Food Habits, Nutrition, and Health
- Food Safety and Quality
- Food Chemistry, Food Microbiology, Food Analysis and Testing
- Food Engineering
- Food Packaging
- Food Waste Management
- Food Entrepreneur
- Food Regulatory
- Post-Harvest Food Management
- Food Supply Chain Management
- Halal Food and Management

# INTERNATIONAL FOOD RESEARCH JOURNAL

[Home](#)[Editorial Board](#)[Author's Guide](#)[Code of Ethics](#)[List of Issues](#)[Contact Us](#)[FOOD Website](#)

## EDITORIAL BOARD

### Editor-in-Chief

#### **Dr. Jinap Selamat**

Professor  
International Food Research Journal (IFRJ)  
Food Safety Research Centre (FOSREC)  
Faculty of Food Science & Technology  
Universiti Putra Malaysia  
MALAYSIA

### Associate Editors

#### **Dr. Tan Chin Ping**

Professor  
Department of Food Technology  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
MALAYSIA

#### **Dr. Mad Nasir Shamsudin**

Professor  
Department of Agribusiness and Bioresource Economics  
Faculty of Agriculture  
Universiti Putra Malaysia  
MALAYSIA

#### **Dr. Faridah Abas**

Associate Professor  
Department of Food Science  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
MALAYSIA

### Board Members

#### **Dr. Nazamid Saari**

Professor  
Department of Food Science  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
MALAYSIA

#### **Dr. Karamatollah Rezaei**

Professor  
Department of Food Science, Engineering and Technology  
Faculty of Agricultural and Engineering Technology  
College of Agriculture and Natural Resources  
University of Tehran  
IRAN

#### **Dr. Olga Martin-Belloso**

Professor  
Department of Food Technology  
University of Lleida  
SPAIN

#### **Dr. Gulum Sumnu**

Professor  
Department of Food Engineering  
Middle East Technical University  
TURKEY

#### **Dr. Stefan Kasapis**

Professor  
School of Science  
RMIT University  
AUSTRALIA

#### **Dr. Hans J.P. Marvin**

Research Coordinator  
RIKILT  
Wageningen University & Research  
THE NETHERLANDS

#### **Dr. Didier Montet**

Professor

#### **Dr. Siree Chaiseri**

Associate Professor

HDR, CIRAD  
FRANCE

Kasetsart University  
THAILAND

**Dr. John Gilbert**  
Professor  
Orta Dogu Teknik Universitesi,  
FoodLife International Ltd  
TURKEY

**Dr. Steve Flint**  
Professor of Food  
Institute of Food, Nutrition and Human Health  
Massey University  
NEW ZEALAND

**Dr. Mansel William Griffiths**  
Professor  
Canadian Research Institute for Food Safety (CRIFS)  
Department of Food Science  
University of Guelph  
CANADA

**Dr. Liu Yang**  
Professor  
Institute of Food Science & Technology  
Chinese Academy of Agricultural Sciences  
CHINA

**Dr. Gregory R. Ziegler**  
Professor of Food Science  
Pennsylvania State University  
University Park  
UNITED STATES OF AMERICA

**Dr. Yonghua Wang**  
Professor in Food Bioprocess  
School of Food Science and Engineering  
South China University of Technology  
CHINA

**Dr. Van Viet Man LE**  
Professor  
Head of Food Technology Department  
Ho Chi Minh City University of Technology  
VIETNAM

**Dr. Yuanfa Liu**  
Professor  
State Key Laboratory of Food Science and Technology  
Synergetic Innovation Center of Food Safety and Nutrition  
School of Food Science and Technology  
Jiangnan University  
CHINA

**Dr. Yin Li**  
Professor  
CAS Key Laboratory of Microbial Physiological and Metabolic  
Engineering  
Institute of Microbiology  
Chinese Academy of Sciences  
CHINA

**Prof. Dr. Franz Berthiller**  
Center for Analytical Chemistry  
Department of Agrobiotechnology (IFA-Tulln)  
University of Natural Resources and Life Sciences  
AUSTRIA

**Prof. Dr. Wan Rosli Wan Ishak**  
Department of Nutrition  
School of Health Science  
Universiti Sains Malaysia  
MALAYSIA

# INTERNATIONAL FOOD RESEARCH JOURNAL

[Home](#)[Editorial Board](#)[Author's Guide](#)[Code of Ethics](#)[List of Issues](#)[Contact Us](#)[FOOD Website](#)

## VOLUME 25 ISSUE 1, 2018

[>> Previous Issues](#)
**International Food Research Journal Volume 25 Issue 1, 2018** **NEW!**

### Mini Review

1. [Adverse effects on consumer's health caused by hormones administered in cattle](#)
  - Senthil Kumar, V., Rajan, C., Divya, P. and Sasikumar, S.
2. [Characteristics and use of electrolyzed water in food industries](#)
  - Athayde, D.R., Flores, D.R.M., Silva, J.S., Silva, M.S., Genro, A.L.G., Wagner, R., Campagnol, P.C.B., Menezes, C.R. and Cichoski, A.J.

### Original articles

3. [The characterization of bioactive peptides of goat milk fermented to activities as anti-hypercholesterolemia](#)
  - Mahdi, C., Untari H., Padaga, M.C. and Raharjo, S.J.
4. [Screening of new bioactive in lebuli beans \(\*Cajanus sp.\*\) of Lombok](#)
  - Mushollaeni, W., Kumalaningsih, S., Wignyanto and Santoso, I.
5. [Genotype impact on antioxidant potential of hull and kernel in Persian walnut \(\*Juglans regia L.\*\)](#)
  - Rahmani, F., Dehghaniasl, M., Heidari, R., Rezaee, R. and Darvishzadeh, R.
6. [Antioxidant activity of roasted and unroasted peanut skin extracts](#)
  - Elsorady, M. E. I. and Ali, S. E.
7. [Additional antioxidants on the determination of quercetin from \*Moringa oleifera\* leaves and variation content from different sources](#)
  - Panya, T., Chansri, N., Sripanidkulchai, B. and Daodee, S.
8. [Antioxidant properties of peels extracts from three varieties of banana \(\*Musa sp.\*\) grown in West Java-Indonesia](#)
  - Fidrianny, I., Anggraeni, N.A.S. and Insanu, M.
9. [Antioxidant capacity and nitrosation inhibition of cruciferous vegetable extracts](#)
  - Amron, N. A. and Konsue, N.
10. [Comparison of phytochemical and antioxidant properties of extracts from flaxseed \(\*Linum usitatissimum\*\) using different solvents](#)
  - Alachaher, F. Z., Dali, S., Dida, N. and Krouf, D.
11. [Evaluation of antioxidant and nutritional properties of sago \(\*Metroxylon sagu\* Rottb.\) and its utilization for direct lactic acid production using immobilized \*Enterococcus faecium\* DMF78](#)
  - Duque, S.M.M., Castro, I.J.L. and Flores, D.M.
12. [Effect of chitosan coating on storage stability of tomatoes \(\*Lycopersicon esculentum\* Mill\)](#)
  - Sucharitha, K. V., Beulah, A. M. and Ravikiran, K.
13. [Effect of edible chitosan coating on combined ultrasound and NaOCl treated kiwi fruits during refrigerated storage](#)
  - Vivek, K. and Subbarao, K.V.
14. [Effect of pre-treatments on the chemical, functional and storage properties of breadfruit \(\*Artocarpus altilis\*\) flour](#)
  - Arinola, S. O. and Akingbala, J. O.
15. [Effects of temperature on postharvest respiratory parameters and quality attributes of ackee \(\*Blighia sapida\* Koenig\) fruit arils during storage](#)
  - Benkeblia, N. and Beaudry, R.M.
16. [Sensory and nutrient quality of wild captured \*Oreochromis shiranus\* \(Boulenger, 1897\) stored at ambient temperature](#)
  - Mchazime, I. and Kapute, F.
17. [Physicochemical characteristics and phytochemical content of jam made from melon \(\*Cucumis melo\*\)](#)
  - Benmeziiane, F., Djermoune – Arkoub, L., Boudraa, A.T. and Bellaagoune, S.
18. [Physicochemical properties of cocoa butter replacers from supercritical carbon dioxide extracted mango seed fat and palm oil mid-fraction blends](#)
  - Jahurul, M.H.A., Zaidul, I.S.M., Sahena, F., Sharifudin, M.S., Norulaini, N.N., Md. Eaqub Ali, Hasmadi, M., Ghafoor, K., Wahidu Zzaman and Omar, A.K.M.

19. Nutritional composition, fatty acids and cholesterol levels in Atlantic white shrimp (*Litopenaeus schimitti*)
  - Pires, D. R., de Moraes, A. C. N., Coelho, C. C. S., Marinho, A. F., Góes, L. C. D. S. A., Augusta, I. M., Ferreira, F. S. and Saldanha, T.
20. Nutrient composition of commonly consumed edible insects in the Lango sub-region of northern Uganda
  - Akullo, J., Agea, J.G., Obaa, B. B., Okwee-Acai, J. and Nakimbugwe, D.
21. Application of modified sorghum flour for improving bread properties and nutritional values
  - Istianah, N., 1Ernawati, L., Anal, A.K. and Gunawan, S.
22. Analysis of nutrient and antinutrient content of the truffle (*Tirmania pinoyi*) from Morocco
  - Bouatia, M., Touré, H. A., Cheikh, A., Eljaoudi, R., Rahali, Y., Oulad Bouyahya Idrissi, M., Khabar, L. and Draoui, M.
23. Kefir ice cream flavored with fruits and sweetened with honey: physical and chemical characteristics and acceptance
  - Januário J.G.B., Oliveira, A.S., Dias, S.S., Klososki, S.J. and Pimentel, T.C.
24. In vitro bioaccessibility of  $\beta$ -carotene in pumpkin and butternut squash subjected to different cooking methods
  - Koh, S.H. Loh, S.P.
25. Optimization of chemical treatment on fresh cut tender jackfruit slices for prevention of browning by using response surface methodology
  - Rana, S. S., Pradhan, R. C. and Mishra, S.
26. Foam-mat drying of passion fruit aril
  - Khamjae, T. and Rojanakorn, T.
27. Volatile chemical profile of cacao liquid smoke
  - Janairo, J.I.B. and Amalin, D.M.
28. Quality performance of protein allergen isolates for allergy diagnostic test (Case: Indonesian soybeans (*Glycine max*) and peanuts (*Arachis hypogaea*))
  - Astuti, R.M., Palupi, N.S. and Zakaria, F.R.
29. Evaluation of ethanol concentration, temperature and shaking time of extracted Thai Jasmine rice on cholinesterase enzyme activity
  - Kukreja, R.K., Sripum, C., Charoenkiatkul, S., Kriengsinyos, W. and Suttisansanee, U.
30. Heavy metal removal ability of *Halomonas elongata* and *Tetragenococcus halophilus* in a media model system as affected by pH and incubation time
  - Asksonthong, R., Siripongvutikorn, S. and Usawakesmanee, W.
31. Hypolipidemic effects of quercetin and kaempferol in human hepatocellular carcinoma (HepG2) cells
  - Yusof, H.M., Sarah, Ng, M.L., Lam, T.W. and Kassim, M.N.I.
32. Evaluation of high pressure treatment for improvement of physicochemical and functional qualities in purple corncobs
  - Itthisoponkul T., Naknan P., Prompun S. and Kasemwong K.
33. Evaluation of caseinolytic, gelatinolytic and milk-clotting activities from germinated seeds of green gram (*Vigna radiata*), chick pea (*Cicer arietinum*), fenugreek (*Trigonellafoenum-graecum*) and moth bean (*Vigna aconitifolia*)
  - Bajulge, P. V., Bhutia, K. C., Awasarmol, A. V., Shaikh, F. K., Panche, A. N. and Harke, S. N.
34. Effect of soaking and autoclaving treatments on structure, properties and resistant starch (RS3) content of edible tapioca pearls
  - Kasote, D. M., Oak, M. D., Nilegaonkar, S. S. and Agte, V. V.
35. Modelling of water loss and solute uptake during osmotic drying of carrots using weibull distribution approach
  - Sahoo, P.K. and Sharma, A.K.
36. Characterization and functional properties of gelatin extracted from goatskin
  - Zilhada, Yahdiana, H., Irwandi, J. and Effionora, A.
37. GC-MS analysis of unpolar fraction from *Ficus carica* L. (fig) leaves
  - Ivanov, I., Dimcheva, I., Budjakov, I., Petkova, N., Denev, P. and Pavlov, A.
38. Formulation and characterization of emergency food based on instan corn flour supplemented by instan tempeh (or soybean) flour
  - Aini, N., Prihananto, V., Wijonarko, G., Sustriawan, B., Dinayati, M., and Aprianti, F.
39. Quality characteristics including formaldehyde content in selected Sea foods of Tuticorin, southeast coast of India
  - Immaculate, J. and Jamila, P.
40. Effect of selected cereal grains on in vitro bioaccessibility of isoflavones in soymilk
  - Ng, C.C. and Loh, S.P.
41. Studies on black tea production from fresh roselle calyxes
  - Aregbesola, O. A., Faborode, M. O. and Hounkanrin, B.A.
42. Glycemic index of *Coleus tuberosus* crackers rich in resistant starch type III
  - Nugraheni, M., Hamidah, S. and Auliana, R.
43. Utilization of soursop leaves as antihyperuricemic in functional beverage 'Herbal Green Tea'
  - Hardoko, Tanudjaja, Y., Mastuti, T.S. and Halim, Y.
44. Potential health risk of heavy metals accumulation in vegetables irrigated with polluted river water
  - Ratul, A.K., Hassan, M., Uddin, M.K., Sultana, M.S., Akbor, M.A. and Ahsan, M.A.

45. **Study on the physicochemical, microbial and sensory characteristics of alcoholic beverage produced by indigenous method**
  - Ahmad, I., Ahmed, S., Yadav, B., Sah, P.P., Alam, J. and Zzaman, W.
46. **Evaluation of the ability of *Micrococcus* sp. isolated from Barada river to biodegradation of vegetable oil wastes**
  - Alsulaiman, A. and Ali Nizam, A.
47. **Evaluation of antimicrobial and antioxidant activities of oily macerates of Algerian dried figs (*Ficus carica* L.)**
  - Debib, A., Tir-Touil, M.A., Meddah, B., Hamaidi-Chergui, F., Menadi, S. and Alsayadi, M.S.
48. **Nutritional and microbiological evaluation on sauces and ketchups available in Bangladesh**
  - Anandsynal, Mumtaz, B., Motalab, M., Jahan, S., Hoque, M.M. and Saha, B. K.
49. **Factors affecting pineapple market supply in Johor, Malaysia**
  - Jaji, K., Man, N. and Nawi, N.M.
50. **Application of a facilitating HACCP system using two innovative methods for the production of Hibiscus syrup by a Senegalese small and medium business (SMBs)**
  - Ndiaye, N.A., Cissé, M., Bonne, R., Sene, B., Kane, N.C. and Montet, D.
51. **Effect of dietary olive leaves on the lipid and protein oxidation and bacterial safety of chicken hamburgers during frozen storage**
  - da Silva, S. L., Marangoni, C., Brum, D. S., Vendruscolo, R. G., Silva, M. S., de Moura, H. C., Rampelotto, C., Wagner, R., de Menezes, C. R., Barin, J. S., Campagnol, P. C. B., Cichoski, A. J.
52. **Microbiological and physicochemical quality of pasteurized milk supplemented with sappan wood extract (*Caesalpinia sappan* L.)**
  - Kimestri, A.B., Indratiningsih and Widodo
53. **Probiotic soft sheep's cheese: evaluation of probiotic survival and its influence on proteolysis and organoleptic characteristics**
  - Cuffia, F., Bergamini, C. and Candioti, M.
54. **Influence on cultivation conditions in the heterotrophic lipid production of the microalga *Chlorella minutissima***
  - Ferreira, S.P., Holz, J.C.P. and Costa, J.A.V.
55. **Comparison between morphophysiological and molecular methods for the identification of yeasts isolated from honey**
  - Rodrigues, A. M. D., Pinheiro, R. E. E., Costa, J. A., Santos, J. T. O., Poli, J. S., Rosa, C. A., Soares, M. J. S., Muratori, M. C. S. and Nóbrega, M. M. G. P.
56. **Detection, identification and phylogenetic analysis of lactic acid bacteria isolated from Tarkhineh, Iranian fermented cereal product, by amplifying the 16s rRNA gene with universal primers and differentiation using rep-PCR**
  - Vasiee, A.R., Mortazavi, A., Tabatabaei-yazdi, F. and Edalatian Dovom, M.R.
57. **Antifungal activity of *Boersenbergia rotunda* (temukunci) extract against filamentous spoilage fungi from vegetables**
  - Zakuan, Z., Mustapa, S. A., Sukor, R. and Rukayadi, Y.
58. **Microbial interaction in selected fermented vegetable condiments in Nigeria**
  - Olanbiwoninu, A. A. and Odunfa, S. A.
59. **Occurrence and PCR identification of *Salmonella* spp. from milk and dairy products in Mansoura, Egypt**
  - Omar, D., Al-Ashmawy, M., Ramadan, H. and El-Sherbiny, M.
60. **In vivo evaluation of snake fruit Kombucha as hyperglycemia therapeutic agent**
  - Zubaidah, E., Apriyadi, T.E., Kalsum, U., Widyastuti, E., Estiasih, T., Srianata, I. and Blanc, P.J.



## Formulation and characterization of emergency food based on instan corn flour supplemented by instan tempeh (or soybean) flour

\*Aini, N., Prihananto, V., Wijonarko, G., Sustriawan, B., Dinayati, M., and Aprianti, F.

Department of Agricultural Technology, Faculty of Agriculture, Jenderal Soedirman University,  
Purwokerto, Central Java-Indonesia, 53123

### Article history

Received: 5 September 2016  
Received in revised form:  
14 November 2016  
Accepted: 25 November 2016

### Abstract

A special disasters emergency food that can be directly consumed, practical and nutritious was formulated. It was developed based on local food materials, namely corn-soybean and corn-tempe flour. The aim of the research is to produce an emergency food formula made from corn-soybean flour and corn-tempe flour that adequately fulfills the standard for emergency food, acceptable and feasibly produced in large quantities. It was found that the best formulation model of emergency food is produced from Srikandi corn flour and tempe flour. Its composition is 42% of corn flour, 20% of tempe flour, 10% of full cream milk powder, 16% of sugar, and 12% of frying oil. Nutrient composition of every piece (each 50 grams of product) was 8.1 g of protein, 20.67 g of lipid, 20.58 g of carbohydrate, and 298.04 kcal of energy. In terms of protein and energy, emergency food already fulfills the adequacy standard, which is the minimum energy 233 kcal and 7.9 to 8.1 g protein. Lipid content of emergency food is too high (9.1 to 11.7 g), while level of carbohydrate is too low (23-35 g). This product is most easily swallowed, most delicious, no after taste, and most preferably compared to other formulas.

© All Rights Reserved

### Keywords

Corn flour  
Emergency food  
Soybean flour  
Tempe flour

### Introduction

The disaster in Indonesia has claimed many lives in a short time, and some of people live at the evacuation site. To avoid a new disaster after natural disasters, namely the emergence of hunger, emergency feeding is absolutely necessary (Rivera and Char, 2004). Emergency food is a special type of food that is consumed during or following an emergency time to fulfill the requirements of daily human consumption (2100 kcal), and is ready to eat and nutritious (Sheu, 2007).

Materials of emergency food should be from local ingredients to raise the potential of local food. Corn is a potential source of food in Indonesia because it contains functional food components such as dietary fiber, vitamin A and iron (Scott and Eldridge, 2005; Aini *et al.*, 2010). Corn was classified as having a moderate glycemic index and being able to be consumed by people with autism (Mulloy *et al.*, 2010).

Another requirement of emergency food is it must be sufficient nutrients. Protein is an essential nutrient that has a role on the absorption of other nutrients, i.e. non-heme iron and calcium. Soybean is one source of vegetable protein. Tempe derived from soybean protein contains high bioavailability and iron (Tyug

*et al.*, 2010). Some disaster conditions often also cause the problem of drinking water availability. Intermediate moisture foods (IMF) is the name given to a category of food that is sufficiently wet to be eaten without rehydration and still remain stable during storage (Furmaniak *et al.*, 2009).

Corn-based IMF has potential as an emergency food in its ready-to-eat form. It contains high nutritional value, especially in providing sufficient energy. To fulfill the quality standards of emergency food, the focus is to supply of protein, vitamins and minerals. It is necessary to improve the nutritional quality of emergency food by combining corn flour with high protein flour (Chen *et al.*, 2006). The addition of soybeans and tempe flour will enrich nutrition. Soybeans and tempe also act as antioxidants, are anti-diarrheal and stimulate growth (Tyug *et al.*, 2010). The existence of the fermentation process will increase the levels of B vitamins, including cobalamin, riboflavin, pyridoxine, niacin, biotin, folate and pantothenic acid compounds (Eklund-Jonsson *et al.*, 2006). Formulation of corn-soybean flour and corn-tempe flour into an emergency food with an intermediate moisture food is expected to be an alternative in the provision of food, especially for refugees of natural disasters. The aim of the research is to produce an emergency food formula made

\*Corresponding author.

Email: [nuraini.1973@gmail.com](mailto:nuraini.1973@gmail.com)

Tel: 02281621004; Fax: 02281628701

from corn-soybean flour and corn-tempe flour that adequately fulfills the standard for emergency food, acceptable and feasibly produced in large quantities.

## Materials and Methods

Three varieties of corn, i.e. Pioneer, Srikandi and Canggal, were purchased from local corn farmers in Temanggung, Central Java. The soybean variety used was Slamet, while tempe was obtained from a producer in the 'Pliken' Village, Banyumas. Additional ingredients were milk powder, refined sugar, cooking oil, sorbitol, glycerol and water.

The experiment was conducted in three steps. The first step was the formulation of emergency food from three varieties of corn and the prediction of the nutritional adequacy of each formula. The second was the determination of the amount of water and a humectant to be added in each formula. It was followed by the production of the emergency food formula. This step also included an analysis of water content, ash, protein, lipids (AOAC, 1995), carbohydrates (by difference) and water activity. The last step was a further analysis of the selected formula for energy value and microbiological properties (total microbial amount and yeasts).

Instant soybean flour was prepared by soaking soybeans for 6 hours and peeling the skin. The soybeans were then washed and blanched for 15 minutes. Afterwards, the soybean seeds were drying, milling and sieving by 80 mesh. The process of making instant corn flour is similar to that of making soybean flour.

Instant tempe flour was prepared by cutting tempe into 1x1x1 cm pieces then steam blanching for 20 minutes. The next step was drying the tempe until thoroughly dried. Finally, the tempe was milled and sieved through 80 mesh.

The calculation of the formula was based on the nutrient requirements of emergency foods, which should contain at least 233 kcal per piece, with 7.9-8.1 g of protein, 9.1-11.7 g of lipids, and 23-35 g of carbohydrates. This value is based on the assumption that one piece is equal to 50 g of dry weight.

Emergency food was made by mixing ingredients until it was homogeneous. In the mixing stage, water (at 80°C) was added while the mixture was stirred. The next stage tasted the product; if it was easy to swallow, a humectant (sorbitol) was added. Sorbitol was added at three concentrations, namely 4, 5 and 6%, to discover the percentage of sorbitol that can cause a bitter taste.

## Results and Discussion

### *Preparation of formula and nutritional adequacy prediction*

Preparation of the emergency food formula is based on the macronutrient content of the raw material. Calculations on the macronutrient content of the instant flour are done through an analysis of lipids, protein and carbohydrates. Instant corn flour has moisture content in the range of 12.2 to 13.59%, while instant tempe flour and soy flour have moisture contents of 11.62 and 11.77%, respectively.

Based on calculations, the emergency food formula from corn-soybean and corn-tempe flours can be seen in Table 1. The composition of corn flour in the emergency food formula ranged from 34.4 to 42%. Formulas II and III have the same composition, which is 42% corn flour, 20% soybean flour, 10% milk powder, 16% icing sugar and 12% cooking oil. Formula I also has almost the same composition as formulas II and III: 40% corn flour, 18% soybean flour, 12% milk powder, 18% sugar and 12% cooking oil. These formulas have the same composition, as they are both made from corn flour and soybean flour with other similar additives.

According to Aini *et al.* (2010), of the three varieties of maize, Srikandi corn has the highest protein content compared to the Canggal and Pioneer varieties. Processing corn into flour will not change its chemical composition. Emergency food based on corn-soybean flour has a less sweet flavor than emergency food from corn-tempe flour, due to fewer added sugars. However, based on its aroma, corn-soybean flour formula has a better aroma than corn-tempe flour formula due to the amount of oil and higher milk content. Lipids provide a delicious aroma for food. Milk powder, which has a lipid content of 25.93%, also contributes to the aroma of products.

Based on proximate analysis, we calculated the percentage composition of the materials for the emergency food formula. The prediction of emergency food nutritional adequacy based on every piece (50 g) can be seen in Table 2. The protein content of the emergency food is in the range of 7.06 to 7.97 g every piece of the product. Based on the nutritional requirements, the emergency food should contain 7.9 to 8.1 g of protein (Zoumas *et al.*, 2002). Based on these requirements, the products that are eligible are formulas III and VI. Formula III has a protein content of 15.86%, equivalent to 7.93 g. Formula VI has a protein content of 15.92%, equivalent to 7.96 g.

The protein content of the emergency food made from Srikandi corn flour is higher (9.86%) than Pioneer (7.35%) and Canggal (7.63%). Tempe flour,

Table 1. Emergency food formulation based on corn flour

| Ingredients   | Formula (%) |    |     |    |      |      |
|---------------|-------------|----|-----|----|------|------|
|               | I           | II | III | IV | V    | VI   |
| Corn flour    | 40          | 42 | 42  | 36 | 36   | 34.4 |
| Soybean flour | -           | -  | -   | 23 | 23.6 | 25   |
| Tempe flour   | 18          | 20 | 20  | -  | -    | -    |
| Milk powder   | 12          | 10 | 10  | 15 | 16   | 16   |
| Icing sugar   | 18          | 16 | 16  | 10 | 8    | 8    |
| Frying oil    | 12          | 12 | 12  | 16 | 16.4 | 16.6 |

Explanation: I = Canggal corn-tempe flour; II = Pioneer corn-tempe flour; III = Srikandi corn-tempe flour; IV = Canggal corn-soy flour; V = Pioneer corn-soy flour; VI = Srikandi corn-soy flour

soybean flour and milk powder have high protein contents, i.e., 50.46, 36.66 and 25.93%, respectively. High levels of protein in these materials increase the protein content of emergency food.

Lipids in emergency food are required to amount of 9.1 to 11.7 g every piece of product. This product has a lipid content predicted at 10.72 to 11.68 g, making it eligible. Lipids as an effective source of energy will maintain the health of the human body. One gram of oils and lipids can produce 9 kcal. Zoumas *et al.* (2002) argue that the source of lipids recommended in the development of emergency food as a source of macro nutrients is the partial hydrogenation of soybean oil, canola oil and sunflower oil. Carbohydrate levels of the emergency food are in the range of 26.29 to 27.11 g for every piece of product. The amount of carbohydrates required for emergency food is 23 to 35 g every piece. Based on the above data, the predicted formula of emergency food has exceeded the minimum recommended caloric value of emergency food.

#### Determination of the amount of water and humectants

Emergency food formulation aims to fulfill the desired energy sufficiency and flavor. Added to the main ingredients of corn-soybean flour and corn-tempe flour, other materials used are whole milk powder, sugar and cooking oil. The water used is preheated to a temperature of 80°C to facilitate the dissolving and mixing processes.

In making emergency food was added water to produce a homogeneous and easy to swallow. Mineral water that has been heated to a temperature of 80°C was gradually added until the dough was homogenous. Homogeneous dough is then subjectively tested by researchers with a limited sensory test to obtain dough that is easy to swallow and does not cause thirst.

Table 2. Prediction of the nutritional adequacy of emergency food formula

| Nutrition           | Formula |        |        |        |        |        |
|---------------------|---------|--------|--------|--------|--------|--------|
|                     | I       | II     | III    | IV     | V      | VI     |
| Protein (g)         | 7.06    | 7.87   | 7.93   | 7.83   | 7.72   | 7.97   |
| Lipid (g)           | 11.20   | 11.68  | 11.41  | 10.72  | 11.36  | 11.27  |
| Carbohydrate (gram) | 27.11   | 26.62  | 26.58  | 27.19  | 26.63  | 26.29  |
| Total energy (kcal) | 241.55  | 243.16 | 240.79 | 237.02 | 239.65 | 238.47 |

Explanation : I = Canggal corn-tempe flour; II = Pioneer corn-tempe flour; III = Srikandi corn-tempe flour; IV = Canggal corn-soybean flour; V = Pioneer corn-soybean flour; VI = Srikandi corn-soybean flour.

The amount of water added varied from 39 to 47.7 g. Differences in the addition of water were affected by the composition of the formula. A formula that has more milk powder and icing sugar requires more added water. Milk powder and icing sugar has a smaller particle size compared with corn flour and soybean flour. According to Aini *et al.* (2010), small particles have a larger surface area that absorbs more water, so formula IV to require more water than the others.

The use of sorbitol as a humectant, which include a polyol group, can affect the texture, flavor and acceptance of the product. Based on these trials, sorbitol was added at a concentration of 5%, or 2.5 g in each of the pieces of emergency food formula. The use of sorbitol is classified as generally recognized as safe (GRAS), but if it is consumed in a greater amount than 50 g/day, it will cause a laxative effect or diarrhea.

#### Nutritional value of emergency food

There are six emergency food products that were produced and had their nutritional values analyzed, as shown in Table 3. The protein content of emergency food was in the range of 12.69 to 16.25%, equivalent to 6.34 to 8.12 g in every piece. Based on the nutritional requirements, the emergency food should contain 7.9 to 8.1 of protein in each piece of the product (Zoumas *et al.*, 2002), so the product that is eligible is formula III, namely Srikandi corn and tempe flour. Every piece of product has a protein content of 16.25%, equivalent to 8.13 g.

High levels of protein in these materials increase the protein content of the emergency food. Srikandi corn flour has a protein content of 9.86%, higher than Pioneer (7.35%) and Canggal (7.63%). Tempe flour

Table 3. Nutrition content of emergency food

| Formula | Ash content<br>(% db) | Lipid content<br>(% db) | Protein total<br>(% db) | Carbohydrate (%)<br>db) |
|---------|-----------------------|-------------------------|-------------------------|-------------------------|
| I       | 1.79                  | 45.12                   | 12.69                   | 40.39                   |
| II      | 1.02                  | 46.79                   | 15.25                   | 36.94                   |
| III     | 1.25                  | 41.34                   | 16.25                   | 41.16                   |
| IV      | 1.85                  | 47.83                   | 14.56                   | 35.76                   |
| V       | 1.07                  | 50.49                   | 13.38                   | 35.06                   |
| VI      | 1.19                  | 44.71                   | 15.26                   | 38.85                   |

Explanation : I = Canggal corn-tempe flour; II = Pioneer corn-tempe flour; III = Srikandi corn-tempe flour; IV = Canggal corn-soybean flour; V = Pioneer corn-soybean flour; VI = Srikandi corn-soybean flour.

and soybean flour also have high protein contents, which amount to 50.36% and 36.66%, thereby increasing the protein content of the emergency food. The addition of milk powder in the amount of 10 to 12% also increases the levels of emergency food proteins. Milk powder has a protein content of 25.93%. According to Zoumas *et al.* (2002), a source of protein recommended for emergency food can be derived from nut products, like isolates or concentrates, and milk powder, such as casein and its derivatives. The protein amino acids must fulfill score of  $\geq 1.0$ .

The quality of protein supplementation can be improved by increasing the levels of limiting amino acids. This can be done by adding limiting amino acids from a pure protein source or mixing two or more types of different protein sources of limiting amino acids. Corn has a limiting amino acid, i.e. lysin and tryptophan, while soybeans have a different limiting amino acid, i.e. methionine. By taking these ingredients together, the deficiency of amino acids from each material can be covered. The use of tempe flour for increase the protein content of the food emergency in this study was appropriate, because in addition to increasing the quantity of protein, it also provided quality of protein.

The emergency food should have lipids in amount of 18.2 to 23.4%. This products have a lipid content of 41.4 to 50.5%, which is greater than the targeted. Corn flour has a lipid content of 0.32 to 1.61%, while tempe flour and soybean flour have lipid contents of 24.7 and 6.74%, respectively. High levels of lipids due to the addition of high amounts of oil, are 12 to 16.6%, and are 10 to 16% due to whole milk powder. To reduce the levels of lipids in the product, skim milk powder should be used, which has low levels of

Table 4. Nutrition composition of selected emergency food

| Types of nutrients     | Composition of product<br>(every 50 grams) | Emergency food<br>nutrition standard<br>(every 50 grams)* |
|------------------------|--------------------------------------------|-----------------------------------------------------------|
| Protein (% of db)      | 8.13                                       | 7.9 – 8.1                                                 |
| Lipid (% of db)        | 20.67                                      | 9.1 – 11.7                                                |
| Carbohydrate (% of db) | 20.58                                      | 23 – 35                                                   |
| Energy (kcal)          | 298.04                                     | Minimal 233                                               |
| Water activity         | 0.93                                       |                                                           |

\*Zoumas *et al.* (2002)

lipids but high protein content, so that the nutritional requirements are achieved.

Carbohydrate values are in the range of 35.2 to 41.2%. The total carbohydrate amount was lower than recommended, i.e. 46 to 70%, with donations valued at 40 to 50% energy. Carbohydrates play an important role in determining the characteristics of the material, such as flavor, color, texture and others. FAO/WHO indicates that the proportional carbohydrate should be of a sufficient quantity in the emergency food to give a sense of function, palatability, stability and metabolic functions.

The presence of sorbitol also affects the carbohydrate content. Sorbitol is a polyol compound belonging to the carbohydrates group that can increase the levels of carbohydrates. Sorbitol is hygroscopic and has a sweet taste, and it can protect the components contained in the carbohydrate content of food.

The addition of sorbitol as a humectant for emergency food has a water activity of 0.6 to 0.8. Water activity is the amount of free water that can be used by microbes for growth. According to Maltini *et al.* (2003), water activity is a thermodynamic concept that is becoming an important factor in the destruction of foodstuff. When water is more strongly bound, it is more difficult to use, both for microbiological activity and chemical hydrolytic activity. The main characteristics of emergency food are easy to swallow without a dry sensation, able to be directly consumed without further preparation, and having a sufficient shelf life because microorganisms cannot grow at specified intervals.

These emergency foods have water activity levels in the range of 0.94 to 0.96. Water activity has not been achieved as expected, i.e. 0.6-0.8. Aw product remains high due to the addition of water in large amounts so that the product is easy to swallow. This is presumably due to the amount of added water



Table 5. Microbiological analysis of selected emergency food

| Week | Total (CFU/gram)   |                    |
|------|--------------------|--------------------|
|      | Yeast              | Microbial          |
| 0    | $1.99 \times 10^2$ | $6.92 \times 10^3$ |
| 1    | $1.99 \times 10^3$ | $3.47 \times 10^4$ |
| 2    | $2.95 \times 10^3$ | $2.29 \times 10^5$ |
| 3    | $2.95 \times 10^3$ | $2.75 \times 10^5$ |
| 4    | $2.95 \times 10^3$ | $4.17 \times 10^5$ |

not being quite right or the addition of sorbitol being less effective in lowering water activity. Addition of sorbitol is expected to lower  $A_w$ , but it was not, so we need research on the use of other humectants.

#### Characterization of selected product

The selection of an emergency food formula is based on the nutritional value, sensory properties, and water activity approaching product. Emergency food from Srikandi corn flour and tempe flour with an additional 2.5% of sorbitol has the lowest water activity, i.e. 0.94. Sensory test results indicate that the product most preferred that fulfills nutritional standards has a composition of 42% Srikandi corn flour, 20% soybean flour, 10% milk powder, 16% icing sugar and 12% cooking oil. The nutrition value of the selected emergency food can be seen in Table 4.

The emergency food contributed an energy total of 298.04 kcal in every piece, up from the prior addition of sorbitol, which was 241.55 kcal. There was an increase in emergency food calories after the addition of humectants. The addition of sorbitol in the amount of 2.5 g can increase calorie emergency food to 2.6 kcal every gram. This is consistent with the results of Ahmad *et al.* (2007), that sorbitol is a carbon source that can provide additional energy to the product.

#### Total microbial and yeasts during storage

Microbiological testing conducted on emergency food was selected to determine the level of product safety. Food safety is one of the main characteristics of emergency food products (Sheu, 2007). Microbiological stability testing is conducted by storing the products in packed aluminum foil, because it has low water vapor permeability. Storage is done for four weeks, and every week, an analysis of microbial and fungal amounts is conducted. The results of microbiological analysis for four weeks can

be seen in Table 5.

The initial microbial count on the emergency food was  $6.92 \times 10^3$  CFU/g. There was microbial growth in the first week of  $3.47 \times 10^4$  CFU/g, whereas after four weeks, it had increased by  $4.17 \times 10^5$  CFU/g. Liu *et al.* (2009) explained that the increase of microbial amounts in food is influenced by intrinsic and extrinsic factors. Intrinsic factors include pH, water activity, relative humidity, nutrient content, structure and biological anti-microbial ingredients, while external factors are influenced by the storage temperature, RH and type and amount of gas in the environment. The water activity level of emergency food is still high, at 0.93, so that the microbial growth is still high.

Referring to yangko and bakpia as intermediate moisture foods, this emergency food has a similar texture and taste. Yangko refers to SNI 01-4325-1996, with a maximum standard value of microorganisms of  $1 \times 10^5$  CFU/g, while bakpia refers to SNI 01-4291-1996 with a maximum value of total microbial as 104 and of yeast as 103. Based on the microbiological standards, the product is safe to consume in the first and second weeks.

Sensory and microbiological characteristics indicate that the product in the third week is not able to be consumed because it has undergone changes in aroma and flavor. According to Chen *et al.* (2010), microbial growth can result in changes in sensory and nutritional properties, which can result in toxicity and mortality.

Total fungi also significantly increased from the first until the fourth week of storage. At the beginning, the total fungi of the product were as much as  $1.99 \times 10^2$  CFU/g. In the first week, the yeast increase by  $1 \times 10^3$  CFU/g and in the fourth week was as much as  $2.95 \times 10^3$  CFU/g. The addition of sorbitol can have a dual role as a barrier to the growth of yeast and a plasticizer of the texture. Sorbitol can also increase the level of dissolved solids in the liquid phase.

Water activity plays an important role in inhibiting the growth of microorganisms (Liu *et al.*, 2009). The water content in the food, especially the free water content, greatly affects the durability of food against invading microorganisms, represented by  $a_w$  (Sabani *et al.*, 2007). Selected products have a high  $a_w$  of 0.93, which allows the bacteria *Lactobacillus* to grow. A high water activity level can cause the growth of *Lactobacillus* due to the composition of a complete nutritional in high quantities. Protein, starches and lipids are a good media for microbial growth; therefore, microbes can grow in considerable numbers (Gidenne *et al.*, 2004).

## Conclusion

The best prototype of emergency food is produced from Srikandi corn flour and tempe flour. The best composition is 42% corn flour, 20% tempe flour, 10% whole milk powder, 16% sugar and 12% frying oil. The nutrient composition of each 50 g of product was 8.1 g of protein, 20.67 g of lipids, 20.58 g of carbohydrate and 298.04 kcal of energy. In terms of protein and energy, emergency food already fulfils the adequacy standard, which is a minimum energy of 233 kcal/piece and 7.9 to 8.1 g protein. The emergency food's lipid content is too high (9.1 to 11.7 g), while the levels are too low for carbohydrates (23-35 g). This product is swallowed easiest, delicious, with no bitter aftertaste and most preferable compared to other formulas.


## Acknowledgement

This research was supported by Directorate of Higher Education, Ministry of Education of Republic Indonesia in the program National Strategic Research in 2013.

## References

- Ahmad, T., Abbasi, N.A., Hafiz, I.A. and Ali, A. 2007. Comparison Of sucrose and sorbitol as main carbon energy sources in micropropagation of peach rootstock GF-677. *Pakistan Journal of Botany* 39(4): 1269-1275.
- Aini, N. and Hariyadi, P. 2010. Gelatinization properties of white maize starch from three varieties of corn subject to oxidized and acetylated-oxidized modification. *International Food Research Journal* 17: 961-968.
- AOAC. 1995. *Official Methods of Analysis*. Association of Official Analytical Chemists's Standard Technical (16<sup>th</sup> ed). Washington DC: USA.
- Chen, L., Remondetto, G.E. and Subirade, M. 2006. Food protein-based materials as nutraceutical delivery systems. *Trends in Food Science and Technology* 17: 272-283.
- Chen, Y.S., Liu, B.L. and Chang, Y.N. 2010. Bioactivities and sensory evaluation of Pu-erh teas made from three tea leaves in an improved pile fermentation process. *Journal Bioscience Bioengineering* 109 (6): 557-563.
- Eklund-Jonsson, C., Sandberg, A.S. and Alminger, M.L. 2006. Reduction of phytate content while preserving minerals during whole grain cereal tempe fermentation. *Journal Cereal Science* 44(2): 154-160.
- Furmaniak, S., Terzyk, A.P., Golembiewski, R., Gauden, P.A. and Czepriski, L. 2009. Searching the most optimal model of water sorption on foodstuffs in the whole range of relative humidity. *Food Research International* 42(8): 1203-1214.
- Gidenne, T., Jehl, N., Lapanouse, A. and Segura, M. 2004. Inter-relationship of microbial activity, digestion and gut health in the rabbit: effect of substituting fibre by starch in diets having a high proportion of rapidly fermentable polysaccharides. *British Journal of Nutrition* 92: 95-104.
- Liu, X., Zhou, P., Tran, A. and Labuza, T.P. 2009. Effects of polyols on the stability of whey proteins in intermediate moisture food model systems. *Journal of Agricultural and Food Chemistry* 57(6): 2339-2345.
- Maltini, E., Torreggiani, D., Venir, E. and Bertolo, G. 2003. Water activity and the preservation of plant foods. *Food Chemistry* 82(1): 79-86.
- Mulloy, A., Lang, R., O'Reilly, M., Sigafoos, J., Lancioni, G. and Rispoli, M. 2010. Gluten-free and casein-free diets in the treatment of autism spectrum disorders: A systematic review. *Research Autism Spectrum Disorders* 4(3): 328-339.
- Rivera, A.F. and Char, D.M. 2004. Emergency department disaster preparedness: Are Regional efforts reaching local front lines? *Annals of Emergency Medicine* 44: S94-S94.
- Sablani, S.S., Kasapis, S. and Rahman, M. 2007. Evaluating water activity and glass transition concepts for food stability. *Journal of Food Engineering* 78(1): 266-271.
- Scott, C. E. and Eldridge, A.L. 2005. Comparison of carotenoid content in fresh, frozen and canned corn. *Journal of Food Composition Analysis* 18(6): 551-559.
- Sheu, J.B. 2007. An emergency logistics distribution approach for quick response to urgent relief demand in disasters. *Trans Rese Part E: Logistics and Transportation Review* 43(6): 687-709.
- Tyug, T. S., Prasad, K. N. and Ismail, A. 2010. Antioxidant capacity, phenolics and isoflavones in soybean by-products. *Food Chemistry* 123(3): 583-589.
- Zoumas, B.L., Armstrong, L.E., Backstrand, J.E., Chenoweth, W.E., Chinachoti, P., Klein, B.P., Lane, H.W., Marsh, K.S. and Tolvanen, M. 2002. *High-Energy, Nutrient Dense Emergency Relief Food Product*. Washington DC: Academy Press.

# International Food Research Journal

|                                                                                 |                                                                                                |                                                                                                                                                                                                       |                                                                                                                                                              |
|---------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------|
|  | <b>SUBJECT AREA AND CATEGORY</b><br><br>Agricultural and Biological Sciences<br>└ Food Science | <b>PUBLISHER</b><br><br>Universiti Putra Malaysia<br><br> University Putra Malaysia in Scimago Institutions Rankings | <b>H-INDEX</b><br><br><b>50</b>                                                                                                                              |
|                                                                                 | <b>ISSN</b><br><br>19854668, 22317546                                                          | <b>COVERAGE</b><br><br>2007-2020                                                                                                                                                                      | <b>INFORMATION</b><br><br><a href="#">Homepage</a><br><a href="#">How to publish in this journal</a><br><a href="mailto:ifrj@upm.edu.my">ifrj@upm.edu.my</a> |



## Proses Pasang Loker Jadi Mudah

Rekrutmen jadi bebas drama, WA ke JobStreet dan kami bantu pasang iklan perusahaan Anda. JobStreet Indonesia

Buka

10451-30467-1-PB.pdf ^


📄 Tesis 14 Februari....docx ^

📁 ScienceDirect\_articl....zip ^



tiles

Science



## Submit Your Manuscript With Us

Accepting Research Focused on Wireless Power Transfer. Publish with Us.

Hindawi

[Open >](#)



### Proses Pasang Loker Jadi Mudah

Rekrutmen jadi bebas drama, WA ke JobStreet dan kami bantu pasang iklan perusahaan Anda. JobStreet Indonesia

[Buka](#)





Hindawi

[Open >](#)

1  
**Food Science and Nutrition**

GBR

**92%**  
similarity

2  
**Carpathian Journal of Food  
Science and Technology**

ROU

**91%**  
similarity

3  
**Journal of Food Processing  
and Preservation**

USA

**89%**  
similarity

4  
**Food Science and  
Technology**

BRA

**89%**  
similarity



## Submit Your Manuscript With Us

Accepting Research Focused on Wireless Power Transfer. Publish with Us.








### Proses Pasang Loker Jadi Mudah

Rekrutmen jadi bebas drama, WA ke JobStreet dan kami bantu pasang iklan perusahaan Anda. JobStreet Indonesia

[Buka](#)

scimagojr.com/journalsearch.php?q=13000154721&tip=sid&clean=0



### Proses Pasang Loker Jadi Mudah

Rekrutmen jadi bebas drama, WA ke JobStreet dan kami bantu pasang iklan perusahaan Anda. JobStreet Indonesia

Buka



Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com/journalsearch.php?q=13000154721&tip=sid&clean=0">
```

## SCImago Graphica

Explore, visually communicate and make sense of data with our **new free tool**.

Get it



## Submit Your Manuscript With Us

Accepting Research Focused on Wireless Power Transfer. Publish with Us.

Hindawi

Open >



## Proses Pasang Loker Jadi Mudah

Rekrutmen jadi bebas drama, WA ke JobStreet dan kami bantu pasang iklan perusahaan Anda. JobStreet Indonesia

Buka



under(s), institution(s) or employer(s) after peer review and before publication.

Nevertheless, IFRJ will remain an 'open access' peer-reviewed journal, and as such, all published reviews and articles are freely available online in perpetuity upon publication through which the author(s) could attain wider readership.

The APC is fixed regardless of manuscript length and page number, and substantially covers the peer review, journal production, online archiving, and copy editing charges. Even so, IFRJ remains without Submission Fee.

IFRJ Editorial would like to extend our appreciation on the unwavering support we received all these years, and hope that you will continue to consider IFRJ as the instrument by which you could circulate your research findings far and wide. For any inquiries, please contact the Editorial at [ifrj@upm.edu.my](mailto:ifrj@upm.edu.my).



**From IFRJ Editorial team.**

Now Indexed in JCR, SCIE, SCOPUS, EBSCO, CHEMICAL ABSTRACT, ProQuest, CABI, AGRICOLA, WORLD ISLAMIC SCIENCE CITATION, MyCite Citation Report

SUBMIT YOUR MANUSCRIPTS - <http://mc.manuscriptcentral.com/upm-ifrj>

**ISSN (Online): 2231 7546**



Our new impact factor is 1.014 (Q4, ranked #125/144 in the category of Food Science and Technology)

# KORESPONDENSI

## Decision Letter (IFRJ16639.R1)

**From:** ifrj@upm.edu.my

**To:** nuraini\_munawar@yahoo.com, nuraini.1973@gmail.com

**CC:**

**Subject:** International Food Research Journal - Decision on Manuscript IFRJ16639.R1

**Body:** 14-Nov-2016

Dear Dr. Aini

Thank you for sending the revised version of this paper ('Formulation and characterization of emergency food based on instan corn flour supplemented by instan tempeh (or soybean) flour'). I have now had the opportunity to examine your revised manuscript and I am pleased to accept it for publication in International Food Research Journal. Kindly refer to the attachment for the acceptance letter.

Please note that all manuscripts must be accompanied by a signed copyright agreement form to enable the manuscript to be processed for publication. You are requested to send the copyright agreement form within three (3) working days from the date of the email. Please print, fill and sign the attached copyright agreement form and send the scanned copy to ifrj@upm.edu.my

Thank you again for your contribution to the Journal.

Sincerely,  
Prof. Chin Ping Tan  
Editor, International Food Research Journal

**Date Sent:** 14-Nov-2016

**File 1:** [\\* IFRJ16-Acceptance-Letter.pdf](#)

**File 2:** [\\* Copyright-Agreement--IFRJ-.pdf](#)

 Close Window





[Home](#)[Author](#)[Review](#)

## Author Dashboard

### 4 Manuscripts with Decisions

[Start New Submission](#)[Legacy Instructions](#)[5 Most Recent E-mails](#)

## Manuscripts with Decisions

| ACTION | STATUS                                                                                                                                                                                                                                | ID         | TITLE                                                                                                          | SUBMITTED   | DECISIONED  |
|--------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|----------------------------------------------------------------------------------------------------------------|-------------|-------------|
|        | ADM: Mohamad Noh, Norhafizah<br><ul style="list-style-type: none"><li>Immediate Reject (16-Jan-2019)</li></ul> <i>Archiving completed on 17-Jan-2020</i><br><a href="#">view decision letter</a><br><a href="#">✉ Contact Journal</a> | IFRJ181426 | The formulation of a cheese analogue using sweet corn extract<br><i>Files Archived</i><br>?                    | 26-Oct-2018 | 16-Jan-2019 |
|        | ADM: Radu, Son<br>ADM: New, Vivian<br><ul style="list-style-type: none"><li>Reject (22-Feb-2017)</li></ul> <i>Archiving completed on 22-Feb-2018</i><br><a href="#">view decision letter</a><br><a href="#">✉ Contact Journal</a>     | IFRJ16997  | Quality evaluation of corn yogurt was packaged polypropylene during cold storage<br><i>Files Archived</i><br>? | 07-Nov-2016 | 22-Feb-2017 |

| ACTION                                       | STATUS                                                                                                                                                                                                                                                                                 | ID           | TITLE                                                                                                                                                                                                                                        | SUBMITTED   | DECISIONED  |
|----------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------|-------------|
|                                              | ADM: Radu, Son<br>ADM: New, Vivian<br><br><ul style="list-style-type: none"> <li>Accept (14-Nov-2016)</li> </ul> <i>Archiving completed on 15-May-2017</i><br><a href="#">view decision letter</a><br><a href="#">✉ Contact Journal</a>                                                | IFRJ16639.R1 | Formulation and characterization of emergency food based on instan corn flour supplemented by instan tempeh (or soybean) flour<br><i>Files Archived</i><br> | 05-Sep-2016 | 14-Nov-2016 |
| a revision has been submitted (IFRJ16639.R1) | ADM: Radu, Son<br>ADM: New, Vivian<br><br><ul style="list-style-type: none"> <li>Minor Revision (13-Aug-2016)</li> <li>a revision has been submitted</li> </ul> <i>Archiving completed on 15-May-2017</i><br><a href="#">view decision letter</a><br><a href="#">✉ Contact Journal</a> | IFRJ16639    | Formulation and characterization of emergency food based on corn flour<br><i>Files Archived</i><br>                                                       | 14-Jun-2016 | 13-Aug-2016 |

Tulis

Kotak Masuk 346

Berbintang

Ditunda

Meet

Rapat baru

Gabung ke rapat

Hangout



Nur



Tidak ada kontak HangoutsCari  
[seseorang](#)

## International Food Research Journal - M



**International Food Research Journal** <onbehalfof+ifrj+upm.edu.i  
kepada nuraini\_munawar, saya, prihananto\_v, goenitp, bsustriawan, in  
05-Sep-2016

Dear Dr. Aini

Your revised manuscript entitled 'Formulation and characterization o  
full consideration for publication in International Food Research Jou

Your manuscript ID is IFRJ16639.R1. Please mention this manuscr

If there are any changes in your street address or email address, ple  
manuscript at any time by checking your Author Centre after logging

Due to high volume of submissions, please expect some delays.

Thank you for submitting your manuscript to International Food Res

Sincerely,  
International Food Research Journal Editorial Office

Balas

Balas ke semua

Teruskan

Tulis

Kotak Masuk 346

Berbintang

Ditunda

Meet

Rapat baru

Gabung ke rapat

Hangout



Nur



Tidak ada kontak HangoutsCari  
[seseorang](#)

## International Food Research Journal - IF



**ifrj@upm.edu.my** lewat manuscriptcentral.com  
kepada nuraini\_munawar, saya

13-Jun-2016

Dear Dr. Aini,

Thank you for submitting your manuscript to International Food Res  
you that your manuscript does not meet the required standards. Ple

Your manuscript is located in the 'Unsubmitted Manuscripts' section  
delete the manuscript.

Please contact the Editorial Office if you have any queries regarding

Comments:

1. Place the \* in front of the author's name
2. Remove 'email:' in the corresponding author information
3. Percentage – Please do not space the unit from the number. i.e.1
4. Bold the word 'keywords' only
5. Maltini et al. (2003) - Remove the (.) after the journal's name
6. Mulloy et al. (2010) - Edit the 'and'
7. Sablani et al. (2007) - Remove the (.) after the journal's name
8. Zoumas et al. (2002) - Location comes first, only then publisher. i.
9. Table 3 - wrong spelling for sampel.
10. Table 4 - Line is missing. Please edit
11. Table 5 - Please change coloni/gram to CFU/g. Week, not week
12. Upload everything in a single MS Word containing title page info



Tulis

Kotak Masuk

346

Berbintang

Ditunda

Meet

Rapat baru

Gabung ke rapat

Hangout



Nur



Tidak ada kontak HangoutsCari

[seseorang](#)

## International Food Research Journal - D



**International Food Research Journal** <onbehalf+ifrj+upm.edu.i

kepada nuraini\_munawar, saya

13-Aug-2016

Dear Dr. Aini

Manuscript IFRJ16639, entitled 'Formulation and characterization of appear below.

Publication is recommended, subject to some minor revision to your

Thank you for submitting your manuscript to International Food Res

Sincerely,

Prof. Chin Ping Tan

Editor, International Food Research Journal

\*\*\*\*\*

Reviewers' Comments to Author:

Reviewer: 1

Comments to the Author

1. As title indicated, the food was formulated using corn flour. Howe paper more accurately. Furthermore, the use terms of instant (for ins
2. References for nutritional composition of the emergency food use

Tulis

**Kotak Masuk** 347

Berbintang

Ditunda

**Meet**

Rapat baru

Gabung ke rapat

**Hangout**



Nur



Tidak ada kontak HangoutsCari  
[seseorang](#)

## International Food Research Journal - D



**International Food Research Journal** <onbehalfof+ifrj+upm.edu.i  
kepada nuraini\_munawar, saya

13-Aug-2016

Dear Dr. Aini

Manuscript IFRJ16639, entitled 'Formulation and characterization of  
appear below.

Publication is recommended, subject to some minor revision to your

Thank you for submitting your manuscript to International Food Res

Sincerely,  
Prof. Chin Ping Tan  
Editor, International Food Research Journal

\*\*\*\*\*

Reviewers' Comments to Author:

Reviewer: 1

Comments to the Author

1. As title indicated, the food was formulated using corn flour. Howe  
paper more accurately. Furthermore, the use terms of instant (for ins
2. References for nutritional composition of the emergency food use

Tulis

Kotak Masuk 347

Berbintang

Ditunda

Meet

Rapat baru

Gabung ke rapat

Hangout



Nur



Tidak ada kontak HangoutsCari  
[seseorang](#)

## International Food Research Journal - D



**International Food Research Journal** <onbehalf+ifrj+upm.edu.i

kepada nuraini\_munawar, saya

14-Nov-2016

Dear Dr. Aini

Thank you for sending the revised version of this paper ('Formulatio  
examine your revised manuscript and I am pleased to accept it for p

Please note that all manuscripts must be accompanied by a signed  
working days from the date of the email. Please print, fill and sign th

Thank you again for your contribution to the Journal.

Sincerely,  
Prof. Chin Ping Tan  
Editor, International Food Research Journal

### 2 Lampiran

