


## Indian Journal Agricultural Research



AGRICULTURAL RESEARCH COMMUNICATION CENTRE

### INDIAN JOURNAL OF AGRICULTURAL RESEARCH

#### **EDITORIAL BOARD**

Chief Editor : T. Mohapatra, Director General

Indian Council of Agricultural Research, New Delhi

Associate Chief Editor : P.C. Sharma,

Director, CSSRI, Karnal

Managing Editor : Gaurav Gupta, A.R.C.C., Karnal

**Editors** 

P.K. Ghosh, Director & V.C., NIBSM, Raipur A. Hemantaranjan, Professor, BHU, Varanasi

S. Alarmelu, Pri. Sci., ICAR-SBI, Coimbatore Sui Kwong YAU, Candian Company, Canada

**MEMBERS** 

Ranvir Singh, New Zealand Gurbachan Singh, Karnal

R.K. Singh, U.A.E. Murari Singh, Jordan

J.S. Chauhan, New Delhi Phil Harris, U.K.

Ramesh Kanwar, U.S.A B.G. Shiva Kumar, Dharwad

C.L. Laxmipathi Gowda, Hyderabad Avijit Sen, Varanasi

Metin Guner, Turkey R.A. Balikai, Dharwad

Ravindra Singh, Ajmer Mahaveer P. Sharma, Indore

Rajendra Kumar, Lucknow N. Ravisankar, Meerut

Pratap Bhattacharyya, Cuttack A.R.G. Ranganatha, Hyderabad

J.S. Mishra, Jabalpur T. Sushila, Hyderabad

Aftabuz Zaman, Paralakhemundi Gyanendra Pratap Singh, Karnal

M. Nedunchezhiyan, Bhubneshwar V. Ramamurthy, Hebbal, Bangalore

Correspondence Address : Managing Editor

**Agricultural Research Communication Centre** 

294, Narsi Village Part 2, Sector 33,

Karnal- 132 001, India.

 $E\text{-mail}: contact@arccjournals.com \; ; \; editor@arccjournals.com \; \\$ 

www.arccjournals.com

#### INDIAN JOURNAL OF AGRICULTURAL RESEARCH

**Print ISSN: 0367-8245** 

**Online ISSN: 0976 - 058X** 

**Print frequency: Bi-monthly** 

National Academy of Agricultural Sciences (NAAS) Rating: 5.20

**Index Copernicus Value (ICV): 110.86** 

Scientific Journal Ranking (SJR): 0.241

#### ARCC Journals are covered in:-

- · BIOSIS Preview
- · ISI Citation Index
- · Biological Abstracts
- · Elsevier (Scopus and Embase)
- · AGRICOLA
- · Google Scholar
- · CrossRef
- · CAB Abstracting Journals
- · Chemical Abstracts
- · Indian Science Abstracts
- EBSCO Indexing Services

#### INDIAN JOURNAL OF AGRICULTURAL RESEARCH

**CONTENT** 

Enhancing the Productivity and Quality of Tomato using Magnetized Water and Humic Acid as

Volume 55 Issue 6, December 2021	NAAS Rating 5.20
----------------------------------	------------------

Amira A. Helaly	645
Microbes Mediated Mitigation of Abiotic Stresses in Agriculture S.S. Bobade, S.P. Dhekane, P.A. Salunke, S.G. Mane, S.S. Dhawan, R.J. Marathe, R.B. Deshmukh, Y.B. Phatake	656
Effect of NPK Fertilizer on Nutrient Uptake, Growth, Yield and Beta-carotene of Cu Groundcherry ( <i>Physalis angulata</i> L.) of Genotypes W.S.D. Yamika, G.P. Simbolon, B. Waluyo, N. Aini	itleaf 667
Management of Insect and Mite Pests of Chili using Botanical and Synthetic Chemicals M. Nasrin, M.R. Amin, M.R.U. Miah, M. Afroz, A.M. Akanda, M.G. Miah, S.J. Suh	674
Assessment of the Genetic Diversity of Joha Rice Germplasms by using Simple Sequence Ro	epeat
Markers P. Saikia, B. Neog, N. Gogoi, D. Baruah	681

Assessment of Genetic Diversity and Character Associations for Yield and Its Traits in Bread Wheat (*Triticum aestivum* L.)

Evaluation of Various Pest Management Modules Against Rice Yellow Stemborer, Scirphopaga

Anuj Kumar, Pooran Chand, Ravi Singh Thapa, Tejbir Singh

695

688

Page No.

Ethno-Botanical Study of Medicinal Plants from Unexplored Area of District Ramban (J&K) India

Munit Sharma, Arvind Kumar Sharma, Munish Sharma

incertulas (Walker) (Crambidae: Lepidoptera)

**Bio-stimulant Agents** 

702

Soil Characterization and Classification in an Upland of Southern Guinea Savannah Zone of Nigeria T.S. Babalola, K.S. Ogunleye, O.J. Omoju, U.C. Osakwe, A.O.A. Ilori 709

Monitoring the Influence of Anthropogenic Pollution on the Quality of Irrigation Water for Market Gardening in Yamoussoukro, Côte d'Ivoire

Tchimonbié Messikely Anoman, Don-Rodrigue Rosin Bi Voko, Dabé Doga,

Aka Niangoran Marie Stephanie Kouadio, Konan Samuel Ahoudjo, Adolphe Zeze

D. Sudha Rani, Ch. Chiranjeevi, T. Madhumathi, S. Krishnam Raju, Sk. Nafeez Umar

715

Reducing Chemical Fertilizer use to Rice through Integrated Nutrient Management in Rice (*Oryza sativa*) Groundnut (*Arachis hypogaea*) Cropping Systems

A. Polthanee, S. Gonkhamdee, M. Srisutham

721

### Indian Journal of Agricultural Research


Control Activity and Antibiotic Gene Detection of Endophytic Bacteria in Suppressing Black Pod Disease ( <i>Phytophthora palmivora</i> Butl.)  Rian Arini, Gusti Ayu Kade Sutariati, Andi Khaeruni, Teguh Wijayanto,  Novita Pramahsari Putri, Tri Joko	Cocoa 727
Effect of Different Crop Sequences on Soil Nutrient Status, Nutrient Uptake and Crop Y Western Himalayas of India  Akashdeep Singh, Pawan Pathania, Tarun Sharma, Sanjay Sharma	ield in
Effect of Application of Different Quality Water through Pitcher Irrigation and Tillage Ty Tomato Production and Soil Properties of Coastal Saline Soil of West Bengal Rahul Adhikary, Arunabha Pal, S.K. De	rpes or
Flood Threat on Cereal Crops Production in Irrigated and Rainfed Agriculture: A Study of Se Indian States  Md Aktar Hussain, Pradyut Guha	elected
The Effect of Aeroponic Potato Seeds from the Tropical Lowland and Highland Based on the Growth and Yield to Become Advanced Seeds by Irrigation Eni Sumarni, Loekas Soesanto, Noor Farid, Ardiansyah	on Size
Effect of Mixed Crops on Soil Plant Parasitic Nematodes in Lettuce Cultivation ( <i>Lactuca sat Lilyana Koleva, Ivanka Mitova</i>	tiva L.) 756
Effect of Palm Oil Mill Effluent on the Growth of Maize (Zea mays) N.C. Ewelike, J.C. Orji, I.E Adieze, V.E. Ogwudire, B.U. Uzoho, A. Ukaoma	761
Yield of Spring Wheat with the Combined Use of Sodium Selenite and Growth Regular Depending on the Conditions of Water Supply  I. Seregina, D.A. Chernyshev, V.I. Trukhachev, S.L. Belopukhov, I.I. Dmitrevskaya, F.I. Zaitsev	ulators 765
Influence of Organic and Inorganic Sources of Fertilizers on Growth and Yield of Maiz mays L.) in Fiji	e (Zea
H.K. Sachan, Deeksha Krishna, A. Prasad  Screening of Pearl Millet Genotypes Suitable for Drought Tolerance at Early Seedling S  R.C. Meena, Supriya Ambawat, C. Tara Satyavathi, Moola Ram, Vikas Khandelwal, S.L. Yadav	769 Stage 773


Esteemed Reviewers of This Issue	776
Authors Index of This Issue	776-777
Keywords Index of This Issue	777-778



## Indian Journal of Agricultural Research







Lampiran 4. List Indian Journal of Agricultural Research di Scopus

В	С	D	E	F	G
Source Title (Medline-sourced journals are indicated in Green)	Print-ISSN	E-ISSN	Active or Inactive	Coverage	Titles discontinued by Scopus due to quality issues
•	.T	•	_	▼	
780 Indian Journal of Agricultural Biochemistry	09706399		Active	2003-ongoing	
781 Indian Journal of Agricultural Economics	00195014		Active	2020-ongoing, 2017-2018, 1978-20	15
782 Indian Journal of Agricultural Research	03678245	0976058X	Active	2012-ongoing, 1990	
783 Indian Journal of Agricultural Sciences	00195022		Active	1993-ongoing, 1987, 1983, 1980-19	981, 1978, 1976, 1974
6784 Indian Journal of Agronomy	0537197X		Active	1993-ongoing	
785 Indian Journal of Anaesthesia	00195049	09762817	Active	2010-ongoing	
786 Indian Journal of Animal Research	03676722	09760555	Active	2008-ongoing	
787 Indian Journal of Animal Sciences	03678318		Active	1996-ongoing, 1987-1988, 1980-19	85, 1971-1978
788 Indian Journal of Applied Economics	09718281		Inactive	1996-1997, 1993	
789 Indian Journal of Biochemistry	00195081		Inactive	1964-1971	
790 Indian Journal of Biochemistry and Biophysics	03011208	09750959	Active	1972-ongoing	
791 Indian Journal of Biotechnology	09725849	09750967	Active	2002-ongoing	
792 Indian Journal of Cancer	0019509X	19984774	Active	1965-ongoing	
793 Indian Journal of Chemical Technology	0971457X	09750991	Active	1994-ongoing	
794 Indian Journal of Chemistry	00195103		Inactive	1983, 1969-1970	
795 Indian Journal of Chemistry - Section A Inorganic, Physical, Theore			Active	1996-ongoing	
796 Indian Journal of Chemistry - Section B Organic and Medicinal Che	m 03764699		Active	1996-ongoing, 1989-1991, 1981, 19	973-1978, 1970
797 Indian Journal of Clinical Biochemistry	09701915		Active	1986-ongoing	
798 Indian Journal of Community Health	09717587	22489509	Active	2012-ongoing	
799 Indian Journal of Community Medicine	09700218	19983581	Active	2009-ongoing	
800 Indian Journal of Computer Science and Engineering	09765166	22313850	Active	2019-ongoing	
6801 Indian Journal of Corporate Governance	09746862	24542482	Active	2008-ongoing	
5802 Indian Journal of Critical Care Medicine	09725229	1998359X	Active	2004-ongoing	
6803 Indian Journal of Dental Research	09709290	19983603	Active	1989-ongoing	
804 Indian Journal of Dermatology	00195154	19983611	Active	2006-ongoing, 1962-1990	
805 Indian Journal of Dermatology and Venereology	00195162		Inactive	1973-1975	
806 Indian Journal of Dermatology, Venereology and Leprology	03786323		Active	2002-ongoing, 1985-1995, 1976-19	182
807 Indian Journal of Earth Sciences	03795128		Inactive	2004-2006, 1979-2002	
808 Indian Journal of Ecology	03045250		Inactive	2013-2021, 1986-1987, 1974-1981	Discontinued by Scopus
6809 Indian Journal of Economics and Business	09725784		Inactive	2019-2021	Discontinued by Sconus
Scopus Sources October 2021 Accepted titles Feb. 2	022 / Disco	ontinued title:	s Feb. 2022	Serial Conf. Proc. with profile	All Conf. Proceedings

## The Effect of Aeroponic Potato

*by* Eni Sumarni

**Submission date:** 06-Dec-2022 08:55PM (UTC+0700)

**Submission ID:** 1973180436

File name: The\_Effect\_of\_Aeroponic\_Potato.pdf (371K)

Word count: 3327

Character count: 16741



# The Effect of Aeroponic Potato Seeds from the Tropical Lowland and Highland Based on Size on the Growth and Yield to Become Advanced Seeds by Irrigation

Eni Sumarni<sup>1</sup>, Loekas Soesanto<sup>2</sup>, Noor Farid<sup>2</sup>, Ardiansyah<sup>1</sup>

10.18805/IJARe.A-514

#### **ABSTRACT**

Background: Potato seed production with aeroponic technology can produce a large number of seeds, besides that it is also healthy because it uses plant seeds from tissue culture. However, how the quality of aeroponic seeds produced to become the next seed (growth capacity) has not been studied and scientigally informed. This research aimed to obtain the effect of aeroponic seed yields from the low and high plains based on their size on the growth and yield into advanced seeds. Potato seeds used were from the highland and lowland using aeroponic technology.

**Methods:** This research was conducted the highland (1000 m above sea level) from April to July 2017. Seeds were classified as  $S \le 1$  mg,  $1 \le Md \le 10$  g and  $L \ge 10$  g. This research used a randomized block design with seven repetitions. The tried factors were follows: 1. The origin of aeroponic seeds (A): A, (Iowland), A<sub>2</sub> (highland); 2. The size of aeroponic seeds (U): U, (S), U, (M), U, (D) Plant growth parameters included plant height, representations and number of tubers.

Result: The data obtained were analyzed using Duncan's multiple range test (DMRT) at the 5% level. Seeds of aeroponic yields of various sizes that are planted using drip irrigation have the potential to become advanced seeds. The L-size aeroponic seeds from the lowland produced an average of 5.9 tubers with an average weight of 68.4 g.

Key words: Aeroponic, Greenhouse, Highland, Lowland, Potato seed, Seed size.

#### INTRODUCTION

Potatoes are tubers that are widely planted throughout the world; they have good nutrition and play a role in food security (Ahmadvand et al., 2009; Karim et al., 2010). According to FAO data (2007), the average production of potato plants in Australia reached 35.9 tons per hectare, while potato production in Indonesia is lower at 18 tons per hectare. The low productivity of potatoes is caused by the limited use of high-grade potato seeds among Indonesian farmers. Such seeds are relatively expensive and seed costs involve the largest proportion of total production costs (Karim et al., 2010). It is estimated that in developing countries, the cost of procuring potato seeds makes up 40 to 70% of the cost of crop production (Masarirambi et al., 2012).

Potato seed production in Indonesia is still low due to climate constraints, pests and plant diseases; it is also due to the limited number of potato seed breeders. Generally, the breeders still use the conventional cultivation method, potato plants face insect and potato cyst nematoda problems in their growing area as well as worldwide (Malik et al., 2018; Mhatre et al., 2019; Priyank et al., 2019). Good-quality potato seeds can be obtained quickly and the result is high enough using aeroponic cultivation. Production technology of aeroponic potato seeds has started to be studied and used by some seed breeders in Indonesia in order to increase the production of potato seeds. Commercial production of aeroponic potato seeds has been developed in Korea and China. In South America, this technology has been used

<sup>1</sup>Department of Agricultural Technology, Faculty of Agriculture, Jenderal Soedirman University, Purwokerto, Indonesia. <sup>2</sup>Department of Agrotechnology, Faculty of Agriculture, Jenderal Soedirman University, Purwokerto, Indonesia.

Corresponding Author: Eni Sumarni, Department of Agricultural Technology, Faculty of Agriculture, Jenderal Soedirman University, Purwokerto, Indonesia. Email: arny0565@gmail.com

How to cite this article: Sum 3 ii, E., Soesanto, L., Farid, N. and Ardiansyah (2021). The Effect of Aeroponic Pol 3 Seeds from the Tropical Lowland and Highland Based on Size of the Growth and Yield to Become Advanced Seeds by Irrigation. Indian Journal of Agricultural Research. 55(6): 751-755. DOI: 10.18805/IJARe.A-514.

Submitted: 09-12-2019 Accepted: 18-08-2021 Online: 06-09-2021

successfully since 2006 (Otazu, 2010). The aeroponic system in Peru's International Potato Center (CIP) produces more than 100 mini bulbs per plant. The production of aeroponic potato tubers reached 1340 bulbs from 100 plants/ m² (Farran et al., 2006). In Indonesia, aeroponic technology has been developed. The result of financial analysis of the potato seed breeding business shows profitability with an R/C ratio value of 2.38, while the result obtained from the farmer breeder value is an R/C ratio equal to 2.08 (Tedy et al., 2011). Aeroponic technology in Indonesia is also being studied fo application in lowlands with root zone cooling (Sumarni, 2013; Sumarni et al., 2013a, b, c; Sumarni et.al., 2019). This technology is able to produce tubers of up to 14 tuber crops (Sumarni et al., 2013b). The production of

Volume 55 Issue 6 (December 2021) 751

aeroponic potato seeds producing 30-40 bulbs per plant in tropical highland (Sumarni et al., 2011). A potato seed tuber produced from aeroponic technology is a mini seed because it is grown inside a greenhouse and the root hangs in the air. Aeroponic potato seed production is profitable enough and produces a healthy seed because it comes from early seeds of tissue culture. However, how the quality of aeroponic seed to be the next seed (growing power) has not been further studied and informed scientifically.

Potato seeds are important in potato production (Rana et al., 2013). Poor potato seeds can reduce yields by 20% (Bentley and Vasques, 1998). The area of potato planting in the highland in Indonesia contains mostly endemic viruses and potato pests. Therefore, if the aeroponic potato seeds from the lowland and highland can produce advanced seeds, they can help the government reach self-sufficiency for seeds. This research aimed to obtain the effect of aeroponic seed yields from the lowland and highland based on their size on the growth and yield into advanced seeds.

#### MATERIALS AND METHODS

This research was conducted in the highland (1000 m above sea level) from April to July 2017. The potato seeds used were derived from the highland and lowland with aeroponic technology. Seeds were classified into sizes S, M and L. Size S were aeroponic seeds with a size of  $\leq$  1 mg, size M:  $1 \leq M \leq 10$  g and size L  $\geq$  10 g. The seeds used passed dormancy three months after harvest. The potato seeds were planted in a *greenhouse* with drip irrigation.

#### Drip irrigation

Drip irrigation system consist of pumps, main pipes, manipol pipes, lateral pipes and emitters, as well as other components such as valves, pressure gauges, filters (sand filter, disk filter and screen filter), faucets (ballvalve) and a nutrient solution tank. The distribution of nutrient solution with drip irrigation (drip system) is an open system; in other words, the nutrient solution that is streamed to the plant is not re-circulated. The pump is the driving force in the drip irrigation system to drain nutrients from the reservoir bucket

to the plants. From the results of measurements that have been carried out, the average discharge using a pump is 1.85 L/hour. The uniformity of dropper discharge for irrigation with a pump was obtained a value of 79%. The drip irrigation scheme is presented in Fig 1.

#### Experimental design and analytical data

The planting of aeroponic potato seeds with drip irrigation in this research used a randomized block design (RBD) with seven repetitions. The drip irrigation used consisted of seven lateral pipes. The distance between plants was 20 cm × 20 cm. The size of the poly bag used was 35 cm × 35 cm in diameter. Each lateral pipe contained 18 plants, so the number of plants was 126 in total. The tried factors were as follows: 1) The origin of aeroponic seed (A): A1 (lowland), A2 (highland), 2) The size of aeroponic seed (U): U1 (S), U2 (M), U3 (L).

The potato seeds planted with drip irrigation were given nutrients with a pH range of 6-6.5. Electric conductivity (EC) at 1–1.5 mS/cm was provided at the 12-ginning of planting to 30 days after planting (HST) and 2 1.5-2 mS/cm for plants aged > 30 HST to harvest (90 HST). Plant growth parameters included plant height, number of leaves and number of tubers. The c11 obtained were analyzed using Duncant's multiple drop test (DMRT) at the 5% level.

#### **RESULTS AND DISCUSSION**

## Air temperature around *greenhouse* for planting aeroponic potato seeds

The air tagperature around the planting location has an important effect on the growth and development of potato crops. The maximum air temperature in the planting greenhouse of potato seeds ranged from 24°C to a minimum of 10°C. The maximum air temperature outside the greenhouse ranged from 23°C to a minimum of 8°C (Fig 2). This air temperature provided an optimal condition for the growth and development of potato crops. High air temperatures (25-32°C) have a negative impact on the potato crop. Such temperatures can cause physiological defects in potato plants (Rykaczewska, 2013). Based on the results




Fig 1: Drip irrigation application in this research.




Fig 2: Air temperature inside and outside of the greenhouse.

of this research, the growth of potato plants occurs quickly at a temperature of 20-25°C, while tuberization and optimal tuber development occur at a temperature of 15-20°C (Rykaczewska, 1993; Van Dam *et al.*, 1996).

#### Size of potato seeds for plant height growth

The size of the aeroponic seeds planted using drip irrigation has a different effect on the plant height and number of leaves. The highest average of potato plants at the age of 10 to 50 days after planting (DAP) is achieved in the size of L-seed potatoes (51.24 cm) and in M-sized seeds at 50 days (46.96 cm). The height of potato plants aged 10 to 30 days at the M size seed was the same as S. On the other hand the M (46.96 cm) size was higher than the size of S (37.45 cm) seed at the age of 40 to 50 days after planting (Table 1).

The use of the L potato seed size produced a higher number of leaves than the M and S sizes, except at the plant age of 40 and 50 HST, which did not differ in the number of leaves between tuber sizes L (94.52 leaves) and M (77.45 leaves) (Table 2).

## The size of the aeroponic potato seeds regarding the advanced seed yield

The number of tubers produced showed that the potato seeds from the lowland produce the largest number of tubers



Fig 3: Advanced seeds produced.

compared to tuber seeds from highland aeroponic yields. The weight per tuber of the yield of potato seed production in lowland was heavier than the seed from the highland (Fig 4 and 5). The largest number of tubers was produced by aeroponic potato seeds from the lowland, with L-size (5.89 tubers) seed equal to the S size (4.86 tubers). Production of potato tuber seeds in the lowland with L size (68.36 g) results in the heaviest weight per tuber than the other sizes. The weight result per tuber for tuber seed production in the lowland for the M-size yield produce the same size as S and is heavier than tuber seed production in the highland for all seed sizes (L, M and S).

Table 1 : Differences in the use of potato seed size on plant height aged 10 to 50 days

7 Treatment			Plant height (cm)		
	10 DAP	20 DAP	30 DAP	40 DAP	50 DAP
L	23.90a	<mark>27</mark> .48a	34.48a	44.83a	51.24a
M	16.71b	20.03b	24.72b	37.96b	46.96a
6	14.51b	16.07b	20.86b	31.27c	37.45b

Table 2: Differences in the use of potato seed size on the number of leaves aged 10 to 50 days.

	·			•	
7 Treatment			Number of leaves		
rreatment	10 DAP	20 DAP	30 DAP	40 DAP	50 DAP
L	68.81a	77.81a	82.81a	86.26a	94.52a
M	50.43b	56.45b	59.45b	72.45a	77.45a
6	47.36b	49.48b	54.15b	66.69b	71.69b

The numbers in the same column followed by the same letters are no different in the DMRT test level of 5%.

The numbers in the same column followed by the same letters are no different in the DMRT test level of 5%.

Volume 55 Issue 6 (December 2021)




Fig 4: Differences in combination between the origin of potato seed production (lowland and highland) with tuber sizes (L, M, and S) on the number of tubers.




Fig 5: Differences in combination between the origin of potato seed production (lowland and highland) with tuber sizes

(L, M and S) on the weight per tuber.

The growth of plant height and the number of leaves for L-size seeds have higher growth potential compared to S- and M-size seeds. These results are consistent with a previous study, in which plants from larger seed potato tubers have higher growth potential than smaller seed tubers. The large and the la 10 r potato tubers show greater efficiency in the overall use of metabolites, as measured by growth and yield components at all levels of plant density, compared to small and medium seeds. This has implications for the duration of bulking, physiological growth and effects on the yield obtained at harvest (Masarirambi et al., 2012).

Mini aeroponic seeds of the largest size (L) are presumed to have decreased water content than M- and S-size seeds. This allows the L-size seed paintain robustness longer in order to produce a greater plant height, number of leaves and tuber yield compared to the M and S sizes. These results are similar to results of a previous study, as the larger microtubers have a shorter dormancy time and show stronger growth capabilities. Starch contact increases in line with increasing microtuber size, with a distribution of about 70-80% dry matter content. The amount of sugar is positively related to the size of the potato. Internal factors such as dry matter and carbohydrate content reveal that microtuber potatoes follow potatoes grown in the field in all

aspects. The result shows that the size of microtubers can be used as an index to assess their quality as seed potatoes, and the size of the microtubers should be at least 0.5 g to be used as potato seeds (Park et al., 2009). From the results of this research, we also found that the aeroponic potato seeds produced in the lowland have higher yield potential than potato seeds that are aeroponically planted in the highland. In this research, aeroponic seeds planted with drip irrigation became advanced seeds, as showed in Fig 3.

#### CONCLUSION

Seeds of aeroponic yields of various sizes that are planted using drip irrigation have the potential to become advanced seeds. The result of statistical analysis shows that the aeroponic seeds from the lowland (modified root) of the L size have higher potency than aeroponic seeds from the highland. The aeroponic seeds of the L size from the lowland produced an average of 5.9 tubers at an average weight of 68.4 g.

#### **ACKNOWLEDGEMENT**

We thank Kemenristekdikti for the National Strategic Grant funding (no.2045/UN23.14/PN/2017) that was given in 2017 so this research could be conducted.

#### REFERENCES

- Ahmadvand, G., Mondani, F and Golzardi, F. (2009). Effect of crop plant density on critical period of weed competition in potato. Scientia Horticulturae. 121: 249-254.
- Bentley, J and Vasques, D. (1998). The Seed Potato System in Bolivia: Organizational Growth and Missing Links. Network Paper No. 85. London, England: Overseas Development International (ODI). 11 p.
- [FAO] Food and Agriculture Organization. (2007). Food and Agricultural Commodities Production. http://faostat.fao.org/ site/339/default.aspx [08 August 2011].
- Farran, I. and Castel, M. (2006). Potato minituber production using aeroponics: Effect of plant density and harvesting intervals. American Journal Potato Research. 83: 47-53.
- Karim, M.R., Hanif, M.M., Shahidullah, S.M., Rahman, M.A., Akanda, A.M. and Khair, A. (2010). Virus free seed potato production through sprout cutting technique under net-house. African Journal Biotechnology. 9: 5852-5858.
- Malik, K., Kumar, P. and Kumar, M. (2018). Potato seed crop will be affected more by vectors under changing climatic conditions. Indian J. Agric. Res. 33(3): 173-176.
- Masarirambi, M.T., Mandisodza, F.C., Mashingaidze, A.B. and Bhebhe, E. (2012). Influence of plant population and seed tuber size on growth and yield components of potato (Solanum tuberosum). International Journal of Agriculture and Biology. ISSN Print: 1560-8530; ISSN Online: 1814-9596.
- Mhatre, P.H., Divya, K.L., Venkatasalam, E.P., Bairwa, A., Sudha, R and Berliner, J. (2019). Potato cyst nematode: A hidden enemy of potato cultivation in hills. Bhartiya Krishi Anusandhan Patrika. 34(1): 50-53.
- Otazu, V. (2010). Manual on Quality Seed Potato Production Using Aeroponics. Lima, Peru: International Potato Center
- Park, S. W., Jeon, J.H., Kim, H.S., Hong, S.J., Swath, C. and Joung, H. (2009). The effect of size and quality of potato microtubers on quality of seed potatoes in the cultivar 'Superior'. Scientia Horticulturae. 120: 127-129.
- Priyank, H.M., Divya, K.L., Venkatasalam, E.P., Bairwa, A., Sudha, R. and Berliner, J. (2019). Potato cyst nematode: A hidden enemy of potato cultivation in hills. Indian J. Agric. Res. 34(1): 50-53.

- Rana, R.K., Sharma, N., Arya, S. Kadian, M.S. and Singh, B.P. (2013). Seed potato utilization pattern and its impact on farmers profitability in Karnataka. Indian J. Agric. Res. 47(6): 488-495.
- Rykaczewska, K. (1993). Effect of temperature during growing season and physiological age of minitubers on potato plant development and yield. Bulletin of the Potato Instatute. 42(1): 39-46.
- Rykaczewska, K. (2013). The impact of high temperature during growing season on potato cultivars with different response to environmental stresses. American Journal of Plant Sciences. 4: 2386-2393.
- Sumarni, E. (2013). Pengembangan zone cooling system untuk produksi benih kentang secara aeroponik di dataran rendah tropika basah. Disertasi, Institut Pertanian Bogor. IPB. Bogor.
- Sumami, E., Ardiansyah and Sumartono, G.H. (2011). Pengembangan produksi benih kentang di sentra Banjarnegara. Laporan Riset Unggulan Daerah. Kerjasama Unsoed-Balai Penelitian dan Pengembangan Propinsi Jawa Tengah.
- Sumarni, E., Herry, S., Kudang, B.S and Saptomo, S.K. (2013a). Aplikasi pendinginan zona perakaran (root zone cooling) pada produksi benih kentang menggunakan sistem aeroponik. Journal Agronomy Indonesia. 41: 154-159.
- Sumarni, E., Herry, S., Kudang, B.S. and Saptomo, S.K. (2013b). Perpindahan panas pada aeroponik chamber dengan aplikasi zone cooling. Jurnal Biofisika. 9: 8-19.
- Sumarni, E., Herry, S., Kudang, B.S and Saptomo, S.K. (2013c). Seed potato production using aeroponics system with zone cooling in wet tropocal lowlands. Proceeding. Second Asia Pacific Symp. on Postharvest Research, Education and Extension. Eds. Acta Hort. 1011. ISHS.
- Sumarni, E., Noor, F., Loekas, S., Darjanto and Ardiansyah (2019).

  Effect of Electrical Conductivity (EC) in the Nutrition
  Solution on Aeroponic Potato Seed Production with Root
  Zone Cooling Application in Tropical Lowland, Indonesia.

  Agricultural Engineering International: CIGR Journal
  21(2): 70-77. Open access, http://www.cigrjournal.org.
- Tedy, S., Endjang, S. and Taemi, F. (2011). Peluang usaha penangkaran bibit kentang di Jawa Barat. Buletin Hasil Kajian 1(1): 1-4.
- Van Dam, J., Kooman, P.L. and Struik, P.C. (1996). Effects of temperature and photoperiod on early growth and final number of tubers in potato (*Solanum tuberosum L.*). Potato Research. 39(1): 51-62. http://dx.doi.org/10.1007/ BF02358206.

Volume 55 Issue 6 (December 2021) 755

## The Effect of Aeroponic Potato

ORIGIN	NALITY REPORT			
SIMIL	4% 10% ARITY INDEX INTERNET SOURCES	10% PUBLICATIONS	2% STUDENT PAPERS	
PRIMA	RY SOURCES			
1	oak.kribb.re.kr Internet Source			3%
2	jos.unsoed.ac.id Internet Source			1 %
3	Eni Sumarni. "Transfer Untuk Produksi Benih Tani Mulya Bersama, k Banjarnegara, Propins Sinergi, 2021	Kentang di Kelo Kabupaten	ompok	<b>1</b> %
4	jurnal.lppm.unsoed.ac	.id		1 %
5	repository.unimal.ac.ic	d		1 %
6	J Jumardi, M Melina, A distribution of the gree Trialeurodes vaporario potato plant canopy", Earth and Environmen	enhouse whiteforum Westwood IOP Conference	fly, d, within e Series:	<b>1</b> %

7	Cindy Eka Puspita, Andriani Eka Prihatiningrum, Saiful Arifin. "The Effect of POC Consetration and Planting Media on the Growth and Production of Pakcoy Aawi Plants (Brassica Rapa L).", IOP Conference Series: Earth and Environmental Science, 2022 Publication	1 %
8	docplayer.net Internet Source	1 %
9	jurnal.fp.unila.ac.id Internet Source	1 %
10	Submitted to CCS Haryana Agricultural University Hisar Student Paper	1 %
11	O. D. Ojo, A. A. Kintomo, E. A. Akinrinde, M. O. Akoroda. "Comparative Effect of Phosphorus Sources for Grain Amaranth Production", Communications in Soil Science and Plant Analysis, 2007 Publication	1 %
12	garuda.ristekbrin.go.id Internet Source	1 %
13	repository.ipb.ac.id Internet Source	1 %
14	R F Lhokitasari, M Hayati, M Rahmawati. "Growth and production of potato mini tubers	1%

(Solanum tuberosum L.) in the aeroponic system by root zone treatment and concentration of leaf-fertilizer", IOP Conference Series: Earth and Environmental Science, 2022

Publication

15

Karseno, T Yanto. "The type and distribution of sap preservatives used by farmers in Kebasen District, Banyumas Regency, Indonesia", IOP Conference Series: Earth and Environmental Science, 2019

Publication

Exclude quotes On

Exclude matches

< 1%

1 %

Exclude bibliography Or

#### **ARTICLE SUBMISSION CERTIFICATE**

Journal Name : Indian Journal of Agricultural Research

Manuscript Title : The effect of aeroponic potato seeds from the tropical lowland and

highland based on size on the growth and yield to become advanced

seeds by drip irrigation

Name of the author(s): Eni Sumarni, Loekas Soesanto, Noor Farid, Ardiansyah

Name and Address of

corresponding author : Eni Sumarni/ Department of Agricultural Technology, Faculty of

Agriculture, Jenderal Soedirman University, Purwokerto, Jl. dr.

Suparno, Karangwangkal. Postal Code. 53123.

Contact # : +62281391396079 Fax # : +62281638791

E-mail : arny0565@gmail.com

#### I (we) affirm that-

1. The manuscript has been prepared in accordance with the latest "Instruction for authors".


- 2. The article is original and has not been published previously, is not under consideration for publication elsewhere, and if accepted, it will not be published elsewhere in the same form, in English or any other language. The submission of the article has the approval of the all the authors and the authorities of the host institute where work had been carried out.
- 3. All the authors have made substantive and intellectual contributions to the article and assume full responsibility for all opinions, conclusion and statements expressed in the articles.
- 4. I (we) agree to abide by the comments of referees/editorial board and will modify the article as per their recommendations for publication.

er) Jenderal Soedirman University, Purwokerto
er) Jenderal Soedirman University, Purwokerto
er) Jenderal Soedirman University, Purwokerto
er) Jenderal Soedirman University, Purwokerto
e

This is certify that the information given above is correct to the best of my knowledge and I have no objection to the publication of the article cited above.

Eni Sumarni Signature and address Korespondensi Indian Journal of Agricultural Research: Vol Vol. 55, Issue 6. No. 751-755. (2021).

The Effect of Aeroponic Potato Seed from the Tropical Lowland and Highland Based on Size on the Growth and Yield to Become Advanced Seed by Irrigation





# AGRICULTURAL RESEARCH COMMUNICATION CENTRE

www.arccjournals.com

Date: 18-08-2021

Reference ID. ARCC/A-514

Eni Sumarni,

Department of Agricultural Technology, Faculty of Agriculture, Jenderal Soedirman University, Purwokerto, Jl. dr. Suparno, Karangwangkal. Postal Code. 53123.

**Acceptance of manuscript** 

Dear Dr. Sumarni,

We are pleased to inform you that your manuscript has been accepted for publication in **Indian Journal of Agricultural Research**. Your submission is a well-thought out piece of writing and follows many of journal guidelines. The editors agreed that your submission showed great writing skills.

Manuscript The effect of aeroponic potato seeds from the tropical Title: lowland and highland based on size on the growth and yield to become advanced seeds by irrigation

Author(s): Eni Sumarni, Loekas Soesanto, Noor Farid, Ardiansyah

Congratulations to you once again on your article acceptance in ARCC Journals, and we look forward to receiving more of your good submissions.

With Best Wishes and Seasonal Greetings,

**Gaurav Gupta** Managing Editor