

9. Characteristics of White Corn Noodle Substitued by Tempeh Flour

Jurnal nasional akreditasi

Jurnal Teknologi dan Industri Pangan

Vol 23 no 2, Desember 2012

1. Cover
2. Editor
3. Daftar isi
4. Artikel
5. Bukti akreditasi (Sinta 2)
6. Bukti koresponden
7. Turnitin

ISSN 1979-7788

Jurnal TEKNOLOGI & INDUSTRI PANGAN

Volume xxiii Nomor 2 Tahun 2012



Publikasi Resmi

Perhimpunan Ahli Teknologi Pangan
Indonesia (PATPI)

bekerjasama dengan

Departemen Ilmu dan Teknologi Pangan
Fakultas Teknologi Pertanian
Institut Pertanian Bogor

Member of



Terakreditasi
No 110/DIKTI/Kep/2009

Jurnal TEKNOLOGI & INDUSTRI PANGAN

ISSN 1979-7788
VOLUME XXIII NUMBER 2
December 2012

Copyright © 2012 Indonesian Association of Food Technologists in collaboration with the Department of Food Science and Technology, IPB

GOVERNING BOARD

Rindit Parabayun, Prof Dr
Chair of Indonesian Association of Food Technologists
(PATPI)

Feri Kusnandar, Dr
Head of Department of Food Science and Technology,
FATETA-IPB

EDITOR IN CHIEF

Harsi D. Kusumaningrum, Dr

NATIONAL EDITORIAL BOARD

Ardiansyah, Dr	Siswa Setyahadi, Dr
Budiatman Satiawihardja, Dr	Sri Widowati, Dr
C. Hanny Wijaya, Prof Dr	Tejasari, Dr
Deddy Muchtadi, Prof Dr	Umar Santoso, Prof Dr
Indah Eprilianti, Dr	Winiati P. Rahayu, Prof Dr
Ratih Dewanti-Hariyadi, Dr	

INTERNATIONAL EDITORIAL BOARD

Irwandi Jaswir, Prof Dr	(Malaysia)
Indrawati Oey, Prof Dr	(New Zealand)

STAFF EDITORS

Anton Rahmadi, MSc
Eko Hari Purnomo, Dr
Nancy Dewi Yuliana, Dr
Nurheni S. Palupi, Dr
Puspo Edi Giriwono, Dr

ADMINISTRATION AND FINANCIAL

Nur Wulandari, MSi
Sutrisno Koswara, Ir
Ifti Hidayah
Vivi Silvia Januar, Amd

Journal of Food Technology and Industry (JFTI) is an Indonesian accredited journal and is a lead reference journal in the field of food science and technology. The journal was established in 1990 and previously known as Bulletin in Research Food of Science and Technology. The name has been updated in 2001 and ever since it became official journal for Indonesian Food Technologist Association (PATPI). The journal is published in collaboration with Department Food Science and Technology, Bogor Agricultural University. The online version is accessible at <http://journal.ipb.ac.id/index.php/jtip> No E-ISSN: 2087-751X.

JFTI is published twice a year, in June and in December. JTIP publishes high quality research articles in food sciences, food technology or its applications in food industry. JFTI accepts Indonesian and English language articles.

Composition of the editorial boards has been restructured since Volume XXIII, issue 2 in 2012 when new chairman of PATPI was elected for the period of 2012-2014. JFTI includes International Editors for internationalization purposes and for improving the performance of the journal.


For further information and correspondence, please contact the Secretariat for Journal of Food Technology and Industry, Department of Food Science and Technology, Faculty of Agricultural Technology, Bogor Agricultural University, IPB Darmaga Campus, PO Box 220, Bogor 16002; Telp/Fax: (0251) 8626725; E-mail: jurnaltip@yahoo.com.



Jurnal
Teknologi dan Industri Pangan
Indonesian Association of Food Technologists (PATPI)
in Collaboration with
Department of Food Science and Technology,
Faculty of Agricultural Technology, Bogor Agricultural University

pISSN 1979-7788
eISSN 2087-751X

Vol. 23 No. 2 (2012): Jurnal Teknologi dan Industri Pangan



Published: 2012-12-30

Research Paper

PEMBUATAN BAHAN BAKU SPREADS KAYA KAROTEN DARI MINYAK SAWIT MERAH MELALUI INTERESTERIFIKASI ENZIMATIK MENGGUNAKAN REAKTOR BATCH [Preparation of Red Palm Oil Based-Spreads Stock Rich in Carotene Through Enzymatic Interesterification in Batch-type Reactor]

. Sugiyono, Mario Wibowo, Soenar Soekopitojo, Nur Wulandari

117

 PDF

AKTIVITAS ANTIBAKTERI EKSTRAK TUMBUHAN LUMUT HATI (*Marchantia paleacea*) TERHADAP BAKTERI PATOGEN DAN PERUSAK PANGAN [Antibacterial Activity of Liverwort (*Marchantia paleacea*) Extract on Pathogenic and Food Spoilage Bacteria]

Reza Fadhillah, Eka Aditya Iskandar, Harsi Dewantari Kusumaningrum

126

 PDF

PRODUKSI DAN EVALUASI KUALITAS SUSU BUBUK ASAL KAMBING PERANAKAN ETTAWA (PE) [Production and Quality Evaluation of Ettawa-Crossbred Goat Milk Powder]

. Widodo, Afina Viyunnur Rachmawati, Rina Chulaila, I Gede Suparta Budisatria

132

 PDF

BIOADSORPSI Hg(II) OLEH PATI SAGU TAUT SILANG FOSFAT [Bioadsorption of Hg(II) by Crosslinked Sago Starch Phosphate]

Jorion Romengga, Tun Tedja Irawadi, Sr Sugiarti

140

 PDF

PENINGKATAN MUTU MI KERING JAGUNG DENGAN PENERAPAN KONDISI OPTIMUM PROSES DAN PENAMBAHAN MONOGLISERIDA [Quality Improvement of Dried Corn Noodle through the Optimization of Processing Conditions and Addition of Monoglyceride]

. Subarna, Tjahja Muhandri, Budi Nurtama, Antung Sima Firlieyanti

146

 PDF

KARAKTERISTIK PENGERINGAN CHIPS MANGGA MENGGUNAKAN KOLEKTOR SURYA KACA GANDA [Characteristics of Mango Chips Drying Using a Double Plated Solar Collector]

. Ansar, . Cahyawan, . Safrani

153

 PDF

POTENSI LACTOBACILLUS YANG DIISOLASI DARI AIR SUSU IBU UNTUK MENCEGAH DIARE [Potential of Lactobacillus Isolated from Breast Milk to Prevent Diarrhea]

Lilis Nuraida, . Hana, Apriliana W. Hartanti, Endang Prangdimurti

158

 PDF

FORMULATION AND PROCESS OPTIMIZATION OF MUFFIN PRODUCED FROM COMPOSITE FLOUR OF CORN, WHEAT AND SWEET POTATO [Formulasi dan Optimasi Proses Produksi Mufin dari Tepung Komposit Jagung, Gandum dan Ubi Jalar]

Eko Hari Purnomo, Azis B. Sitanggung, Denny S. Agustin, Purwiyatno Hariyadi, Stefani Hartono
165

 PDF

STRUCTURAL ANALYSIS OF SPRAY-DRIED COCONUT SHELL LIQUID SMOKE POWDER [Analisis Struktural Bubuk Asap Cair Batok Kelapa Hasil Pengeringan Semprot]

Satrijo Saloko, Purnama Darmadji, Bambang Setiaji, Yudi Pranoto
173

 PDF

CHARACTERISTICS OF WHITE CORN NOODLE SUBSTITUED BY TEMPEH FLOUR [Karakteristik Mi Jagung Putih dengan Substitusi Tepung Tempe]

Nur Aini, V. Prihananto, S. Joni Munarso
179

 PDF

SURVIVAL OF *Cronobacter sakazakii* IN SKIM MILK DURING SPRAY DRYING, STORAGE AND RECONSTITUTION [Ketahanan Hidup *Cronobacter sakazakii* dalam Susu Skim selama Proses Pengeringan Semprot, Penyimpanan dan Rekonstitusi]

Ratih Dewanti Hariyadi, Fenny Larasati, Lilis Nuraida
186

 PDF

QUANTIFICATION OF *Salmonella Typhimurium* REDUCTION DURING COLD STORAGE OF RAW SHRIMPS IN THE PRESENCE OF SODIUM METABISULFITE [Kuantifikasi Reduksi *Salmonella Typhimurium* pada Udang Segar selama Penyimpanan Dingin dengan Penambahan Natrium Metabisulfit]

Harsi Dewantari Kusumaningrum, Andiarto Yanuardi
193

 PDF

PHYSICOCHEMICAL PROPERTIES OF INSTANT PUMPKIN JAVANESE NOODLE GRAVY [Sifat Fisikokimia Saus Bubuk Mi Jawa Instan dari Labu Kering]

Ng Kim Yee, Yusnita Hamzah
199

 PDF

EFFECT OF PRETREATMENTS ON CHEMICAL AND ANTIOXIDANT PROPERTIES OF SKY FRUIT (*Swietenia macrophylla*) SEED OIL [Pengaruh Pra-perlakuan Terhadap Sifat Kimia dan Antioksidan Minyak Biji Buah Tunjuk Langit (*Swietenia macrophylla*)]

Lau Chiew Ping, Nor Hayati Ibrahim, Hayati Mohd Yusof

205



EXTRACTION AND CHARACTERIZATION OF MUCILAGE FROM LEAVES OF *Pereskia bleo* (ROSE CACTUS) [Ekstraksi dan Karakterisasi Getah Daun Kaktus Mawar (*Pereskia bleo*)]

Ng Tze Hong, Nor Hayati Ibrahim

210



Short Communication

AKTIVITAS ANTIBAKTERI FRAKSI-FRAKSI EKSTRAK SIRIH HIJAU (*Piper betle* Linn) TERHADAP PATOGEN PANGAN [Antibacterial Activity of Fractionated Green Sirih (*Piper betle* Linn) Extract Against Food Pathogenic Bacteria]

. Suliantari, Betty S. L. Jenie, Maggy T. Suhartono

217



Citedness Scopus:



Sidebar Menu:

[Editorial Team](#)

[Peer Reviewer](#)

[Abstracting and Indexing](#)

[Focus and Scope](#)

[Ethic in Publishing](#)

[Author Guideline](#)

Journal Template:



Accreditation



Indexed by:





Publisher:



Department of Food Science and Technology, Faculty of Agricultural Technology and Engineering,
IPB University

In Collaboration with



Indonesian Association of Food Technologists (IAFT)

Member of IUFOST:



Plagiarism Check:



Google Scholar Citation:



Citation Indices	All	Since 2015
Citations	2125	1687
h-index	22	20

I10-index10	63	49
-------------	----	----

SINTA Score:

S2 Sinta Score	
22	19
H-Index	HS-Index
2093	1657
Citations	5 Year Citations

Current Issue



[Visitor Statistic](#)

[pISSN: 1979-7788](#)

[eISSN: 2087-751X](#)



This journal is published under the terms of the is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](#)

Platform &
workflow by
OJS / PKP

CHARACTERISTICS OF WHITE CORN NOODLE SUBSTITUED BY TEMPEH FLOUR

[Karakteristik Mi Jagung Putih dengan Substitusi Tepung Tempe]

Nur Aini^{1)*}, V. Prihananto¹⁾ and S. Joni Munarso²⁾

¹⁾ Department of Agricultural Technology, Jenderal Soedirman University, Purwokerto

²⁾ Center for Estate Crops Research and Development, Bogor

Submitted November 08th 2012 / Accepted December 07th 2012

ABSTRACT

Different corn type and processing of corn flour can produce flour with different physical and chemical characteristics. Processing of such flour into noodle will also result in different properties of noodle. While substitution of corn flour with tempeh flour can improve the protein content of the noodle it also impair its sensory characteristics. The objectives of this research were to determine the best combination of corn flour-type, corn flour processing and proportion of corn:tempeh flour to produce the best corn noodles based on physical, chemical and sensory properties. The observed variables of this study were moisture, ash, soluble protein, fat, elongation, colour, flavour, taste, and preferences. The results showed that based on physical, chemical and sensory properties, the best corn noodle could be made from flour made from waxy corn soaked for 24 hours and a ratio of corn:tempeh flour of 80:20. The noodle produced had 12.1% of total protein content, 2.8% of soluble protein, 8.4% of fat, 5.9% of moisture, 3.6% of ash, 69.9% of carbohydrate, 17.6% of elongation, yellow colour, slight flavour of corn/soybean, and 2.6 of preferences.

Keywords: corn, noodle, soaking, tempeh, waxy

ABSTRAK

Jenis jagung (waxy dan non waxy) serta proses pengolahan tepung menghasilkan perbedaan sifat tepung jagung, dan apabila diaplikasikan pada pembuatan mi akan menghasilkan sifat mi yang berbeda. Substitusi tepung tempe pada pembuatan mi jagung meningkatkan kadar proteinnya, tetapi dapat mengubah sifat sensori mi. Penelitian bertujuan menentukan kombinasi perlakuan terbaik (antara jenis jagung, proses pengolahan tepung dan proporsi tepung jagung:tempe) yang dapat menghasilkan mi jagung terbaik (berdasar sifat fisik, kimia dan sensoris). Variabel yang diukur meliputi kadar air, kadar abu, protein terlarut, lemak, pemanjangan/elongasi, warna, aroma, rasa dan kesukaan. Hasil penelitian menunjukkan bahwa berdasar sifat fisik, kimia dan sensoris, mi jagung terbaik dibuat dari tepung jagung waxy yang dibuat melalui proses perendaman selama 24 jam, dan proporsi tepung jagung: tempe yang digunakan pada pembuatan mi 80:20 (V₂P₁T₂). Karakteristik mi adalah sebagai berikut: kadar protein total 12,1%, protein terlarut 2,8%, lemak 8,4% b, air 5,9% bk, abu 3,6%bk, karbohidrat 69,9% bk, elongasi 17,6%, warna kuning, aroma jagung/tempe yang agak terasa, dan kesukaan 2,6.

Kata kunci: jagung, mi, perendaman, tempe, waxy

INTRODUCTION

White corn has high carbohydrate, so it potentially to be used as staple food like noodle. Before processed into a noodle, white corn flour was made first. Corn flour manufacturing consist of seed polishing, washing, soaking, grinding and sieving proces. Soaking in the corn flour manufacture has several functions, namely to separate the corn from germ, skin, and soften texture of to facilitate corn milling (Eneje *et al.*, 2004) and as one of the natural fermentation techniques that can produce certain functional properties of corn flour (Aini *et al.*, 2010) (Yuan *et al.*, 2008). Difference characteristics of the flour caused by soaking will affect characteristics of corn noodles, presumably.

Based on the amylosa-amylopectin levels, white corn can be divided in to non-waxy and waxy corn. Amylose and amylopectin fractions have different properties that will make a difference product characteristic when applied to the food product. Protein content of corn noodles do not meet the noodle standards quality according to SNI 01-2974-1996, which is less than the minimum 8% wb. The corn noodles protein content can be improved by other materials substitution that have a high protein content, such as tempeh flour (Rodriguez *et al.*, 2004). Protein content of tempeh flour reaches 41.64% db. Presence of tempeh flour can increase the protein content but change the sensory properties of corn noodles.

The objectives of this research were to determine the best combined treatment (between corn flour-type, corn flour processing and proportion of corn:tempeh flour) to produce the best corn noodles (based on physical, chemical and sensory properties).

Paper was Presented at International Conference on "Future of Food Factors", October 3-4, 2012, Jakarta, Indonesia.

*Correspondence Author: email nuraini@unsoed.ac.id; nuraini.1973@gmail.com /nuraini_munawar@yahoo.com, phone+62 281 621094

MATERIALS AND METHODS

Materials

The materials used in this study were waxy corn from Gorontalo, non-waxy white corn from Temanggung, tempeh, corn starch, additional materials for the noodles and chemicals for analysis.

Making of corn flour

The processing of white corn flour was conducted by modification method of Aini *et al.* (2009). Modification was done by soaking treatments (soaking for 24 hours and without soaking). Corn flour then sifted 100 mesh.

Making of corn noodles

To make corn noodles, first the corn flour, tempeh and corn starch were mixed. Then, salt was added gradually while stirring until form a semi solid noodles dough. Once dissolved, the solution was mixed with some warm water. Once the dough was semi-solid form, the yolks and ovalet were added. After all the ingredients mixed, added the remaining water into the mixture then poured in the pan. The next process, it was steaming for 3 minutes to form a sheet of dough that was ready to be molded into noodles. The noodles sheets was trimmed and molded, then dried with cabinet dryer at $\pm 60^{\circ}\text{C}$ until dry.

Variables

The observed variables of this study were moisture content (AOAC, 2000), ash (AOAC, 2000), soluble protein (AOAC, 2000), fat (AOAC, 2000) and elongation (Muhandri *et al.*, 2011). In addition, the analysis of total protein and carbohydrate (by difference) was also performed for the best treatment. The observed sensory variables include colour, texture (firmness), flavour, taste, and preferences were measured by the scoring method.

Design experimental

This research use Completely Randomized Design (CRD) with 20 factorial combinations treatment repeated 2 times. The tested factors were the types of corn (V) i.e. non-waxy (V1) and waxy (V2); the soaking process in the making of corn flour (P) i.e. without soaking (Po) and 24-hour soaking (P1); and the proportion of corn:tempeh flour (T) which consists of five levels, namely 100:0(T0); 90:10(T1); 80:20(T2); 70:30(T3) and 60:40(T4).

Statistical analysis

The obtained physical and chemical data were analyzed with "F" test on 5 percent and 1% significant level. Duncan's Multiple Range Test (DMRT) was conducted if there any significant diversity to determine the difference effects between single and combination treatment level. The sensory variables data were analyzed by the Friedman test followed by multiple comparisons as a further test if there any significant differences.

RESULTS AND DISCUSSION

Moisture content

Moisture content is an important food constituent. It can be expressed in the form of water stating the amount of absolute moisture content in food as a food component in the various number. The types of corn affect to the noodle water content, significantly. Non-waxy corn noodles had a higher water content ($7.53 \pm 0.32\%$) than waxy corn noodles ($5.16 \pm 0.14\%$) as shown in Figure 1. The water content differences of the corn noodle due to the raw materials differences, mainly corn flour. Non-waxy corn flour has higher moisture content (11.43%) than waxy flour (8.5%).




Figure 1. The moisture content of corn noodles

The soaking did not significantly affect the moisture content of corn noodles. The water content of maize flour noodles without soaking corn kernels (P0) is $5.46 \pm 0.22\%$ and the soaking seeds (P1) yields $5.75 \pm 0.32\%$ dry basis moisture. Noodles was made from corn flour without undergoing the process of soaking corn kernels, have a higher water content, but not significantly different. It is due to the ability of the flour to absorb the water associated with higher levels of amylose.

According to Aini and Hariyadi (2010), a straight-chain amylosa has the ability to absorb water is higher than the branched chain amylopectin. The higher levels of amylosa, the lower water absorption capacity of corn flour. Corn flour that produced without soaking of corn grits have amylosa content 3.69% dry basis. While corn flour that produced from soaking of corn grits have amylosa content 2.17% dry basis, because soaking corn kernels for 24 hours will solute of the amorphous granules. Most of the amylosa present in the amorphous regions, so the leaching section during soaking reduces amorphous amylosa.

Tempeh flour substitution did not significantly affect to the moisture content of corn noodles as well as the interaction between treatments. The moisture content of the corn noodles that have a proportion of corn : soybean flour 100:0(T0) is $6.03 \pm 0.29\%$ db, 90:10(T1) is $6.25 \pm 0.21\%$ db, 80:20(T2) is $5.60 \pm 0.31\%$ db, 70:30(T3) is $5.49 \pm 0.21\%$ db, and 60:40(T4) is $4.64 \pm 0.19\%$ db.

Corn noodle had moisture content under 10%. It was lower than the water content of yellow corn noodles according Yuan *et al.* (2008). The maximum moisture content of corn noodles according to SNI 01-2974-1996 is 8% for the first quality and 10% for second quality.

Fat content

Fat has important functions in food processing i.e. as a source of energy, play a role in the formation of texture and sensory quality of food products, medium heat in the frying processes well as the solvent for the essential fat-soluble vitamin.

The types of corn significantly affect to the fat content of corn noodles. Fat content of noodles from waxy corn flour was higher ($10.18 \pm 0.41\%$) than non-waxy corn ($6.78 \pm 0.27\%$) (Figure 2). This was due to waxy corn has a higher fat content (2.94%) than non-waxy corn (1.58%). According to Lu *et al.* (2003), diversity levels of fat in the product due to fat content variations of raw materials.





Figure 2. Fat content of corn noodles

The manufacturing techniques of corn flour gave significantly difference to the levels of noodles fat. Corn noodle produced from the immersed corn flour had a lower fat content ($8.02 \pm 0.32\%$) than corn noodles made for non immersed corn flour ($8.94 \pm 0.43\%$) as shown in Figure 3. This is consistent to the statement Belyea *et al.* (2004) that wet milling of rice flour to produce a smaller amount of fat.



Gambar 3. Effect of soaking in the processing of corn flour to the fat content of corn noodles

The difference in fat content as well as soaking in accordance with Aini *et al.* (2009) that corn fermentation techniques with soaking resulted partly dissolved germ. Fat is a macro chemical component that dominates the germ, so the dissolved germ can reduce the fat content of corn flour. It is suspected as the cause of reduced fat noodles by soaking corn seeds. Fat content of corn flour made through and without soaking proces were $2.43 \pm 0.07\%$ db and $2.94 \pm 0.06\%$ db, respectively.

The proportion of soybean flour significantly affect the fat content of corn noodles, as shown in Figure 4. Increased levels

of fat noodle flour substitution with increasing proportions due to the fat content of soybean tempeh flour is quite high (23.88% db).

Soluble protein

Soluble protein is the percent protein contained in the materials that can be extracted by or dissolved in the water under specified conditions (Young *et al.*, 2004). Foodstuffs that have high levels of soluble protein has a good quality because it is more easily utilized by the body.

Type of maize significantly affect soluble protein noodles. Noodles from waxy maize has higher soluble protein content ($2.52 \pm 0.01\%$ db) compared to corn noodles non-waxy ($1.97 \pm 0.08\%$ db). Variations in soluble protein content of corn noodles due to differences in the base material, which is waxy maize (1.64% db) has a higher protein content than non-waxy maize (1.16% db).




Figure 4. Fat content of corn noodles that influenced by the proportion of corn and tempeh flour




Figure 5. Soluble protein content of different types corn noodle

Soaking in the processing of corn flour can significantly affect on the soluble protein of corn noodles. Noodles made from corn flour with soaking (P1) has a higher soluble protein ($2.85 \pm 0.11\%$ db) than corn noodles from corn flour without soaking (P0), which had $2.13 \pm 0.14\%$ db soluble protein levels (Figure 6). It was influenced by the levels of corn flour dissolved protein. Corn flour made without soaking and soaking had 9.28% and 3.59% of soluble protein respectively. Increased levels of soluble protein of immersed corn flour thought to be caused by a protease enzyme activity during fermentation which can break down proteins into small molecular weight amino acids that soluble in the water (Abdel-Aal, 2008).

The higher tempe flour in the noodles, the higher levels of noodle soluble protein (Figure 7). This was due to tempeh flour had higher levels of soluble protein (12.89%) than corn starch. During fermentation of soybeans into tempeh, *Rhizopus oligosporus* produces protease enzymes (Rodríguez *et al.*, 2006). Protein complexes occur reshuffle into simpler compounds. Protein content expressed as total nitrogen levels did not change during fermentation, but the levels of soluble protein and free amino acid was increased.

Soluble protein is the percent protein contained in the materials that can be extracted by or dissolved in the water under specified conditions (Rodríguez *et al.*, 2004). Foodstuffs that have high levels of soluble protein has a good quality because it is more easily utilized by the body.




Figure 6. The soaking process effect upon the corn noodles soluble protein content




Figure 7. Soluble protein of cor noodles affected by tempeh flour proportion

Texture

Texture of noodle after cooking is one of important indicator to determine the quality of noodles. Texture conducted in an objective measurement using the texture analyzer to determine the noodle elongation (elongation). Elongation show a maximum length of noodles before breaking. Elongation is one of the most important parameters of noodles (Bellido and Hatcher, 2009). High percentage of elongation showed the noodle are not easily broken. This is important characteristic because the noodle is not expected to crumble when eaten (Muhandri and Subarna, 2009).

The interaction between type of corn, soaking in the manufacture of corn flour and tempeh flour substitution significantly affect the noodle elongation. Elongation is a level of the material to stretch or pull before the material to be broken.

Elongation measurements performed to determine the noodles resistance againsts the pull. The good noodle have high tensile strength, characterized by a high percentage of noodles elongation.

The noodle from non-waxy unsoaked corn flour with 20% of tempeh (V1P0T2) had the lowest elongation 8.68% (Figure 8). This is presumably its because the noodles had the highest water content and cooking loss. The more water that was bound by the components in the noodles, the weaker bond was formed. The high cooking loss was causing noodle to be easily broken up with noodle heavy shrinkage. It can lower the elongation.




Figure 8. Interaction of the corn type, processing of corn flour and proportion of corn:tempeh flour to the noodle elongation (V₁=non-waxy corn, V₂=waxy corn, P₀= processing of flour without soaking P₁= 24-hour soaking, T = proportion of corn:tempeh flour, T₀= 100:0, T₁= 90:10, T₂= 80:20, T₃= 70:30, T₄= 60:40)

Non-waxy corn flour noodle had a higher elongation than waxy corn noodles. Non-waxy maize contains more straight-chain amylose so it make easier to change shape due to the pull. Waxy corn has higher levels of amylopectin and the presence of branched structures causing the noodles produced less elongation.

Percentage of corn noodles extension ranged from 8.679 to 28.205%, similar to rice flour noodles ranged from 27.84 to 30.91% (Lu *et al.*, 2003). This range was similar to the extension percentage of noodles brand Myojo (28.66%) or Pop Mi (22.5%). However, the research results of Bellido and Hatcher (2009) showed that the percent elongation of rehydrated wet wheat noodles was 107.35%.

The low elongation of corn noodles or other noodles allegedly due to the substantial rehydration during processing of dried noodles into a wet noodle (Bellido and Hatcher, 2009). The solids loss during cooking caused by dispersion of amylose in hot water can be occurred. Substantial solids loss causing the increased of rehydration so that the noodles texture to be less compact and easily broken.

Incompletely retrogradation was causing the low percent of elongation. According to Aini *et al.* (2010), at gelatinization, amylose leach from starch granules and form fat-amylose

inclusion complexes. This complex can reduce the tendency of amylose to bind, to form a gel and to retrograde. It causes the formed hydrogen bonds less compact and produce easy to broken noodles. In addition, alternately steaming process causes the dough through a omission stage that allow to the precipitation.

In the absence of gluten, corn noodle texture was more delicate than wheat noodles. Functional properties of soy protein may help to improve the texture of noodles, but the functional properties of carbohydrates in the dough was more dominant in the binding process water, viscosity and dough development. Functional properties of carbohydrates was determined by several factors including the shape and size of starch granules (Moorthy, 2002). Texture strength was supported by the interaction of crystalline starch inter molecular bonds that were less powerful.

Sensory properties

Sensory properties were important properties that will affect the consumer's decision to buy or not. Analyzed sensory properties of corn noodles were include the colour, flavour, taste and preferences.

Colour

Combination treatment of corn types, soaking and tempeh flour substitution had significantly affect to the colour of corn noodles. The result of sensory test showed that the corn noodle colour varies from white to yellow (score of 1 to 4.87) (Figure 9). Variations of corn noodles colour is influenced by raw material, namely white and yellow corn flour. Moreover, soaking process also resulted in the colour of noodles differences. According to Aini *et al.* (2009), the degree of corn white flour that made from soaking corn is higher than non-soaking.




Figure 9. Interaction between corn type, processing of corn flour and proportion of corn:tempeh flour to the colour of noodle (V₁=non-waxy corn, V₂=waxy corn, P₀= processing of flour without soaking P₁= 24-hour soaking, T= proportion of corn:tempeh flour, T₀= 100:0, T₁= 90:10, T₂= 80:20, T₃=70:30, T₄ = 60:40)

Increasing of the proportion of tempeh flour make the colour more brown. The high protein content of tempeh flour resulted in

non-enzymatic browning reactions during drying so that the colour becomes golden brown. In addition, a grayish-white mycelia formed by the mold in the tempeh flour will affect the colour of the flour. This will affect the resulting colour of the corn noodles. In the non-waxy corn noodles, the addition of tempeh flour resulted in increased intensity of the brown-gray colour, while the noodles from waxy corn, the addition of tempeh flour makes better corn noodle colour (more yellow).

Colour is a very important on food. Before other factors considered, the visual colour factors was appearing first and sometimes determines the selection of food products (De Groot and Kimenju, 2008). At corn noodles, the higher value of the resulting indicates the better colour of the noodles with the highest colour is yellow. Usually the panelists preferred colour for the noodles is opaque to yellow (Chung *et al.*, 2012). The colours on the noodles are formed by the heating during drying. During the heating process, the amino acids in soy protein were primarily reacted with a reducing sugar of carbohydrate formed a brownish yellow pigment (melanoidin).

Flavour

Combination of corn types, soaking and tempeh flour substitution significantly affect to the corn noodles flavour. The flavour of corn/tempeh were detected in corn noodles ranged from strong-stronger (2.72) to slightly strong (3.56). Effect of interaction between corn types, processing of corn flour and proportion of corn:tempeh flour to the flavour of corn noodles can be seen in Figure 10.




Figure 10. Interaction between the corn type, processing of corn flour and proportion of corn:tempeh flour to the flavour of noodle (V₁=non-waxy corn, V₂=waxy corn, P₀= processing of flour without soaking P₁= 24-hour soaking, T= proportion of corn:tempeh flour, T₀= 100:0, T₁= 90:10, T₂= 80:20, T₃=70:30, T₄ = 60:40)

The expected flavour of corn noodles was not strong (Inglett *et al.*, 2005). Combination treatment of local corn types, non-soaking and non-substitution of tempeh flour (V1P0T0) is the best combination. There were some other treatments that also had same value. Combination of local corn types with soaking

and 20% tempeh flour substitution (V2P1T2) had the strongest flavour, although there is some combination of treatments that had similar flavour.

The presence of corn/tempeh flavour allegedly due to the fermentation process during soaking and the manufacture of substituted tempeh. The flavour of fermented corn is more acidic than the unfermented (Maurel *et al.*, 2003). Tempeh flour substitution also gives pungent flavour, because during the fermentation of soybeans into tempeh was increasing the number of amino acids, especially phenylalanine and tryptophan increased 20%. Tryptophan is a volatile amino because it has an flavouric side chain, so the its increase is to strengthen the distinctive flavour of tempe.

Taste

The addition of tempeh flour as a substituted was lowered the taste of produced corn noodles. Combination of soaking and tempeh flour substitution gave a significant effect to the corn/tempe taste of corn noodles. Sense of taste is referred to the assessment of corn, tempeh or other foreign undesirable taste contained in the corn noodles. Effect of interaction between corn types, processing of corn flour and proportion of corn:tempeh flour to the taste of corn noodles can be seen in Figure 11.




Figure 11. Interaction between the corn type, processing of corn flour and proportion of corn:tempeh flour to the taste of noodle (V₁=non-waxy corn, V₂=waxy corn, P₀= processing of flour without soaking P₁= 24-hour soaking, T= proportion of corn:tempeh flour, T₀= 100:0, T₁= 90:10, T₂= 80:20, T₃=70:30, T₄= 60:40)

According to Chelule *et al.* (2010), corn fermentation caused a bitter taste, however it wasn't happened in the 24-hour soaking fermentation. The corn noodles of soaking corn tend to had a better taste characterized by a higher taste score than the noodles of unsoaked corn flour. Unusual characteristic tempeh taste found in the noodles increased as the increase of the proportion of tempeh flour substitution.

Preference

Preference was the conclusion of the quality attributes assessment such as colour, texture, and flavour to determine the accepting of presented material (Van-Kleef *et al.*, 2005). Noodle was made from waxy corn, non-soaking and 10 % proportion of tempeh flour % (V2P0T1) was the most preferred product, but not significantly different from V1P1T1, V1P1T0, V2P1T2, and V2P0T4 corn noodles (Table 1). The high value of the combined treatment was caused by its texture (the elasticity) and the best colour, presumably. The most favorite corn noodles had yellow colour, less strong of tempeh flavour and taste, and chewy texture.

The effectiveness index test results show that the best noodle was produced from waxy corn flour made from soaking corn and 20% substitution of tempeh flour. Analysis of nutrient content of the best treatment noodles compared with the Indonesian National Standard (SNI) of noodle SNI 01-2974-1996 can be seen in Table 1. From Table 1 it can be seen that the product can virtually meet the SNI criteria, except for ash content that was worth of 3.18%, still slightly higher than the SNI, 3%.




Figure 12. Interaction between the corn type, processing of corn flour and proportion of corn:tempeh flour to the preference of noodle (V₁=non-waxy corn, V₂=waxy corn, P₀= processing of flour without soaking P₁= 24-hour soaking, T= proportion of corn:tempeh flour, T₀= 100:0, T₁= 90:10, T₂= 80:20, T₃=70:30, T₄= 60:40)

Table 1. Nutritional value of best corn noodle comparison with SNI

Component	SNI	Corn Noodle
Water content (% wb)	Max 10	5.67
Soluble protein (%db)	-	2.89
Total Protein (%wb)	Min 8	12.12
Fat (%db)	-	8.76
Ash (%wb)	Max 3	3.16
Carbohydrate (%)		70.19
Energi (kkal)		310.7
Texture (N)		1.76
Elongation (%)		12.17

CONCLUSION

Based on physical, chemical and sensory properties, the best corn noodle was made from waxy corn flour in the combined treatment 24 hours of soaking with the substitution ratio of corn flour:tempeh flour ($V_2P_1T_2$). The characteristics was 12.1% of total protein content, 2.8% of soluble protein, 8.4% of fat, 5.9% of moisture, 3.6% of ash, 69.9% of carbohydrate, 17.6% of elongation, yellow colour, slightly flavour of corn or soybean, and 2.6 of preferences.

ACKNOWLEDGEMENTS

Thanks to Indonesian Agency for Agricultural Research and Development (IAARD) that has funded this research through the Agricultural Research and Higher Education (KKP3T) Partnership in 2010.

REFERENCES

- Abdel-Aal ESM, Rabalski I. 2008. Effect of baking on nutritional properties of starch in organic spelt whole grain products. *Food Chem* 111: 150-156. DOI: [10.1016/j.foodchem.2008.03.050](https://doi.org/10.1016/j.foodchem.2008.03.050).
- Aini N, Hariyadi P, Muchtadi TR, Andarwulan N. 2009. Hubungan sifat kimia dan rheologi tepung jagung putih dengan fermentasi spontan butiran jagung. *Forum Pasca Sarjana* 32: 33-43.
- Aini N, Hariyadi P, Muchtadi TR, Andarwulan N. 2010. Hubungan antara waktu fermentasi grits jagung putih dengan sifat gelatinasi tepung jagung putih yang dipengaruhi ukuran partikel. *J Teknol dan Industri Pangan* 21: 18-24.
- Aini N, Hariyadi P. 2010. Gelatinization properties of white maize starch from three varieties of corn subject to oxidized and acetylated-oxidized modification. *Int Food Res J* 17: 961-968.
- [AOAC] American Association of Cereal Chemists. 2000. Approved methods of the AACC, 10th ed. Methods 22-12, 46-12, 54-10, 54-21, 76-30A. St Paul MN: The Association.
- Belyea RL, Rausch KD, Tumbleson ME. 2004. Composition of corn and distillers dried grains with solubles from dry grind ethanol processing. *Bioresource Technol* 94: 293-298. DOI: [10.1016/j.biortech.2004.01.001](https://doi.org/10.1016/j.biortech.2004.01.001).
- Bellido GG, Hatcher DW. 2009. Stress relaxation behaviour of yellow alkaline noodles: Effect of deformation history. *J Food Eng* 93: 460-467. DOI: [10.1016/j.jfoodeng.2009.02.013](https://doi.org/10.1016/j.jfoodeng.2009.02.013).
- Chelule PK, Mbongwa HP, Carries S, Gqaleni N. 2010. Lactic acid fermentation improves the quality of amahewu, a traditional South African maize-based porridge. *Food Chem* 122: 656-661. DOI: [10.1016/j.foodchem.2010.03.026](https://doi.org/10.1016/j.foodchem.2010.03.026).
- Chung JH, Cho A, Lim ST. 2012. Effect of heat-moisture treatment for utilization of germinated brown rice in wheat noodle. *LWT Food Sci Tech* 47: 342-347. DOI: [10.1016/j.lwt.2012.01.029](https://doi.org/10.1016/j.lwt.2012.01.029).
- De Groote H, Kimenju SC. 2008. Comparing consumer preferences for colour and nutritional quality in maize: Application of a semi-double-bound logistic model on urban consumers in Kenya. *Food Policy* 33: 362-370. DOI: [10.1016/j.foodpol.2008.02.005](https://doi.org/10.1016/j.foodpol.2008.02.005).
- Eneje LO, Ogu EO, Aloh CU, Odibo, FJC, Agu RC, Palmer GH. 2004. Effects of stepping and germination time on malting performance of Nigerian white and yellow maize varieties. *Proc Biochem* 39: 1013-1016. DOI: [10.1016/S0032-9592\(03\)00202-4](https://doi.org/10.1016/S0032-9592(03)00202-4).
- Ingllett GE, Peterson SC, Carriere CJ, Maneepun S. 2005. Rheological, textural, and sensory properties of Asian noodles containing an oat cereal hydrocolloid. *Food Chem* 90: 1-8. DOI: [10.1016/j.foodchem.2003.08.023](https://doi.org/10.1016/j.foodchem.2003.08.023).
- Lu ZH, Li LT, Cao W, Li ZG, Tatsumi E. 2003. Influence of natural fermentation on physico-chemical characteristics of rice noodles. *Int J of Food Sci Tech* 38: 505-510. DOI: [10.1046/j.1365-2621.2003.00701.x](https://doi.org/10.1046/j.1365-2621.2003.00701.x).
- Maurel BV, Orliac O, Christen P. 2003. Sensors and measurements in solid state fermentation: A review. *Process Biochem* 38: 881-896. DOI: [10.1016/S0032-9592\(02\)00093-6](https://doi.org/10.1016/S0032-9592(02)00093-6).
- Moorthy SN. 2002. Physicochemical and functional properties of tropical tuber starches: A Review. *Starch* 54: 559-592. DOI: [10.1002/1521-379X\(200212\)54:12<559::AID-STAR2222559>3.0.CO;2-F](https://doi.org/10.1002/1521-379X(200212)54:12<559::AID-STAR2222559>3.0.CO;2-F).
- Muhandri, T, Subarna. 2009. Pengaruh kadar air, NaCl, dan jumlah passing terhadap karakteristik reologi mi jagung. *J Teknol dan Industri Pangan* 20: 71-77.
- Muhandri T, AB Ahza, R Syarif, Sutrisno. 2011. Optimasi proses ekstrusi mi jagung dengan metode permukaan respon. *J Teknol dan Industri Pangan* 22: 97-104.
- Rodríguez CEO, Carrillo MJ, Escobedo MR, Valenzuela COG, Moreno RC. 2004. Quality protein maize (*Zea mays* L.) tempeh flour through solid state fermentation process. *LWT Food Sci Tech* 37: 59-67. DOI: [10.1016/S0023-6438\(03\)00134-8](https://doi.org/10.1016/S0023-6438(03)00134-8).
- Rodríguez CEO, Montoya VNM, Bejarano API, Carrillo MJ, Escobedo MR, Pérez BLA, Tiznado GJA, Moreno RC. 2006. Nutritional properties of tempeh flour from quality protein maize (*Zea mays* L.). *LWT Food Sci Tech* 39: 1072-1079. DOI: [10.1016/j.lwt.2005.07.003](https://doi.org/10.1016/j.lwt.2005.07.003).
- Van-Kleef E, van Trijp HCM, Luning P. 2005. Consumer research in the early stages of new product development: a critical review of methods and techniques. *Food Quality and Preference* 16: 181-201. DOI: [10.1016/j.foodqual.2004.05.012](https://doi.org/10.1016/j.foodqual.2004.05.012).
- Young H, Borrel A, Holland D, Salama P. 2004. Public nutrition in complex emergencies. *The Lancet* 364: 1899-1909. DOI: [10.1016/S0140-6736\(04\)17447-3](https://doi.org/10.1016/S0140-6736(04)17447-3).
- Yuan, ML, Z.-H. Lu ZH, Cheng, YQ, Li LT. 2008. Effect of spontaneous fermentation on the physical properties of corn starch and rheological characteristics of corn starch noodle. *J of Food Eng* 85: 12-17. DOI: [10.1016/j.jfoodeng.2007.06.019](https://doi.org/10.1016/j.jfoodeng.2007.06.019).

Journal Profile

Jurnal Teknologi dan Industri Pangan

eISSN : 2087751X | pISSN : 19797788

Engineering

Institut Pertanian Bogor



S2

Sinta Score



Indexed by GARUDA

27

H-Index

22

H5-Index

3481

Citations

2487

5 Year Citations

Jurnal

TEKNOLOGI & INDUSTRI PANGAN

Volume 24 Nomor 2 Februari 2012





Pembinaan/Revisi

Prof. Dr. Ir. H. M. M. Teknologi Pangan
Indonesia (IPATI)

and Equivalents dengan

Departemen Ilmu dan Teknologi Pangan
Fakultas Teknologi Pertanian
Institut Pertanian Bogor

Member of



Perhimpunan Ahli Teknologi Pangan Indonesia (PATPI) yang diterbitkan bekerjasama dengan Departemen Ilmu dan Teknologi Pangan, Fakultas Teknologi Pertanian, Institut Pertanian

[Website](#) | [Editor URL](#)

Address:

Departemen Ilmu dan Teknologi Pangan, Fateta-IPB Gedung PAU Lt. 2, P.O. Box 220 Bogor 16680 Kampus IPB Darmaga
Bogor Telp/Fax (0251) 8626725
KAB. BOGOR

Email:

jurnaltip@gmail.com

Phone:

(0251) 8626725

Last Updated :

2022-01-31

2017 2018 2019 2020 2021 2022 2023 2024 2025

 Accreditations

Search..





Page 1 of 69 | Total Records : 685


Publications	Citation
<u>KAPASITAS ANTIOKSIDAN DAN INHIBITOR ALFA GLUKOSIDASE EKSTRAK UMBI BAWANG DAYAK (Antioxidant and Alpha-Glucosidase Inhibitory Properties of Bawang Dayak Bulb Extracts)</u> AE Febrinda, M Astawan, T Wresdiyati, ND Yuliana Jurnal Teknologi Dan Industri Pangan 24 (2), 161	93
<u>OPTIMASI PEMURNIAN ASAP CAIR DENGAN METODA REDISTILASI</u> P Darmadji Jurnal Teknologi dan Industri Pangan 13 (3), 267	87
<u>AKTIVITAS ANTIOKSIDAN EKSTRAK DAN FRAKSI EKSTRAK BIJI ATUNG (Parinarium glaberrimum Hassk.) (Antioxidant Activities of Parinarium glaberrimum Hassk Extracts and their Fractions)</u> D Sarastani, ST Soekarto, TR Muchtadi, D Fardiaz, A Apriyantono Jurnal Teknologi dan Industri Pangan 13 (2), 149-149	84
<u>Aktivitas antioksidan ekstrak dan fraksi ekstrak biji atung (Parinarium glaberrimum Hassk.)</u> D Sarastani, ST Soekarto, TR Muchtadi, D Fardiaz, A Apriyantono jurnal teknologi dan industri pangan, 149-156	84

Publications	Citation
<p><u>Pemanfaatan ubi jalar ungu (<i>Ipomoea batatas</i> L. Poir) sebagai pengganti sebagian tepung terigu dan sumber antioksidan pada roti tawar</u></p> <p>Handoko, L Hendarto, TM Siregar Jurnal Teknologi Dan Industri Pangan 21 (1), 25</p>	85
<p><u>EKSTRAKSI DAN STABILITAS ZAT PEWARNA ALAM DARI KATUL BERAS KETAN HITAM (<i>Oryza sativa glutinosa</i>). EXTRACTION AND STABILITY OF NATURAL COLORANT FROM RED GLUTINOUS RICE BRAN (<i>Oryza ...</i></u></p> <p>T Hanum Jurnal Teknologi dan Industri Pangan 11 (1), 17-17</p>	59
<p><u>PENGEMBANGAN BERAS ANALOG DENGAN MEMANFAATKAN JAGUNG PUTIH (Development of White Corn-Based Rice Analogues)</u></p> <p>S Noviasari, F Kusnandar, S Budijanto Jurnal Teknologi Dan Industri Pangan 24 (2), 194</p>	59
<p><u>PENDUGAAN UMUR SIMPAN PRODUK BISKUIT DENGAN METODE AKSELERASI BERDASARKAN PENDEKATAN KADAR AIR KRITIS (Accelerated Shelf-life Testing of Biscuits Using a Critical Moisture ...</u></p> <p>F Kusnandar, DR Adawiyah, M Fitria Jurnal Teknologi Dan Industri Pangan 21 (2), 117</p>	56
<p><u>EKSTRAKSI DAN STABILITAS ZAT PEWARNA ALAM DARI KATUL BERAS KETAN HITAM (<i>Oryza sativa glutinosa</i>). EXTRACTION AND STABILITY OF NATURAL COLORANT FROM RED GLUTINOUS RICE BRAN (<i>Oryza ...</i></u></p> <p>T Hanum Jurnal Teknologi dan Industri Pangan 11 (1), 17-17, 2000</p>	56
<p><u>Karakteristik es krim hasil modifikasi dengan formulasi bubuk timun suri (<i>Cucumis melo</i> L.) dan sari kedelai</u></p> <p>S Oksilia, E Lidiasari Jurnal Teknologi dan Industri Pangan 23 (1), 17-22</p>	56

Page 1 of 69 | Total Records : 685



Citation Statistics



KORESPONDENSI

HOME MAIL NEWS FINANCE SPORTS ENTERTAINMENT LIFE SHOPPING YAHOO PLUS MORE Upgrade Now

Find messages, documents, photos or people



Home

Compose

← Back ↶ ↷ → Archive ↗ Move Delete Spam



nuraini... 119

nuraini.1973

Inbox 119

Unread

Starred

Drafts 7

Sent

Archive

Spam

Trash

^ Less

Views Hide

Photos

Documents

Subscriptions

Deals

Groceries

Receipts

Travel

Folders Hide

+ New Folder

Notes

Nur aini



nuraini munawar

To:
jurnaltip@live.co

Tue, Nov 13, 2012 at 10:59 AM

Kepada Redaksi Jurnal Teknologi dan Industri Pangan

Terima kasih atas kesempatan yang telah diberikan
Berikut saya kirimkan naskah untuk Jurnal TIP yang saya
perbaiki sesuai format

terima kasih

Nur Aini

From: Jurnal TIP <jurnaltip@yahoo.com>
To: nuraini munawar <nuraini_munawar@yahoo.com>
Sent: Thursday, November 8, 2012 3:42 PM
Subject: Re: artikel untuk jurnal TIP

Yth. Ibu Nuraini Munawar

Baik, naskah sudah kami terima.
Setelah kami lihat dan cek kembali naskah belum sesuai
dengan Pedoman JTIP yang telah di berikan.

Berikut kami lampirkan kembali naskah untuk dapat
diperbaiki lebih lanjut.
Perbaikan naskah dapat dikirimkan pada tanggal **13
November 12** untuk kami proses lebih lanjut.

Atas perhatian dan kerjasamanya kami ucapkan terima
kasih.

salam,

=====
[Redaksi Jurnal Teknologi & Industri Pangan](#)
[Departemen Ilmu & Teknologi Pangan](#)
[Fakultas Teknologi Pertanian](#)
[Institut Pertanian Bogor](#)
[Kampus IPB Darmaga,](#)
[PO Box 220, Bogor 16002](#)
[Telp/Faks. 62251 3426715](#)

From: nuraini munawar <nuraini_munawar@yahoo.com>
To: "jurnaltip@yahoo.com" <jurnaltip@yahoo.com>

Find messages, documents, photos or people



Home

Compose

← Back ↩️ ⏪ ⏩ 📁 Archive 📁 Move 🗑️ Delete 🛡️ Spam



nuraini_... 119

nuraini.1973 ⚠️

...

Inbox 119

Unread

Starred

Drafts 7

Sent

Archive

Spam

Trash

^ Less

Views Hide

Photos

Documents

Subscriptions

Deals

Groceries

Receipts

Travel

Folders Hide

+ New Folder

Notes

Nur aini

jadi terlambat

Terima kasih atas kesempatan yang diberikan

Nur Aini

----- Forwarded Message -----

From: nuraini munawar <nuraini_munawar@yahoo.com>**To:** Jurnal TIP <jurnaltip@live.com>**Sent:** Tuesday, November 13, 2012 10:59 AM**Subject:** Re: artikel untuk jurnal TIP

Kepada Redaksi Jurnal Teknologi dan Industri Pangan

Terima kasih atas kesempatan yang telah diberikan
Berikut saya kirimkan naskah untuk Jurnal TIP yang saya
perbaiki sesuai format

terima kasih

Nur Aini

From: Jurnal TIP <jurnaltip@yahoo.com>**To:** nuraini munawar <nuraini_munawar@yahoo.com>**Sent:** Thursday, November 8, 2012 3:42 PM**Subject:** Re: artikel untuk jurnal TIP

Yth. Ibu Nuraini Munawar

Baik, naskah sudah kami terima.
Setelah kami lihat dan cek kembali naskah belum sesuai
dengan Pedoman JTIP yang telah di berikan.

Berikut kami lampirkan kembali naskah untuk dapat
diperbaiki lebih lanjut.


Perbaikan naskah dapat dikirimkan pada tanggal **13
Nopember 12** untuk kami proses lebih lanjut.

Atas perhatian dan kerjasamanya kami ucapkan terima
kasih.

salam,

Global Trade Starts Here

Alibaba.com

fat tire popular 200
electric scooter

IDR 2.6M

[Visit Site](#)

Find messages, documents, photos or people



Home

Compose

← Back ↶ ↷ → Archive ↗ Move 🗑 Delete 🛡 Spam



nuraini... 119
nuraini.1973 ⚠

Inbox 119

Unread

Starred

Drafts 7

Sent

Archive

Spam

Trash

^ Less

Views Hide

Photos

Documents

Subscriptions

Deals

Groceries

Receipts

Travel

Folders Hide

+ New Folder

Notes

Nur aini

• Permintaan abstrak Bahasa Indonesia naskah JTIP (segera)

Yahoo/Inbox ★



• jurnaltip@yahoo



Thu, Dec 20, 2012 at 5:52 PM ★

To: nuraini@unsoed.a

Cc: nuraini.1973@gm

, nuraini_munawar@

Yth. Ibu Nur Aini

Berikut kami lampirkan naskah yang telah di lakukan editing.

Mohon untuk dapat membuat Abstrak dalam Bahasa Indonesia dari Abstrak yang sudah ada.

Penambahan abstrak pada naskah mohon dapat dikirimkan pada tanggal **21 Des 12** untuk kami proses lebih lanjut.

Atas perhatian dan kerjasamanya kami ucapkan terima kasih.

salam,

=====
[Redaksi Jurnal Teknologi & Industri Pangan](#)
[Departemen Ilmu & Teknologi Pangan](#)
[Fakultas Teknologi Pertanian](#)
[Institut Pertanian Bogor](#)
[Kampus IPB Darmaga,](#)
[PO Box 220, Bogor 16002](#)
[Telp/Faks. 62251 3426715](#)



jurnaltip@yahoo.com 🔍

[jurnaltip@yahoo.com](#)[Edit contact](#)

Find messages, documents, photos or people

Home

Compose

Back Reply Reply All Forward Archive Move Delete Spam

Calendar Lists Help



nuraini... 119
nuraini.1973

Inbox 119
Unread
Starred
Drafts 7
Sent
Archive
Spam
Trash
Less

Views Hide

Photos
Documents
Subscriptions
Deals
Groceries
Receipts
Travel

Folders Hide

+ New Folder
Notes
Nur aini

Persetujuan Penerbitan & Proofread-JTIP(Segera) 3 Yahoo/Inbox

jurnaltip@yahoo.com 0 Fri, Jan 11, 2013 at 5:29 PM

nuraini munawar <jurnaltip@yahoo.co> To: jurnaltip@yahoo.co

Yth
Redaksi Jurnal TIP

Bersama ini kami sampaikan sedikit perbaikan pada ucapan terima kasih dan afiliasi penulis
Terima kasih atas kesempatan yang diberikan

Nur Aini

From: "jurnaltip@yahoo.com" <jurnaltip@yahoo.com>
To: "nuraini_munawar@yahoo.com" <nuraini_munawar@yahoo.com>
Cc: "nuraini@unsoed.ac.id" <nuraini@unsoed.ac.id>; "nuraini.1973@gmail.com" <nuraini.1973@gmail.com>
Sent: Friday, January 11, 2013 2:29 AM
Subject: Persetujuan Penerbitan & Proofread-JTIP(Segera)

Nomor : 010/ SK. Naskah
Disetujui/JTIP/01/2013
Lamp. : 1 Berkas
Hal : Persetujuan Penerbitan

Kepada Yth. Diterima : 07
Ibu Nur Aini November 2012
Di Tempat No. Naskah :
56/JTIP/11/12

Dengan hormat,

Dengan ini kami memberitahukan bahwa naskah:

Find messages, documents, photos or people



Home

Compose

Archive Move Delete Spam



nuraini... 119

nuraini.1973

...

Inbox 119

Unread

Starred

Drafts 7

Sent

Archive

Spam

Trash

^ Less

Views Hide

Photos

Documents

Subscriptions

Deals

Groceries

Receipts

Travel

Folders Hide

+ New Folder

Notes

Nur aini



Journal TIP 'edoman JTIP yang ' Thu, Nov 8, 2012 at 3:42 PM



nuraini munawar Tue, Nov 13, 2012 at 10:59 AM

To:
jurnaltip@live.co

Kepada Redaksi Jurnal Teknologi dan Industri Pangan

Terima kasih atas kesempatan yang telah diberikan
Berikut saya kirimkan naskah untuk Jurnal TIP yang saya
perbaiki sesuai format

terima kasih

Nur Aini

From: Jurnal TIP <jurnaltip@yahoo.com>
To: nuraini munawar <nuraini_munawar@yahoo.com>
Sent: Thursday, November 8, 2012 3:42 PM
Subject: Re: artikel untuk jurnal TIP

Yth. Ibu Nuraini Munawar

Baik, naskah sudah kami terima.
Setelah kami lihat dan cek kembali naskah belum sesuai
dengan Pedoman JTIP yang telah di berikan.

Berikut kami lampirkan kembali naskah untuk dapat
diperbaiki lebih lanjut.
Perbaikan naskah dapat dikirimkan pada tanggal **13
Nopember 12** untuk kami proses lebih lanjut.

Atas perhatian dan kerjasamanya kami ucapkan terima
kasih.

salam,

=====
Redaksi Jurnal Teknologi & Industri Pangan
Departemen Ilmu & Teknologi Pangan
Fakultas Teknologi Pertanian
Institut Pertanian Bogor
Kampus IPB Darmaga,
PO Box 220, Bogor 16002
Telp/Faks. 62251 3426715

Find messages, documents, photos or people

Home

Compose

Back Reply Reply All Forward Archive Move Delete Spam

Calendar Lists Help



nuraini_muna... 119
nuraini.1973

Inbox 158

Unread

Starred

Drafts 10

Sent

All Mail

Spam

Trash

Less

Views

Photos

Documents

Subscriptions

Deals

Groceries

Receipts

Travel

Folders

+ New Folder

[Gmail]/Penting 126

Kerjaan

Notes

nur.aini@unso... 41

Perjalanan

Pribadi

Resi

4

jurnaltip@yahoo.com Thu, Dec 20, 2012 at 5:52 PM

nur aini Kepada Yth Redaksi J Thu, Dec 20, 2012 at 6:50 PM

nur aini Iesia naskah JT Thu, Dec 20, 2012 at 6:54 PM

jurnaltip@yahoo.com To: nur aini Fri, Dec 21, 2012 at 8:58 AM

Yth. Ibu Nur Aini

Baik, terima kasih.
Abstrak sudah kami terima. Dapat kami informasikan sebelumnya mohon untuk alamat e-mail **jurnaltip@live.com** untuk di hapus. Kami hanya menggunakan alamat e-mail : jurnaltip@yahoo.com

Demikian yang dapat kami sampaikan. Mohon menunggu hasil proses naskah selanjutnya.

Atas perhatian dan kerjasamanya kami ucapkan terima kasih.

salam,
=====
Redaksi Jurnal Teknologi & Industri Pangan
Departemen Ilmu & Teknologi Pangan
Fakultas Teknologi Pertanian
Institut Pertanian Bogor
Kampus IPB Darmaga,
PO Box 220, Bogor 16002
Telp/Faks. 62251 3426715

Show original message



Reply, Reply All or Forward

Send



Find messages, documents, photos or people



Home

Compose

← Back ↶ ↷ → Archive ↗ Move Delete Spam



nuraini_muna... 119
nuraini.1973 ⚠

Inbox 158
Unread
Starred
Drafts 10
Sent
All Mail
Spam
Trash
^ Less

Views Hide
Photos
Documents
Subscriptions
Deals
Groceries
Receipts
Travel

Folders Hide
+ New Folder
[Gmail]/Penting 126
Kerjaan
Notes
nur.aini@unso... 41
Perjalanan
Pribadi
Resi

perbaiki naskah 56-JTIP-11-12 nuraini.1973/Inbox ★
3

nur aini iatif ini 56-JTIP-11-12 revisi. Fri, Dec 7, 2012 at 6:13 AM ★

Jurnal TIP :kah 56-JTIP-11-12 Kepad Fri, Dec 7, 2012 at 9:29 AM ★

Jurnal TIP <jurnaltip@yahoo.c 📧 Fri, Dec 7, 2012 at 9:40 AM ★
To: nur aini

Yth. Ibu Nur Aini

Baik, naskah hasil perbaikan sudah akmi terima.
Akan kami tindaklanjuti kembali.
Mohon kesabarannya menunggu hasil proses naskah selanjutnya.

Atas perhatian dan kerjasamanya kami ucapkan terima kasih.

salam,


=====
[Redaksi Jurnal Teknologi & Industri Pangan](#)
[Departemen Ilmu & Teknologi Pangan](#)
[Fakultas Teknologi Pertanian](#)
[Institut Pertanian Bogor](#)
[Kampus IPB Darmaga,](#)
[PO Box 220, Bogor 16002](#)
[Telp/Faks. 62251 3426715](#)

> Show original message



[Reply](#), [Reply All](#) or [Forward](#)

Send



jurnaltip@yahoo.com 🔍

[jurnaltip@yahoo.com](#)
[Edit contact](#)

Find messages, documents, photos or people



Home

Compose

← Back ↶ ↷ → Archive ↗ Move Delete Spam



nuraini_muna... 119
nuraini.1973

Inbox 158
Unread
Starred
Drafts 10
Sent
All Mail
Spam
Trash
^ Less

Views Hide
 Photos
 Documents
 Subscriptions
 Deals
 Groceries
 Receipts
 Travel

Folders Hide
+ New Folder
[Gmail]/Penting 126
Kerjaan
Notes
nur.aini@unso... 41
Perjalanan
Pribadi
Resi

• Permintaan abstrak Bahasa Indonesia naskah JTIP (segera) nuraini.1973/Inbox ★

4



• **jurnaltip@yahoo.com** Thu, Dec 20, 2012 at 5:52 PM ★



• **nur aini** <nuraini.1973@...> Thu, Dec 20, 2012 at 6:50 PM ★
To: jurnaltip@yahoo.com

Kepada Yth Redaksi Jurnal TIP

Bersama ini kami kirimkan perbaikan naskah yang sudah ditambah abstrak bahasa Indonesia
Mohon maaf, saya sudah beberapa kali mengemailnya, yg pertama salah email lagi dan yg kemarin sepertinya belm terkirim

Atas perhatian dan kesempatan yang diberikan kami ucapkan terima kasih

Nur Aini

----- Forwarded Message -----
From: nuraini munawar <nuraini_munawar@yahoo.com>
To: Jurnal TIP <jurnaltip@yahoo.com>
Sent: Tuesday, December 18, 2012 1:17 PM
Subject: Fw: penambahan abstrak

----- Forwarded Message -----
From: nuraini munawar <nuraini_munawar@yahoo.com>
To: Jurnal TIP <jurnaltip@yahoo.com>
Sent: Tuesday, December 18, 2012 1:14 PM
Subject: Fw: penambahan abstrak

----- Forwarded Message -----
From: nuraini munawar <nuraini_munawar@yahoo.com>
To: Jurnal TIP <jurnaltip@live.com>
Sent: Monday, December 17, 2012 1:49 PM
Subject: penambahan abstrak

Kepada Yth Redaksi Jurnal TIP

Bersama ini kami kirimkan perbaikan naskah yang sudah ditambah abstrak bahasa Indonesia
Atas perhatian dan kesempatan yang diberikan kami ucapkan terima kasih

Nur Aini

Find messages, documents, photos or people



Home

Compose

nuraini_muna... 119

nuraini.1973

...

Inbox 158

Unread

Starred

Drafts 10

Sent

All Mail

Spam

Trash

^ Less

Views Hide

Photos

Documents

Subscriptions

Deals

Groceries

Receipts

Travel

Folders Hide

+ New Folder

[Gmail]/Penting 126

Kerjaan

Notes

nur.aini@unso... 41

Perjalanan

Pribadi

Resi

← Back Archive Move Delete Spam



To: nuraini_munawar@y
 Cc: nuraini@unsoed.ac.
 ,
 nuraini.1973@gmai

Nomor : 010/ SK. Naskah
 Disetujui/JTIP/01/2013
 Lamp. : 1 Berkas
 Hal : Persetujuan Penerbitan

Kepada Yth.	Diterima	: 07
Ibu Nur Aini		
Di Tempat	November 2012	
	No. Naskah	:
	56/JTIP/11/12	

Dengan hormat,

Dengan ini kami memberitahukan bahwa naskah:

No : 56/JTIP/11/12

Judul : **CHARACTERISTICS OF WHITE
 CORN NOODLE SUBSTITUED BY TEMPEH
 FLOUR**

Penulis : **Nur Aini¹⁾, V. Prihananto¹⁾, and S.
 Joni Munarso²⁾**

¹⁾ Department of Agricultural
 Technology, Jenderal Soedirman University, Purwokerto

²⁾ Research and Development Center of Plantation,
 Bogor

telah disetujui pada tanggal : 07 Desember 2012

Naskah tersebut telah disetujui untuk diterbitkan dalam Jurnal
 Teknologi Industri Pangan Volume XXIII No. 2 Tahun 2012.

Bersama ini juga kami sampaikan contoh cetak untuk
 naskah tersebut. Kami mohon dapat menyampaikan koreksian
 dari contoh cetak halaman dalam waktu **lima hari kerja**
 setelah menerima surat ini. Jika masih terdapat perbaikan atau
 koreksian mohon untuk mencoret dan memperbaikinya
 langsung pada naskah contoh cetak yang kami lampirkan.

Atas perhatian dan kerjasamanya kami ucapkan terimakasih



jurnaltip@yahoo.com

jurnaltip@yahoo.com
 Edit contact