MENU

SSDM2023

SSDM2023

ⓘ
✕

AIP Publishing **Conference Proceedings**

f ⟋

HOME   BROWSE   INFO   FOR AUTHORS   🔔 SIGN UP FOR ALERTS

FOR ORGANIZERS

AIP Conference Proceedings has been a trusted publishing partner for more than 40 years, delivering fast, affordable, and versatile publishing for maximum exposure of your meeting's key research. Our conference proceedings program reports the findings presented at scientific meetings from large international conferences to small specialist workshops. Subject areas span the physical sciences, including physics, math, chemistry, materials science, and engineering.

Why publish with us:

- Indexed in leading databases — Web of Science, Scopus, Inspec, Chemical Abstracts Service (CAS), and Astrophysics Data System (ADS)
- Author-friendly license agreement — you retain copyright of your own work
- Global reach — nearly 4,000 institutions in 190+ countries
- Customized publishing options including print-on-demand — best value and flexibility

Most Read

Most Cited

FEB 02 2018

A solid waste management survey in Davao del Sur (school and household waste management survey)

52038 VIEWS

Find out why AIP Conference Proceedings is the right choice for your next event.

Request a QUOTE today.

Most Recent

MAR 31 2023

Preface: 1st International & 4th Local Conference for Pure Science (ICPS-2021)

MAR 31 2023

Wastewater: A study of its sources, methods of treatment, and impact on environmental pollution

MAR 31 2023

Molecular detection of canine distemper virus using one-step RT-QPCR in domestic dog in Iraq

MAR 31 2023

Comparative study of ICP-OES, μ-XRF and P-XRF of trace elements analysis in various Sasanian and Parthian artworks found in the Garmian region of Iraq

MAR 31 2023

Feature and utility in diagnosis of papillary thyroid carcinoma by immunohistochemistry CD56

OCT 03 2017

A review of the motivation theories in learning

47 CITATIONS

95784 VIEWS

MAR 10 2017

Integrated management system: The integration of ISO 9001, ISO 14001, OHSAS 18001 and ISO 31000

10 CITATIONS

32107 VIEWS

Active Topics

Students

Learning and learning models

Machine learning

Computer simulation

Thin films

MAR 31 2023

Indoor radon concentration measurements for University of Karabük building using Cr-39 detector

MAR 31 2023

Computational approaches for solving fuzzy linear fractional models

MAR 31 2023

Fetal age estimation using ultrasonography in dairy cattle

Browse All Topics

SIGN UP NOW

About AIP Publishing

AIP Publishing is a wholly owned not-for-profit subsidiary of the American Institute of Physics (AIP). Our portfolio comprises highly regarded, peer-reviewed journals, including a growing portfolio of Open Access titles, that cover all areas of the physical sciences.

The research published in these titles paves the way for new fields of study, gives rise to new techniques, and provides inspiration to contemporary researchers.

### Applied Physics Letters

The Journal of Chemical Physics

Physics Today

Review of Scientific Instruments

Physics of Fluids

Physics of Plasmas

Journal of Applied Physics

AIP Advances

VISIT PUBLISHER'S WEBSITE

VISIT ALL PUBLICATIONS

Latest Physics Jobs

VIEW ALL JOBS FROM PHYSICS TODAY

Visiting Assistant Professor of Earth and Climate Sciences and Physics | Bates College

Fri, 31 Mar 2023 14:23:59 -0400 Lewiston, Maine, The Department of Earth and Climate Sciences (EACS) and the Department of Physics and Astronomy (PA) at Bates College invite applications for a joint full-time (five courses) Visiting Assistant Professor for the academic year 2023-24. Candidates with a demonstrated commitment to the success of historically underrepresented, marginalized and first-generation students are encouraged to apply. The successful candidate must demonstrate potential for teaching excellence in a vigorous undergraduate program, have a d knowledge of/experience with

Professorship in Experimental Quantum Nanophysics and/or Quantum Materials (open rank) | University of Basel

Thu, 30 Mar 2023 08:37:35 -0400 Basel, Switzerland, The Department of Physics at the University of Basel, Switzerland, invites applications for a professorship (open rank) in the areas of experimental quantum nanophysics and/or quantum materials. The University of Basel is the home institution of the Swiss Nanoscience Institute (SNI) and of the Swiss National Centre of Competence in Research (NCCR) "SPIN: Spin Qubits in Silicon". Candidates at the tenure-track assistant professorship level are particularly encouraged to apply. In exceptional cases, candidates at other levels (associate or full

Physical Scientist GS-1301-15 | Department of Energy Office of Science

Thu, 30 Mar 2023 13:55:49 -0400 Germantown, Maryland, Overview   The two job opportunities include one program manager in the Atomic, Molecular, and Optical Sciences (AMOS) program, and another program manager to manage the Quantum Information Sciences (QIS) portfolio in CSGB. BES CSGB has an opening for a program manager to manage the portfolio of fundamental research projects in the Atomic, Molecular, and Optical Sciences (AMOS) Program. The AMOS Program currently focuses on fundamental, hypothesis-driven research in ultrafast chemical sciences. The program portfolio supports

Medical Imaging Physics Fellowship | National Cancer Institute/Molecular Imaging Branch

Thu, 30 Mar 2023 09:41:35 -0400 Bethesda, Maryland, The Molecular Imaging Branch (MIB) at the Center for Cancer Research (CCR) of the National Cancer Institute (NCI) is looking for a highly motivated individual to join the MIB in a multi-organizational effort to develop paired radiation detectors that can accurately locate annihilation events in space using only differences in photon arrival times (time-of-flight). Applicants should have received a Ph.D. degree within the last 5 years in physics or engineering (electrical or biomedical) and are expected to be familiar with the full range of contemporary radiation detection methods and

inclusive pedagogies, and welcome the opportunity to mentor serious students and advise senior theses. Field-based, computational, and analytical laboratory projects, and collaboration with departmental ...

professor) will also be considered.   Your position The new professor will be expected to conduct an excellent research ...

basic experimental and theoretical research aimed at understanding the structural and dynamical properties of atomic and molecular systems. The research targets ...

computer codes simulating these methods. This individual will be ...

Resources

AUTHOR

LIBRARIAN

ADVERTISER

General Information

ABOUT

CONTACT

HELP

PRIVACY POLICY

TERMS OF USE

FOLLOW AIP PUBLISHING:

Scitation

AIP Publishing   **Conference Proceedings**

f   ⋒

HOME   BROWSE   INFO   FOR AUTHORS   🔔 **SIGN UP FOR ALERTS**

FOR ORGANIZERS

Browse Volumes

2553   ⌄

Submit

Browse Volumes

Table of Contents

⟨ PREV

NEXT ⟩

PROCEEDINGS OF THE 16TH JOINT CONFERENCE ON CHEMISTRY (JCC 2021): Visualizing the latest up-front of chemistry research and education for better future

Conference date: 9 September 2021
Location: Purwokerto, Indonesia
ISBN: 978-0-7354-4281-8
Editors: Uyi Sulaeman, Hassan Y. Aboul-Enein, Ahman Zuhairi Abdullah, Zaheer Ul-Haq Qasmi, Warakorn Limbut, Jas Raj Subba and Mardiyah Kurniasih
Volume number: 2553
Published: Nov 22, 2022

DISPLAY :   20   50   100   all

PRELIMINARY

🔒 No Access . November 2022

Preface: The 16th Joint Conference on Chemistry 2021 (16th JCC)

AIP Conference Proceedings **2553**, 010001 (2022);
https://doi.org/10.1063/12.0012151

🔒 No Access . November 2022

Committee: The 16th Joint Conference on Chemistry 2021 (16th JCC)

AIP Conference Proceedings **2553**, 010002 (2022);
https://doi.org/10.1063/12.0015105

ARTICLES

🔒 No Access . November 2022

🔒 No Access . November 2022

Effect of cetyltrimethylammonium bromide on hydrophobicity of silica thin layer and xerogel derived from sodium silicate and methyltrimethoxysilane

Riza Fithrotul Maghfiroh, Choiril Azmiyawati and Adi Darmawan

AIP Conference Proceedings **2553**, 020045 (2022); https://doi.org/10.1063/5.0104042

SHOW ABSTRACT

🔒 No Access . November 2022

Molecular docking approach for prediction of enantioseparation of miconazole using cyclodextrin derivatives as chiral selector

Dadan Hermawan, Ainaya Halwa Lathufa, Cacu, Ponco Iswanto, Uyi Sulaeman and Hassan Y. Aboul-Enein

AIP Conference Proceedings **2553**, 020046 (2022); https://doi.org/10.1063/5.0107144

SHOW ABSTRACT

🔒 No Access . November 2022

Synthesis of citric acid-crosslinked chitosan membrane with zeolite filler and its application as Cu(II) ion separation membrane

Khabibi, Nor Basid Adiwibawa Prasetya, Retno Ariadi Lusiana, Linda Suyati, Rahmad Nuryanto, Lailatul Rohmah and Ika Aprilia Khoirunnisa

SHOW ABSTRACT

No Access . November 2022

Activating cashew nut shell carbon (CNSC) produced by pyrolysis process using microwave and its application on the methylene blue adsorption

Mashuni, La Ode Ahmad, M. Jahiding, Fitri Handayani Hamid, Irmawati and Andi Muhammad Naufal Khaeri

SHOW ABSTRACT

1   2   >

Molecular docking approach for prediction of enantioseparation of miconazole using cyclodextrin derivatives as chiral selector

Dadan Hermawan, Ainaya Halwa Lathufa, Cacu, et al.

View Online   Export Citation

Molecular Docking Approach for Prediction of Enantioseparation of Miconazole Using Cyclodextrin Derivatives as Chiral Selector

Dadan Hermawan[1, a], Ainaya Halwa Lathufa[1], Cacu[1], Ponco Iswanto[1], Uyi Sulaeman[1], and Hassan Y. Aboul-Enein[2]

[1]*Department of Chemistry, Faculty of Mathematics and Natural Sciences, Jenderal Soedirman University, Jl. Dr. Soeparno, Karangwangkal, Purwokerto 53122, Indonesia*

[2]*Department of Pharmaceutical and Medicinal Chemistry, Pharmaceutical and Drug Industries Research Division, National Research Centre, Dokki, Giza 12622, Egypt*

[a]*Corresponding author: dadan.hermawan@unsoed.ac.id*

Abstract. Enantioseparation of miconazole has been successfully predicted based on the molecular docking approach using cyclodextrin derivatives such as sulfated-β-cyclodextrin (S-β-CD), hydroxypropyl-β-cyclodextrin (HP-β-CD) and hydroxypropyl-γ-cyclodextrin (HP-γ-CD) as chiral selectors. Molecular docking was performed using AutoDock Vina software and the root mean square deviation (RMSD) was calculated using PyMol software. Molecular docking shows that R-Miconazole forms more stable interaction with all cyclodextrin derivatives than S-Miconazole forms, suggesting that S-miconazole will be eluted earlier than R-miconazole. In addition, the stability level of cyclodextrin derivatives as chiral selector was HP-β-CD > S-β-CD > HP-γ-CD.

INTRODUCTION

Miconazole is a class of imidazole compounds that are used as an antifungal agent. This drug has been used to treat fungal skin infections such as ringworm, water fleas, tinea versicolor, and candidiasis. This compound has one chiral center and two different enantiomers. The enantiomers have identical physicochemical characteristics but can show differences in pharmacological and toxicological [1]. Based on several reports, there is only one enantiomer that gives good effects while the other one may cause negative effects [2], therefore, the enantiomeric separation of chiral compounds is important to do. One of the most possible methods to separate enantiomers of chiral compounds is High Performance Liquid Chromatography (HPLC).

Chiral separation can be achieved using chiral selectors which clearly distinguish the two enantiomers. Chiral selector changes one of the two enantiomers at different rates into a new compound (kinetics enantioselective) or established a labile molecule at different stability (thermodynamics enantioselective). One of the most widely used chiral selectors is cyclodextrin due to its excellent chiral recognition abilities [3]. However, the HPLC chiral separation can only separate a limited chiral range. Therefore, a preliminary schematic was needed to analyze the three-dimensional structure of compounds to predict the characteristics and functions of these compounds and analyze the actual interactions at the molecular level [4].

Molecular docking is a computational method that has been used to predict the molecular interactions of ligands and receptors. The molecular docking prediction increases accuracy and precision in chiral recognition and also reduces research time since it can be predicted quickly [5]. The molecular docking method can save time, effort, and the use of solvents or chemicals that pollute the environment [6]. This method was performed by placing the ligands systematically on the active site of receptors. The molecular docking method aims to achieve the optimal complex conformation and to predict the interactions of drugs/ligands and receptors, therefore, the best geometry of ligands

and receptors complex was obtained. In addition, a more effective ligand can be predicted by calculated the energy interactions of the different ligands [7]. Based on this report, the aims of this study are to analyzing the structure of the miconazole complex, calculated the binding energy, and determined the most suitable chiral selector to predicting separation of the enantiomers of miconazole. Chiral selectors commonly used are cyclodextrin and its derivatives. Cyclodextrin is an oligosaccharide composed of six (α-CD), seven (β-CD), and eight (γ-CD) glucose units through α-1,4 glycoside bonds [8]. Cyclodextrin can form an inclusion complex with drugs by inserting drug molecules into the central cavity of cyclodextrin. The inclusion complex formed can improve the solubility, dissolution, stability, and bioavailability of the guest molecule [9]. Cyclodextrin shows good enantioselectivity for a wide range of analytes, transparent to UV light and has good water solubility. In addition, cyclodextrins are available in various ranges, generally, give fast kinetics for the complex formation and breakdown enantiomers, and are relatively cheap [10]. In this study, miconazole acts as ligands, and cyclodextrin acts as receptors. Cyclodextrins used for this study are sulfated-β-cyclodextrin (S-β-CD), hydroxypropyl-β-cyclodextrin (HP-β-CD) and hydroxypropyl-γ-cyclodextrin (HP-γ-CD).

EXPERIMENTAL DETAILS

Materials

The crystal structure of S-β-CD, HP-β-CD, and HP-γ-CD were obtained from PubChem (CID: 12049144, 14049689, and 119174) in sdf format. The R-Miconazole and S-Miconazole crystal structures were obtained from PubChem (CID: 1150344 and 969503).

Instrumentation

A computer with an Operating System (OS) *Windows 10 Home 64-bit* with *processor Intel® Core™ i3-6006U CPU @ 2.0GHz (4 CPUs)*, 1.99GHz and external hard disk 500 GB, RAM 4 GB was utilized in this experiment. The CDs structures were converted into pdb file format using Avogadro software. The obtained structures of miconazole were optimized using the PM3 method implemented in HyperChem. The optimized structures of miconazole were used for the molecular docking using Autodock Vina software. The docking results were calculated RMSD using PyMol software and visualized using Discovery Studio software.

Geometry Optimization

The structure of miconazole was geometrically optimized using HyperChem software. The density function used is semiempirical with PM3 as the calculation method. The structure of miconazole and its enantiomers are shown in Fig. 1.

**FIGURE 1.** Structure of R-Miconazole and S-Miconazole

Molecular Docking

Molecular docking was carried out using automated docking program, AutoDock Vina. Water molecules on the receptor structures are removed to avoid interference during the molecular docking process. Polar hydrogen atoms are added to the structure as only polar atoms will interact with ligands while non-polar hydrogen atoms will be hidden

to increase molecular docking calculation speed. The grid box was determined according to the coordinates of the active site of receptors. This study was conducted using a blind docking process as the grid box parameters are not yet known. The large grid box was arranged so that ligands can rotate freely to find the most stable site on the receptors.

Exhaustiveness is a parameter of Autodock Vina that controls how comprehensive its prediction is. The increase of the exhaustiveness value will be slow down the docking process. However, the higher the exhaustiveness, the higher the probability of a good result of docking. The default value of Autodock Vina exhaustiveness is 8 [7,11]. In this study, exhaustiveness of 264 was used to obtain a more consistent docking result. The S-β-CD structure was docked with coordinate value of X = -3.160, Y = -2.483, Z = -1.514, Grid Box size of 52 x 52 x 52 and Grid Spacing 0.375. The HP-β-CD structure was docked with coordinate value of X = -0.421, Y = -5.367, Z = -2.059, Grid Box size of 60 x 60 x 60 and Grid Spacing 0.375. The HP-γ-CD structure was docked with coordinate value of X = 7.235, Y = 14.629, Z = 2.358, Grid Box size of 52 x 64 x 60 and Grid Spacing 0.375. The docking results should have Root Mean Square Deviation (RMSD) of less than 3 Å [12].

RESULTS AND DISCUSSION

Geometry Optimization

The first step is ligands and receptors preparation using Avogadro software to changes the file format into pdb. Miconazole as a ligand was optimized using HyperChem software by the semiempirical PM3 method, therefore, the most stable structure that has the lowest binding energy was obtained. Miconazole enantiomers structures before and after optimization were shown in Fig. 2 and Fig. 3.

**FIGURE 2.** The structure of R-Miconazole before and after optimized

FIGURE 3. The structure of S-Miconazole before and after optimized

Molecular Docking

The running process was performed using Autodock Vina software. The RMSD values of inclusion complexes obtained are shown in Table 1. All of the RMSD values obtained are below 3 Å, this indicates that the ligand structure before and after docking is almost accurate.

The results of molecular docking are shown the stability of the inclusion complex based on its binding energy (ΔG). Fig. 4 shows the structure of S-β-CD/R-Miconazole and S-β-CD/S-Miconazole complexes. The complex which the miconazole is not included in the S-β-CD has ΔG value of S-β-CD/R-Miconazole in the range -4.6 to -4.2 Kcal/mol and S-β-CD/S-Miconazole in the range -4.4 to -3.9 Kcal/mol. The S-β-CD/R-Miconazole complex has a lower binding energy value (-4.6 Kcal/mol) compared to S-β-CD/S-Miconazole (-4.4 Kcal/mol). Miconazole that does not include the S-β-CD may be caused by the size of the S-β-CD cavity, therefore, the complexes between miconazole and S-β-CD could be trapped. The ΔG value of S-β-CD/R-Miconazole and S-β-CD/S-Miconazole docked using Autodock Vina software are shown in Table 2.

TABLE 1. RMSD value of the docking results

Inclusion complex	RMSD value
S-β-CD/R-Miconazole	2.227
S-β-CD/S-Miconazole	1.893
HP-β-CD/R-Miconazole	2.383
HP-β-CD/S-Miconazole	2.389
HP-γ-CD/R-Miconazole	2.941
HP-γ-CD/S-Miconazole	1.964

TABLE 2. The ΔG value of S-β-CD/R-Miconazole and S-β-CD/S-Miconazole

Inclusion complex	ΔG (Kcal/mol)	|ΔΔG| (Kcal/mol)
S-β-CD/R-Miconazole	-4.6	0.2
S-β-CD/S-Miconazole	-4.4	

FIGURE 4. Structure of S-β-CD/R-Miconazole and S-β-CD/S-Miconazole

The more negative the ΔG value, the higher affinity between ligands and receptors, therefore, the inclusion complex obtained was more stable. The enantiomer with the highest affinity will spend the most time in the mobile phase while the enantiomer with the least affinity will be eluted first from the column. The |ΔΔG| value is the difference ΔG value of R-Miconazole and S-Miconazole that describes the enantiomer separation ability to miconazole compounds using chiral selector cyclodextrin. The value of |ΔΔG| obtained is 0.2 Kcal/mol. The higher |ΔΔG| value of |ΔΔG|, the better separation of chiral compounds. The value of |ΔΔG| indicates that cyclodextrin can be used as a chiral selector to separating the enantiomer of miconazole compounds. In this study, the inclusion complex of R-Miconazole has lower ΔG values compared to S-Miconazole. It causes S-Miconazole will be eluted first from the cyclodextrin chiral selector because the stability of S-Miconazole is weaker than R-Miconazole.

TABLE 3. The ΔG value of HP-β-CD/R-Miconazole and HP-β-CD/S-Miconazole

Inclusion complex	ΔG (Kcal/mol)	|ΔΔG| (Kcal/mol)
HP-β-CD/R-Miconazole	-5.1	0.3
HP-β-CD/S-Miconazole	-4.8	

Following the same approach, the ΔG value of the inclusion complex was obtained from the docking process using Autodock Vina. The ΔG value of HP-β-CD/R-Miconazole is in the range -5.1 to -4.5 Kcal/mol and HP-β-CD/S-Miconazole in the range -4.8 to -4.3 Kcal/mol. HP-β-CD/R-Miconazole complex has a lower binding energy value (-5.1 Kcal/mol) compared to HP-β-CD/S-Miconazole (-4.8 Kcal/mol). Similar to the S-β-CD/Miconazole complex, this inclusion complex has |ΔΔG| value in the range 0.3 Kcal/mol. In addition, HP-β-CD/R-Miconazole has a lower ΔG value compared to S-Miconazole, therefore, S-Miconazole will be eluted first from HP-β-CD chiral selector. Fig. 5 shows the structure of HP-β-CD/R-Miconazole and HP-β-CD/S-Miconazole. The ΔG value of HP-β-CD/R-Miconazole and HP-β-CD/S-Miconazole docked using Autodock Vina software are shown in Table 3.

FIGURE 5. Structure of HP-β-CD/R-Miconazole and HP-β-CD/S-Miconazole

The ΔG value of HP-γ-CD/Miconazole complex was obtained from the docking process using Autodock Vina. The ΔG value of HP-γ-CD/R-Miconazole is in the range -4.4 to -4.0 Kcal/mol and HP-γ-CD/S-Miconazole in the range -4.3 to -3.8 Kcal/mol. HP-γ-CD/R-Miconazole complex has a lower binding energy value (-4.3 Kcal/mol) compared to HP-γ-CD/S-Miconazole (-4.4 Kcal/mol). This complex has |ΔΔG| value in the range 0.1 Kcal/mol. In addition, HP-γ-CD/R-Miconazole has a lower ΔG value compared to S-Miconazole, therefore, S-Miconazole will be eluted first from HP-γ-CD chiral selector. Fig. 6 shows the structure of HP-γ-CD/R-Miconazole and HP-γ-CD/S-Miconazole. The ΔG value of HP-γ-CD/R-Miconazole and HP-γ-CD/R-Miconazole docked using Autodock Vina software are shown in Table 4.

TABLE 4. The ΔG value of HP-γ-CD/R-Miconazole and HP-γ-CD/S-Miconazole

| Inclusion complex	ΔG (Kcal/mol)	|ΔΔG| (Kcal/mol)
HP-γ-CD/R-Miconazole	-4.4	0.1
HP-γ-CD/S-Miconazole	-4.3	

FIGURE 6. Structure of HP-γ-CD/R-Miconazole and HP-γ-CD/S-Miconazole

The complexing ability of cyclodextrin was modified by the presence of hydroxypropyl (HP) substituent, therefore, the interaction and complexation energies for HP-β-CD are higher than other cyclodextrin derivatives. In contrast to HP-γ-CD, the complexation energies, deformation, and interaction in HP-γ-CD are weak as the large cavity in HP-γ-CD prevents the fit between host and guest molecules [13-16]. Based on the research that has been done, the use of HP-β-CD as a chiral selector for chiral separation of miconazole obtained the lowest ΔG value with high affinity and stable complex. It indicates that HP-β-CD performed better than S-β-CD and HP-γ-CD as a chiral selector for chiral separation of miconazole compounds.

CONCLUSION

Cyclodextrin derivatives can be used to predict chiral separation of miconazole compounds, specifically sulfated-β-cyclodextrin (S-β-CD), hydroxypropyl-β-cyclodextrin (HP-β-CD), and hydroxypropyl-γ-cyclodextrin (HP-γ-CD). The results of molecular docking using Autodock Vina shows that R-Miconazole has a more stable interaction with all of the cyclodextrin derivatives compared to S-Miconazole based on its lower binding energies. These results

indicate that S-Miconazole will be eluted first followed by R-Miconazole. In addition, the stability level of cyclodextrin derivatives as a chiral selector to predicting the separation of miconazole is HP-β-CD>S-β-CD>HP-γ-CD.

ACKNOWLEDGMENTS

REFERENCES

1. E. Sanganyado, Z. Lu, Q. Fu, D. Schlenk, and J. Gan, Water Res. **124**, 527–542 (2017).
2. R. Chang. *General Chemistry: The Essential Concepts, 3rd Edition*. (McGraw-Hill, New York, 2014).
3. C. S. Nascimento, J. F. Lopes, L. Guimarães, and K. B. Borges, Analyst. **139**, 3901-3910 (2014).
4. D. K. Pratoko, Chem. Prog, **5**, 31-36 (2012).
5. S. R. Arsad, H. Maarof, W. A. W. Ibrahim, and H. Y. Aboul-Enein, Chirality. **28**, 209-214 (2016).
6. Y. Liu, L. Cai, J. Lun, M. Zhao, and X. Guo, New J. Chem., **44**, 18337-18346 (2020).
7. H. Setiawan and M. I. Irawan, Jurnal Sains dan Seni ITS. **6**, 68-72 (2017).
8. T. Rostinawati, Y. C. Sumirtapura, Elfahmi, and D. S. Retnoningrum, Pharmauho. **2**, 1-5 (2016).
9. I. Makmur, A. Halim, M. D. Oktavia, and H. Anggraeni, Jurnal Farmasi Higea, **10**, 60-75 (2018).
10. S. Wren, T. A. Berger, K.S. Boos, H. Engelhardt, E. R. Adlard, I. W. Davies, K. D. Altria, and R. Stock. Chromatographia, **54**, 59-77 (2001).
11. S. Forli, R. Huey, M. E. Pique, M. F. Sanner, D. S. Godsell, and A. J. Olson, Nat. Prot. **11**, 905-919 (2016).
12. A. N. Jain and A Nicholls, J. Comput. Aided. Mol. **22**, 133-139 (2008).
13. V. Barillaro, G. Dive, P. Bertholet, B. Evrard, L. Delattre, Z. Eric, and G. Piel, Int. J. Pharm. **342**, 152-160 (2007).
14. S. Yusri, H. Sutanto, and M.Nasikin, AIP Conf Proc. 020006-1-020006-6 (2021).
15. S. Priya, B. S. Bharathi, G. B. Veerakanellore, T. Stalin, J. Mol. Struct. **1160**, 1–8 (2018).
16. N. R. Azhari, N. Yahaya, F. Bukhari, M. Suah, S. Prabu, B. Y. Hui, M. S. Shahriman, N. N. M. Zain, Chirality. **44**, 18337–18346 (2020).

SJR   Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

Home   Journal Rankings   Country Rankings   Viz Tools   Help   About Us

AIP Conference Proceedings

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
United States	Physics and Astronomy Physics and Astronomy (miscellaneous)	American Institute of Physics	**75**
Universities and research institutions in United States			
Media Ranking in United States			

| PUBLICATION TYPE	ISSN	COVERAGE	INFORMATION
Conferences and Proceedings	0094243X, 15517616	1973-1978, 1983-1984, 1993, 2000-2001, 2003-2021	Homepage How to publish in this journal confproc@aip.org

SCOPE

Today, AIP Conference Proceedings contain over 100,000 articles published in 1700+ proceedings and is growing by 100 volumes every year. This substantial body of scientific literature is testament to our 40-year history as a world-class publishing partner, recognized internationally and trusted by conference organizers worldwide. Whether you are planning a small specialist workshop or organizing the largest international conference, contact us, or read these testimonials, to find out why so many organizers publish with AIP Conference Proceedings.

🔍 Join the conversation about this journal

SJR

Total Documents

Total Cites   Self-Cites

Citations per document

- Cites / Doc. (4 years)
- Cites / Doc. (3 years)
- Cites / Doc. (2 years)

External Cites per Doc   Cites per Doc

% International Collaboration

Citable documents   Non-citable documents

Cited documents   Uncited documents

AIP Conference Proceedings

Not yet assigned quartile

SJR 2021
0.19

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

`<a href="https://www.scima`

SCImago Graphica

Explore, visually communicate and make sense of data with our