



ScienceDirect®

Ecotoxicology and Environmental Safety

Open access

10.1

CiteScore

7.129

Impact Factor

[Submit your article](#)

[Guide for authors](#)



Menu



Search in this journal

Latest
issue

Volume 256

In progress • May 2023

About the journal

Announcement: From **January 2021** *Ecotoxicology and Environmental Safety* will become an open access journal. Authors who publish in *Ecotoxicology and Environmental Safety* will be able to make their work immediately, permanently, and freely accessible.

Ecotoxicology and Environmental Safety continues ...

[View full aims & scope](#)

\$2700 ⓘ

Article Publishing Charge
for open access

4.3 weeks

Time to First Decision

6.7 weeks

Review Time

0.9 weeks

Publication Time



View all insights

Co-Editors-in-Chief | [View full Editorial Board](#)

Richard Handy

University of Plymouth, Plymouth, United Kingdom

Bing Yan, PhD

Guangzhou University, Guangzhou, China

Articles

Latest published	Top cited	Most downloaded	Most popular
------------------	-----------	-----------------	--------------

Research article *Open access*

Influence of exposure to pesticides on telomere length and pregnancy outcome: Diethylphosphates but not Dimethylphosphates are associated with accelerated telomere attrition in a Palestinian cohort



Ecotoxicology and Environmental Safety

Co-Editors-in-Chief

Richard D. Handy
University of Plymouth
Department of Biology
Plymouth, Devon PL4 8AA
United Kingdom

Bing Yan
Shandong University,
Jinan, China

Associate Editors

Fernando Barbosa Jr
University of Sao Paulo, Sao
Paulo, Brazil
Yong Liang
Jiangnan University,
Wuhan, China
Sijin Liu
State Key Laboratory of
Environmental Chemistry and
Ecotoxicology, Research Center
for Eco-Environmental Sciences,
Chinese Academy of Sciences

Ruth Pereira
Universidade do Porto, Porto,
Portugal
Paul Sibley
School of Environmental Sciences
University of Guelph
Guelph, Ontario
N1G 2W1
Canada

Hyo-Bang Moon
Associate Professor
Marine Environment Analytical
Laboratory (MEAL)
Dept. of Marine Sci. and
Convergent Tech.,
Hanyang University,
Korea

Editorial Board

Eduardo Alves de Almeida
Shashi Bhushan Agrawal
Fernando Barbosa Jr
Jingwen Chen
Swaran J. S. Flora
Xiangang Hu
Taisen Iguchi
Hisato Iwata
Kelvin Sze-Yin Leung
Yi-Fan Li
Daohui Lin
Weiping Liu
Megharaj Mallavarapu

Ann Miracle
Haruhiko Nakata
Jae-Sung Rhee
Susan Shaw
Jianbo Shi
Maoyong Song
Jennifer Stauber
Louis Tremblay
Lei Wang
Po-Keung Wong
Tao Zhang
Hao Zhu

Ecotoxicology and Environmental Safety

Open access

Submit your article

Menu

Search in this journal

Volume 170

Pages 1-804 (15 April 2019)

 Download full issue

[< Previous vol/issue](#)

[Next vol/issue >](#)

Receive an update when the latest issues in this journal are published

Sign in to set up alerts

Full text access

Editorial Board

Page ii

 [View PDF](#)

FEEDBACK 

Ecotoxicology and Environmental Safety

Open access

Submit your article

Menu 

Michael A. Hart, Tom D. Miller, Charles R. Kanukiev

Pages 1-8

Article preview 

Research article ☐ Abstract only

Effects of tris(1,3-dichloro-2-propyl) phosphate (TDCPP) and triphenyl phosphate (TPP) on sex-dependent alterations of thyroid hormones in adult zebrafish

Xiaoshan Liu, Yi Cai, Yao Wang, Suhua Xu, ... Kyungho Choi

Pages 25-32

Article preview 

Research article ☐ Abstract only

Food up-take and reproduction performance of *Daphnia magna* under the exposure of Bisphenols

Yanhua Liu, Zhengyu Yan, Ling Zhang, Zhujiangcai Deng, ... Ruixin Guo

Pages 47-54

Article preview 

Research article ☐ Abstract only

Soil pollution and toxicity in an area affected by emissions from a bauxite processing plant and a power plant in Gardanne (southern France)

Rahime Oral, Giovanni Pagano, Antonietta Siciliano, Maria Toscanesi, ... Marco Trifuoggi

Pages 55-61

Article preview 

Research article ☐ Abstract only

FEEDBACK 

Ecotoxicology and Environmental Safety

Open access

Submit your article

Menu 

Research article ☐ Abstract only

Foliar spraying of melatonin confers cadmium tolerance in *Nicotiana tabacum* L

Meng Wang, Shuhui Duan, Zhicheng Zhou, Shibao Chen, Duo Wang

Pages 68-76

Article preview 

Research article ☐ Abstract only

Iron mediated hematological, oxidative and histological alterations in freshwater fish *Labeo rohita*

Mamta Singh, Anindya Sundar Barman, Asem Lembika Devi, Ayam Gangarani Devi, Pramod Kumar Pandey

Pages 87-97

Article preview 

Research article ☐ Abstract only

Allantoin attenuates cadmium-induced toxicity in cucumber plants

Sławomir Dresler, Barbara Hawrylak-Nowak, Jozef Kováčik, Martyna Pochwatka, ... Magdalena Wójciak-Kosior

Pages 120-126

Article preview 

Research article ☐ Abstract only

Responses of duckweed (*Lemna minor* L.) to aluminum stress: Physiological and proteomics analyses

Chunlei Su, Yuji Jiang, Yaru Yang, Wei Zhang, Qinsong Xu

Pages 127-140

FEEDBACK 

Ecotoxicology and Environmental Safety

Open access

Submit your article

Menu 

Article preview 

Research article ☐ Abstract only

Using Nrf2/antioxidant response element-dependent signaling to assess the toxicity potential of fly ash particles

Hongyan Wang, Yiping Chen, Jingwen Zhang, Xiuwen Tang, Xiu Jun Wang

Pages 172-179

Article preview 

Research article ☐ Abstract only

Insight into occurrence, profile and spatial distribution of organochlorine pesticides in soils of solid waste dumping sites of Pakistan: Influence of soil properties and implications for environmental fate

Marriya Sultan, Sidra Waheed, Usman Ali, Andrew James Sweetman, ... Riffat Naseem Malik

Pages 195-204

Article preview 

Research article ☐ Abstract only

Developmental neurotoxicity of maneb: Notochord defects, mitochondrial dysfunction and hypoactivity in zebrafish (*Danio rerio*) embryos and larvae

Fangjie Cao, Christopher L. Souders, Pengfei Li, Sen Pang, ... Christopher J. Martyniuk

Pages 227-237

Article preview 

Research article ☐ Abstract only

FEEDBACK 

Ecotoxicology and Environmental Safety

Open access

Submit your article

Menu



Research article ☐ Abstract only

A potential involvement of plant systemic response in initiating genotoxicity of Ag-nanoparticles in *Arabidopsis thaliana*

Ting Wang, Jingjing Wu, Shaoxin Xu, Chenguang Deng, ... Po Bian

Pages 324-330

Article preview

Research article ☐ Abstract only

Feminization, altered gonadal development, and liver damage in least killifish (*Heterandria formosa*) exposed to sublethal concentrations of 17 α -ethinylestradiol

Latonya M. Jackson, Bruce E. Felgenhauer, Paul L. Klerks

Pages 331-337

Article preview

Research article ☐ Abstract only

Scavenging effect of oxidized biochar against the phytotoxicity of lead ions on hydroponically grown chicory: An anatomical and ultrastructural investigation

Mostafa F. El-Banna, Ahmed Mosa, Bin Gao, Xianqiang Yin, ... Zahoor Ahmad

Pages 363-374

Article preview

Research article ☐ Abstract only

A novel computational solution to the health risk assessment of air pollution *via* joint toxicity prediction: A case study on selected PAH binary mixtures in particulate matters

Xian Liu, Huazhou Zhang, Wenxiao Pan, Qiao Xue, ... Aiqian Zhang

FEEDBACK

Ecotoxicology and Environmental Safety

Open access

Submit your article

Menu



Vera Kovacevic, André J. Simpson, Myrna J. Simpson

Pages 468-478

Article preview

Research article ☐ Abstract only

The response of maize protoplasts to cadmium stress mitigated by silicon

Karin Kollárová, Zuzana Kusá, Zuzana Vatehová-Vivodová, Desana Lišková

Pages 488-494

Article preview

Research article ☐ Abstract only

Metabolic perturbation, proliferation and reactive oxygen species jointly contribute to cytotoxicity of human breast cancer cell induced by tetrabromo and tetrachloro bisphenol A

Chao Zhao, Zhi Tang, Arthur Chi Kong Chung, Hailin Wang, Zongwei Cai

Pages 495-501

Article preview

Research article ☐ Abstract only

EDTA-facilitated toxic tolerance, absorption and translocation and phytoremediation of lead by dwarf bamboos

Mingyan Jiang, Shiliang Liu, Yangfan Li, Xi Li, ... Qibing Chen

Pages 502-512

Article preview

FEEDBACK

Ecotoxicology and Environmental Safety

Open access

Submit your article

Menu 

Article preview 

Research article ☐ Abstract only

Identifying sublethal endpoints for evaluating neurotoxic compounds utilizing the fish embryo toxicity test

Julie C. Krzykwa, Asal Saeid, Marlo K. Sellin Jeffries

Pages 521-529

Article preview 

Research article ☐ Abstract only

Accumulation of trace elements by corn (*Zea mays*) under irrigation with treated wastewater using different irrigation methods

Khaskhoussy Khawla, Kahlaoui Besma, Misle Enrique, Hachicha Mohamed

Pages 530-537

Article preview 

Research article ☐ Abstract only

Revisiting fish toxicity of active pharmaceutical ingredients: Mechanistic insights from integrated ligand-/structure-based assessments on acetylcholinesterase

Nikola Minovski, Melek Türker Saçan, Elif Merve Eminoğlu, Safiye Sağ Erdem, Marjana Novič

Pages 548-558

Article preview 

Research article ☐ Abstract only

Effects on tadpole snail gene expression after exposure to vinclozolin

FEEDBACK 

Ecotoxicology and Environmental Safety

Open access

Submit your article

Menu 

ultrastructural changes and antioxidative metabolism in lettuce genotypes under Cd stress

José Lavres, Flávio Henrique Silveira Rabêlo, Flávia Regina Capaldi, André Rodrigues dos Reis, ... Neusa de Lima Nogueira

Pages 578-589

Article preview 

Research article ☐ Abstract only

Lipidomic alteration and stress-defense mechanism of soil nematode *Caenorhabditis elegans* in response to extremely low-frequency electromagnetic field exposure

Yongyan Sun, Xiaomei Huang, Yahong Wang, Zhenhua Shi, ... Peng Cai

Pages 611-619

Article preview 

Research article ☐ Abstract only

Ambient PM_{2.5} causes lung injuries and coupled energy metabolic disorder

Xia Ning, Xiaotong Ji, Guangke Li, Nan Sang

Pages 620-626

Article preview 

Research article ☐ Abstract only

Development of bisphenol A (BPA)-sensing indicator *Arabidopsis thaliana* which synthesizes anthocyanin in response to BPA in leaves

DongGwan Kim, Ramin Bahmani, Jae-Heung Ko, Seongbin Hwang

Pages 627-634

Article preview 

FEEDBACK 

Ecotoxicology and Environmental Safety

Open access

Submit your article

Menu 

Article preview 

Research article ☐ Abstract only

Ecotoxic effects of loratadine and its metabolic and light-induced derivatives

Maria Rosaria Iesce, Margherita Lavorgna, Chiara Russo, Concetta Piscitelli, ... Marina Isidori
Pages 664-672

Article preview 

Research article ☐ Abstract only

The MEK/ERK/CREB signaling pathway is involved in atrazine induced hippocampal neurotoxicity in Sprague Dawley rats

Jianan Li, Xueting Li, Haoran Bi, Baixiang Li
Pages 673-681

Article preview 

Research article ☐ Abstract only

Tissue-specific transcriptional regulation of seven heavy metal stress-responsive miRNAs and their putative targets in nickel indicator castor bean (*R. communis* L.) plants

Özge Çelik, Enes Yağız Akdaş
Pages 682-690

Article preview 

Research article ☐ Abstract only

Carbon black induced DNA damage and conformational changes to mouse hepatocytes and DNA molecule: A combined study using comet assay and multi-spectra methods

FEEDBACK 

Ecotoxicology and Environmental Safety

Open access

Submit your article

Menu 

and metal uptake

Munyaradzi Mtisi, Willis Gwenzi

Pages 750-762

Article preview 

Research article ☐ Abstract only

Microbial reduction fate of chromium (Cr) in aqueous solution by mixed bacterial consortium

Linlin Ma, Jinming Xu, Nan Chen, Miao Li, Chuanping Feng

Pages 763-770

Article preview 

Research article ☐ Abstract only

Species-species interactions modulate copper toxicity under different visible light conditions

Giulia Cheloni, Valérie Gagnaux, Vera I. Slaveykova

Pages 771-777

Article preview 

Research article ☐ Abstract only

Sub-lethal effects of waterborne copper in early developmental stages of rainbow trout (*Oncorhynchus mykiss*)

Shannon Weeks Santos, Jérôme Cachot, Pierre-Yves Gourves, Christelle Clérandeau, ... Patrice Gonzalez

Pages 778-788

Article preview 

FEEDBACK 

Ecotoxicology and Environmental Safety

Open access

Submit your article

Menu



Jie Meng, Xiaoyan Lin, Jian Zhou, Ruigang Zhang, ... Xuegang Luo

Pages 9-17

Article preview

Research article ☐ Abstract only

A pilot study on using biochars as sustainable amendments to inhibit rice uptake of Hg from a historically polluted soil in a Karst region of China

Ying Xing, Jianxu Wang, Jicheng Xia, Zhenmei Liu, ... Wanli Wei

Pages 18-24

Article preview

Research article ☐ Abstract only

Photodegradation of fluazaindolizine in aqueous solution with graphitic carbon nitride nanosheets under simulated sunlight illumination

Nannan Pang, Hongfang Lin, Jiye Hu

Pages 33-38

Article preview

Review article ☐ Abstract only

Adsorption of gaseous elemental mercury on soils: Influence of chemical and/or mineralogical characteristics

Andres J. Montoya, Jorge C. Lena, Cláudia C. Windmüller

Pages 98-106

Article preview

FEEDBACK

Ecotoxicology and Environmental Safety

Open access

Submit your article

Menu 

Article preview 

Research article ☐ Abstract only

Immobilized bacterial biosensor for rapid and effective monitoring of acute toxicity in water

Hendri Wasito, Amin Fatoni, Dadan Hermawan, Sri Sutji Susilowati

Pages 205-209

Article preview 

Research article ☐ Abstract only

Combined nitrogen fertilizer and wheat straw increases the cadmium phytoextraction efficiency of *Tagetes patula*

Xinxin Ye, Hongxiang Hu, Hongying Li, Qizhong Xiong, Hongjian Gao

Pages 210-217

Article preview 

Research article ☐ Abstract only

Study on the influence of soil microbial community on the long-term heavy metal pollution of different land use types and depth layers in mine

Xingqing Zhao, Jian Huang, Jin Lu, Yu Sun

Pages 218-226

Article preview 

Research article ☐ Abstract only

Oxidative degradation of chlorpyrifos using ferrate(VI): Kinetics and reaction mechanism

FEEDBACK 



Immobilized bacterial biosensor for rapid and effective monitoring of acute toxicity in water

Hendri Wasito^{a,c,*}, Amin Fatoni^{b,c}, Dadan Hermawan^b, Sri Sutji Susilowati^a

^a Department of Pharmacy, Faculty of Health Sciences, Jenderal Soedirman University, Purwokerto 53123, Indonesia

^b Department of Chemistry, Faculty of Mathematics and Natural sciences, Jenderal Soedirman University, Purwokerto 53123, Indonesia

^c Biosensory Technology Division, Center for Maritime Biosciences Studies, Jenderal Soedirman University, Purwokerto 53123, Indonesia

ARTICLE INFO

Keywords:

Water
Toxicant
Environment
Biosensor

ABSTRACT

The use of biosensors by using microorganisms such as bacteria have short life cycles and provide other advantages. One colorimetric biosensor technique that has been developed is the use of a biosensor utilizing the incorporation of Prussian blue formation reactions mediated by *E. coli* bioreactors with ferricyanide. Immobilization is a method that allows the bacteria can be used for long-term without reducing its ability as bioreceptor. This study aimed to develop a novel and rapid immobilized bacterial biosensor for the detection of toxic compound in water and to evaluate their analytical performances. Immobilization of *E. coli* performed by trapping method using alginate material support. The bacterial suspension was mixed with sodium alginate (1:1 v/v), and the mixture was continuously dropped in CaCl_2 solution to be a form of beads. The beads were used as bioreceptor to detect toxicants regarding cadmium, arsenic, mercury, chromium and lead solutions with Prussian blue as a colorimetric indicator. The linearity and sensitivity of detection of beads to the toxicants were tested, the stability of repeated use and storage were evaluated as well. The results showed that *E. coli* could be immobilized using alginate with response value was correlated with toxic concentration. The developed biosensor was more stable when used repeatedly and could be stored in a long time. The immobilization of *E. coli* in calcium alginate bead was successfully performed as a biosensor system for monitoring acute toxicity in water.

1. Introduction

Research and publications related to the development and manufacture of biosensors that are simple, fast, easy to make and low cost to monitor pollution and detect the contaminants in water are growing rapidly due to increasing the amount of pollution and industrial activity (Cao et al., 2018; Neufeld et al., 2006; Rodriguez-Mozaz et al., 2006). Several studies used biosensor for toxicant sensing in water with various kind of biosensing and detectors (Axelrod et al., 2016; Homaei, 2017; Mohseni et al., 2018; Rensing and Maier, 2003; Tsai et al., 2015). Each of them had advantages and limitations related to their application.

The use of organisms such as fish, shrimp, or animals living in the water to monitor water quality take a long time and need a sophisticated procedure and high cost. Thus, the use of biosensors by using microorganisms such as bacteria, become a solution to solve these problems, because the microorganisms have short life cycles and can provide an excellent response to pollution or toxins that can be used within the scope of health research that related ecosystem or

environment (Bentley et al., 2001; Di Gennaro et al., 2011). Detection technique using a change of color, colorimetric, is widely used in the detection of biomolecules, metal ions, and the presence of other compounds because the response can be seen directly with the naked eye without the need for specialized instruments. Detection system by using colorimetric more simple and without using sophisticated instruments, furthermore this could reduce the costs significantly (Ali et al., 2008; Duke et al., 2008; Kim et al., 2012; Yuan et al., 2012).

Colorimetric biosensor for detection of toxicant in water can be used as an early warning sign to monitor water toxicity in the environment. One colorimetric biosensor technique that has been developed was the use of a microbial biosensor utilizing the incorporation of Prussian blue formation reactions mediated by bioreactors with ferricyanide (Zhai et al., 2013). That study used *E. coli* as a model organism and the formation of Prussian blue with degradation of distinctive blue color as an indicator of the presence of contaminants or toxicant in water. The biosensor developed method has also been successfully used to detect the presence of toxicant in waters with the indicator of color change that can be observed visually with the naked eye. This study was still

* Corresponding author at: Department of Pharmacy, Faculty of Health Sciences, Jenderal Soedirman University, Purwokerto 53123, Indonesia.

E-mail address: hendri.wasito@unsoed.ac.id (H. Wasito).

not satisfactory for practical application because the method was using fresh microbial cell suspensions that could only be used once time and the time of microbial preparation was quite long. Therefore, there is considerable interest in using immobilized microbes to solve the problem.

Immobilization is a technique that allows the bacteria can be used for long-term without reducing its ability as bioreceptor (D'souza, 2001). Among the immobilization techniques, trapping method is one method which can be used for cell immobilization. In this method, the bacteria will be in a trap in the form of a matrix polymer, for example in the form of a bead. Polymers commonly used are agarose, acrylamide, chitosan and alginate (Dong et al., 2014a). Alginate has the advantage among others because it has a good biocompatibility and low toxicity (Barin et al., 2009; Ruan et al., 2018). In this study, we developed immobilized *E. coli* bacteria as a biosensor for monitoring and detecting toxicity in waters. *E. coli* was entrapped in the form of beads using calcium alginate and evaluated their analytical performances regarding the sensing ability as a toxicant biosensor, reuse cycles of beads and storage stability. Finally, the immobilized bacterial biosensor was applied for monitoring toxicants in groundwater samples.

2. Material and methods

2.1. Chemical

Nutrient Broth (Merck, KgaG) was obtained from Darmstadt, German and used as a medium growth for *E. coli*. Sodium alginate (Sigma Aldrich, USA) and calcium chloride (Merck, Darmstadt, Germany) were used for the preparation of immobilized microbial beads. Potassium ferricyanide ($K_3(FeCN_6)$), sodium chloride, ferric chloride ($FeCl_3$), sulfuric acid, hydrochloric acid, disodium hydrogen phosphate, sodium dihydrogen phosphate, were prepared as a reagent for sensing toxicant in the sample. Standard of cadmium ($Cd(NO_3)_2$), arsenic (H_3AsO_4), mercury ($Hg(NO_3)_2$), chromium ($Cr(NO_3)_3$) and lead ($Pb(NO_3)_2$) in HNO_3 solution were purchased from Merck (Darmstadt, Germany). The groundwater from our laboratory was chosen for the real sample analysis without any treatment. All other chemicals were of analytical grade.

2.2. Instruments

Autoclave YXQ-SG 41–280 (Ningbo, China), shaker incubator Memmert EN 60529 (Schwabach, Germany), and incubator IN-601 (Taipei, Taiwan) were used for beads preparation. The UV–Vis spectrophotometer for sensing measurement of OD_{600} was performed with a Cecil CE 3021 UV–visible Spectrophotometer (Cambridge, England). Scanning electron microscopes (SEM) TM-3000 (Hitachi, Japan) was used for bead characterization. All sensing measurements were carried out at room temperature.

2.3. Microbial culture

E. coli strain ATCC 25922 that subcultured from Laboratory of Microbiology, Faculty of Medicine, Jenderal Soedirman University, Indonesia was used as a microbial biosensor. It was added to 100 mL of Nutrient broth, pH 7.2. Bacterial medium subsequently incubated in a rotary shaker with a speed of 20 rcf for 24 h at 37 °C and transferred into 1.5 mL tubes for centrifuged at 50,000 rcf for 10 min. Part of pellets and the bacterial medium was separated by pipetting supernatant then the pellets washed with 9.0 mg/mL NaCl (w/v) sterile solution. Obtained pellets then resuspended in 10 mL of sterile saline, followed by stored at 4 °C. The incubation time for culturing bacteria found to be 25 h and used to obtain the desired bacterial concentration. The calculation of the concentration of bacteria was performed using the McFarland standard (Sood et al., 2011). Absorbance value is measured using spectrophotometer at a wavelength of 600 nm.

2.4. Microbial immobilization in calcium alginate

Immobilization of *E. coli* performed by trapping method using alginate material support. Immobilization of bacterial cells adopting from Rehn et al. (2012) The amount of 7.38×10^{11} cells/mL ($OD_{600} = 2.5$) suspension of *E. coli* added to a solution of sodium alginate with a ratio of 1:1 (v/v) in beaker glass. During the addition, stirring was performed by shaking slowly. The mixture was dripped continuously using a micropipette into 0.4 mol/L $CaCl_2$ solution to form immobilized bacterial beads. Bacterial Beads consecutively washed with a solution of 0.1 mol/L phosphate buffer, pH 7 and stored at 4 °C in 0.04 mol/L $CaCl_2$ solution. Surface topography of beads was examined by scanning electron microscope (SEM).

2.5. Sensing performance

Determination of the ability of immobilized bacteria as a biosensor was adopted from Zhai et al. (2013). The method was performed by 30 beads added 250 μ L of 450 mmol/L $K_3(FeCN_6)$, 500 μ L of deionized water and 250 μ L of toxicants standard solution with concentrations of 10–500 μ g/mL followed by incubating for 30 min at 37 °C. A 400 μ L of solution was taken and mixed with 400 μ L of 5 mmol/L $FeCl_3$ in 3 mol/L HCl and 1200 μ L of deionized water. The absorbance was measured using spectrophotometers at 700 nm wavelengths and using deionized water as a blank. Measurement performed in three replicated and shown as relative activity between absorption intensity beads with toxicant and absorption intensity beads without toxicant with deionized water as a control. The stability of reuse cycles of beads was performed by measure the relative activity of beads that repeatedly used for toxicant detection. Beads washed with phosphate buffer solution three times and incubated for 15 min before subsequent use. The stability of storage of beads assayed by the evaluated relative activity of beads that stored in 0.04 mol/L $CaCl_2$ solution at 4 °C for 20 days. Application of biosensor for assays of groundwater samples was conducted using groundwater sample, and spiked groundwater that added Cd and As standard solutions.

3. Results and discussion

3.1. Immobilization of *E. coli* in calcium alginate

The bacteria of *E. coli* was harvested at the end of the log phase period and subsequently immobilized using calcium alginate matrixes in the form of beads. Bacteria cell was harvested and determined as the optimum incubation time because, in this phase, the bacteria achieved maximum growth (Odaci et al., 2009). The concentration result of *E. coli* suspension that calculated based on the McFarland approaches using spectrophotometer at OD_{600} 2.5 was 7.38×10^{11} cells/mL. The light emitted in the spectrophotometer will hit the bacterial cell resulted in some of the light will be absorbed and partly passed on. The amount of light absorbed was proportional to the concentration of bacterial cells (Domínguez et al., 2001; Koch, 2007; Rescott et al., 1988).

E. coli trapped in alginate beads when a solution of 5% alginate mixed with bacteria was dropped into the $CaCl_2$ solution. Beads were formed due to ionic interactions between Ca^{2+} bivalent cations with carboxylate groups of alginates (Bajpai and Sharma, 2004). The bead of immobilized *E. coli* was a round shape (Fig. 1). The results of this study were accordance with other research (Dong et al., 2014b) which states that the use of alginate with a concentration of 50 mg/mL can produce hard and round shape beads. The higher concentration of alginate would make the smaller pore size of the beads followed by in reduced efficiency of immobilization. The occurrence of diffusion detention whereas low alginate concentrations will result in fragile beads and large porous this will lead to the release of bacterial cells from beads (Bibi et al., 2015; Bilal and Asgher, 2015). Increased $CaCl_2$




Fig. 1. Calcium alginate beads containing immobilized cells of *E. coli*. (A) The visual picture of beads (B) Scanning electron microscopy image of the bead with a diameter of 3–4 mm.

concentrations strengthen Ca-bonds with alginate chains thus the strength of the beads will also increase, and 0.4 mol/L CaCl_2 would produce stronger beads (Chai et al., 2004; Donati and Paoletti, 2009). In this study, 1000 μL of alginate-bacterial suspensions roughly formed 30 beads with a diameter of 3–4 mm.

3.2. Sensing performance of the immobilized microbial biosensor

Prussian blue was a colorimetric indicator of the process of reduction of ferricyanide to ferrocyanide during bacterial respiration with the addition of FeCl_3 and can be measured using spectrophotometer (Fig. 2). Consequently, selecting the wavelength that gave the optimum absorption in spectrophotometry was essential to prove appropriate measurement results (Wasito et al., 2018). The maximum wavelength of Prussian blue was 700 nm, this result was in accordance with related work that shows similar UV–Vis spectra (Zhai et al., 2013). This wavelength then used to measure the response of biosensors for this study. The feasibility of the novel immobilized bacterial biosensor in detecting toxicant was performed with the toxic metal standard solution of Cd, As, Hg, Cr, and Pb as the model toxicants. The absorbance data obtained than converted into relative activity. Calibration curves between




Fig. 2. UV–vis absorption spectra of Prussian blue formed by reagents that incubated using beads. Beads with *E. coli* could produce Prussian blue that has maximum absorption spectra of 700 nm, on the other hand, beads without *E. coli* that incubated with reagents, not interference the absorption spectra of Prussian blue formed.

Table 1

Values of regression coefficient between concentration of toxicant and relative activity for bacterial beads after incubated with various toxicants.

Toxicant	Regression equation	R^2	LOD ($\mu\text{g/mL}$)
Cd	$y = -0.099x + 84.197$	0.9349	61
As	$y = -0.077x + 88.871$	0.8265	8
Hg	$y = -0.101x + 80.997$	0.8915	82
Cr	$y = -0.062x + 74.045$	0.9562	51
Pb	$y = -0.066x + 91.268$	0.9058	76

the concentration of toxicant and relative activity (Table 1) shown that there was a linear relationship with the square of correlation coefficients of 0.8265–0.9562 over the concentration range of 10–500 $\mu\text{g/mL}$. Based on the standard deviation of the response and the slope of the calibration curve, limit of detection (LOD) was different among toxicants from 8 to 82 $\mu\text{g/mL}$.

The relative activity of bacterial beads for monitoring various toxicants with different concentration levels (Fig. 3) proved that the higher the toxicant concentration, the smaller the response value. The higher the toxicant concentration, the less the amount of $\text{K}_3[\text{Fe}(\text{CN})_6]$ converted to $\text{K}_4[\text{Fe}(\text{CN})_6]$ during bacterial respiration process resulted in the amount of Prussian blue produced was also decreased because toxicant inhibits bacterial respiration (Nwachukwu and Pulford, 2011; Zhai et al., 2013). Hg was seemed to be more toxic than other toxins. On the other hand, Pb appeared less toxic for *E. coli*. Each toxicant had different response depended on their characteristics. For instance, toxicants of Cd and As have different binding sites on *E. coli*; Cd has two binding sites whereas As has three binding sites. The difference between binding sites affects the toxicity and selectivity of toxicant (Chen and Rosen, 2014). This result was in accordance with the previous




Fig. 3. The effect of Cd, As, Hg, Cr, and Pb with various concentrations on the relative activity of bacterial biosensor beads for monitoring toxicants in water.




Fig. 4. Profile of reuse cycles of bacterial biosensor beads for monitoring of toxicants in water.




Fig. 5. The stability of bacterial biosensor beads after different storage times at 4 °C for monitoring toxicants in water.

study which reported that As not inhibit bacterial respiration significantly (Zhai et al., 2013).

The reuse cycles assay of bacterial beads was performed to determine the stability of bacterial beads biosensor when used more than one time for toxicant detection (Fig. 4). In general, it could be seen that immobilized of *E. coli* in beads after used four times could retain its response ability up to 85% and decreased gradually. The stability of beads on reuse cycles assay showed that immobilization of *E. coli* with calcium alginate proved the stability of beads form and bacterial entrapment. This result was in accordance with a previous study using other microorganisms of immobilized *Aspergillus* in calcium alginate which reported the ability to maintain the biosensor response of 91% after three times of use (El-Ghonemy, 2015). The gradual decrease in response activity could be caused by a loss of microbial cells during the washing process after each recycle used (Bibi et al., 2015).

Storage beads stability tests were performed to evaluate the stability of beads during storage in 4 °C for various periods (Fig. 5). Biosensor microbial beads seemed to be stable when it was stored up to 20 days. This stability of beads simplified the assay process for biosensing toxicant compared using free cells microbial biosensor. Microbial biosensor using free cells should be used on the day of harvest, so it is not possible to be stored (Zhai et al., 2013). This research found that the microbial biosensor with immobilized *E. coli* has higher stability than the free cell. According to the previous study with other microorganisms which reported that the immobilization of *A. fischeri* in alginate could be stored up to 10 weeks of storage and a longer storage time of 200 days immobilization of *E. coli* in alginate still shows stability response (Futra et al., 2014; Rehn et al., 2012).

The next step in this study was to apply the immobilized microbial biosensor developed to monitor some toxicants in groundwater samples. *E. coli* can be applied for toxic detection in groundwater and the response of relative activity correlated with increasing of toxicant




Fig. 6. Application of bacterial biosensor beads for monitoring toxicants in groundwater with and without added of toxicants.

concentrations (Fig. 6). In the assay using groundwater samples showed that no toxicant detected indicated by the relative activity value found to be 100% or the same as the blank sample. The assay was also performed with the toxicant of Cd, and As at the concentrations of 10 µg/mL and 100 µg/mL respectively, that added to the groundwater sample. On the addition of Cd or As, the relative activity was lower than the sample without toxicant; this result indicated that the biosensor beads was able to monitor the toxicant in groundwater samples where the higher the added toxicant concentration, the lower the relative activity biosensor response. This result proved that biosensors with immobilized bacterial of *E. coli* in calcium alginate proved a promising approach for early risk warning of acute water toxicity in groundwater.

4. Conclusion

The immobilized microbial biosensor of *E. coli* in calcium alginate beads was successfully developed and have appropriate sensing performances. The proposed microbial biosensor was novel, rapid and convenient for detection of acute toxicity and water monitoring effectively.

Acknowledgments

The authors would like to thank the Jenderal Soedirman University for financial support by the Institutional Research Program and the Department of Pharmacy, Faculty of Health Sciences, Jenderal Soedirman University for research facilities. Acknowledgments were also conveyed to Aliyah and Ratna Mutiara who had assisted technically in the laboratory in this study.

Conflict of interest

Authors declare no conflict of interest.

References

- Ali, H.D.P., Kruger, P.E., Gunnlaugsson, T., 2008. Colorimetric “naked-eye” and fluorescent sensors for anions based on amidourea functionalized 1, 8-naphthalimide structures: anion recognition via either deprotonation or hydrogen bonding in DMSO. *New J. Chem.* 32, 1153–1161.
- Axelrod, T., Eltzov, E., Marks, R.S., 2016. Bioluminescent bioreporter pad biosensor for monitoring water toxicity. *Talanta* 149, 290–297. <https://doi.org/10.1016/j.talanta.2015.11.067>.
- Bajpai, S.K., Sharma, S., 2004. Investigation of swelling/degradation behavior of alginate beads crosslinked with Ca^{2+} and Ba^{2+} ions. *React. Funct. Polym.* 59, 129–140. <https://doi.org/10.1016/j.reactfunctpolym.2004.01.002>.
- Barin, M., Otadi, M., Khorasheh, F., Kheirloomoom, A., 2009. Effect of cell concentration on the acylation of penicillin G enzymatic reaction in immobilized cells. *Sci. Iran. Trans. C Chem. Chem. Eng.* 16, 69.
- Bentley, A., Atkinson, A., Jezek, J., Rawson, D.M., 2001. Whole cell biosensors — electrochemical and optical approaches to ecotoxicity testing. *Toxicol. Vitro* 15, 469–475. [https://doi.org/10.1016/S0887-2333\(01\)00052-2](https://doi.org/10.1016/S0887-2333(01)00052-2).

- Bibi, Z., Qader, S.A.U., Aman, A., 2015. Calcium alginate matrix increases the stability and recycling capability of immobilized endo- β . *Extremophiles* 19, 819–827.
- Bilal, M., Asgher, M., 2015. Dye decolorization and detoxification potential of Ca-alginate beads immobilized manganese peroxidase. *BMC Biotechnol.* 15, 111.
- Cao, Y., Lei, K., Zhang, X., Xu, L., Lin, C., Yang, Y., 2018. Contamination and ecological risks of toxic metals in the Hai River, China. *Ecotoxicol. Environ. Saf.* 164, 210–218. <https://doi.org/10.1016/j.ecoenv.2018.08.009>.
- Chai, Y., Mei, L., Wu, G., Lin, D., Yao, S., 2004. Gelation conditions and transport properties of hollow calcium alginate capsules. *Biotechnol. Bioeng.* 87, 228–233.
- Chen, J., Rosen, B.P., 2014. Biosensors for inorganic and organic arsenicals. *Biosensors* 4, 494–512.
- Di Gennaro, P., Bruzzese, N., Anderlini, D., Aiossa, M., Papacchini, M., Campanella, L., Bestetti, G., 2011. Development of microbial engineered whole-cell systems for environmental benzene determination. *Ecotoxicol. Environ. Saf.* 74, 542–549. <https://doi.org/10.1016/j.ecoenv.2010.08.006>.
- Domínguez, M.C., de la Rosa, M., Borobio, M.V., 2001. Application of a spectrophotometric method for the determination of post-antibiotic effect and comparison with viable counts in agar. *J. Antimicrob. Chemother.* 47, 391–398. <https://doi.org/10.1093/jac/47.4.391>.
- Donati, I., Paoletti, S., 2009. Material properties of alginates. In: *Alginates: Biology and Applications*. Springer, pp. 1–53.
- Dong, Y., Zhang, Y., Tu, B., Miao, J., 2014a. Immobilization of ammonia-oxidizing bacteria by calcium alginate. *Ecol. Eng.* 73, 809–814.
- Dong, Y., Zhang, Y., Tu, B., Miao, J., 2014b. Immobilization of ammonia-oxidizing bacteria by calcium alginate. *Ecol. Eng.* 73, 809–814. <https://doi.org/10.1016/j.ecoleng.2014.09.020>.
- D'souza, S., 2001. Microbial biosensors. *Biosens. Bioelectron.* 16, 337–353.
- Duke, R.M., O'Brien, J.E., McCabe, T., Gunnlaugsson, T., 2008. Colorimetric sensing of anions in aqueous solution using a charge neutral, cleft-like, amidothiourea receptor: tilting the balance between hydrogen bonding and deprotonation in anion recognition. *Org. Biomol. Chem.* 6, 4089–4092. <https://doi.org/10.1039/B807579D>.
- El-Ghonemy, D.H., 2015. Immobilization and characterization of a thermostable β -glucosidase from *Aspergillus terreus* NRRL 265. *J. Microbiol. Biotechnol. Food Sci.* 4, 287–291.
- Futra, D., Heng, L.Y., Surif, S., Ahmad, A., Ling, T.L., 2014. Microencapsulated *Aliivibrio fischeri* in alginate microspheres for monitoring heavy metal toxicity in environmental waters. *Sensors* 14, 23248–23268.
- Homaei, A., 2017. Immobilization of *Penaeus merguensis* alkaline phosphatase on gold nanorods for heavy metal detection. *Ecotoxicol. Environ. Saf.* 136, 1–7. <https://doi.org/10.1016/j.ecoenv.2016.10.023>.
- Kim, H.N., Ren, W.X., Kim, J.S., Yoon, J., 2012. Fluorescent and colorimetric sensors for detection of lead, cadmium, and mercury ions. *Chem. Soc. Rev.* 41, 3210–3244. <https://doi.org/10.1039/C1CS15245A>.
- Koch, A.L., 2007. Growth Meas. 172–199. <https://doi.org/10.1128/9781555817497.ch9>.
- Mohseni, M., Abbaszadeh, J., Maghool, S.-S., Chaichi, M.-J., 2018. Heavy metals detection using biosensor cells of a novel marine luminescent bacterium *Vibrio* sp. MM1 isolated from the Caspian Sea. *Ecotoxicol. Environ. Saf.* 148, 555–560. <https://doi.org/10.1016/j.ecoenv.2017.11.002>.
- Neufeld, T., Biran, D., Popovtzer, R., Erez, T., Ron, E.Z., Rishpon, J., 2006. Genetically engineered pfabA pfabR bacteria: an electrochemical whole cell biosensor for detection of water toxicity. *Anal. Chem.* 78, 4952–4956. <https://doi.org/10.1021/ac052096r>.
- Nwachukwu, O., Pulford, I., 2011. Microbial respiration as an indication of metal toxicity in contaminated organic materials and soil. *J. Hazard. Mater.* 185, 1140–1147.
- Odaci, D., Timur, S., Telefoncu, A., 2009. A microbial biosensor based on bacterial cells immobilized on chitosan matrix. *Bioelectrochemistry* 75, 77–82. <https://doi.org/10.1016/j.bioelechem.2009.01.002>.
- Rehn, G., Grey, C., Branneby, C., Lindberg, L., Adlercreutz, P., 2012. Activity and stability of different immobilized preparations of recombinant *E. coli* cells containing ω -transaminase. *Process Biochem.* 47, 1129–1134. <https://doi.org/10.1016/j.procbio.2012.04.013>.
- Rensing, C., Maier, R.M., 2003. Issues underlying use of biosensors to measure metal bioavailability. *Ecotoxicol. Environ. Saf. Spec. Issue Methodol. Assess. Expo. Met.: Speciat. Bioaccess. Bioavailab. Environ. Food Feed* 56, 140–147. [https://doi.org/10.1016/S0147-6513\(03\)00057-5](https://doi.org/10.1016/S0147-6513(03)00057-5).
- Rescott, D., Nix, D., Holden, P., Schentag, J., 1988. Comparison of two methods for determining in vitro postantibiotic effects of three antibiotics on *Escherichia coli*. *Antimicrob. Agents Chemother.* 32, 450–453.
- Rodríguez-Mozaz, S., de Alda, M.J.L., Barceló, D., 2006. Biosensors as useful tools for environmental analysis and monitoring. *Anal. Bioanal. Chem.* 386, 1025–1041.
- Ruan, B., Wu, P., Chen, M., Lai, X., Chen, L., Yu, L., Gong, B., Kang, C., Dang, Z., Shi, Z., Liu, Z., 2018. Immobilization of *Sphingomonas* sp. GY2B in polyvinyl alcohol-alginate-kaolin beads for efficient degradation of phenol against unfavorable environmental factors. *Ecotoxicol. Environ. Saf.* 162, 103–111. <https://doi.org/10.1016/j.ecoenv.2018.06.058>.
- Sood, S., Singhal, R., Bhat, S., Kumar, A., 2011. 2.13 - Inoculum preparation. In: Moo-Young, M. (Ed.), *Comprehensive Biotechnology*, Second edition. Academic Press, Burlington, pp. 151–164.
- Tsai, H.-F., Tsai, Y.-C., Yagur-Kroll, S., Palevsky, N., Belkin, S., Cheng, J.-Y., 2015. Water pollutant monitoring by a whole cell array through lens-free detection on CCD. *Lab. Chip* 15, 1472–1480. <https://doi.org/10.1039/c4lc01189a>.
- Wasito, H., Buranaphalin, S., Sratthaphut, L., Suntrornsuk, L., Wilairat, P., Phechkrajang, C.M., 2018. Comparison of mean centering of ratio spectra based spectrophotometric approach and HPLC method for quantitative determination of pirenixine in the presence of methylparaben and propylparaben. *Indian J. Pharm. Educ. Res.* 52, 284–292.
- Yuan, J., Gaponik, N., Eychmüller, A., 2012. Application of polymer quantum dot-enzyme hybrids in the biosensor development and test paper fabrication. *Anal. Chem.* 84, 5047–5052. <https://doi.org/10.1021/ac300714j>.
- Zhai, J., Yong, D., Li, J., Dong, S., 2013. A novel colorimetric biosensor for monitoring and detecting acute toxicity in water. *Analyst* 138, 702–707.