

Welcome...

Author and Reader



Current Issue

Vol. 4 No. 3 (2023): Journal Health Sains



Published: 2023-03-20

Supported & Tools

 Chat via WhatsApp



tools by






About journal


[Editorial Team](#)

[Reviewers](#)


[Contact](#)

[Focus and scope](#)

 ilovepdf_merged (8).pdf ^

 ilovepdf_merged (8).pdf ^

 DeteksiPembawaSi....pdf
Removed ^

 ilovepdf_merged (8).pdf
Removed ^

Show all



[Home](#) / Editorial Team

Editorial Team

Editor in Chief

Cucu Herawati, Sekolah Tinggi Ilmu Kesehatan Cirebon, Indonesia [[Scopus](#)] [[Sinta](#)] [[Google Scholar](#)]

Journal Manager

[Dian Wuri Astuti](#), STIKES Guna Bangsa Yogyakarta, Indonesia [[Scopus](#)] [[Sinta](#)] [[Google Scholar](#)]

Editorial Board

1. [Valen Fridolin Simak](#), Universitas Sam Ratulangi, Indonesia [[Scopus](#)] [[Sinta](#)] [[Google Scholar](#)]
2. [Dyah Suryani](#), Universitas Ahmad Dahlan, Indonesia [[Scopus](#)] [[Sinta](#)] [[Google Scholar](#)]
3. [Prasojo Pribadi](#), Universitas Muhammadiyah Magelang, Indonesia [[Scopus](#)] [[Sinta](#)] [[Google Scholar](#)]
4. [Jonathan Necor Tariga](#), Quirino State University, Filipina [[Scopus](#)] [[Google Scholar](#)]
5. [Fahrur Nur Rosyid](#), Universitas Muhammadiyah Surakarta, Indonesia [[Scopus](#)] [[Google Scholar](#)]
6. [Putri Ayu Ika Setiyowati](#), Universitas Muhammadiyah Lamongan, Indonesia [[Sinta](#)] [[Google Scholar](#)]
7. [Luky Dwiantoro](#), Universitas Dipenogoro, Jawa Tengah Indonesia [[Scopus](#)] [[Google Scholar](#)]
8. [Danik Iga Prasiska](#), Dinas Kesehatan Kab. Nganjuk, Indonesia [[Scopus](#)] [[Google Scholar](#)]
9. [Azza Moustafa Fahmy](#), Theodore Bilharz Research Institute, Egypt [[Scopus](#)] [[Google Scholar](#)]
10. [Sri Sumarni](#), Politeknik Kesehatan Semarang Jawa Tengah, Indonesia [[Scopus](#)] [[Google Scholar](#)]

Supported & Tools

[Home](#) / [Archives](#) / Vol. 3 No. 12 (2022): Jurnal Health Sains

Vol. 3 No. 12 (2022): Jurnal Health Sains



DOI: <https://doi.org/10.46799/jhs.v3i12>

Published: 2022-12-14

Articles

The Effect of Educational Intervention of Infant Massage Through Android Application on The Growth and Development of Infants Aged 6 to 12 Months at Anny Rahardjo Main Clinic in 2022

Elin Puspita, Rasumawati Rasumawati, Vini Yuliani, Yusuf Kristianto
1752-1761

[pdf](#)[html](#)

Faktor-Faktor yang Mempengaruhi Penggunaan Kontrasepsi Hormonal Suntik 3 Bulan

Fira Nur Hasna, Is Susiloningtyas, Friska Realita
1794-1807

[PDF](#)[html](#)

Analysis And Management Of Polychlorinated Biphenyls (Pcbs) At Temporary Storage Of Hazardous And Toxic Waste Of Pln Up3 Tanjung Karang

Romi Sepsriza, Dikpride Despa, FX. Arinto Setyawan, Rinawati Rinawati

1770-1776

 PDF html

Cytomorphological Characteristics Of Ameloblastoma By Fine Needle Aspiration Biopsy Procedure Confirmed With Histopathological Examination

Dewi Safnita, Noza Hilbertina, Yessy Setiawati

1733-1743

 PDF html

Pengaruh Kelas Remaja Putri Dan Integrasi Model Aplikasi Android "My Periode Tracker" Untuk Menurunkan Gejala Kecemasan Premenstrual Syndrome

Alice Leiwakabessy, Henny Novita, Endah Dian Marlina, Ayi Tansah Rohaeti, Jusuf Kristianto

1714-1723

 pdf html

Innovation of Elderly in Care Sticks in Maintaining Elderly Walking Balance and Speed

Jusuf Kristianto, Ni Made Riasmini, Eviana Sumarti Tambunan, Sri Djuwitaningsih, Nur Rachmat, Fajar Susanti

1808-1815

 PDF HTML

Monitoring Baby's Motor Development Through Android-Based Baby GYM Application

Nurhayati Nurhayati, Masita Masita, Husnul Khatimah, Tutik Iswanti, Jusuf Kristianto

1744-1751

 pdf html

Promosi Gizi Bayi Balita Bagi Orang Tua Anak Penderita Stunting di Puskesmas Alak Kota Kupang Tahun 2022

Loriana L. Manalor, Martina F. Diaz, Jane A. Peni

1785-1793

 PDF HTML

Hubungan Pengendalian Diabetes Melitus Type 2 Dengan Kejadian Retinopati Diabetes Di Rsud Dr Chasan Boesoirie Ternate

Eko Sudarmo Dahad Prihanto, Yetrina Yetrina, Muhammad Sultan Firman Syah, Andri W Johan Imbar

1762-1769

 PDF html

Jurnal Health Sains

The Jurnal Health Sains is a double blind peer-reviewed academic journal and open access to scientific fields. The Jurnal Health Sains monthly once by CV. Syntax Corporation Indonesia.

The Jurnal Health Sains provides a means for sustained discussion of relevant issues that fall within the focus and scopes of the journal which can be examined empirically covering health sciences, which includes: Medical sciences, Psychology, Pharmacy, Public Health, Environmental Health that belong to the health sciences context.

Journal has become a member of Crossref (Prefix:[10.46799](#)) with Online ISSN [2548-1398](#) and Print ISSN [2723-4339](#)

Journal title	Jurnal Health Sains
Initials	JHS
Abbreviation	JHS
Frequency	12 issues per year (monthly)
DOI	prefix 10.46799
Online ISSN	2723-4339
Print ISSN	2723-6927
Editor-in-chief	Cucu Herawati
Publisher	Syntax Corporation Indonesia
Citation Analysis	Google Scholar

Permanent Call for Papers

Jurnal Health Sains invites the authors to submit their manuscript for forthcoming issues. Before submitting, we encourage you to read and confirm your manuscript to the [Author Guidelines](#) and [Manuscript Template](#). Articles are normally published in Indonesia and English. Submit your manuscript to [Submission Menu](#). If you have any questions, please feel free to contact the Editor.

Jurnal Health Sains is indexed and abstracted in the following databases:

- [EBSCO](#)
- [Base](#)
- [Google Scholar](#)
- [Garuda](#)
- [Copernicus](#)
- [Road](#)
- [Scilit](#)
- [Neliti](#)
- [Crossref](#)
- [dimension](#)

THE RELATIONSHIP BETWEEN ESBLs PRODUCTION AND VIRULENCE FACTORS GENE FIMA AND PAPC IN UROPATHOGENIC ESCHERICHIA COLI ISOLATED FROM A PRIVATE HOSPITAL IN BANYUMAS REGION IN CENTRAL JAVA, INDONESIA - A CROSS SECTIONAL STUDY

Ade Hasan Basri¹, Anang Kurniawan², Joko Setyono³, Dwi Utami Anjarwati⁴

Faculty of Medicine, Jenderal Soedirman University, Purwokerto, Central Java, Indonesia

Email: adehasanbasri83@gmail.com, Kuriaku1234@gmail.com

joko.setyono1972@unsoed.ac.id, dwi.anjarwati@unsoed.ac.id

INFO ARTIKEL	ABSTRACT
Diterima 29 November 2022 Direvisi 12 December 2022 Disetujui 13 December 2022	<i>Escherichia coli</i> is one of the opportunistic pathogenic that occupies the highest position causing the incidence of UTI. Fimbriae, particularly type 1 and P fimbriae, are the most commonly implicated bacterial cell surface virulence factors. The production of ESBL and virulence factors in <i>E.coli</i> bacteria causes chronicity, persistence, and recurrence of infections that cause high morbidity and mortality. Therefore, this study was conducted to explain the relationship between ESBL production and its virulence factors in <i>E.coli</i> bacteria. The design of this research is analytic observational with a cross-sectional approach was conducted from March to May 2021. A total of 40 <i>E. coli</i> strains were isolated and collected from urine samples of UTI patients who were admitted to the hospital. in a private Hospital in Banyumas Region in Central Java, Indonesia. The HiChrome ESBL Agar Base media was used to screen for ESBL-Producing <i>E. coli</i> . Identification of <i>fimA</i> and <i>ppC</i> genes was performed by using the PCR method. All urine samples diagnosed with UTI were examined for ESBL production. As many as 25% of <i>E.coli</i> were ESBL-production. All isolates showing positive <i>E.coli</i> ESBL results were then analyzed for <i>fimA</i> and <i>papC</i> genes using the PCR method. The results obtained 100% <i>fimA</i> gene and 80% <i>papC</i> gene. The conclusion is that there is a strong relationship between ESBL production with <i>fimA</i> and <i>papC</i> genes.
Keywords: <i>ESBL, fimA, papC, Escherichia coli, virulence factors</i>	

Introduction

Infection of the urinary tract is caused by the proliferation of bacteria in the urinary system of humans (Andersen et al., 2022). Bacteria, viruses, and fungi can cause Urinary Tract Infections. The type of bacteria that causes UTI is anaerobic Gram-negative bacteria commonly found in the digestive tract (Enterobacteriaceae) (Terlizzi et al., 2017). *Escherichia coli* is opportunistic pathogenic. Enterobacteriaceae bacteria occupy the highest

position, causing UTI incidence (Prasetya et al., 2019). *E.coli* can transform from flora in the intestine to pathogens in the urinary system, where they can flourish and persist. This pathogen has various virulence factors and tactics that allow them to infect and illness the urinary tract. This strain is a uropathogenic *E.coli* (UPEC) because it is persistently linked to uropathogenic infections (Shah et al., 2019).

Fimbriae, especially type 1 and P fimbriae, are the most frequently implicated

How to cite:	Ade Hasan Basri, Anang Kurniawan, Joko Setyono, Dwi Utami Anjarwati (2022). The Relationship Between Esbls Production And Virulence Factors Gene Fima And Papc In Uropathogenic Escherichia Coli Isolated From A Private Hospital In Banyumas Region In Central Java, Indonesia - A Cross Sectional Study. <i>Jurnal Health Sains</i> , 3(12). https://doi.org/10.46799/jhs.v3i12.695
E-ISSN:	2722-5356
Published by:	Ridwan Institute

The Relationship Between Esbls Production And Virulence Factors Gene Fima And Papc In Uropathogenic Escherichia Coli Isolated From A Private Hospital In Banyumas Region In Central Java, Indonesia - A Cross Sectional Study

bacterial cell surface virulence factors (Emody et al., 2003). Fimbriae type 1 is an essential UPEC virulence factor that can stabilize bacteria's adhesion to different cell types in the urinary system (Parvez & Rahman, 2018). P fimbriae are connected to the carbohydrate complex alpha-D-Galp-(1-4)-beta-D-Galp and are pyelonephritogenic. They adhere tightly to Bowman's capsule, glomerulus, and endothelial cells that line blood channel walls in the kidney. PapC protein is the most significant protein with 80 KD, aiding this process by transporting subunits outside the cell (Wullt et al., 2000).

UTIs will be challenging to treat when experiencing antibiotic resistance problems. E.coli is one of the bacteria that can become resistant to antibiotic drugs in UTI because it can produce the Extended-Spectrum -lactamase (ESBL) enzyme, and E.coli is the highest producer of ESBL (Prasetya et al., 2019). Extended- Spectrum Beta-lactamase is an enzyme that breaks down Beta-lactams into ineffective. Beta-lactams are a class of antibiotics that work to inhibit and damage the cell walls of Gram-negative bacteria (Vickers, 2017). The production of ESBL and virulence factors in E.coli bacteria causes chronicity, persistence, and recurrence of infections that cause high morbidity and mortality (Dumaru et al., 2019). Understanding the relationship between virulence genes and ESBL production in E.coli is critical in developing successful UTI prevention and management strategies and actions, particularly for severe, recurrent, and complicated UTIs (Katongole et al., 2020). Therefore, this study was conducted to explain the relationship between ESBL production and its virulence factors in E.coli bacteria

Methods

This cross-sectional study was conducted from March to May 2021 in Central Java. E. coli isolates were isolated and collected from urine specimens of UTI patients admitted to a private Hospital in Banyumas Region in Central Java, Indonesia. The microorganisms were stored in TSB (Tryptic soy broth) containing 15% glycerol at -70°C (Fattahi et al., 2015).

Determination of ESBL-producing E. coli isolates

The screening of ESBL-producing E. coli was performed using the HiCrome™ ESBL Agar Base media. Inoculate related samples directly on the plate and incubate for 18-24 hours in aerobic conditions at 35-37 °C. Pink to purple colonies showed a positive result, namely E.coli producing ESBL (Grohs et al., 2013).

DNA extraction and PCR method

The DNA extraction kit was used to extract total genomic DNA from ten E. coli isolates. The DNA of the fimA gene in the chromosome was extracted using Presto Mini gDNA Bacteria Kit Geneid and PapC gene. DNA in Plasmid was extracted using Presto Mini Plasmid Kit Geneid according to the manufacturer's directions. Specific primers were used for amplification of the fimA and papC genes (Table 1).

Table 1.
PCR primers

Gene	Primers (5'-3')	Size of product (bp)
fimA	F:GTTGTTCTGTCGGCTCTGTC	400
	R:ATGGTGTTGGTTCCGTTATCC	
papC	F:GACGGCACTGCTGCAGGGTGTGGCG	328
	R:ATATCCTTTCTGCAGGGATGCAATA	

PCR primers adapted from Zamani and Salehzadeh (Zamani & Salehzadeh, 2018)

papC temperature 63°C for 30 seconds; elongation at 72°C for 1 minute and repeated 30 cycles; final elongation performed at 72°C for

5 minutes. The reaction was stopped at 4°C (Zamani & Salehzadeh, 2018).

Statistical analysis

SPSS program for Windows, version 16, was used for statistical analysis (SPSS 16.0).

Results and discussion

All urine samples from patients diagnosed with UTI were 40 samples. The samples were then tested for ESBL production. The results obtained were 10 isolates of *E.coli* ESBL (Figure 1) or 25% of *E.coli* ESBL, 18% other than *E.coli* (Figure 2). All isolates showing positive *E.coli* ESBL were then identified with *fimA* and *papC* genes using the PCR method. The results were 10 isolates (100%) positive for

The association between the variables was assessed using the Chi-square or Fisher's exact test. The level of significance at $p < 0.05$.

fimA gene (Figure 3), and *papC* gene 8 isolates (80%) positive, 2 isolates (20%) negative (Figure 4). Based on statistical tests to see the relationship between ESBL with *fimA* and *papC* genes, the results were $p < 0.01$ for the *fimA* gene, $p < 0.02$ for the *papC* gene (Table 2). These results indicate that there is a strong relationship between ESBL production with *fimA* and *papC* genes.




Figure 1. Isolated on HiCrome™ ESBL Agar Base media

The Relationship Between Esbls Production And Virulence Factors Gene Fima And Papc In Uropathogenic Escherichia Coli Isolated From A Private Hospital In Banyumas Region In Central Java, Indonesia - A Cross Sectional Study




Figure 2. positive ESBL percentage




Figure 3. fimA gene PCR results




Figure 4. PapC gene PCR results

Table 2.
Relationship between ESBL production with fimA and papC genes

Gene	Positive	Negative	ESBL	P values
fimA	10	0	10	0.01
papC	8	2	10	0.02

One of the most frequent bacterial illnesses is urinary tract infection (UTI), and UPEC is the culprit that causes more than half of nosocomial UTIs. The virulence factors of UPEC strains can cause an inflammatory response in UTI (Bien et al., 2012). This study aimed to determine the relationship between ESBL production and its virulence factors in *E.coli* bacteria. In this study, out of 40 isolated urine samples, *E. coli* was the highest producer of ESBL, namely 10 isolates (25%). The possible reason was *E.coli* has a plasmid that can encode resistance genetic mutation factors. The mechanism of ESBL resistance in *E.coli* is genetically inherited by new intrinsically resistant strains (Grohs et al., 2013). The results of this study are in accordance with the study of (Prasetya et al., 2019) in East Java, which explained that *E.coli* is one of the opportunistic pathogenic Enterobacteriaceae bacteria that occupy the highest position causing the incidence of UTIs and is the highest ESBL-producing bacteria. The results of a study conducted by (Shrestha et al., 2019) in Nepal showed the same results, namely *E.coli*, which was positive for ESBL more than 20% of the total sample examined.

The fimA and papC genes appeared in isolate *E. coli* ESBL. The result indicates that

fimA and papC genes can occur in all *E.coli* strains (Bien et al., 2012). fimA is the most abundant protein produced by type 1 fimbriae and functions at the time of adhesin (Parvez & Rahman, 2018). PapC genes were detected to be associated with pyelonephritis and were found in 60% of *E.coli* strains. The ability to colonize the urinary tract epithelium is known to be linked to the presence of this gene. The papC gene produces an outer membrane protein that regulates the development of P fimbriae (Winberg, 1984). Most infections caused by *E.coli* are closely related to virulence factors with the pathogenicity of *E.coli* in urinary tract infections. Several important virulence genes of UPEC strains associated with severe urinary tract infection are aseptic adhesin (AfaI), hemolysin (HLY), cytotoxic necrosis factor (cnf 1), aerobin (aer), S-pilus (sfa), P-pilus (porridge), type 1 fimbriae (fimA) (Winberg, 1984).

The results of this study indicate that there is a strong relationship between ESBL production and the virulence factors of the fimA and papC genes. These results are consistent with the study of (Shah et al., 2019), which explains that there is a significant relationship between virulence factors and ESBL resistance.

The Relationship Between Esbls Production And Virulence Factors Gene Fima And Papc In Uropathogenic Escherichia Coli Isolated From A Private Hospital In Banyumas Region In Central Java, Indonesia - A Cross Sectional Study

In the urinary system, virulence factors play several roles in the development and colonization process (Winberg, 1984). Colonization ability depends on the expression of other pilus adhesins. The virulence factor involved in the adhesion process is type 1 pili, which is essential for the adhesion process (Emody et al., 2003). They produce erythrocyte hemagglutination when they enter the urinary system and cause bacteriuria. They also enable bacteria to overcome the epithelial barrier to enter the circulation (Connell et al., 1996). For entry into urinary tract host cells, type 1 fimbriae play a significant role. Fimbriae type 1 is a highly versatile UPEC virulence factor that can stabilize bacterial attachment to various cell types throughout the urinary tract (Al-Amiery et al., 2016). UPEC strain 99% can encode genes present in type 1 fimbriae (Vigil et al., 2011), consisting mainly of the protein FimA along with FimF, FimG, and FimH (Klemm & Schembri, 2000). Another virulence factor is P fimbriae, which *E. coli* expresses. They produce erythrocyte hemagglutination when they enter the urinary system and cause bacteriuria. They also enable bacteria to overcome the epithelial barrier to enter the circulation (Riegman et al., 1988). This type of

fimbriae is encoded by the pap gene (Wullt et al., 2000). The pap gene cluster contains at least nine genes, each with two restriction sites on each end. Another papC protein, the largest at 80 KD, assists in this process by transporting subunits outside the cell (Collinson et al., 1992).

ESBL production is a common resistance mechanism of UPEC (Talbot et al., 2006). UTIs caused by ESBL-producing *E. coli* are becoming more widespread, and ESBL-producing *E. coli* are found in various Asian nations (Heffernan et al., 2009). Multidrug resistance makes selecting an antibiotic agent difficult. There is a growing link between the creation of ESBLs and multidrug resistance. The emergence of multidrug-resistant UPEC poses a serious threat to managing UTIs as medical costs increase (Neupane et al., 2016). UPEC strains that acquire potential virulence factors can improve their ability to adapt to novel environments, colonize and invade host tissues, elude immune responses, and collect resources from the host (Köhler & Dobrindt, 2011).

Conclusions

The conclusion of this study is that a strong association between ESBL production and the virulence factors of the fimA and papC genes. These findings will undoubtedly aid in understanding the pathogenicity of UTIs and their effective management, thereby reducing the inappropriate use of antibiotics. Therefore, increased physician vigilance and increased testing with Laboratory tests are needed to reduce treatment failure and prevent the spread of ESBL-producing *E. coli*.

References

- Al-Amiery, A. A., Binti Kassim, F. A., Kadhum, A. A. H., & Mohamad, A. B. (2016). Synthesis and characterization of a novel eco-friendly corrosion inhibition for mild steel in 1 M hydrochloric acid. *Scientific Reports*, 6(1), 1–13.
- Andersen, S., Nawrocki, A., Johansen, A. E., Herrero-Fresno, A., Menéndez, V. G., Møller-Jensen, J., & Olsen, J. E. (2022). Proteomes of Uropathogenic Escherichia coli Growing in Human Urine and in J82 Urinary Bladder Cells. *Proteomes*, 10(2), 15.
- Bien, J., Sokolova, O., & Bozko, P. (2012). Role of uropathogenic Escherichia coli virulence factors in development of urinary tract infection and kidney damage. *International Journal of Nephrology*, 2012.
- Collinson, S. K., Emödy, L., Trust, T. J., & Kay, W. W. (1992). Thin aggregative fimbriae from diarrheagenic Escherichia coli. *Journal of Bacteriology*, 174(13), 4490–4495.
- Connell, I., Agace, W., Klemm, P., Schembri, M., Märdil, S., & Svanborg, C. (1996). Type 1 fimbrial expression enhances Escherichia coli virulence for the urinary tract. *Proceedings of the National*

- Academy of Sciences*, 93(18), 9827–9832.
- Dumaru, R., Baral, R., & Shrestha, L. B. (2019). Study of biofilm formation and antibiotic resistance pattern of gram-negative Bacilli among the clinical isolates at BPKIHS, Dharan. *BMC Research Notes*, 12(1), 1–6.
- Emody, L., Kerenyi, M., & Nagy, G. (2003). Virulence factors of uropathogenic *Escherichia coli*. *International Journal of Antimicrobial Agents*, 22, 29–33.
- Fattahi, S., Kafil, H. S., Nahai, M. R., Asgharzadeh, M., Nori, R., & Aghazadeh, M. (2015). Relationship of biofilm formation and different virulence genes in uropathogenic *Escherichia coli* isolates from Northwest Iran. *GMS Hygiene and Infection Control*, 10.
- Grohs, P., Tillevold, B., Caumont-Prim, A., Carbonnelle, E., Day, N., Podglajen, I., & Gutmann, L. (2013). Comparison of five media for detection of extended-spectrum Beta-lactamase by use of the wasp instrument for automated specimen processing. *Journal of Clinical Microbiology*, 51(8), 2713–2716.
- Heffernan, H. M., Woodhouse, R. E., Pope, C. E., & Blackmore, T. K. (2009). Prevalence and types of extended-spectrum β -lactamases among urinary *Escherichia coli* and *Klebsiella* spp. in New Zealand. *International Journal of Antimicrobial Agents*, 34(6), 544–549.
- Katongole, P., Nalubega, F., Florence, N. C., Asimwe, B., & Andia, I. (2020). Biofilm formation, antimicrobial susceptibility and virulence genes of Uropathogenic *Escherichia coli* isolated from clinical isolates in Uganda. *BMC Infectious Diseases*, 20(1), 1–6.
- Klemm, P., & Schembri, M. A. (2000). Bacterial adhesins: function and structure. *International Journal of Medical Microbiology*, 290(1), 27–35.
- Köhler, C.-D., & Dobrindt, U. (2011). What defines extraintestinal pathogenic *Escherichia coli*? *International Journal of Medical Microbiology*, 301(8), 642–647.
- Neupane, S., Pant, N. D., Khatriwada, S., Chaudhary, R., & Banjara, M. R. (2016). Correlation between biofilm formation and resistance toward different commonly used antibiotics along with extended spectrum beta lactamase production in uropathogenic *Escherichia coli* isolated from the patients suspected of urinary tract infections visiting Shree Birendra Hospital, Chhauni, Kathmandu, Nepal. *Antimicrobial Resistance and Infection Control*, 5(1), 1–5.
- Parvez, S. A., & Rahman, D. (2018). Virulence Factors of Uropathogenic *E. coli*. *Microbiology of Urinary Tract Infections-Microbial Agents and Predisposing Factors*, 7–21.
- Prasetya, Y. A., Nisya, K., & Amanda, E. R. (2019). Aktivitas Nanoemulsi Minyak Lengkuas (*Alpinia galanga* [L] Willd) Dalam Menghambat Bakteri *Escherichia coli* Penghasil Extended Spectrum Beta Lactamases (ESBLs).
- Riegman, N., Van Die, I., Leunissen, J., Hoekstra, W., & Bergmans, H. (1988). Biogenesis of F71 and F72 fimbriae of uropathogenic *Escherichia coli*: influence of the FsoF and FstFG proteins and localization of the Fso/FstE protein. *Molecular Microbiology*, 2(1), 73–80.
- Shah, C., Baral, R., Bartaula, B., & Shrestha, L. B. (2019). Virulence factors of uropathogenic *Escherichia coli* (UPEC) and correlation with antimicrobial resistance. *BMC Microbiology*, 19(1), 1–6.
- Shrestha, R., Khanal, S., Poudel, P., Khadayat, K., Ghaju, S., Bhandari, A., Lekhak, S., Pant, N. D., Sharma, M., & Marasini, B. P. (2019). Extended spectrum β -lactamase producing uropathogenic *Escherichia coli* and the correlation of biofilm with antibiotics resistance in Nepal. *Annals of Clinical Microbiology and Antimicrobials*, 18(1), 1–6.
- Talbot, G. H., Bradley, J., Edwards Jr, J. E., Gilbert, D., Scheld, M., & Bartlett, J. G.

The Relationship Between Esbls Production And Virulence Factors Gene Fima And Papc In Uropathogenic Escherichia Coli Isolated From A Private Hospital In Banyumas Region In Central Java, Indonesia - A Cross Sectional Study

- (2006). Bad bugs need drugs: an update on the development pipeline from the Antimicrobial Availability Task Force of the Infectious Diseases Society of America. *Clinical Infectious Diseases*, 42(5), 657–668.
- Terlizzi, M. E., Gribaudo, G., & Maffei, M. E. (2017). UroPathogenic Escherichia coli (UPEC) infections: virulence factors, bladder responses, antibiotic, and non-antibiotic antimicrobial strategies. *Frontiers in Microbiology*, 8, 1566.
- Vickers, N. J. (2017). Animal communication: when i'm calling you, will you answer too? *Current Biology*, 27(14), R713–R715.
- Vigil, P. D., Alteri, C. J., & Mobley, H. L. T. (2011). Identification of in vivo-induced antigens including an RTX family exoprotein required for uropathogenic Escherichia coli virulence. *Infection and Immunity*, 79(6), 2335–2344.
- Winberg, J. (1984). P-fimbriae, bacterial adhesion, and pyelonephritis. *Archives of Disease in Childhood*, 59(2), 180.
- Wullt, B., Bergsten, G., Connell, H., Röllano, P., Gebretsadik, N., Hull, R., & Svanborg, C. (2000). P fimbriae enhance the early establishment of Escherichia coli in the human urinary tract. *Molecular Microbiology*, 38(3), 456–464.
- Zamani, H., & Salehzadeh, A. (2018). Biofilm formation in uropathogenic Escherichia coli: Association with adhesion factor genes. *Turkish Journal of Medical Sciences*, 48(1), 162–167. <https://doi.org/10.3906/sag-1707-3>

Copyright holder :

Ade Hasan Basri, Anang Kurniawan, Joko Setyono, Dwi Utami Anjarwati (2022)

First publication right :

Jurnal Health Sains

This article is licensed under:

