

Indonesian Journal of Cancer Chemoprevention

Online ISSN: 2355-8989, Print ISSN: 2088-0197

Indonesian Journal of Cancer Chemoprevention (IJCC) is an open access, peer-reviewed, triannual journal devoted to publishing articles on Cancer Chemoprevention including Experimental and Clinical Pharmacology, especially concerning Anti-Oxidants, Anti-Aging, Anti-Inflammation, Anti-Angiogenesis, and Anti-Carcinogenesis; Cancer Detection; Stem Cell Biology; Immunology; *in vitro* and *in silico* Exploration of Chemopreventive Mechanism; and Natural Products.

IJCC is an official journal of the Indonesian Society for Cancer Chemoprevention. All IJCC's materials are free to be copied and redistributed in any medium or format. However, appropriate credit should be given. The material may not be used for commercial purposes. This content licensing is in accordance with a CC license: CC-BY-NC.

IJCC has been accredited by the Ministry of Research, Technology and Higher Education of the Republic of Indonesia.



DOAJ DIRECTORY OF
OPEN ACCESS
JOURNALS



Announcements

USER

Username
Password
☐ Remember me

JOURNAL CONTENT

Search
Search Scope
All

Browse

- » By Issue
- » By Author
- » By Title

INFORMATION

- » For Readers
- » For Authors
- » For Librarians

CURRENT ISSUE

INDONESIAN JOURNAL OF CANCER CHEMOPREVENTION

[HOME](#) [ABOUT](#) [LOGIN](#) [REGISTER](#) [SEARCH](#) [CURRENT](#) [ARCHIVES](#) [ANNOUNCEMENTS](#) [ARTICLE IN PRESS](#)

[Home](#) > [About the Journal](#) > **Editorial Team**

Editorial Team

Editor in Chief

Ferry Sandra, Department of Biochemistry and Molecular Biology, Faculty of Dentistry, Universitas Trisakti, Indonesia

Board of Editors

Ari Widiyanto, Chemistry Study Program, Faculty of Mathematics and Natural Science, Universitas Tanjungpura, Indonesia, Indonesia
 Adam Hermawan, Faculty of Pharmacy, Universitas Gadjah Mada, Indonesia
 Evamarie Hey-Hawkins, Department of Chemistry and Mineralogy, Institute of Inorganic Chemistry, Universität Leipzig, Germany
 Ines Atmosukarto, College of Medicine, Biology & Environment, Australian National University, Australia
 Masashi Kawaichi, Laboratory of Gene Function in Animals, Graduate School of Biological Sciences, Nara Institute of Science and Technology, Japan
 Taifo Mahmud, Department of Pharmaceutical Science, Oregon State University, The USA, Indonesia
 Muhammad Da'i, Faculty of Pharmacy, Universitas Muhammadiyah Surakarta, Indonesia
 Nunuk Aries Nurulita, Faculty of Pharmacy, Universitas Muhammadiyah Purwokerto, Indonesia
 Restu Syamsul Hadi, Faculty of Medicine, Universitas Yarsi, Indonesia
 Solachuddin Jauhari Arief, Dentistry Programme PAPRSB Institute of Health Sciences Universiti Brunei Darussalam Jalan Tungku Link, Gadong Brunei Darussalam, BE1410, Malaysia
 Sri Handayani, Indonesian Institute of Sciences, Indonesia
 Yashwant Pathak, Department of Pharmaceutical Sciences, College of Pharmacy, University of South Florida, United States

Board of Reviewers

Adek Zamrud Adnan, Faculty of Pharmacy, Universitas Gadjah Mada, Indonesia
 Arry Yanuar, Universitas Indonesia, Indonesia
 Daryono Hadi Tjahjono, Institut Teknologi Bandung, Indonesia
 Edy Meiyanto, Cancer Chemoprevention Research Center, Faculty of Pharmacy, Universitas Gadjah Mada, Indonesia
 Elfahmi Elfahmi, Department of Pharmaceutical Biology, College of Pharmacy, Institut Teknologi Bandung, Indonesia
 Indwiani Astuti, Faculty of Medicine, Universitas Gadjah Mada, Indonesia
 Jun-ya Kato, Tumor Cell Biology Laboratories, Graduate School of Biological Sciences, Nara Institute of Science and Technology, Japan
 Koji Itahana, Duke-NUS, a school of National University of Singapore, Singapore
 Mitsunori Kiriha, Research Center for Boron Neutron Capture Therapy, Osaka Prefecture University, Japan
 Nico P.E. Vermeulen, Amsterdam Institute for Molecules, Medicine and System, Vrije Universiteit Amsterdam, Netherlands
 Prof. Ratna Asmah Susidarti, Faculty of Pharmacy, Universitas Gadjah Mada, Indonesia
 Rolf F. Barth, Department of Pathology, The Ohio State University, United States
 Sukardiman Sukardiman, Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Universitas Airlangga, Indonesia
 Wahyu Widowati, Medical Research Center, Faculty of Medicine, Maranatha Christian University, Indonesia

Officers

Beni Lestari, Cancer Chemoprevention Research Center, Faculty of Pharmacy, Universitas Gadjah Mada, Yogyakarta, Indonesia
 Dian Anita, Cancer Chemoprevention Research Center, Faculty of Pharmacy, Universitas Gadjah Mada, Indonesia
 Hilyatul Fadliyah, Cancer Chemoprevention Research Center, Faculty of Pharmacy, Universitas Gadjah Mada, Indonesia
 Nur Dina Amalina, Semarang State University, Indonesia
 Nurrani Mustika Dewi, Faculty of Pharmacy, Universitas Padjadjaran, Indonesia
 Rohmad Yudi Utomo, Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Universitas Gadjah Mada, Indonesia
 Yonika Arum Larasati, Cancer Chemoprevention Research Center, Faculty of Pharmacy, Universitas Gadjah Mada, Indonesia

Indexed by:



Indonesian Society for Cancer Chemoprevention



USER

Username

Password

☐ Remember me

JOURNAL CONTENT

Search

Search Scope

All

Browse

- » [By Issue](#)
- » [By Author](#)
- » [By Title](#)

INFORMATION

- » [For Readers](#)
- » [For Authors](#)
- » [For Librarians](#)

INDONESIAN JOURNAL OF CANCER CHEMOPREVENTION

[HOME](#) [ABOUT](#) [LOGIN](#) [REGISTER](#) [SEARCH](#) [CURRENT](#) [ARCHIVES](#) [ANNOUNCEMENTS](#) [ARTICLE IN PRESS](#)

[Home](#) > [Archives](#) > **Vol 12, No 3 (2021)**

Vol 12, No 3 (2021)

Front and Back Matters

[View or download the Front and Back Matters](#)

[PDF](#)

Table of Contents

Research Article

Citrus Flavonoids from Citrus reticulata Peels Potentially Target an Autophagy Modulator, MAP1LC3A, in Breast Cancer

Bayu Anggoro, Dennaya Kumara, Dhella Angelina, Muthi Ikawati

[PDF](#)
114-122

Association between Vitamin D and Calcium Level and BMD Alteration in Post-Menopausal Osteoporosis Patients Treated with Bisphosphonate Therapy for at Least 1 Year in Saiful Anwar Hospital Malang

Hamzah Hamzah, Thomas Erwin Christian Junus Huwae, Chodidjah Chodidjah

[PDF](#)
123-129

Addition of Beetroot Extract to Neoadjuvant Adriamycin Cyclophosphamide Regimen Increased Tumor Cell Apoptosis in Mammary Adenocarcinoma Rats

Sri Susilowati, Neni Susilningsih, Catharina Suharti

[PDF](#)
130-136

Phytochemical and Bioinformatic Studies of Citrus Flavonoids as Chemopreventive Agents Targeting GGPS1 for Liver Cancer

Ratih Kurnia Wardani, I Made Rhamandana, Christina Mutiara Putri Gono, Muthi Ikawati

[PDF](#)
137-147

Inhibition of Human Epidermal Growth Factor Receptor-2 (HER-2) from Pomelo (Citrus maxima) Flavonoid Compounds: an In Silico Approach

Roihatul Mutiah, Tanaya Jati Dharma Dewi, Arief Suryadinata, Kesimira Qonita

[PDF](#)
148-160

Bioinformatics Analysis of Rho GTP-ase Activating Protein 35 (ARHGAP35) in Breast Cancer Migration

Dicky Rizky Febrian, Joko Setyono, Muhamad Salman Fareza, Nur Amalia Choironi, Arif Fadlan, Sarmoko Sarmoko

[PDF](#)
161-169

Review Article

A Mini-Review: Possible Mechanisms of Hepatoprotective Effect of Aloe Vera Gel

Sri Handayani, Dina Aprilia, Khoirun Nisa, Vita Taufika Rosyida, Martha Purnami Wulanjati, Anjar Windarsih, Cici Darsih, Andri Frediansyah, Sari Haryanti

[PDF](#)
170-179

Indexed by:



Indonesian Society for Cancer Chemoprevention



USER

Username

Password

☐ Remember me

JOURNAL CONTENT

Search

Search Scope

All


Browse

- » [By Issue](#)
- » [By Author](#)
- » [By Title](#)

INFORMATION

- » [For Readers](#)
- » [For Authors](#)
- » [For Librarians](#)

CURRENT ISSUE



[OPEN JOURNAL SYSTEMS](#)

Bioinformatics Analysis of Rho GTP-ase Activating Protein 35 (*ARHGAP35*) in Breast Cancer Migration

Dicky Rizky Febrian¹, Joko Setyono², Muhamad Salman Fareza¹, Nur Amalia Choironi¹,
Arif Fadlan³, Sarmoko^{1*}

¹Department of Pharmacy, Faculty of Health Sciences, Jenderal Soedirman University, Jl. Dr. Soeparno Karangwangkal, Purwokerto, Central Java, Indonesia 53123

²Laboratory of Biochemistry, Faculty of Medicine, Jenderal Soedirman University, Jl. Gumbreg No 1 Purwokerto, Central Java, Indonesia, 53146

³Department of Chemistry, Faculty of Science and Data Analytics, Institut Teknologi Sepuluh Nopember, East Java, Indonesia, 60111

Abstract

Breast cancer is a second deadly cancer after lung cancer worldwide. Progression of cancer is driven by mutated cancer drive gene such as *ARHGAP35*. This study aims to analyze the role of *ARHGAP35* in the growth and development of breast cancer cells. *ARHGAP35* expression level was analyzed using Oncomine (p -value $<1E-4$; gene rank top 10%). Overall survival (OS) and disease-free survival (DFS) were evaluated by using GEPIA (median cutoff; HR displayed with 95% CI). STRING was used for analyzing the protein-protein interaction network, while WEBGESTALT for KEGG pathway and gene ontology (GO) of *ARHGAP35* and associated proteins and cBioPortal for gene mutation. *ARHGAP35* was overexpressed in several types of breast cancer, namely invasive ductal breast carcinoma (IDC), invasive ductal and lobular breast carcinoma (IDLC), invasive lobular breast carcinoma (ILC), male breast carcinoma, and mixed ductal and lobular carcinoma (MDLC). High expression of *ARHGAP35* had significantly lower OS ($p=0.045$) compared to low expression of *ARHGAP35* and the difference in DFS was not significant ($p=0.98$). *ARHGAP35* interacted with RHOA, RHOB, RHOC, RHOD, RASA1, RND1, RAC1, CDC42, FYN and SRC. KEGG pathway and GO analysis showed that these proteins are highly involved in actin-based processes through adherent junction, axon guidance, focal adhesion, regulation of actin cytoskeleton, and tight junction. Mutation rate analysis showed 34 missense, 29 truncating, 3 fusion, and 1 in frame on *ARHGAP35*. Taken together, *ARHGAP35* may involve in the growth and development of breast cancer through regulation of actin cytoskeleton pathway.

Keywords: *ARHGAP35*, breast cancer, KEGG pathway, mutation rate, actin cytoskeleton.

INTRODUCTION

Breast cancer is the second most common cancer that causes death after lung cancer. In 2004, breast cancer was the most common cancer in women globally. In 2012, 13.8% of all cancer cases were breast cancer. Based on the Global Trends of

Breast Cancer Mortality Rate, mortality from breast cancer increases significantly every year, especially

Submitted: January 1, 2022

Revised: February 22, 2022

Accepted: February 24, 2022

*Corresponding author: sarmoko@unsoed.ac.id

in developing countries and low-income areas (Azamjah, *et al.*, 2019). In addition, breast cancer patients have poor survival and prognosis (Wang, *et al.*, 2015).

Breast cancer develops due to DNA damage and genetic mutations. In some cases, DNA defects and oncogenes can be inherited, so family history is a risk factor for breast cancer (Feng, *et al.*, 2018). In addition, genetic mutations are also essential factors that can disrupt cell regulatory pathways and promote the initiation and progression of breast cancer. A study states that *ARHGAP35* has the potential as a breast cancer gene driver, which supports breast cancer growth (Rajendran and Deng, 2017).

Small GTPase (guanosine triphosphatase) of Rat sarcoma (Ras) is an essential protein in cell signaling pathways. One of the Ras groups, namely Homologous Ras (Rho), has a vital role in regulating processes involving the actin cytoskeleton involved in adhesion, migration, and cell division. GTPase switch from an active to inactive state or vice versa, which in turn activate downstream signaling (Heraud, *et al.*, 2019). GTPase is active when it binds to GTP and is inactive when it binds to GDP. GEF and GAP control GTPase activation regulation. GEF supports the active state of GTP on GTPase, and GAP inhibits GTPase by increasing GTP hydrolysis. *ARHGAP35* acts as a GAP, a negative regulator for Rho GTPase (Heraud, *et al.*, 2019; Chen, *et al.*, 2019).

The *ARHGAP35* is thought to be involved in the growth and development of cancer cells. It is based on the role of *ARHGAP35* involved in cell cycle events, migration, and invasion (Heraud, *et al.*, 2019). The deregulation of *ARHGAP35* in breast cancer can lead to increased Brk/*ARHGAP35* interactions, cell proliferation, migration, and invasion, as well as increased tumorigenicity in mouse models (Shen, *et al.*, 2008).

Bioinformatics is an interdisciplinary field that mainly involves molecular biology, computers, mathematics, and statistics (Can, 2014). Chen, *et al.*

(2019) conducted a bioinformatics analysis on the *ARHGAP* gene group in breast cancer, namely *ARHGAP4*, *ARHGAP6*, *ARHGAP7*, *ARHGAP8*, *ARHGAP9*, *ARHGAP10*, *ARHGAP11A*, *ARHGAP14*, *ARHGAP15*, *ARHGAP18*, *ARHGAP19*, *ARHGAP23*, *ARHGAP29*, and *ARHGAP2924*, and *ARHGAP30*. The study used the Oncomine database, Kaplan-Meier Plotter, bcGenExMiner, and cBioportal to determine gene expression levels, patient survival, clinical parameters, and protein-networks. The *ARHGAP9*, *ARHGAP15*, *ARHGAP19*, and *ARHGAP30* were found to be promising targets for breast cancer treatment. According to these data, the *ARHGAP* gene family impacts breast cancer prognosis (Chen, *et al.*, 2019). However, the analysis of *ARHGAP35* on the development of breast cancer has never been carried out, thus prompting us to analyze the role of *ARHGAP35* in the growth and development of breast cancer cells.

MATERIAL AND METHODS

This study was an *in silico* study to reveal the role of *ARHGAP35* on the growth and development of breast cancer cells. The databases used were: Oncomine to analyze the expression level of *ARHGAP35* in breast cancer cells and normal cells, GEPIA to analyze overall survival (OS) and disease-free survival (DFS) due to the effect of *ARHGAP35* expression, STRING to analyze the protein-protein interaction network associated with *ARHGAP35*, WEBGESTALT to analyze GO and KEGG pathway, and cBioPortal to analyze the mutation rate of *ARHGAP35*.

Expression Level of *ARHGAP35* in Breast Cancer

The expression of *ARHGAP35* in cancer and normal cells was searched on Oncomine (www.oncomine.org) by using “*ARHGAP35*” keyword through “cancer vs normal analysis” filter. The threshold used was the *p*-value of 1E-4, the fold change of 2, and the top 10% of gene rank. The

search resulted in a comparison of the expression levels of *ARHGAP35* in normal and cancer cells. The data were taken from the TCGA Breast dataset which analyzed 593 samples and 20,423 genes. The data were exported as box plot data.

Survival Analysis of Breast Cancer Affected by *ARHGAP35*

Survival analysis was performed using GEPIA database (www.gepia.cancer-pku.cn) with the keyword “*ARHGAP35*”. Next, the “survival” and “Survival Plots” tabs were selected in the new page. The “overall survival” to analyze the overall survival and “disease free survival” to analyze DFS were then selected. The intersection point was set in the median, while the hazard ratio was displayed with a 95% confidence interval (CI) along with the log *p* value. The type of cancer was entered in the “datasets” column and the samples were evaluated using the GEPIA survival plot. The data obtained was then exported in pdf file and converted into a jpg format.

Protein-protein Networking Analysis

Protein-protein interaction network associated with *ARHGAP35* was searched using STRING database (www.string-db.org). The keyword “*ARHGAP35*” was applied in the search field and *Homo sapiens* was selected as organism. The results showed a network of protein-protein interactions that have a possible involvement with *ARHGAP35* which was exported as portable network graphic file.

A schematic of the *ARHGAP35* associated protein-protein interaction network consisted of nodules and lines. Each nodule represented an interaction of protein with *ARHGAP35*. Nodules were colored and colorless (white) indicating direct and indirect interaction of protein with *ARHGAP35*. Lines depicted interactions of nodules (proteins) connected by the lines. The light blue and purple lines represent known interactions based on the databases and experimental studies; the green,

blue and red lines represent predictable interactions through gene similarity, gene fusion and co-occurrence; and the yellow, black, and white lines depict the interactions obtained through text mining, co-expression and protein similarity.

Gene Ontology dan KEGG Pathway Analysis

Gene ontology and KEGG pathway were analyzed using WEBGESTALT (www.webgestalt.org) by *Homo sapiens* organisms and the ORA method. In the functional database column, the selected gene ontology included biological processes (BP), cellular components (CC), and molecular functions (MF). In the same column, the KEGG pathway was also selected, and proteins involved with *ARHGAP35* based on protein-protein interaction network analysis was entered in the gene list column. The genome was selected as the reference set and submitted to start the analysis. The analysis produced a bar graph of the enrichment ratio and it was exported in a zip file.

Mutation of *ARHGAP35* Analysis

The *ARHGAP35* mutation rate was analyzed by using cBioPortal (www.cbioportal.org). The “breast cancer” was selected for the “query by genes” analysis and *ARHGAP35* for “gene list”. The mutations occurred in *ARHGAP35* was viewed by selecting the “mutation” tab. The analysis resulted in a mutation curve of the *ARHGAP35* gene including the type of mutation, the number of mutated samples, and the location of the mutation in the gene. The data was exported as png and tsv files.

RESULTS

The *ARHGAP35* Expression Level in Breast Cancer

The *ARHGAP35* expression level in breast cancer and in normal cells were analyzed using Oncomine. The results showed that *ARHGAP35* was overexpressed in several types of breast cancer, namely invasive ductal breast carcinoma (IDC), in-




Figure 1. Expression level of *ARHGAP35*. (a) IDC, (b) IDLC, (c) ILC, (d) MBC, (e) MDLC. 1=normal cells, 2=cancer cells.

vasive ductal and lobular breast carcinoma (IDLC), invasive lobular breast carcinoma (ILC), male breast carcinoma, and mixed ductal and lobular carcinoma (MDLC) with fold change values of 2.950, 4.039, 2.889, 2.400, and 4.017 with p -value $< 1E-4$ (Figure 1 and Table 1).

Overall Survival (OS) and Disease-Free Survival (DFS) of Breast Cancer

The effect of *ARHGAP35* expression on OS and DFS was analyzed using GEPIA. The OS and DFS analysis used samples on breast invasive carcinoma (BRCA) with a cut point in the median. The analysis produced a Kaplan-Meier graph of survival versus time. The high expression of *ARHGAP35* showed a significantly lower OS (Figure 2). The OS values of BRCA patients decreased significantly at 100 to 150 months. Meanwhile, the DFS analysis showed a non-significant difference before

150 months. It showed a significant decrease in DFS up to < 0.4 in high expression of *ARHGAP35* in the mid-month between 150 and 200. The decrease in breast cancer survival percentage may be related to higher *ARHGAP35* expression levels thereby increasing the growth of breast cancer cells.

Protein-protein Interaction Network of *ARHGAP35*

The protein-protein interaction network of *ARHGAP35* was analyzed using the STRING database. *ARHGAP35* interacted with ten proteins namely RHOA, RHOB, RHOC, RHOD, RASA1, RND1, RAC1, CDC42, FYN, and SRC (Figure 3). The analysis of KEGG pathway and GO with the WEBGESTALT database was carried out to determine the involvement and role of these proteins.

KEGG pathway analysis described the interaction pathways that occur from proteins ob-

Table 1. Fold change expression level of *ARHGAP35* in breast cancer cells vs normal cells.

Breast cancer types	Fold change	p-value
Invasive ductal breast carcinoma (IDC)	2.950	283E-21
Invasive ductal and lobular breast carcinoma (IDLC)	4.039	287E-10
Invasive lobular breast carcinoma (ILC)	2.889	334E-13
Male breast carcinoma (MBC)	2.400	3.42E-5
Mixed ductal and lobular carcinoma (MDLC)	4.017	4.63E-6

tained by the STRING analysis. The enrichment ratio level represents the evidence related pathways. The KEGG pathway analysis denoted that there are pathways related to actin-based processes, namely adherens junction, axon guidance, focal adhesion, regulation of the actin cytoskeleton, and tight junctions (Figure 4).

The GO analysis described the characteristics of a gene. Gene ontology consists of three categories, namely biological process (BP), cellular component (CC), and molecular function (MF). The BP analysis showed actin-based processes such as the formation or maintenance of the polarity of the actin cytoskeleton, regulation of cell shape, regulation of cell morphogenesis, and the formation of actin filaments. CC analysis describes the localization sites of the analyzed protein. The CC analysis reported that the analyzed proteins were available at the site of cell division, cleavage

furrow, cell cortex, ruffle, endosome, and actin cytoskeleton. The MF analysis reflects the molecular functions involving the analyzed protein. The MF analysis pointed out that the molecular functions related to GTPase activity were GTPase activity, GTP binding and amino acids (Figure 5).

Mutation Rates of *ARHGAP35*

Mutation rates in *ARHGAP35* were analyzed using cBioPortal. The number of analyzed samples was 9502 from 16 studies published in Nature, TCGA, Firehose Legacy, PanCancer, PLoS Med, and NPJ Breast Cancer. The results showed that the *ARHGAP35* mutation involved 34 missenses, 29 truncating, 3 fusions, and 1 inframe in breast cancer. The nonsense mutation caused a premature stop codon, namely in R997*, which causes *ARHGAP35* losing its ability to bind to the active RhoA. The nonsense mutation R997* in




Figure 2. Overall survival (OS) and disease-free survival in breast cancer (BRCA) related to *ARHGAP35* expression.




Figure 3. *ARHGAP35*-associated protein-protein interaction network.

ARHGAP35 resulted in the absence of a GAP domain in the *ARHGAP35* structure. In addition, there are also missense mutations such as the Y742H and R832Q that do not have an impact on the protein activity.

Mutations in the pG2 domain showed a strong impact on protein activity. This can be seen from the S866F mutation and in-frame deletion del865-870 which showed an increase in *ARHGAP35* activity. The enhanced function of *ARHGAP35* has been reported to support random cell migration and to reduce the ability of cancer cells to migrate directionally (Binamé, *et al.*, 2016). The pG2 domain is located at amino acid number 766-958 (Héraud, *et al.*, 2019). Based on the re-

sults of cBioPortal analysis, there are three mutations that occur in the pG2 domain, namely non-sense mutation Q805*, missense mutation R832Q and frame shift deletion mutation K945Rfs*3. The missense mutation R832Q has been shown to have no effect on *ARHGAP35* activity, but no studies of nonsense mutation Q805* and frame shift deletion K945Rfs*3 have reported an effect on *ARHGAP35*.

DISCUSSION

The *ARHGAP35* is a GAP protein coding gene which acts as a Rho GTPase negative regulator. The *ARHGAP35* implicates in the growth and development of breast cancer cells due to its role in cell migration and invasion (Héraud, *et al.*, 2019; Chen, *et al.*, 2019). The *ARHGAP35* has the potential to be a driver gene that can support cancer growth (Rajendran and Deng, 2017). The *ARHGAP35* is overexpressed in several types of breast cancer, namely IDC, IDLC, ILC, MBC and MDLC. High expression of *ARHGAP35* had a lower survival rate. Protein-protein interaction network analysis showed that *ARHGAP35* interacted with ten proteins, namely RHOA, RHOB, RHOC, RHOD, RASA1, RND1, RAC1, CDC42, FYN and SRC. The results of the KEGG pathway and GO analysis showed that these proteins were widely involved in cell growth and development.




Figure 4. KEGG pathway of *ARHGAP35*.




Figure 5. Gene ontology analysis of *ARHGAP35*. (a) biological process (BP), (b) cellular component (CC), and molecular function (MF).

The Oncomine analysis for gene expression level showed increasing *ARHGAP35* expression in IDC, IDLC, ILC, MBC, and MDLC. Overexpression of *ARHGAP35* may be associated with genetic and epigenetic changes (Axelsen, *et al.*, 2007). Upregulation of *ARHGAP35* has been linked to the development of breast cancer because it can enhance cell proliferation, migration, and

invasion. Regarding the function of *ARHGAP35* as a GAP, phosphorylation of *ARHGAP35* at residue Y1105 by breast tumor kinase (Brk) promotes the formation of the *ARHGAP35*/p120RasGAP complex which can stimulate *ARHGAP35* function. This can result in RhoA inactivation. In addition, disruption of the *ARHGAP35*/p120RasGAP complex with a dominant-negative *ARHGAP35* moiety

on cells may reduce the stimulatory effect of Brk on cell proliferation, migration, and invasion (Shen, *et al.*, 2008).

The survival analysis by using GEPIA indicated that high expression of *ARHGAP35* had a significantly lower OS than low expression of *ARHGAP35* ($p=0.045$). A similar analysis with DFS displayed a non-significant result ($p=0.98$), but the DFS value dropped drastically in the middle of the 150th to 200th months. These findings showed that a higher *ARHGAP35* expression level has a faster growth of breast cancer cells. Upregulation of *ARHGAP35* has been shown to promote cell proliferation, migration, and invasion, as previously described (Shen, *et al.*, 2008).

The KEGG pathway analysis exhibited that *ARHGAP35* was heavily involved in the actin cytoskeleton regulatory pathway required for RhoA regulation (Heraud, *et al.*, 2019). RhoA, a Rho GTPase, is a major regulator of the actin cytoskeleton pathway. The active Rho GTPase interacts with effector actin binding protein (ABP) which in turn regulates the rearrangement of actin filament. ABP can be categorized into two, namely G- and F-actin. The former interacts with actin monomers and the latter associates with actin filaments (Lee and Dominguez, 2010).

The *ARHGAP35* as a major RhoA regulator is involved in cell migration, entosis, and cell division. At the cell-cell junction, inhibition of RhoA activity is required to reduce actomyosin activity at cell-cell contacts and facilitate coordinated cell movement. The interaction between claudin-11 and *ARHGAP35* in the p120RasGAP domain can reduce the level of RhoA at the cell-to-cell junction so that the stability of cell-to-cell contact is maintained and facilitates the spread of tumor cells (Li, *et al.*, 2019).

The *ARHGAP35* is known to be involved in the mechanism of entosis based on its involvement in the attenuation of cell contractility. Entosis is described as a non-apoptotic cell killing program for cancer cells, in which one cell can eat the adjacent cells. This competition between cells

results in differences in cell contractility. One of the cells shows high mechanics in deformation, while the other cells accumulate actin and myosin in the cell cortex which generated tension in the cell. The increase of tension in the cell is regulated by Rho/ROCK. Rho regulates cofilin activity via Rho kinase (ROCK) which phosphorylates LIM kinase (LIMK), which in turn phosphorylates cofilin. This signal transduction pathway modulates actin accumulation in response to extracellular stimuli. The localization of *ARHGAP35* in cells that accumulate actin and myosin can polarize RhoA activity and cause propagation of cell tension to promote entosis (Lee and Dominguez, 2010; Sun, *et al.*, 2014).

This *in silico* research was conducted by utilization of the data in the database. Therefore, the *in vivo* and *in vitro* studies is necessarily needed regarding the effect of *ARHGAP35* on breast cancer based on its role as the main regulator of RhoA through the regulatory pathway of the actin cytoskeleton.

CONCLUSION

The *ARHGAP35* may involve in the growth and development of breast cancer through regulation of actin cytoskeleton pathway.

ACKNOWLEDGEMENTS

This work was supported by the Ministry of Education and Culture, the Republic of Indonesia, under the Jenderal Soedirman University Competence-Based Research Grant 2020.

DECLARATIONS OF INTEREST

None

REFERENCES

- Axelsen, J.B., Lotem, J., Sachs, L., and Domany, E., 2007, Genes overexpressed in different human solid cancers exhibit different tissue-specific

- expression profiles, *Proc. Natl. Acad. Sci. USA.*, **104**(32), 13122-13127.
- Azamjah, N., Soltan-Zadeh, Y., and Zayeri, F., 2019, Global Trend of Breast Cancer Mortality Rate: A 25-Year Study, *Asian Pac. J. Cancer Prev.*, **20**(7), 2015-2020.
- Binamé, F., Bidaud-Meynard, A., Magnan, L., Piquet, L., Montibus, B., Chabadel, A., *et al.*, 2016, Cancer-associated mutations in the protrusion-targeting region of p190RhoGAP impact tumor cell migration, *J. Cell Biol.*, **214**(7), 859-873.
- Can, T., 2014, Introduction to bioinformatics, *Methods Mol. Biol.*, **1107**, 51-71.
- Chen, W-X., Lou, M., Cheng, L., Qian, Q., Xu, L-Y., Sun, L., *et al.*, 2019, Bioinformatics analysis of potential therapeutic targets among ARHGAP genes in breast cancer, *Oncol. Lett.*, **18**(6), 6017-6025.
- Feng, Y., Spezia, M., Huang, S., Yuan, C., Zeng, Z., Zhang, L. *et al.*, 2018, Breast cancer development and progression: Risk factors, cancer stem cells, signaling pathways, genomics, and molecular pathogenesis, *Genes Dis.*, **5**(2), 77-106.
- Héraud, C., Pinault, M., Lagrée, V., and Moreau, V., 2019, p190RhoGAPs, the ARHGAP35- and ARHGAP5-Encoded Proteins, in Health and Disease, *Cells*, **8**(4), 351.
- Lee, S.H., and Dominguez, R., 2010, Regulation of actin cytoskeleton dynamics in cells, *Mol. Cells.*, **29**(4), 311-325.
- Li, C-F., Chen, J-Y., Ho, Y-H., Hsu, W-H., Wu, L-C., Lan, H-Y., *et al.*, 2019, Snail-induced claudin-11 prompts collective migration for tumour progression, *Nat. Cell Biol.*, **21**(2), 251-262.
- Rajendran, B.K. and Deng, C-X., 2017, Characterization of potential driver mutations involved in human breast cancer by computational approaches., *Oncotarget.*, **8**, 50252-50272.
- Shen, C-H., Chen, H-Y., Lin, M-S., Li, F-Y., Chang, C-C., Kuo, M-L., *et al.*, 2008, Breast tumor kinase phosphorylates p190RhoGAP to regulate rho and ras and promote breast carcinoma growth, migration, and invasion, *Cancer Res.*, **68**(19), 7779-7787.
- Sun, Q., Cibas, E.S., Huang, H., Hodgson, L., and Overholtzer, M., 2014, Induction of entosis by epithelial cadherin expression, *Cell Res.*, **24**, 1288-1298.
- Wang, Y., Zhang, Y., Pan, C., Ma, F., and Zhang, S., 2015, Prediction of poor prognosis in breast cancer patients based on microRNA-21 expression: a meta-analysis, *PLoS One*, **10**(2), e0118647.