BMC Pregnancy and Childbirth

HomeAboutArticlesCollectionsSubmission GuidelinesJoin The Editorial Board

About ▼

Contact

Editorial Board

In Review

Editorial Board

Tovah Honor Aronin, Ph.D. - Senior Editor, BMC series



Tovah Honor Aronin has been an Editor for BMC since 2016 and has been working on *BMC Pregnancy and Childbirth* since 2017. Before moving into publishing, Tovah received her Ph.D. from Johns Hopkins University, Baltimore, USA, where she studied calcium signaling in yeast, developing a novel microscopy probe for real-time measurement of calcineurin activity. An early fascination with Punnett squares lead to a focus on genetics, which then broadened into an interest in the communication and

application of scientific research across disciplines. Tovah is an editor for the <u>BMC series blog</u> and is interested in promoting best practices in health care.

Senior Editorial Board Members

Ishag Adam - Qassim University, Saudi Arabia



Prof Ishag Adam is a full Professor at Unaizah Medical College, Qassim University in the Kingdom of Saudi Arabia.

Professor Ishag is a holder of MD in Obstetrics and Gynaecology as well as PhD in Public Health, Division of Epidemiology and Biostatistics. Ishag's areas of interest, where he published extensively, included maternal and perinatal epidemiology.

Professor Ishag has been author or co-author on over 375 articles and 6 chapters that have appeared in peer-reviewed journals. He spent most of his carrier investigating communicable diseases during pregnancy, anemia, preeclampsia and he is known in the safety of artemisinins during pregnancy. Prof Ishag and his collaborators succeeded to move Praziquantel during pregnancy from unsafe zoon (X) to be safely used during pregnancy as per WHO quidelines.

Amir Aviram - University of Toronto, Canada



Dr. Amir Aviram graduated medical school in the Sackler Faculty of Medicine, Tel-Aviv University, Tel-Aviv, Israel. He finished is post-graduate training in Obstetrics and Gynecology in Israel, followed by Maternal-Fetal Medicine Fellowship program at the University of Toronto. He currently holds an academic position as an Associate Professor at the Faculty of Medicine, University of Toronto, and practices as a staff physician at the Division of Maternal-Fetal Medicine at Sunnybrook Health Sciences Centre, Toronto, ON, Canada.



Paolo Ivo Cavoretto MD PhD is a senior consultant in Obstetrics and Gynaecology at IRCCS San Raffaele Hospital, Milan, Italy and Senior Board Member of *BMC Pregnancy and Childbirth*. He is also active member of the Editorial Board of *Ultrasound in Obstetrics and Gynaecology (ISUOG)*, Associate Editor for *Archives of Gynecology and Obstetrics* and past Advisory Member for *ISUOG*. Dr Cavoretto is a member of the *International Fetoscopy Working Group* and Coordinator of the *First Trimester Committee of Italian Society of Obstetrics and*

Gynaecology (SIEOG). He is involved at editorial level or as a reviewer with other international Journals as well as in several national and international multicenter scientific projects.

He attended extensive sub-speciality training as a research fellow in King's College Hospital (London, UK). He achieved the Fetal Medicine Foundation (FMF) Diploma in Fetal Medicine and the Italian Scientific Qualification of Associate Professor in Obstetrics-Gynaecology. Dr Cavoretto has commitment in clinical activity, teaching and clinical research with a broad range of interests within the obstetric field (fetal-maternal medicine and prenatal diagnosis with a focus on ultrasound; fetal echocardiography and congenital heart defects; high-risk obstetrics including fetal monitoring in labour as well as obstetric procedures and fetal surgery for spina bifida). Since the year 2006 he serves as ultrasound trainer for the FMF.

Christian Göbl - Medical University of Vienna, Austria



Christian Göbl is associate professor and senior consultant at the Department of Obstetrics and Gynaecology, Medical University of Vienna. He graduated in *Human Medicine* in 2008 and holds a Ph.D. degree in *Endocrinology and Metabolism* as well as a M.Sc. degree in *Medical Biometry and Biostatistics*. He is the head of the metabolic research group with strong research focus in gestational diabetes mellitus,

obesity as well as pregnancy in women after bariatric surgery. Another research activity is in the assessment of novel methods to describe the major components of glucose metabolism. He joined *BMC pregnancy and Childbirth* in 2017 and is a Senior Board Member since 2020.

Vishnu Khanal - Nepal Development Society, Nepal



Dr. Vishnu Khanal is a public health researcher in Menzies School of Health Research, Australia. He has completed his PhD (public health) from Curtin University, Western Australia. Prior to PhD, he has completed Maser of Public Health (Applied Epidemiology and Biostatistics with distinction), Master of Sociology and Anthropology and Bachelor of Public Health. In his current research, he is working towards increasing access to Comprehensive Primary Health

Care using digital technologies among Indigenous Australians living in remote and very remote communities. He has significant experience in maternal and child health, primary health care, breastfeeding, international health and nutrition from his previous work with UNICEF, CARE International, and the Ministry of Health and Population Nepal. Vishnu's areas of interest, which he has published extensively, include inequity in the access and utilisation of health services, global health, and maternal and child health.

In addition to his role at Menzies School of Health Research, Vishnu is a founding member of the Nepal Development Society which is one of the leading research organizations in non-communicable disease prevention in Nepal. He joined *BMC Pregnancy and Childbirth* in 2014 as an Associate Editor and has been contributing since then.

Ashraf Nabhan - Ain Shams University, Egypt



Ashraf Nabhan is a Professor of Obstetrics and Gynecology at Ain Shams University, Egypt. He is a consultant of Obstetrics and Gynecology at the University Hospitals. His university hospital is one of the largest maternity hospitals in the Middle East and Africa with more than 16,500 births per annum. Nabhan's areas of interest, where he published extensively, include high risk pregnancy and operative obstetrics. He is a world expert in synthesized evidence and in

developing evidence-based clinical practice guidelines. Among many credentials, He is the founder and director of Egyptian Center of Evidence Based Medicine (ECEBM), renowned as a national and regional center of excellence in Evidence Based Healthcare. His center is a member of Global Evidence Synthesis Initiative (GESI) network and The Partnership for Reproductive, Maternal, Newborn, Child & Adolescent Health (PMNCH). His center produces and disseminates systematic reviews of research evidence in health care, supports policy making and guideline development and provides training to strengthen research capacity and provides peer mentoring in evidence synthesis and guideline development.

Hora Soltani - Sheffield Hallam University, UK



Professor Hora Soltani holds the Chair of Maternal and Infant Health in Sheffield Hallam University. She is a Senior Board Member for *BMC Pregnancy & Childbirth*, Co-Chair of the International Confederation of Midwives - Research Standing Committee (ICM-RSC) with a history of working as an advisor for the World Health Organization on several guideline developments, and as a member of Human Reproduction Programme Alliance for Research capacity Strengthening and Steering

Committee. Through her 20 year research experience, she has led interdisciplinary projects in collaboration with a wide range of disciplines from "Design and Technology, Psychology and Nutrition Sciences" whilst retaining a thematic integrity with the core aim of enhancing care standards and reducing health

inequalities for mothers and their babies. Using both qualitative and quantitative approaches, her main areas of research include postpartum haemorrhage, weight and physical activity during pregnancy and maternity care models particularly for vulnerable groups of women (e.g. those with obesity, of young age or from ethnic minorities).

Valerie Smith - University of Dublin, Trinity College Dublin, Ireland



Prof Valerie Smith is a Professor in Midwifery at the School of Nursing and Midwifery, Trinity College Dublin. Valerie's research interests extend to all areas of maternity care, with specific topic expertise in the areas of assessing fetal wellbeing (fetal heart rate monitoring and fetal movements), preterm birth, caesarean section, and intrapartum care. Valerie has extensive methodological expertise in systematic reviews, randomised and non-randomised trials, survey research, and qualitative research methods. Valerie has

been author or co-author on ~130 peer reviewed publications, including three book chapters and two co-authored textbooks. Valerie has contributed to national and international (World Health Organisation) guidelines, is a member of the Steering Committee of Evidence Synthesis Ireland and was a member of a previous EU COST Action (ISO907) Childbirth Cultures, Concerns and Consequences, Creating a Dynamic EU Framework for Optimal Maternity Care.

Jos van Roosmalen - Athena Institute VU University Amsterdam and Department of Obstetrics, Leiden University Medical Center, Leiden, the Netherlands



Jos van Roosmalen is emeritus professor of Safe Motherhood and Health Systems at VU University Amsterdam and a retired consultant obstetrician in the department of Obstetrics, Leiden University Medical Center, the Netherlands.

He published extensively on Safe Motherhood issues in low and high income countries and is a (co-)promotor of 25 Safe Motherhood theses, written by students of both the Northern and Southern hemisphere. He is a Senior Board Member of *BMC Pregnancy and*

Childbirth.

Among many credentials, he is a founder of the Dutch Working Party for International Safe Motherhood and Reproductive Health (since 1987), he was the Chairman of the Maternal Mortality Audit Committee of the Netherlands Society of Obstetrics and Gynecology (2001-2016) and a co-founder of the International Network for Obstetric Survey Systems (INOSS) in 2010. He was a board member of the Netherlands Society for Tropical Medicine and International Health (1992-1998). He acted as technical advisor to WHO for Maternal Near Miss (2015) and Maternal Sepsis (2016). He works as a consultant for CCBRT Capacity building for emergency obstetric care in Dar es Salaam, Tanzania and for WLF's Maternity care project Thamini Uhai in Tanzania.

Senior Editorial Board Members

Ishag Adam, University of Khartoum, Sudan
Amir Aviram, University of Toronto, Canada
Paolo Ivo Cavoretto, San Raffaele Scientific Institute, Italy
Christian Göbl, Medical University of Vienna, Austria
Eran Hadar, Rabin Medical Center, Israel
Vishnu Khanal, Menzies School of Health Research, Australia and Nepal
Development Society, Nepal
Ashraf Nabhan, Ain Shams University, Egypt
Valerie Smith, Trinity College Dublin, Ireland
Hora Soltani, Sheffield Hallam University, UK

Editorial Board Members

Raymond Aborigo, Navrongo Health Research Centre, Ghana

Sumaiya Adam, University of Pretoria, South Africa

Akilew Adane, Murdoch University, Australia

Emmanuel Adewuyi, Edith Cowan University, Australia

Agani Afaya, School of Nursing and Midwifery, University of Health and Allied

Sciences, Ho, Ghana

Kingsley Agho, University of Western Sydney, Australia

Ki Hoon Ahn, Korea University Anam Hospital, Republic of Korea

Ashraful Alam, The University of Sydney School of Public Health, Australia

Souhail Alouini, Centre Hospitalier Régional d'Orleans, France

Negash Wakagri Amanta, Ambo University, Ethiopia

Charles Ameh, Liverpool School of Tropical Medicine, United Kingdom

Carmen Amezcua-Prieto, *Preventive Medicine and Public Health, University of Granada, Spain*

Mary Amoakoh-Coleman, University of Ghana, Ghana

Melania Amorim, Instituto de Medicina Integral Prof. Fernando Figueira, Brazil

Carla Andreucci, Federal University of Sao Carlos, Brazil

Anteneh Asefa, Institute of Tropical Medicine, Belgium

Nina Ayala, Alpert Medical School of Brown University, USA

Elie Azria, Paris University - Paris Saint Joseph Hospital, France

Aduragbemi Banke-Thomas, University of Greenwich, UK

Jacob Bar, Edith Wolfson Medical Centre, Israel

Eran Barzilay, Samson Assuta Ashdod University Hospital, Israel

Theresa Bastain, University of Southern California, USA

Wubet Alebachew Bayih, Debre Tabor University, Ethiopia

Artur Beke, Semmelweis University, Hungary

Mireille Bekker, University Medical Center Utrecht, Netherlands

Pamela Berens, McGovern Medical School, University of Texas at Houston, USA

Alexandra Berezowsky, North York General Hospital and Sunnybrook Health

Sciences Centre, University of Toronto, Canada

Yifru Berhan, St Paul's Hospital Millennium Medical College, Ethiopia

Lorena Binfa, University of Chile, Chile

Hilary Brown, University of Toronto, Canada

Danilo Buca, Center for Fetal Care and High Risk Pregnancy, University of Chieti, Italy

Wally Carlo, University of Alabama at Birmingham, USA

Piya Chaemsaithong, Ramathibodi Hospital, Mahidol University, Thailand

Christos Chatzakis, Aristotle University of Thessaloniki, Greece

Leonardo Chavane, Eduardo Mondlane University, Mozambique

Min Chen, Guangzhou Medical University, China

Xiaoyan Chen, Shenzhen Baoan Women's and Children's Hospital, China, People's Republic

Christian Chisholm, University of Virginia, USA

Etienne Ciantar, Leeds Teaching Hospitals NHS Trust, United Kingdom

Ricardo Cobucci, Federal University of Rio Grande do Norte, Brazil

Patricia Coffey, PATH, USA

Denise Cornelius, University of Mississippi Medical Center, United States

Raquel Costa, Universidade Lusófona, Portugal

Berihun Dachew, Curtin University, Australia

Andrea Dall'Asta, University of Parma, Italy / Imperial College London, UK

Edward Dassah, School of Public Health, Kwame Nkrumah University of Science and Technology, Ghana

Angela Dawson, University of Technology Sydney, Australia

Ank De Jonge, Amsterdam University Medical Center, Netherlands

Jill Demirci, University of Pittsburgh, USA

Erhan Demirdag, Gazi University School of Medicine, Turkey

Deepa Dongarwar, Baylor College of Medicine, USA

Guillaume Ducarme, Centre Hospitalier Departemental, France

Pratibha Dwarkanath, St. John's Research Institute, Bangalore, India

Kristina Edvardsson, La Trobe University, Australia

Erika Edwards, University of Vermont, USA

Samantha Ehrlich, University of Tennessee, Knoxville, United States

Tariku Ersado, Wachemo University Durame campus, Ethiopia

Fabio Facchinetti, *University of Modena and Reggio Emilia, Department of Medical and Surgical Sciences for Mother Child and Adult, Italy*

Alessandra Familiari, Fondazione Policlinico Agostino Gemelli, Italy

Dazhi Fan, Foshan Institute of Fetal Medicine, Affiliated Foshan Maternity & Child

Healthcare Hospital, Southern Medical University, China

Ilaria Fantasia, I.R.C.C.S. Materno Infantile Burlo Garofolo, Italy

Alex Farr, Medical University of Vienna, Austria

Shohreh Farzan, University of Southern California, USA

Vicki Flenady, Mater Research - University of Queensland, Australia

Kazumichi Fujioka, Kobe University, Japan

Romy Gaillard, Erasmus University Medical Center, Netherlands

Jewel Gausman, Harvard TH Chan School of Public Health, USA

Ismail Guler, Gazi University, Turkey

Amanuel Gebremedhin, Curtin School of Population Health, Curtin University,

Australia

Tamirat Getachew, Haramaya University, Ethiopia

Solmaz Ghanbari-Homayi, Tabriz University of Medical Sciences, Iran

Elvira Grandone, University of Foggia, Italy

Hisham Haggag, Cairo University, Egypt

Amr Hamza, University of Saarland, Germany

Claudia Hanson, Karolinska Institutet, Sweden

Christian Haslinger, University Hospital Zurich, Switzerland

Abigail Hatcher, University of North Carolina at Chapel Hill, USA

Melanie Hayman, CQUniversity, Australia

Jianrong He, Guangzhou Women and Children's Medical Center, China

Alexander Heazell, University of Manchester, UK

Amanda Henry, University of New South Wales, Australia

Kamran Hessami, Harvard Medical School, USA

Liran Hiersch, Lis Maternity Hospital, Tel Aviv Sourasky Medical Center and Sackler

Faculty of Medicine, Tel Aviv University, Israel

Larry Hinkson, Charité University Hospital, Berlin, Germany

Sara Holton, Deakin University, Australia

Shigeko Horiuchi, St Luke's International University, Japan

Bo Huang, Huazhong University of Science and Technology, China, People's Republic

Jialyu Huang, Jiangxi Maternal and Child Health Hospital, China

Evelyn Huhn, Department of Obstetrics and Perinatal medicine, University Hospital

Basel, Switzerland and Department of Obstetrics and Perinatal medicine,

University Hospital Hamburg, Germany

Peiyin Hung, University of South Carolina Arnold School of Public Health, USA

Lisa Hurt, Cardiff University School of Medicine, UK

Nik Hazlina Nik Hussain, Universiti Sains Malaysia, Malaysia

Chiara Ionio, Dipartimento di Psicologia, Universita Cattolica, Milano, Italy

Shayesteh Jahanfar, Tufts University School of Medicine USA,

Elaine Jefford, University of the Sunshine Coast, Australia

Hong Jiang, Fudan University, China, People's Republic

Sadhana Joshi, Interactive Research School For Health Affairs, Bharati Vidyapeeth University, India

Ixora Kamisan Atan, National University of Malaysia, Malaysia

Leila Katz, Institute of Integral Medicine Prof Fernando Figueira IMIP, Brazil

Anish Keepanasseril, *Jawaharlal Institute of Postgraduate Medical Education and Research, India*

Hussein Kidanto, Aga Khan University Medical College, Tanzania

Eyal Krispin, Boston Children's Hospital, Harvard Medical School, USA

Sailesh Kumar, University Of Queensland, Australia

Sonia Lansky, Health Department Belo Horizonte City, Brasil

Olav Lapaire, University Hospital Basel, Switzerland

Osaheni Lawani, Alex Ekwueme Federal University Teaching Hospital, Ebonyi

State, Nigeria

Debora Leite, Federal University of Pernambuco, Brazil

Kwok-yin Leung, Gleneagles Hospital Hong Kong, Hong Kong SAR

Gabriel Levin, Hadassah Medical Center, Israel

Cheng Li, Obstetrics & Gynecology Hospital, Fudan University, China

Hong-tian Li, Peking University Institute of Reproductive and Child Health, China, People's Republic

Pei Shan Lim, Universiti Kebangsaan Malaysia, Malaysia

Yuan Lin, Nanjing Medical University, China

Helena Lindgren, Karolinska Institutet, Sweden

Jufen Liu, Peking University, China, People's Republic

Xitong Liu, Assisted reproduction center, China

Ambrogio P Londero, Academic Unit of Obstetrics and Gynaecology, Department

of Neuroscience, Rehabilitation, Ophthalmology, Genetics Maternal and Infant

Health, University of Genoa, Genova, Italy

Ahmed Maged, Cairo University, Egypt

Moke Magoma, EngenderHealth, Tanzania

Zaleha Abdullah Mahdy, The National University of Malaysia, Malaysia

Juan Miguel Martinez Galiano, University of Jaen, Spain

Daniele Di Mascio, Sapienza University of Rome, Italy

Salvatore Andrea Mastrolia, Department of Obstetrics and Gynecology, Ospedale

dei Bambini "Vittore Buzzi", University of Milano, Italy

Jiji Mathews, Christian Medical College Vellore, India

Mitsuaki Matsui, Nagasaki University School of Tropical Medicine and Global

Health, Japan

Shinya Matsuzaki, Osaka University, Japan and Osaka International Cancer

Institute, Japan

Shahla Meedya, University of Wollongong, Australia

Tarek Meguid, Child Health Unit, Department of Paediatrics and Child Health,

University of Cape Town, Cape Town, South Africa

Elad Mei-Dan, North York General Hospital, Canada

Philippa Middleton, South Australian Health and Medical Research Institute,

Australia

Attila Molvarec, Semmelweis University, Hungary

Maddalena Morlando, Department of Woman, Child and General and Specialized

Surgery, Obstetrics and Gynecology Unit, University of Campania "Luigi Vanvitelli",

Naples, Italy

Cheryl Moyer, University of Michigan, USA

Achenef Muche, University of Gondar and EPHI, Ethiopia

David Mukunya, Busitema University, Uganda

Mona Nabulsi, American University of Beirut, Lebanon

Ahmed Nassr, Baylor College of Medicine amd Texas Children's Hospital, USA Ilona Nenko, Jagiellonian University Medical College, Poland

Sutapa Bandyopdhyay Neogi, International Institute of Health Management

Research (IIHMR) Delhi, India

Tanya Nippita, University of Sydney, Australia

Theoneste Nkurunziza, Technical University of Munich, Germany

Roseli Nomura, Universidade Federal de Sao Paulo and Universidade de São

Paulo, Brazil

Mohd Noor, Norhayati Universiti Sains Malaysia, Malaysia

Rhona O'Connell, *University College Cork, Ireland*

Friday Okonofua, Centre of Excellence in Reproductive Health Innovation,

University of Benin, Nigeria

Rossana Orabona, Spedali Civili of Brescia, Italy

Tiago Miguel Pinto, Lusófona University, Portugal

Shuby Puthussery, University of Bedfordshire, Maternal and Child Health Research

Centre, UK

Paola Quaresima, Magna Graecia University of Catanzaro, Italy

Maggie Redshaw, University of Oxford, UK

Jianzhi Ren, Xiamen University Affiliated Chenggong Hospital, China

Elisha Riggs, Murdoch Childrens Research Institute, Australia

Fredrick J. Rosario, University of Colorado, USA

Amihai Rottenstreich, Hadassah Medical Center, Israel

Marilza Rudge, Sao Paulo State University -Medical school, Botucatu, Brazil

Stephen Rulisa, University of Rwanda, Rwanda

Gabriele Saccone, University of Naples Federico II, Italy

Emma Sacks, John Hopkins School of Public Health, USA

Hidayet Sal, Araklı Bayram Halil State Hospital, Turkey

Srividhya Sankaran, Guy's & St Thomas' NHS Foundation Trust, UK

Laura Sarno, University of Naples Federico II, Italy

Dietmar Schlembach, Vivantes Network for Health Berlin, Germany

Hen Sela, Shaare Zedek Medical Center in Jerusalem Affiliated with the Hebrew

University School of Medicine, Israel

Katherine Semrau, Ariadne Labs and Harvard Medical School, USA

Endalew Gemechu Sendo, Addis Ababa University, College of Health Sciences, Ethiopia

Loïc Sentilhes, Bordeaux Univeristy Hospital, France

Mehmet Baki Senturk, Namik Kemal University, Turkey

Steven W. Shaw, Chang Gung Memorial Hospital, Taiwan R.O.C.

Songying Shen, Guangzhou women and children's medical center, Guangzhou medical university, China

Andrew Shennan, King's College London, UK

Makino Shintaro, Juntendo University Urayasu Hospital, Japan

Filomena Giulia Sileo, University of Modena and Reggio Emilia, Italy

Angelo Sirico, AORN Caserta Hospital, Caserta, Italy

Renato Souza, University of Campinas, Brazil

Jelle Stekelenburg, University of Groningen, Netherlands

Zaharah Sulaiman, Universiti Sains Malaysia, Malaysia

Linda Tahaineh, Jordan University of Science and Technology, Jordan

Maira Takemoto, Botucatu Medical School, São Paulo State University (UNESP), Brazil

Hannah Tappis, Jhpiego - an affiliate of Johns Hopkins University, USA

Chiara Tersigni, Fondazione Policlinico A. Gemelli IRCCS, Italy

Gill Thomson, University of Central Lancashire, UK

Rachel Tinius, Western Kentucky Univeristy, USA

Stefania Triunfo, S. Paolo University Hospital, Italy

Akaninyene Eseme Ubom, Department of Obstetrics, Gynaecology & Perinatology,

Obafemi Awolowo University Teaching Hospitals Complex, Ile-Ife, Osun State,

Nigeria

Akihiko Ueda, Kyoto University, Japan

Thomas van den Akker, Leiden University Medical Center, Netherlands

Paige van der Pligt, Deakin University, Institute of Physical Activity and Nutrition,

Geelong, Victoria, Australia & Western Health, Sunshine Hospital, St Albans,

Victoria, Australia

Marleen van Gelder, Radboud University Medical Center, Netherlands

Edwin van Teijlingen, Bournemouth University, UK

Corine Verhoeven, VUMC, Netherlands

Angela Vinturache, *Grande Prairie Regional Hospital, Alberta, Canada*Bellington Vwalika, *University of Zambia, School of Medicine, Zambia*Dongqing Wang, *George Mason University, USA*Xiao Wanqing, *Guangzhou Women and Children's Medical Center, China*Tuangsit Wataganara, *Faculty of Medicine Siriraj Hospital, Mahidol University*,

Xiaozhong Wen, State University of New York at Buffalo, USA
Aleksandra Wesołowska, Warsaw Medical University, Poland
Jiang-Nan Wu, Obstetrics and Gynecology Hospital of Fudan University, China
Karen Wynter, Deakin University, Australia
Jeanine Young, University of the Sunshine Coast, Australia
Agnieszka Zawiejska, Poznan University of Medical Sciences, Poland
Depeng Zhao, Shenzhen Maternity and Child Healthcare Hospital, Southern
Medical University, China
Yeyi Zhu, Kaiser Permanente Division of Research, USA
Fabrizio Zullo, University of Rome "La Sapienza", Italy

Editorial Advisors

Thailand

Howard Berger, *University of Toronto, St. Michael's Hospital, Canada* Luis Gabriel Cuervo, *Pan American Health Organization, USA* Pisake Lumbiganon, *Khon Kaen University, Thailand*



Submit manuscript

Important information

Editorial board

For authors

For editorial board members

For reviewers

Contact Us

Annual Journal Metrics

Citation Impact

3.105 - <u>2-year Impact Factor</u> (2021)

3.853 - <u>5-year Impact Factor</u> (2021)

1.518 - <u>SNIP</u> (Source Normalized Impact per Paper)

0.960 - SJR (SCImago Journal Rank)

Speed

41 days to first decision for all manuscripts (Median) 64 days to first decision for reviewed manuscripts only (Median)

Usage

7,560,102 Downloads (2021) 4,644 Altmetric mentions (2021)

More about our metrics

Peer-review Terminology

The following summary describes the peer review process for this journal:

Identity transparency: Single anonymized

Reviewer interacts with: Editor

Review information published: Review reports. Reviewer Identities reviewer opt in. Author/reviewer

communication

More information is available here

Follow



BMC Pregnancy and Childbirth

ISSN: 1471-2393

Contact us

Submission enquiries: bmcpregnancyandchildbirth@biomedcentral.com

General enquiries: ORSupport@springernature.com

Read more on our blogs Policies Support and Contact

Receive BMC newsletters Accessibility Leave feedback

Manage article alerts Press center Careers

Language editing for authors

Scientific editing for authors

Follow BMC







By using this website, you agree to our <u>Terms and Conditions</u>, <u>California Privacy Statement</u>, <u>Privacy statement</u> and <u>Cookies</u> policy. <u>Manage cookies/Do not sell my data</u> we use in the preference centre.

SPRINGER NATURE

© 2023 BioMed Central Ltd unless otherwise stated. Part of Springer Nature.

BMC Pregnancy and Childbirth

HomeAboutArticlesCollectionsSubmission GuidelinesJoin The Editorial Board

Articles ▼

Collections

Supplements

Reviewer acknowledgements

Articles



Joint effect of maternal pre-pregnancy body mass index and folic acid supplements on gestational diabetes mellitus risk: a prospective cohort study

The joint effect of folic acid (FA) supplements and maternal pre-pregnancy body mass index (BMI) on gestational diabetes mellitus (GDM) has not been fully addressed. This study aimed to examine the joint effec...

Minyu Li, Lijiang Wang, Zhanhui Du, Qianqian Shen, Lu Jiang, Lun Sui, Nan Zhang, Hong Wang and Guoju Li

BMC Pregnancy and Childbirth 2023 23:202

Research | Published on: 23 March 2023

> Full Text> PDF

Lifestyle and the hypertensive disorders of pregnancy in nulliparous women in the United States: a secondary data analysis of the nuMom2b

Hypertensive disorders of pregnancy are a leading cause of maternal and fetal morbidity and mortality and a significant risk factor for future cardiovascular disease development in women. This study aimed to e...

Elizabeth Mollard and Constance Cottrell

BMC Pregnancy and Childbirth 2023 23:201

Research | Published on: 23 March 2023

Brief and long co-incubation of sperm and oocytes for in vitro fertilization: a meta-analysis of randomized controlled trials

There is still no consensus on the optimal time of oocyte–sperm co-incubation during in vitro fertilization and embryo transfer (IVF-ET). The aim of this meta-analysis was to compare the effects of brief (1-6 ...

Yiyue Fan, Zeyu Wu, Fang Peng, Hongyao Peng, Xin Liang and Shaomi Zhu

BMC Pregnancy and Childbirth 2023 23:200

Research | Published on: 23 March 2023

> Full Text> PDF

Histological endometrial dating: a reliable tool for personalized frozen-thawed embryo transfer in patients with repeated implantation failure in natural cycles

To evaluate the clinical availability and stability of histological endometrial dating as a tool for personalized frozen-thawed embryo transfer (pFET) in patients with repeated implantation failure (RIF) in na...

Yuan Li, Quan Wen, Jing Hu, Jingnan Liao, Xiangxiu Fan, Huijun Chen, Qi Zhao, Guang-Xiu Lu, Ge Lin and Fei Gong

BMC Pregnancy and Childbirth 2023 23:199

Research | Published on: 22 March 2023

Acceptability and user experiences of a patient-held smart card for antenatal services in Nigeria: a qualitative study

Poor maternal, newborn and child health outcomes remain a major public health challenge in Nigeria. Mobile health (mHealth) interventions such as patient-held smart cards have been proposed as effective soluti...

Ijeoma Uchenna Itanyi, Juliet Iwelunmor, John Olajide Olawepo, Semiu Gbadamosi, Alexandra Ezeonu, Adaeze Okoli, Amaka Grace Ogidi, Donaldson Conserve, Byron Powell, Chima Ariel Onoka and Echezona Edozie Ezeanolue

BMC Pregnancy and Childbirth 2023 23:198

Research | Published on: 22 March 2023

> Full Text> PDF

Women's experiences of trauma, the psychosocial impact and health service needs during the perinatal period

Traumatic events are associated with psychological and physical health problems for women in the perinatal period (i.e., pregnancy-12-months after childbirth).

Despite the negative impact of trauma on perinata...

Essence Perera, Sharon Chou, Nicole Cousins, Natalie Mota and Kristin Reynolds

BMC Pregnancy and Childbirth 2023 23:197

Research | Published on: 21 March 2023

The feasibility and acceptability of Project POWER: a mindfulness-infused, cognitive-behavioral group intervention to address mental and sexual health needs of young pregnant women in Liberia

Following 14 years of civil war in Liberia, war exposure, gender-based violence, and extreme poverty have been identified as key challenges affecting the mental and sexual health of young pregnant women and th...

Tamora A. Callands, Kandyce Hylick, Alethea Desrosiers, Shantesica M. Gilliam, Erica N. Taylor, Josalin J. Hunter and Nathan B. Hansen

BMC Pregnancy and Childbirth 2023 23:196

Research | Published on: 20 March 2023

> Full Text> PDF

Lessons from digital technology-enabled health interventions implemented during the coronavirus pandemic to improve maternal and birth outcomes: a global scoping review

Timely access to essential obstetric and gynecologic healthcare is an effective method for improving maternal and neonatal outcomes; however, the COVID-19 pandemic impacted pregnancy care globally. In this glo...

Imelda K. Moise, Nicole Ivanova, Cyril Wilson, Sigmond Wilson, Hikabasa Halwindi and Vera M. Spika

BMC Pregnancy and Childbirth 2023 23:195

Research article | Published on: 20 March 2023

Using household survey data to explore the effects of the domiciliary environment on weight at birth: a multilevel mixed-effects analysis of the 2016 Ethiopian Demographic Health Survey

Low birth weight (LBW) is associated with infant mortality and postpartum health complications. In previous studies, overall LBW has been found to be significantly associated with several sociodemographic fact...

Aiggan Tamene, Aklilu Habte, Mihretu Tagesse, Zablon Wale Sewalem and Abel Afework

BMC Pregnancy and Childbirth 2023 23:194

Research | Published on: 20 March 2023

> Full Text> PDF

Trophoblast retrieval from the cervical canal to predict abnormal pregnancy early in gestation: a pilot study

The current detection of fetal chromosomal abnormalities by non-invasive prenatal testing (NIPT) mainly relies on the cell free DNA(cfDNA) in the maternal blood. However, a gestational age of less than 12 week...

Xiaoke Yang, Liuyezi Du, Yue Li, Lin Liang, Linlin Ma and Shaowei Wang

BMC Pregnancy and Childbirth 2023 23:193

Research | Published on: 18 March 2023

Non-adherence to WHO's recommended 8-contact model: geospatial analysis of the 2017 Maternal Health Survey

Evidence shows that most women in Ghana do not meet the minimum 8-contact model for antenatal care as recommended by WHO with only 31.2%-41.9% of them meeting the recommendation. To the best of our knowledge, ...

Kwamena Sekyi Dickson, Ebenezer N. K. Boateng, Kenneth Setorwu Adde, Edward Kwabena Ameyaw and Michelle L. Munro-Kramer

BMC Pregnancy and Childbirth 2023 23:192

Research | Published on: 18 March 2023

> Full Text> PDF

Self-management, care needs and clinical management of primiparous mothers during early labour – a qualitative content analysis

Childbearing women face the problem of managing spontaneous onset of labour without professional support. It is their responsibility to diagnose and react to early labour and subsequently recognise the right t...

Antonia N. Mueller and Susanne Grylka-Baeschlin

BMC Pregnancy and Childbirth 2023 23:191

Research article | Published on: 18 March 2023

Analysis of the clinical outcomes of microbial contamination caused by environmental contamination of the embryology laboratory during IVF-ET treatment cycles

Bacterial contamination may cause loss of or damage to cultured oocytes or embryos, resulting in the lack of transplantable embryos during IVF embryo culture. However, there are few reports about IVF embryo co...

Tingting Zheng, Qinhua Li, Ningjing Chen, Peiyao Du and Hong Ye

BMC Pregnancy and Childbirth 2023 23:190

Research | Published on: 18 March 2023

> Full Text> PDF

A novel clinical prediction model of severity based on red cell distribution width, neutrophil-lymphocyte ratio and intra-abdominal pressure in acute pancreatitis in pregnancy

Acute pancreatitis in pregnancy (APIP) with a high risk of death is extremely harmful to mother and fetus. There are few models specifically designed to assess the severity of APIP. Our study aimed to establis...

Wenyan Liao, Guangwei Tao, Guodong Chen, Jun He, Chunfen Yang, Xiaohua Lei, Shuo Qi, Jiafeng Hou, Yi Xie, Can Feng, Xinmiao Jiang, Xin Deng and Chengming Ding

BMC Pregnancy and Childbirth 2023 23:189

Research | Published on: 18 March 2023

Effects of maternal pre-pregnancy body mass index and gestational weight gain on antenatal mental disorders in China: a prospective study

Maternal obesity is the most common medical condition among women of reproductive age worldwide. The pre-pregnancy body mass index and gestational weight gain have been suggested to be associated with maternal...

Xuan Zhou, Lin Rao, Dongjian Yang, Tong Wang, Hong Li and Zhiwei Liu

BMC Pregnancy and Childbirth 2023 23:188

Research | Published on: 18 March 2023

> Full Text> PDF

The experience of pregnant women in contexts of vulnerability of prenatal primary nursing care: a descriptive interpretative qualitative study

Prenatal primary nursing care contributes to improving the health outcomes of mothers and unborn babies. Some pregnant women in contexts of vulnerability experience prenatal nursing care in a positive way, whi...

Émilie Hudon, Maud-Christine Chouinard, Édith Ellefsen, Jérémie Beaudin and Catherine Hudon

BMC Pregnancy and Childbirth 2023 23:187

Research | Published on: 18 March 2023

A new continuous glucose monitor for the diagnosis of gestational diabetes mellitus: a pilot study

Gestational Diabetes Mellitus (GDM) incidence and adverse outcomes have increased globally. The validity of the oral glucose tolerance test (OGTT) for GDM diagnosis has long been questioned, with no suitable s...

Daria Di Filippo, Amanda Henry, Chloe Bell, Sarah Haynes, Melissa Han Yiin Chang, Justine Darling and Alec Welsh

BMC Pregnancy and Childbirth 2023 23:186

Research | Published on: 18 March 2023

> Full Text> PDF

Determinants of health facility delivery among young mothers aged 15 – 24 years in Nigeria: a multilevel analysis of the 2018 Nigeria demographic and health survey

Young mothers aged 15 to 24 years are particularly at higher risk of adverse health outcomes during childbirth. Delivery in health facilities by skilled birth attendants can help reduce this risk and lower mat...

Tope Olubodun, Semiu Adebayo Rahman, Oluwakemi Ololade Odukoya, Ifeoma P. Okafor and Mobolanle Rasheedat Balogun

BMC Pregnancy and Childbirth 2023 23:185

Research | Published on: 17 March 2023

Self-management system for postpartum women with hypertension disorders: an eHealth application intervention study

Hypertension disorders are relatively common in pregnant women and often persist in the postpartum period. Few studies are available regarding the self-management of postpartum hypertension via the eHealth sys...

Chung-Wei Chang, Yi-Jing Tsai, Yu-Yun Hsu and Ting-Wei Hou

BMC Pregnancy and Childbirth 2023 23:184

Research | Published on: 16 March 2023

> Full Text> PDF

Patient perspectives on noninvasive prenatal testing among black women in the United States: a scoping review

Advances in reproductive health technologies such as noninvasive prenatal testing (NIPT) are changing the landscape of prenatal care and maternal health. NIPT, made clinically available in the United States (U...

Shameka P. Thomas, Madison A. Keller, Tiara Ranson and Rachele E. Willard

BMC Pregnancy and Childbirth 2023 23:183

Research | Published on: 16 March 2023

Comparison of the incidence of fetal prolonged deceleration after induction of labor analgesia between dural puncture epidural and combined spinal epidural technique: a pilot study

Abnormal cardiotocogram (CTG) tracing may appear after induction of neuraxial labor analgesia. Non-reassuring fetal status (NRFS) indicated by severely abnormal tracings, such as prolonged deceleration (PD) or...

Shoko Okahara, Rie Inoue, Yumi Katakura, Hitomi Nagao, Saori Yamamoto, Shuko Nojiri, Jun Takeda, Atsuo Itakura and Hiroyuki Sumikura

BMC Pregnancy and Childbirth 2023 23:182

Research | Published on: 16 March 2023

> Full Text> PDF

A simulation and experiential learning intervention for labor and delivery providers to address HIV stigma during childbirth in Tanzania: study protocol for the evaluation of the MAMA intervention

The experience of HIV stigma during intrapartum care can impact women's trust in the health care system and undermine their long-term commitment to HIV care engagement. Delivery of respectful maternity care (R...

Melissa H. Watt, Linda M. Minja, Mariam Barabara, Pendo Mlay, Maya J. Stephens, Gaudensia Olomi, Janeth Mlay, Virginie Marchand, Blandina T. Mmbaga, Olivia R. Hanson and Susanna R. Cohen

BMC Pregnancy and Childbirth 2023 23:181

Study Protocol | Published on: 16 March 2023

Closed-loop insulin delivery in pregnant women with type 1 diabetes (CRISTAL): a multicentre randomized controlled trial – study protocol

Despite increasing use of continuous glucose monitoring (CGM) and continuous subcutaneous insulin infusion (CSII, insulin pumps) in type 1 diabetes (T1D) in pregnancy, achieving recommended pregnancy glycaemic...

Kaat Beunen, Nancy Van Wilder, Dominique Ballaux, Gerd Vanhaverbeke, Youri Taes, Xavier-Philippe Aers, Frank Nobels, Joke Marlier, Dahae Lee, Joke Cuypers, Vanessa Preumont, Sarah E. Siegelaar, Rebecca C. Painter, Annouschka Laenen, Pieter Gillard, Chantal Mathieu...

BMC Pregnancy and Childbirth 2023 23:180

Study Protocol | Published on: 16 March 2023

> Full Text> PDF

Exploring the antenatal care challenges faced during the COVID-19 pandemic in rural areas of Indonesia: a qualitative study

The COVID-19 pandemic affected almost all healthcare services in Indonesia, including antenatal care (ANC). Pregnant women were a vulnerable group during the pandemic since the Indonesian government's policies...

Mekar Dwi Anggraeni, Rahmi Setiyani, Endang Triyanto, Asep Iskandar, Desiyani Nani and Amin Fatoni

BMC Pregnancy and Childbirth 2023 23:179

Research | Published on: 16 March 2023

Fatherhood and high-risk pregnancy: a scoping review

The experience of expectant parenthood is commensurate of relative angst and nervousness albeit one of overall excitement and joy. However, when the pregnancy is regarded as high-risk, this experience changes ...

Kyle Jackson, Erika Erasmus and Ntobizodwa Mabanga

BMC Pregnancy and Childbirth 2023 23:168

Research | Published on: 16 March 2023

> Full Text> PDF

Application value of Early-Follicular Phase Long-Acting Gonadotropin-Releasing Hormone Agonist Long Protocol in patients with resistant ovary syndrome

Resistant ovarian syndrome(ROS) is a rare disease. It is difficult to diagnose and treat. Most of the literature reports on assisted pregnancy treatment for ROS patients are individual case reports. In this pa...

Fan Zhang, Huixia Zhang, Hui Du, Xin Li, Haixia Jin and Gang Li

BMC Pregnancy and Childbirth 2023 23:178

Research article | Published on: 15 March 2023

Vitamin D in pregnancy (GRAVITD) – a randomised controlled trial identifying associations and mechanisms linking maternal Vitamin D deficiency to placental dysfunction and adverse pregnancy outcomes – study protocol

The prevalence of vitamin D deficiency is high among pregnant women. Vitamin D deficiency in pregnancy is associated with increased risk of adverse pregnancy outcomes especially complications related to placen...

Anna Louise Vestergaard, Martin Christensen, Mette Findal Andreasen, Agnete Larsen and Pinar Bor

BMC Pregnancy and Childbirth 2023 23:177

Study protocol | Published on: 15 March 2023

> Full Text> PDF

Overall childbirth experience: what does it mean? A comparison between an overall childbirth experience rating and the Childbirth Experience Questionnaire 2

In clinical settings and research studies, childbirth experience is often measured using a single-item question about overall experience. Little is known about what women include in this rating, which complica...

Frida Viirman, Susanne Hesselman, Inger Sundström Poromaa, Agneta Skoog Svanberg and Anna Wikman

BMC Pregnancy and Childbirth 2023 23:176

Research | Published on: 14 March 2023

Testing effects of partner support and use of oral contraception during relationship formation on severity of nausea and vomiting in pregnancy

A recent study focusing on dietary predictors of nausea and vomiting in pregnancy (NVP) found that women with higher levels of partner support, and those who had used oral contraception (OC) when they met the ...

Kateřina Roberts, Jan Havlíček, Šárka Kaňková, Kateřina Klapilová and S. Craig Roberts

BMC Pregnancy and Childbirth 2023 23:175

Research article | Published on: 14 March 2023

> Full Text> PDF

Additional single dose GnRH agonist during luteal phase support may improve live birth rate in GnRHa-HRT frozen-thawed embryo transfer cycle: a retrospective cohort study

GnRH agonist (GnRHa) has been reported to have direct effects and functional roles in the endometrium and embryos. Several meta-analyses have shown that GnRHa administration in the luteal phase improved the li...

Wei-Shan Chang, Pei-Hsuan Lin, Chia-Jung Li, Chyi-Uei Chern, Yu-Chen Chen, Li-Te Lin and Kuan-Hao Tsui

BMC Pregnancy and Childbirth 2023 23:174

Research | Published on: 14 March 2023

The effect of perceived spousal support on childbirth self-efficacy on pregnant women in turkey

Spousal support during pregnancy reduces the anxiety and stress of the pregnant women and increases the ability to cope with the problems experienced by the pregnant women. The study aimed to examine the effec...

Burcu Kucukkaya and Sukran Basgol

BMC Pregnancy and Childbirth 2023 23:173

Research | Published on: 14 March 2023

> Full Text> PDF

Data collection systems for active safety surveillance of vaccines during pregnancy in low- and middle-income countries: developing and piloting an assessment tool (VPASS)

There is an urgent need for active safety surveillance to monitor vaccine exposure during pregnancy in low- and middle-income countries (LMICs). Existing maternal, newborn, and child health (MNCH) data collect...

Maria Belizán, Federico Rodriguez Cairoli, Agustina Mazzoni, Erin Goucher, Sabra Zaraa, Sarah Matthews, Verónica Pingray, Andy Stergachis, Xu Xiong, Mabel Berrueta and Pierre Buekens

BMC Pregnancy and Childbirth 2023 23:172

Research | Published on: 13 March 2023

The assessment in patients with acute fatty liver of pregnancy (AFLP) treated with plasma exchange: a cohort study of 298 patients

To assess the prevalence, risk factors, clinical characteristics of Acute fatty liver of pregnancy (AFLP) patients, and outcomes of AFLP patients treated with plasma exchange (PE).

Lingxia Li, Dengchao Huang, Jing Xu, Miaojing Li, Juan Zhao, Qindong Shi and Qinyue Guo

BMC Pregnancy and Childbirth 2023 23:171

Research | Published on: 13 March 2023

> Full Text> PDF

Live birth in woman with premature ovarian insufficiency and 46, XY karyotype after chemotherapy and bone marrow transplant: a case report

Premature ovarian insufficiency (POI) is a clinical syndrome defined by loss of ovarian function before the age of 40 years, characterized by elevated serum gonadotropin levels and decreased estrogen levels wi...

Yaojia Zhang, Haiyan Wang and Xiaoqin Pan

BMC Pregnancy and Childbirth 2023 23:170

Case report | Published on: 13 March 2023

Environmental factors influencing women's childbirth experiences in labor–delivery–recovery–postpartum unit: a qualitative cross-sectional study

Women's experiences of birth environment influence their mental health and that of their families. Identifying women's childbirth experiences in the labor-delivery-recovery-postpartum unit (LDRP) unit can help...

Ashraf Kazemi, Marjan Beigi and Hajar Enteshary Najafabadi

BMC Pregnancy and Childbirth 2023 23:169

Research | Published on: 13 March 2023

> Full Text> PDF

Implementation of a women's reproductive behavioral health telemedicine program: a qualitative study of barriers and facilitators in obstetric and pediatric clinics

Perinatal Mood and Anxiety Disorders and Substance Use Disorders are common and result in significant morbidities and mortality. Despite evidence-based treatment availability, multiple barriers exist to care d...

Katherine R. Sterba, Emily E. Johnson, Edie Douglas, Rubin Aujla, Lisa Boyars, Ryan Kruis, Rebecca Verdin, Rachel Grater, Kathryn King, Dee Ford and Constance Guille

BMC Pregnancy and Childbirth 2023 23:167

Research | Published on: 11 March 2023

Perceptions on acceptability of the 2016 WHO ANC model among the pregnant women in Phalombe District, Malawi – a qualitative study using Theoretical Framework of Acceptability

The World Health Organization introduced a new model of care, 'The 2016 WHO ANC Model' to overcome challenges encountered during the implementation of the Focused Antenatal Care Approach. For any new intervent...

Prince Nyumwa, Agatha Kapatuka Bula and Alinane Linda Nyondo-Mipando

BMC Pregnancy and Childbirth 2023 23:166

Research | Published on: 11 March 2023

> Full Text> PDF

Body perception in pregnant women: a qualitative study

Dramatic body changes in pregnancy cause severe concerns among pregnant women about their appearance. Therefore, this study aimed to explore body perception in pregnant women.

Zahra Sohrabi, Ashraf Kazemi, Ziba Farajzadegan and Mojgan Janighorban

BMC Pregnancy and Childbirth 2023 23:165

Research | Published on: 11 March 2023

Clinical guidelines for the management of weight during pregnancy: a qualitative evidence synthesis of practice recommendations across NHS Trusts in England

Women who enter pregnancy with a Body Mass Index above 30 kg/m² face an increased risk of complications during pregnancy and birth. National and local practice recommendations in the UK exist to guide healthcare ...

Lucy Goddard, Nerys M. Astbury, Richard J. McManus, Katherine Tucker and Jennifer MacLellan

BMC Pregnancy and Childbirth 2023 23:164

Research | Published on: 11 March 2023

> Full Text> PDF

Does benefits-of-breastfeeding language or risks-of-formula-feeding language promote more-positive attitudes toward breastfeeding among midwives and nurses?

Midwives and nurses are crucial in breastfeeding support. Few studies have explored appropriate language for nursing education on breastfeeding. We assessed the impact of the language used on breastfeeding att...

Ayumi Toda, Keiko Nanishi and Akira Shibanuma

BMC Pregnancy and Childbirth 2023 23:163

Research | Published on: 11 March 2023

Adverse neonatal outcome and veno-arterial differences in umbilical cord blood pH (Δ pH) at birth: a population-based study of 108,629 newborns

Umbilical cord blood gases are routinely used by midwives and obstetricians for quality assurance of birth management and in clinical research. They can form the basis for solving medicolegal issues in the ide...

Tiia-Marie Sundberg, Nana Wiberg, Karin Källén and Mehreen Zaigham

BMC Pregnancy and Childbirth 2023 23:162

Research | Published on: 11 March 2023

> Full Text> PDF

Complication experience during pregnancy and place of delivery among pregnant women: a cross-sectional study

Unlike other causes such as abortion, obstetric complications like hemorrhage, and hypertensive disorders of pregnancy, which are difficult to resolve for women who give birth out of health facilities are pers...

Bekelu Teka Worku, Misra Abdulahi, Merertu Tsega, Birtukan Edilu, Rahma Ali, Mahilet Berhanu Habte, Samira Awel, Masrie Getnet, Yabsira Melaku, Radiet Kidane, Abonesh Taye and Meseret Tamirat

BMC Pregnancy and Childbirth 2023 23:161

Research | Published on: 11 March 2023

Association of delivery mode and number of pregnancies with anorectal manometry data in patients with postpartum constipation

To explore the association of delivery mode and the number of pregnancies with anorectal manometry data in patients with postpartum constipation.

Yan Yin, Yumin Zhang and Cheng Qian

BMC Pregnancy and Childbirth 2023 23:160

Research | Published on: 11 March 2023

> Full Text> PDF

Healthcare providers perceptions regarding the presence of Birth Companion during childbirth at a tertiary care hospital in India

Despite an increase in institutional births and a fall in maternal mortality, the satisfaction of women with their birthing experience in public health institutions is low. Birth Companion (BC) is an important...

Tanvi Sarwal, Yamini Sarwal, Shakun Tyagi and Rakesh Sarwal

BMC Pregnancy and Childbirth 2023 23:159

Research | Published on: 10 March 2023

Association between very advanced maternal age women with gestational diabetes mellitus and the risks of adverse infant outcomes: a cohort study from the NVSS 2014–2019

To evaluate the association between gestational diabetes mellitus (GDM) and infant outcomes in women of very advanced maternal age (vAMA) (≥45 years).

Lin Lu, Lidan He, Jifen Hu and Jianhua Li

BMC Pregnancy and Childbirth 2023 23:158

Research | Published on: 10 March 2023

> Full Text> PDF

A feasible method to evaluate fetal palate: sequential sector-scan through oral fissure

Prenatal diagnosis of cleft palate is still challenging. To describe a practical and efficient method, sequential sector-scan through oral fissure (SSTOF), to evaluate palate.

Xiaofeng Lu, Yi Zhou, Min Fan, Wanyan Li and Chaoxue Zhang

BMC Pregnancy and Childbirth 2023 23:157

Research | Published on: 9 March 2023

Application of machine learning to identify risk factors of birth asphyxia

Developing a prediction model that incorporates several risk factors and accurately calculates the overall risk of birth asphyxia is necessary. The present study used a machine learning model to predict birth ...

Fatemeh Darsareh, Amene Ranjbar, Mohammadsadegh Vahidi Farashah, Vahid Mehrnoush, Mitra Shekari and Malihe Shirzadfard Jahromi

BMC Pregnancy and Childbirth 2023 23:156

Research | Published on: 8 March 2023

> Full Text> PDF

Foetal death due to extensive extra-abdominal umbilical vein Varix with umbilical vein thrombosis: a case report

Foetal anaemia and umbilical vein thrombosis are rare pregnancy complications that can increase the risk of perinatal adverse events, which, in severe cases, can lead to foetal death. During pregnancy, umbilic...

Qing-Yun Song and Ying Tang

BMC Pregnancy and Childbirth 2023 23:155

Case report | Published on: 8 March 2023

Spontaneous skull base cerebrospinal fluid leak during pregnancy: a case report and review of the literature

Idiopathic intracranial hypertension can lead to dural defects and spontaneous leakage of cerebrospinal fluid (CSF) from the skull base. Skull base CSF leaks are rarely reported in pregnancy but pose unique ch...

Lauren Michelle, Rebecca J. Post, Edward C. Kuan and Michael P. Nageotte

BMC Pregnancy and Childbirth 2023 23:154

Case report | Published on: 8 March 2023

> Full Text> PDF

Pre-pregnancy complications - associated factors and wellbeing in early pregnancy: a Swedish cohort study

Many couples experience difficulties to become pregnant or carry a pregnancy to term due to unknown causes. Here we define pre-pregnancy complications as having prior recurrent pregnancy loss, prior late misca...

Unnur Gudnadottir, Juan Du, Luisa W. Hugerth, Lars Engstrand, Ina Schuppe-Koistinen, Eva Wiberg Itzel, Emma Fransson and Nele Brusselaers

BMC Pregnancy and Childbirth 2023 23:153

Research | Published on: 8 March 2023



Awful Bad COK Good Great Send feedback

How was your experience today?



Submit manuscript

Important information

Editorial board

For authors

For editorial board members

For reviewers

Manuscript editing services

Contact Us

Annual Journal Metrics

Citation Impact

3.105 - 2-year Impact Factor (2021)

3.853 - <u>5-year Impact Factor</u> (2021)

1.518 - SNIP (Source Normalized Impact per Paper)

0.960 - SJR (SCImago Journal Rank)

Speed

41 days to first decision for all manuscripts (Median)

64 days to first decision for reviewed manuscripts only (Median)

Usage

7,560,102 Downloads (2021) 4,644 Altmetric mentions (2021)

More about our metrics

Peer-review Terminology

The following summary describes the peer review process for this journal:

Identity transparency: Single anonymized

Reviewer interacts with: Editor

Review information published: Review reports. Reviewer Identities reviewer opt in. Author/reviewer

communication

More information is available here

Follow



Sign up for article alerts and news from this journal

BMC Pregnancy and Childbirth

ISSN: 1471-2393

Contact us

Submission enquiries: bmcpregnancyandchildbirth@biomedcentral.com

General enquiries: ORSupport@springernature.com

Read more on our blogs Policies Support and Contact

Receive BMC newsletters Accessibility Leave feedback

Manage article alerts Press center Careers

Language editing for authors

Scientific editing for authors

Follow BMC







By using this website, you agree to our <u>Terms and Conditions</u>, <u>California Privacy Statement</u>, <u>Privacy statement</u> and <u>Cookies</u> policy. <u>Manage cookies/Do not sell my data</u> we use in the preference centre.

SPRINGER NATURE

© 2023 BioMed Central Ltd unless otherwise stated. Part of <u>Springer Nature</u>.

also developed by scimago:





Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

Home

Journal Rankings

Country Rankings

Viz Tools

Help

About Us



Research Paper Editing

A Nature Research service. Expert scientific English ed

Nature Research Editing

0

BMC Pregnancy and Childbirth 3

COUNTRY SUBJECT AREA AND CATEGORY

PUBLISHER

H-INDEX

United Kingdom

Universities and research institutions in United Kingdom

Media Ranking in United

Medicine

Obstetrics and Gynecology

BioMed Central Ltd.

92

PUBLICATION TYPE

Kingdom

ISSN

COVERAGE

INFORMATION

Journals

14712393

2001-2021


Homepage

How to publish in this journal

sparhandy=



Plan P Open Science - Transform Peer Review






health, maternity care, trends and sociological aspects of pregnancy and childbirth.

Q Join the conversation about this journal



BMC Pregnancy and Childbirth



Metrics based on Scopus® data as of April 2022

Rawan A Rahim Alsayed 2 years ago

hello

I have a case report

title: Spontaneous pregnancy and live birth to a patient with premature ovarian failure I want to publish it in your journal

what are the requirements?

is it any fees?

thanks

reply



Melanie Ortiz 2 years ago

Dear Rawan,

thank you for contacting us.

We are sorry to tell you that SCImago Journal & Country Rank is not a journal. SJR is a portal with scientometric indicators of journals indexed in Elsevier/Scopus.

Unfortunately, we cannot help you with your request, we suggest you visit the journal's homepage (See submission/author guidelines) or contact the journal's editorial staff, so they could inform you more deeply.

Best Regards, SCImago Team

T tadele 3 years ago

your posts are very important.

reply

SCImago Team

Tatyana 4 years ago

Hello! I plan to publish a clinical case. Is this possible in your journal?

reply



Melanie Ortiz 4 years ago

SCImago Team

Dear Tatyana,

thank you for contacting us.

Sorry to tell you that SCImago Journal & Country Rank is not a journal. SJR is a portal with scientometric indicators of journals indexed in Elsevier/Scopus. For the publication of an article you must submit your manuscript to a journal.

SJ&CR is an Open Access tool. We suggest you use the SJ&CR search options to check if a journal is indexed or not. We make public all the information we have sent us Scopus and if there are journals indexed in Scopus that do not appear in the SJ&CR is because we have not received the data from Elsevier/Scopus.

http://www.scimagojr.com/

Greetings from Spain and thank you for using the SCImago products, SCImago Team

Leave a comment

Name

Email

(will not be published)

l'm not a robot
reCAPTCHA
Privacy - Terms

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular

articles, maintain the dialogue through the usual channels with your editor

Developed by:

Powered by:





Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2022. Data Source: Scopus®

EST MODUS IN REBUS Horatio (Satire 1,1,106)

Cookie settings

Cookie policy

RESEARCH **Open Access**



Exploring the antenatal care challenges faced during the COVID-19 pandemic in rural areas of Indonesia: a qualitative study

Mekar Dwi Anggraeni^{1*}, Rahmi Setiyani¹, Endang Triyanto¹, Asep Iskandar¹, Desiyani Nani¹ and Amin Fatoni²

Abstract

Introduction The COVID-19 pandemic affected almost all healthcare services in Indonesia, including antenatal care (ANC). Pregnant women were a vulnerable group during the pandemic since the Indonesian government's policies at the time influenced the delivery of ANC services, particularly in rural areas. Investigating the ANC challenges faced during the pandemic from the perspectives of pregnant women and healthcare providers is important for our understanding of ANC provision. This study, therefore explores barriers to ANC appointments faced during the COVID-19 pandemic in rural areas of Indonesia from the perspectives of pregnant women and health care providers.

Methods This was a qualitative exploratory descriptive study involving 31 participants, consisting of 25 pregnant women and six healthcare providers who were selected via a purposive sampling method. Thadeus and Maine's Three Delays Model was used as the theoretical framework. Data were collected between March and August 2021, through two focus group discussions (FGDs), ten in-depth interviews, and field notes. Data were analyzed using a thematic analysis method.

Results Three themes describing barriers to ANC during the COVID-19 pandemic in rural areas of Indonesia emerged from this study. Those themes were: (1) The fear of being infected with COVID-19, related to anxiety, perceived vulnerability, and the desire to protect oneself and loved ones; (2) The stay-at-home policy, related to transport barriers and restricted social activity; and (3) Re-designed ANC services, related to ANC adjustments, high-risk pregnancies, insufficient information, and adherence to COVID-19 preventive behaviors.

Conclusion Based on the Three Delays Model, several challenges to carrying out ANC during the COVID-19 pandemic in rural areas of Indonesia were identified. These findings demonstrate the need to formulate and implement ANC packages to facilitate pregnant women's access to health care services.

Keywords Antenatal care, COVID-19, Indonesia, Pregnancy, Rural areas

*Correspondence: Mekar Dwi Anggraeni mekar.anggraeni@unsoed.ac.id ¹Nursing Department, Faculty of Health Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia

²Chemistry Department, Faculty of Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia



Introduction

In 2020, the COVID-19 pandemic was declared an international public health problem [1]. The virus spread massively to all regions of Southeast Asia, with the total number of confirmed cases and deaths in Indonesia up to September 3rd 2021 reaching 4,116,890 and 134,930 respectively [2]. These figures meant that Indonesia ranked 13 globally in terms of total COVID-19 confirmed cases [3]. The pandemic affected all aspects of life in Indonesia, including routine health services such as antenatal care (ANC). Pregnant women were one of the most vulnerable groups during the pandemic [4], and, despite the disruption, they were still encouraged to engage in ANC during the period because of the massive health benefits associated with ANC. Despite this, ANC coverage in Indonesia has decreased over the past two years. Antenatal care coverage among pregnant Indonesian women was 88%, 88.5%, and 84.6% in 2018, 2019, and 2020 respectively [5–7]. The noticeable decline in ANC coverage in 2020 might be due to the pandemic. The Indonesian government reported the first case in Indonesia on March 2nd, 2019. Following that, the Indonesian Obstetrics and Gynecologists Association announced that 20% of MMR cases in the country between the beginning of 2020 and August 2021 were related to COVID-19 [8]. A similar study in Brazil found that 13.19% of maternal deaths in 2020 were linked to COVID-19 [9].

ANC is essential care that should be provided to all pregnant women. The main aim of ANC is to ensure that every pregnancy ends in the delivery of a healthy infant without any undesirable effects on the health of women, and this is achieved through health promotion, disease prevention, early detection, and management of complications and existing diseases [10]. There is less ANC coverage for pregnant women living in developing countries than those in developed countries, particularly in rural areas that lack healthcare providers and facilities [11] Irregular ANC is associated with complications during pregnancy, delivery, and puerperium for both mother and infant, including preeclampsia, eclampsia, anemia, preterm birth, low birth weight, and stillbirth [12]. Occasionally, pregnancy triggers diseases such as pregnancy-induced hypertension, gestational diabetes, pre-eclampsia, and eclampsia [13].

Although most maternal deaths linked to COVID-19 have occurred in pregnant women with comorbidities, previous studies have confirmed that COVID-19 in pregnant women may cause severe outcomes such as abortion [14] and increased maternal morbidity and mortality, preeclampsia, and preterm birth [15]. One study carried out in India found that during the pandemic, the number of deliveries taking place in institutional hospitals reduced by 45.1%, while high-risk pregnancies increased by 7.2%, admission to intensive care units saw a 2.5-fold increase,

and one-third of pregnant women experienced inadequate ANC. The authors stated that the principal reasons for delayed health-seeking behaviors were lockdown and fear of transmission, and this resulted in 44.7% of pregnancies developing complications during that period [16]. Another study found that access to and continuity of ANC, particularly with regard to the early detection of COVID-19 symptoms, is highly recommended [17]. This could be because pregnant women tend to be relatively immunodeficient which could worsen an infection and lead to bad outcomes for mothers and fetuses [18]. It is clear to see that ANC is hugely important among pregnant women.

Attending ANC appointments was more challenging during the COVID-19 pandemic worldwide. One study carried out in India found that of 144 pregnant women, two thirds reported at least one barrier to ANC during the pandemic, with one third reporting fear, and more than three quarters refused ultrasound assessments and blood tests [19]. Another study identified the main barriers to ANC during the pandemic as being health facility-related barriers, poor quality of care, government regulations, anxiety, and fear of infection [20]. Another study added closing hospital/healthcare centers, the involvement of private health sectors, lack of transport, and fear of transmission from health care providers to be the major barriers to ANC during the pandemic [21]. The authors of one study cited lack of social support from husband, restricted face-to face interaction with health professionals, lack of information on COVID-19, and high uncertainty about pregnancy and birth during the pandemic as being the main ANC barriers [22]. Psychosocial challenges such as social isolation, lack of information about neonatal care, lack of public transport, and lack of up-to-date information related to COVID-19, particularly among women in low-and middle income countries were cited as the main factors in another study [23]. Pregnant women from low socio-economic classes with pre-existing pregnancy complications were more likely to experience barriers to ANC [19]. To date, little is known about the ANC challenges faced by Indonesian women during the pandemic, particularly those in rural areas who already faced a lot of barriers to ANC. Since ANC challenges during the COVID-19 pandemic in rural areas of Indonesia have not yet been sufficiently explored from the perspective of pregnant women and their healthcare providers, the knowledge provided in this study could be used to develop an appropriate and safe ANC program for pregnant women living in rural areas.

Methods

Aim

The aim of this study was to explore the ANC challenges during the COVID-19 pandemic in rural areas of Indonesia.

Study design

A qualitative exploratory descriptive design was applied since it was deemed the best way of answering the who, what, and where of experiences and of obtaining insights from participants about an inadequately understood phenomenon [24, 25]. The qualitative method emphasizes deep understanding, complexity, and details of the phenomena under study, and the researcher was actively involved in the research process. The Three Delays Model developed by Thaddeus and Maine was used as the framework for this study [26].

Setting

Indonesia is the most populous nation in Southeast Asia. It also has one of the highest MMR rates in the region [27], with around 305 MMR incidents occurring per 100,000 live births [28]. Indonesia is divided into 38 provinces and is home to over 300 ethnic groups. The biggest ethnicity in Indonesia is Javanese which accounts for around 41.71% of the total population, the majority of whom live in rural areas [29]. The third highest number of MMR cases in the country are found in Central Java Province. Data were collected in Banyumas District, the area with the sixth highest rate of MMR cases in the province. There are 17 hospitals and 30 primary health centers (PHCs) in Banyumas District, however, ANC coverage there remains lower than the national average. The district is one of the lowest 10 districts for ANC coverage in Central Java Province [30]. The study was conducted between April and August 2021. Most pregnant women attend ANC in a primary health center according to Ministry of Health policies, but high-risk pregnancies may be referred to secondary or tertiary hospitals, depending on the severity of their pregnancy-related problems.

Participants

Included in this study were pregnant women and health-care providers living or working in rural areas. A purposive sampling technique was used to select the study participants. The inclusion criteria for pregnant women were those in their second or third trimester who had attended ANC at least three times in a PHC. Participants were recruited through the PHC list of pregnant women and all those meeting the inclusion criteria who agreed to participate in the study were invited for data collection. Women whose pregnancy was deemed high-risk were excluded from the study. The inclusion criteria for healthcare providers were those who worked in a PHC

providing ANC that had had at least three years' experience. Healthcare providers were selected purposively and sent a formal invitation to participate in the study by post. A total of 31 participants including 25 pregnant women and six healthcare providers agreed to participate in the study and were invited to attend focus group discussions (FGDs) and in-depth interviews.

Data collection

Data collection only began after the researchers received approval from the Institutional Review Board (IRB) of the Faculty of Medicine, Universitas Jenderal Soedirman. Data were collected by the researchers, all of whom are faculty members of the nursing department in the Faculty of Health Sciences, Universitas Jenderal Soedirman. Most researchers were of Javanese ethnicity and familiar with the principles of qualitative research and interview guidelines. Prospective participants were selected with consideration given to their social, economic, and demographic characteristic, including age, level of education, employment status, and parity for pregnant women, and length of work experience for healthcare providers, since a greater diversity of study participants allows for a more in-depth exploration and analysis of a phenomena. To adhere the Declaration of Helsinki for studies involving human subjects and ethical issues, each participant was informed of the study's purpose, procedure, and risks, and told that participation was voluntary and that they could quit at any time without facing any consequences. Pregnant women choosing to withdraw from the study knew that they would continue to receive ANC in their PHCs as usual. The researchers also guaranteed confidentiality and anonymity of participants' information.

Two FGDs were conducted for the purpose of this study. Twelve pregnant women participated in first FGD and nine pregnant women took part in the second. The study was conducted between March and August 2021. Because the COVID-19 situation worsened in Indonesia in July 2021, it was not possible to conduct another FGD and so the data collection method switched to in-depth interviews conducted through online interviews. At this point, the researchers conducted ten in-depth interviews to four pregnant women and six healthcare providers using video calls. The researchers initially met each participant in the PHC. There they introduced themselves, informed the participants of the study's aims, and shared contact information. The researchers then contacted each proposed participant via WhatsApp to gain their permission to participate in the study. Each participant was provided with an electronic informed consent form before starting the interviews, and any participant struggling to fill out the electronic version was provided with a hard copy. Researchers verbally told all participants that the interviews would be recorded and transcribed.

Information related to demographic characteristics was collected before the FGDs began. The FGDs were conducted in the Bahasa Indonesia language using interview guidelines by a researcher who was an expert in community health nursing and familiar with the FGD process. The researcher leading the FGDs holds a doctoral degree in community health and has experience in carrying out qualitative research. The first FGD was conducted in the meeting room at the PHC and the second FGD was conducted in the meeting room of the Sub-District Government Office. The in-depth interviews were conducted by a researcher holding a magister degree in community health and has experience in carrying out qualitative research. The researchers introduced themselves as faculty members in Universitas Jenderal Soedirman and let participants know they were free to talk with them.

All discussions and interviews were audio recorded for verbatim transcription. During the FGDs, researchers encouraged participants to share their knowledge, feelings, and experiences regarding ANC coverage during the COVID-19 pandemic, and another researcher observed and took notes to supplement the audio recorded FGDs. The observations were used to confirm the results from the FGDs and in-depth interviews to enrich the data. Only researchers and participants attended the FGD sessions to ensure confidentiality and privacy.

In a qualitative study, data saturation is reached when extending the interviews delivers no new data and all founded codes are recurrent [24]. In this study, saturation was achieved with the 31st participant. At this point, the researchers ended the data collection, meaning there were 31 participants were involved in this study, ten of whom met on an individual basis and the others of whom attended the two focus groups. Each interview and FGD lasted around 60 and 90 min respectively. No repeat interviews were carried out and two prospective participants declined to participate, citing busy schedules.

Research instrument

Data were collected using a semi-structured interview guideline that was developed based on the literature review on ANC in the COVID-19 pandemic situation. A panel of experts consisting of a maternity nurse and two lecturers from the maternity department reviewed the interview guidelines for their relevance to the study's purpose. The interview guidelines were also verified and pilot tested on five pregnant women not involved as study participants. The FGDs and in-depth interviews used semi-structured questions, and probes were used to elicit further descriptions of challenges and experiences.

Data analysis

Data were analyzed using a thematic analysis method included six phase framework for doing a thematic

analysis included familiarized themselves with the data, produced initial codes, searched for themes, evaluated the themes, defined and named the themes, and made the study report [31]. Three researchers analyzed the data independently. These researchers analyzed the data manually. First, the data were transcribed the day after collection and were analyzed by the researchers to be familiar with the data. Data collection and data analysis were conducted simultaneously. The transcribed verbatim was carefully read and re-read by the researchers, and the researchers provided the participants with the opportunity to check the verbatim. Second, all the transcribed verbatim were carefully and precisely examined line by line to find initial codes, and then were categorized based on the meaning similarity. Data was cleaned by eliminating all distinguishable information, and the researchers created initial codes before interviewing the next participant. Third, the researchers developed sub-themes and themes based on the categorized data. They categorized the codes with associated meanings into one group called sub-themes and considered their significance. They then categorized all codes and categories into central categories called themes. Fourth, researchers worked together to evaluate the draft of themes in biweekly meetings. The entire research team then discussed the coding, categories, sub-themes, and themes and organized them into charts. Fifth, each member of the research team agreed upon the names of the themes and sub-themes, and some participants were asked to read the study results and give feedback about whether the themes accurately portrayed their views. Finally, the research team discussed and agreed upon the names of the final themes that emerged from this study before making the final report. The last phase is write-up study report following the consolidated criteria for reporting qualitative research (COREQ) [32]. The demographic characteristics of the participants were presented in frequency, percentage, median, minimum, and maximum.

Trustworthiness

To assess the quality of the study data, the researchers used Lincoln and Guba's trustworthiness criteria [24]. Most researchers in this study were Javanese and native residents of the region under investigation, and this helped gain participants' confidence and familiarity. The study included pregnant women and healthcare providers varying in age, parity, level of education, working status, and working experience to help identify different views and concepts and heighten credibility. During the study's execution, a qualitative research specialist observed the data collection and processing, and two qualitative researchers analyzed the data independently. The data collection, data analysis, and theory generation process can be audited, which heightens the study's

dependability. The researchers attempted to avoid subjective prejudices by recording all FGDs and interviews, keeping accurate field notes, and avoiding interfering with the data analysis results, leading to better confirmability within the study. The results were shared with participants who were asked to confirm whether they accurately reflected their experiences, leading to better transferability within the study.

Rigor

The interview guidelines were based on a literature review focusing on ANC coverage during the COVID-19 pandemic. The draft of interview guidelines was checked for clearness and modified based on suggestions made by a panel of experts and pregnant women independent of the study. It was then tested on five pregnant women and two healthcare providers. The data were translated into English at each stage of the data analysis process. Participants were given the opportunity to carry out member checking to validate the data and were asked for their opinions regarding the themes. The entire research team read and discussed the data to confirm the right information was reported. Three researchers involved in this study had previous experience in conducting and publishing qualitative studies, which further helped to ensure the rigor of the qualitative data.

Ethical consideration

Ethical approval for this study was granted by the IRB of the Medical Faculty, Universitas Jenderal Soedirman No 1204/KEPK/III/2017. Permissions were also granted by district assemblies and the District Ministry of Health. Researchers explained the research purposes, procedures, risks, participants' expected role, and the voluntary nature of participation in the study to each participant. Participants were also informed that they were free to leave the study any time and that this would not affect any healthcare services received by them. Either a written or electronic informed consent form was obtained by each participant before involvement in the study and agreement to audio record the discussion or in-depth conversation was also obtained. All participants' data were documented with codes to guarantee anonymity and only the research team had access to the research data to ensure confidentiality.

Results

The majority of pregnant women included in the study were between 20 and 35 years old (52%), multiparous (68%), housewives (68%), and had graduated from senior high school (40%). The healthcare providers included in the study consisted of three midwives (50%), a nurse (16.7%), a head of the PHC (16.7%), and a doctor (16.7%). The mean age of the healthcare providers was 41.83

Table 1 The participant's demographic data

Table 1 The participant's demographic data				
Variables	Frequency (%)	Median (Min-Max)		
Pregnant women				
Age (years)		31.5 (19–43)		
<20	1 (4%)			
20–35	13 (52%)			
> 35	11 (44%)			
Parity				
Primiparous	8 (32%)			
Multiparous	17 (68%)			
Occupation				
Seller	1 (4%)			
Teacher	1 (4%)			
Housewife	15 (60%)			
Private employee	6 (24%)			
Government employee	2 (8%)			
Education				
Elementary School	3 (12%)			
Junior High School	6 (24%)			
Senior High School	10 (40%)			
University	6 (24%)			
Healthcare providers				
Gender				
Male	1 (16.7%)			
Female	5 (83.3%)			
Age (years)		41 (34–52)		
20–35	1 (16.7%)			
> 35	5 (83.3%)			
Working experience				
5–10 years	2 (33.3%)			
>11 years	4 (66.7%)			
Education				
Bachelor of Public Health	1 (16.7%)			
Medical Doctor	1 (16.7%)			
Diploma of Midwifery	3 (50%)			
Bachelor of Nursing	1 (16.7%)			
Occupation				
Head of PHC	1 (16.7%)			
Medical Doctor	1 (16.7%)			
Midwives	3 (50%)			
Nurse	1 (16.7%)			

years, with a range of 34–52 years. The majority of healthcare providers had more than 11 years' work experience (66.7%). Half of healthcare providers had completed either a bachelor's degree or diploma of midwifery (Table 1). The details of participant's demographic data were provided in Table 2.

Themes

The themes emerged out of findings from in-depth interviews, focus group discussions, and field notes which were acquired from observations at the Primary Health Centers. Using the Three Delays Model by Thaddeus and Maine as the study framework, the researchers identified. three themes: (1) Fear of being infected with COVID-19, (2) The stay-at-home policy, and (3) Re-designed ANC services. Each theme was broken into sub-themes

Table 2 Participants demographic data

Participant	Age	Parity	Education level	Occupation
P1	36	Multiparous	Senior High School	Housewife
P2	29	Multiparous	Bachelor's degree	Private employee
P3	36	Primiparous	Senior High School	Housewife
P4	31	Multiparous	Bachelor's degree	Government employee
P5	41	Multiparous	Elementary School	Housewife
P6	28	Primiparous	Senior High School	Housewife
P7	27	Primiparous	Bachelor's degree	Government employee
P8	39	Multiparous	Senior High School	Private employee
P9	24	Primiparous	Senior High School	Housewife
P10	32	Multiparous	Elementary School	Housewife
P11	22	Primiparous	Bachelor's degree	Private employee
P12	40	Multiparous	Junior High School	Seller
P13	19	Primiparous	Senior High School	Housewife
P14	30	Multiparous	Junior High School	Housewife
P15	36	Multiparous	Bachelor's degree	Private employee
P16	23	Primiparous	Senior High School	Private employee
P17	39	Multiparous	Junior High School	Housewife
P18	34	Multiparous	Junior High School	Housewife
P19	37	Multiparous	Junior High School	Housewife
P20	40	Multiparous	Senior High School	Private employee
P21	27	Primiparous	Senior High School	Housewife
P22	38	Multiparous	Bachelor's degree	Teacher
P23	29	Multiparous	Junior High School	Housewife
P24	31	Multiparous	Senior High School	Housewife
P25	43	Multiparous	Elementary School	Housewife
Participant	Age	Gender	Education	Occupation
P26	34	Female	Bachelor of Nursing	Nurse
P27	52	Male	Bachelor of Public Health Science	Head of PHC
P28	45	Female	Medical Doctor	Doctor
P29	38	Female	Diploma of Midwifery	Midwife
P30	39	Female	Diploma of Midwifery	Midwife
P31	43	Female	Diploma of Midwifery	Midwife

Table 3 Summary of themes and sub-themes from the transcribed data

Themes	Sub-themes
1. Fear of being infected with COVID-19	Anxiety
	Perceived vulnerability
	The desire to protect self and loved ones
2. The stay-at-home policy	Transport barriers to healthcare
	Social activity restriction
3. Re-designed ANC services	ANC adjustments
	Focus on high-risk pregnancy
	Insufficient information
	COVID-19 preventive behaviors adherence

and the descriptive quotes for each sub-theme are illustrated in Table 3.

1. Fear of being infected with COVID-19

According to Three Delays Models, the first delay is related to the decision to seek care. During the pandemic time, pregnant women often prefer to delay their ANC visit because they are afraid of being infected with COVID-19. This theme is comprised of three subthemes: anxiety, perceived vulnerability, and the desire to protect self and loved ones.

Anxiety

Almost all of participants stated that the pandemic had made them feel anxious. Interacting with people outside the home induced anxiety of being infected with COVID-19.

...I am very worried since I am pregnant ... my office

colleague was confirmed as having COVID-19 so I feel very anxious to have contact with other people including healthcare providers ... (P8, multiparous, 39 years old).

Many participants worried about their health condition because of their weak immunity status and knew that their risk of contracting the virus increased when they visited healthcare facilities for ANC.

... I am scared because the immunity status of pregnant women is decreased and the COVID-19 virus spread very quickly, I really feel fear when I go to the PHC to attend ANC so, I prefer to delay my next ANC schedule ... (P23, multiparous, 29 years old).

Perceived vulnerability

Another participant expressed having mixed feelings about getting pregnant at this time. She felt grateful because she had been trying to get pregnant for a long time. However, she knew that the fact she was pregnant during the COVID-19 pandemic made her more vulnerable and this hindered her ANC attendance.

... I'm grateful and happy for my first pregnancy after having waited for a long time ... however I'm a vulnerable person because I am pregnant during the COVID-19 pandemic, so I do not attend ANC regularly ... I hope to contact healthcare providers using social media ... (P3, primiparous, 36 years old).

The desire to protect self and loved ones

Some participants lived with members of other vulnerable groups such as the elderly, children, and people with co-morbidities. Some participants feared that engaging in activities outside the home increased their chances of being infected and they tried to minimize visiting places they deemed high-risk, such as clinics where ANC was carried out. Some participants preferred to delay ANC to prevent themselves or their children and parents from getting ill.

... I have to minimize visiting high-risk transmission places because I have children and my parents have co-morbidities. We have to take care of each other ... I will delay my ANC until the COVID-19 situation improves ... (P8, multiparous, 39 years old).

2. The stay-at-home policy

According to Three Delays Models, the second delay is related to identify and reach the health facility. During

the COVID-19 pandemic, pregnant women prefer to delay their ANC visit due to Large Social Scale Restriction policy by Indonesia Government. This theme is comprised of two sub-themes: transport barriers to health care and restriction of social activities.

Transport barriers to health care

Participants disclosed that the COVID-19 pandemic had created several barriers to their daily activities. The Indonesian government's policy encouraged people to stay at home during the pandemic. Participants therefore faced public transport barriers to attending their ANC.

... I faced difficulties attending ANC during the COVID-19 pandemic because of a lack of public transport due to the government's stay at home policy ... it would be really helpful if I could have my healthcare consultations over social media ... (P14, multiparous, 30 years old).

Social activity restriction

Almost all participants described experiencing social activity restrictions during the pandemic due to government policies. Participants were encouraged to stay at home and not to attend activities involving a lot of people to prevent transmission.

... My husband did not allow me to join social gatherings, recitations, or religious meetings during the pandemic ... I had to follow the government's regulations and stay at home, and so I decided not to attend ANC this month ... (P6, multiparous, 28 years old).

3. Re-designing ANC services

According to Three Delays Models, the third delay is related to receive appropriate treatment at the health facility. Health care providers re-design ANC services during the COVID-19 pandemic. This theme is comprised of four sub-themes: ANC adjustments; focus on high-risk pregnancy; insufficient information; and adherence to COVID-19 preventive behaviors.

ANC adjustments

All participants experienced some ANC services being modified during the pandemic. Many services were redesigned to reduce COVID-19 transmission. This meant ANC sessions were shortened and pregnancy classes were cancelled.

... They asked me some questions such as whether I had flu-like symptoms or fever, then they measured

my blood pressure and weight and assessed the fetus in my womb very quickly ... the midwife did a quick physical examination ... (P9, primiparous, 24 years old).

Healthcare providers confirmed that ANC procedures were adjusted to prevent transmission. The Indonesia Ministry of Health ANC adaptation guidance called for early screening, the separation of healthy women suspected of being pregnant, and modified seating and room arrangement.

... We provide ANC using a new adaptation era guidance from the Ministry of Health. We take pregnant women's' temperatures, assess symptoms such as coughs and colds, and ask their history before carrying out the physical examination ... we adjusted the ANC procedures and duration to prevent COVID-19 transmission ... we also canceled prenatal classes during the pandemic ... (P30, midwife, 39 years old).

Focus on high-risk pregnancy

Healthcare providers said that the ANC examinations during the pandemic focused on patients' health problems and complaints, meaning that pregnant women with no problems were more likely to delay ANC to minimize risk. Healthcare providers made home visits to women with high-risk pregnancies to keep them safe.

... We re-arranged ANC schedules for low-risk pregnancies to prevent infection and we made home visits to high-risk pregnant women to keep them at home ... (P31, midwife, 43 years old).

... The midwife came to my house and did a physical examination. She explained that I should stay at home during this time and told me she would visit again in the following months to keep me safe ... (P20, multiparous, 40 years old).

Insufficient information

All pregnant participants said they had received insufficient information during their ANC appointments. They maintained that ANC assessments were short in duration and that there was no discussion or health promotion given by healthcare providers. They were given no chance to express their feeling or ask questions related to their pregnancies, and they said they used social media to facilitate discussions with their healthcare providers and get pregnancy-related advice.

... when I had my ultrasound examination, the doc-

tor did it quickly and did not explain the results in detail ... he just said it was good and normal. The doctor didn't give me the opportunity to express my feelings or ask questions about my pregnancy ... (P10, multiparous, 32 years old).

... I met the midwife in the ANC room for a very short time ... she did a quick physical examination, did not give much information about the examination results, and did not explain about my fetus in detail ... maybe I can use social media to communicate with healthcare providers and get some pregnancy-related advice ... (P11, primiparous, 22 years old).

Adherence to COVID-19 preventive behaviors

All participants said that they adhered to COVID-19 preventive behaviors when going outside the home. They wore masks, washed their hands, and observed physical distancing when attending ANC. There is a big banner containing information about COVID-19 preventive measures in front of the PHC office.

... I always wear a mask, wash my hands in the wash basin which placed in front of the PHC, and sit down not too close to other patients in the PHC ... there is a person in charge who always reminds us to wear a mask, wash our hands, and keep a safe distance from other people ... (P14, multiparous, 30 years old).

Healthcare providers confirmed that there is a banner in front of the PHC that informs patients about COVID-19 preventive behaviors. They maintained that they frequently reminded patients to follow the preventative measures, provided a hand washing basin outside the PHC, and wore additional personal protective equipment.

.... We provide healthcare services in the PHC using a high standard of COVID-19 preventive measures ... There is a big banner in front of the PCH informing patients of the COVID-19 preventive behaviors ... we always remind patients to wear a mask, keep at least two meters distance from others, and wash their hands using water and soap before and after entering the PHC ... healthcare providers should wear additional personal protective equipment ... (P2, head of PHC, 52 years old).

Discussion

To our knowledge, this is the first Indonesian study to investigate the barriers to ANC from the perspectives of pregnant mothers and healthcare providers during the COVID-19 pandemic. Despite facing many challenges and restrictions throughout the pandemic, healthcare providers have had to continue providing a full range of ANC services to the pregnant population. Some services were modified in order to prevent COVID-19 transmission [33]. The number of ANC visits decreased during the pandemic, particularly among pregnant women in rural areas [33]. This might have been due to fear of infection, transport problems, the fact that women lived far away from health facilities, the government's COVID-19 policies, changing healthcare services, or lack of awareness about the benefits of ANC.

Our findings show that many pregnant women were afraid of COVID-19 infection. Pregnant woman are particularly vulnerable to viruses due to changes to maternal immune responses and increased sensitivity to respiratory pathogens [18]. Most pregnant women in this study attended ANC in PHCs however, this made them feel anxious. Our results are similar to those of a previous study that found that pregnant women during the pandemic missed ANC visits and were afraid of giving birth in health facilities due to a fear of contracting COVID-19 [34]. Our results also correlate to those of another study that discovered that while women desired to selfmonitor their pregnancies, they did not have sufficient knowledge, and that they felt the need to strictly isolate at home and expected pregnancy tele-consultations [35]. In our study, fear was the most important factor in reduced ANC during the pandemic, and this correlates to the findings of a study carried out in Israel, whereby women refused to visit healthcare centers for ANC out of fear of viral infection [36]. A large survey found that pregnant women were particularly fearful of restrictions to family visits post-delivery, the possibility of their babies being infected COVID-19, the lack of support offered during delivery, and delivery plans changing [37]. Anxiety related to COVID-19 transmission was also experienced by healthcare workers in Finland, particularly among women and young people [38]. Unsurprisingly, our results show that the combination of pregnancy and working status increased the fear of viral transmission among Indonesian working women who conceived during the pandemic.

Our study shows that Indonesian pregnant women were not only scared of COVID-19 infecting themselves but also worried that they would transmit the virus to family members with comorbidities since most participants lived with their extended families. Pregnant women in Iran also experienced fear, obsession, boredom, nervousness, and despair during the pandemic due

to the highly contagious nature of the virus [39]. Living with the extended family is common for people living in rural areas of Indonesia. The majority of participants in this study lived with their parents, parents-in-law, or grandparents, many of whom had comorbidities. Fear of COVID-19 transmission among people living in rural areas may have been caused by low adherence to healthcare protocols. Participants in this study said they felt uncomfortable wearing masks and wanted to join their neighbors for social interactions. Urban dwellers in other studies found it easier to share knowledge about COVID-19 [40], advised others to carry out preventive behaviors [40], and were found to have less negative attitudes regarding preventive behaviors [41]. The low adherence of COVID-19 preventive behaviors among people in rural areas might be due to low health literacy and lack of information. One study found that people with adequate knowledge of the virus were more likely to engage in appropriate preventive behaviors and that urban dwellers were more likely to practice good preventive behaviors than rural dwellers [42].

The qualitative findings of this study further illustrate that the pandemic affected public transport and restricted social activities. The Indonesian government restricted non-essential travel and group activities to prevent the spread of COVID-19 [43]. Participants in this study reported that public transport was limited due to government policies and so they found it difficult to attend their ANC appointments. These findings are similar to those reported in a previous study that also found that pregnant women faced difficulty accessing public transport pandemic, resulting in missed ANC appointments [44]. Social support from family members, friends, and significant others contributes to developing resilience among pregnant women, positively affecting perinatal mental health and obstetric outcomes [45]. Social activities are part of daily life for people living in Indonesia's rural areas. People are accustomed to visiting neighbors, praying together in the mosque, and attending funerals, recitations, wedding ceremonies, and social gatherings.

The Ministry of Health launched perinatal and newborn health service guidelines to guide healthcare providers in the delivery of healthcare services during the pandemic. The guidelines included information regarding general precautions, health facility readiness, perinatal care recommendations, and health services for maternal and newborns [33]. Healthcare providers in this study described having to redesign ANC programs, limit prenatal examinations, carry out more home visits, and limit the number of prenatal class participants. These ANC program adjustments were made to protect both healthcare providers and pregnant women from COVID-19 transmission, particularly during July 2021

when daily cases in Indonesia reached the highest level [3]. Unsurprisingly, the situation of maternal health services during a pandemic is more precarious in developing countries due to insufficient infrastructure and resources, the collapse of healthcare systems, decreased workforce, access decline, and fewer ANC visits [46]. Consistent with a study in Kenya that reported ANC program modifications including limiting ANC hospital appointments and increasing networking between community health workers and communities [47]. Participants in this study reported that they feel unsatisfied because they did not have sufficient time to discuss their pregnancies with healthcare providers during their ANC appointments. Participants in another study reported that if they stated dissatisfaction with ANC services due to changes caused by the pandemic, they had their ANC appointments and pregnancy classes cancelled or suspended [48]. One study found that pregnant women were not examined in detail during ANC appointments, and that this may have led to health professionals missing important aspects in their pregnancy [44]. Participants in the same study also used social media or written documents to obtain information from their healthcare providers during the pandemic. Pregnant women in another study declared social media a useful tool in providing antenatal care and support during the pandemic, particularly with regard to obtaining pregnancy-related information, managing feelings of isolation, service specific issues, and routine care [49]. The use of social media is an interesting way to communicate and deliver health promotion to pregnant women, as identified in this study.

In another study, good preventive behavior among pregnant women was found to be significantly associated with fear of transmission and good knowledge of COVID-19 preventive behaviors [50]. Participants in this study understand that pregnant women are a vulnerable group due to their lowered immune response, and they rigorously implemented preventive measures such as wearing masks, washing hands with water and soap, and observing physical distancing, even while attending ANC appointments in the PHC. Such careful adherence to COVID-19 preventive behaviors may be due to the government's massive publicity campaign.

Strengths and limitations

A strength of this study is that it used open-ended questions which provided rich data for qualitative analysis. Its findings are still relevant to improving maternal health services. Despite its strengths, this study also has limitations. Its principal limitation is that the findings are of limited generalizability since all participants were married women of Javanese ethnicity who attended ANC appointments during the pandemic. Further research is required to examine the utility of social media as way of

communicating between healthcare providers and pregnant women.

Conclusion

This study highlights the substantial impact COVID-19 had on ANC in rural areas of Indonesia. Its findings contribute to the growing body of evidence outlining the challenges faced in providing ANC in rural areas during pandemics. Pregnant women faced several difficulties in seeking antenatal care, reaching antenatal care facilities, and receiving satisfactory services at the health care facilities during the COVID-19 pandemic.

Acknowledgements

We would like to express our deepest gratitude to participants in this study to share their valuable experiences. We would like to extend our sincere thanks to data collection team. We also could not have undertaken this study without funding support from Universitas Jenderal Soedirman.

Author Contribution

MDA conceptualized and designed the study, wrote, and revised the final manuscript. RS analyzed the data and critically reviewed and revised the manuscript. ET designed the study, analyzed the data, and critically reviewed and revised the manuscript. Al contributed to collected data and assisted with data analysis, critically reviewed, and revised the manuscript. DN analyzed the data, and critically reviewed and revised the manuscript. AF contributed to collected data, critically reviewed, and revised the manuscript.

Funding

This study was supported by a research grant from Universitas Jenderal Soedirman.

Data Availability

The datasets generated during and/or analyzed during the study are available from the corresponding author on reasonable request.

Declarations

Ethical approval and consent

All methods were carried out in accordance with relevant guidelines and regulations. All procedures were approved by the Institutional Review Board, Faculty of Medicine, Universitas Jenderal Soedirman. Informed consent was obtained from all subjects and/or their legal guardian(s).

Consent to publish

Not applicable.

Conflict of Interest

There is no conflict of interest to declare.

Received: 28 July 2022 / Accepted: 3 March 2023 Published online: 16 March 2023

References

- World Health O. COVID 19 Public Health Emergency of International Concern (PHEIC). Global research and innovation forum: towards a research roadmap. 2020.
- WHO. WHO coronavirus disease (COVID-19) dashboard with vaccination data: Indonesia 2021 [Available from: https://covid19.who.int/region/searo/country/id.
- Worldmeter. Worldmeter's COVID-19 data: Indonesia 2021 [Available from: https://worldmeters.info/coronavirus/country/indonesia/.

- Read S, Wild SH. What are the risks of COVID-19 infection in pregnant women? 2020; 395:[760-2 pp.].
- 5. Health IMo. Indonesia Health Profile. In: Health Mo, editor. Jakarta 2018.
- 6. Health IMo. Indonesia Health Profile. In: Health Mo, editor. Jakarta 2019.
- 7. Health IMo. Indonesia Health Profile. In: Health Mo, editor. Jakarta 2020.
- Nasional BKdKB. Kematian ibu hamil selama pandemi tinggi, Kepala BKKBN apresiasi kecepatan vaksinasi di DKI Jakarta2021. Available from: https:// www.bkkbn.go.id/detailpost/kematian-ibu-hamil-selama-pandemi-tinggikepala-bkkbn-apresiasi-kecepatan-vaksinasi-di-dki-jakarta.
- Carvalho-Sauer R, Costa M, Teixeira M, Nascimento E, Silva E, Barbosa M et al. Impact of COVID-19 pandemic on time series of maternal mortality ration in Bahia, Brazil: analysis of period 2011–2020.BMC pregnancy and childbirth. 2021;21.
- Tola W, Negash E, Sileshi T, Wakgari N. Late initiation of antenatal care aand associated factors among pregnant women attending antenatal clinic of Ilu Ababor Zone, Southwest Ethiopia: a cross-sectional study. PLoS ONE. 2021;16(1):1–11.
- 11. UNICEF. Antenatal Care. In: UNICEF, editor. 2021.
- Abbas A, Rabeea M, Hafiz H, Ahmed N, editors., editors. Effects of irregural antenatal care attendace in primiparas on the perinatal outcomes: a cross sectional study. Proceedings in Obstetrics and Gynecology; 2017.
- 13. Mckinney E. Maternal-child nursing. Canada: Elsevier; 2018.
- Kazemi SN, Hajikhani B, Didar H, Hosseini SS, Haddadi S, Khalili F et al. COVID-19 and cause of pregnancy loss during the pandemic: A systematic review. Plos One. 2021:1–10.
- Westgren M, Pettersson K, Hagberg H, Ganesh A. Severe maternal morbidity and mortality associated with COVID-19: the risk should not be downplayed. Acta Obstet Gynecol Scand. 2020;99(7):815–6.
- Goyal M, Singh P, Singh K, Shashank S, Agrawal N, Misra S. The effect of COVID-19 pandemic on maternal health due to delay in seeking health care: experience from a tertiary center. International Journal of Gynecology & Obstetrics: 2020.
- Organization) PPAH. Epidemiological alert: COVID-19 during pregnancy Washington D.C2020 [Available from: https://www.paho.org/en/news/4-8-2020-paho-director-warns-disruptions-regular-health-services-duecovid-19.
- Liu H, Wang L-L, Zhao S-J, Kim JK, Mor G, Liao A-H. Why are pregnant women susceptible to COVID-19? An immunological viewpoint. J Reprod Immunol. 2020:139:1–4.
- Zacharias P, Joseph M, Jacob A, Viji CLM, Josy N, Lydia S, et al. Barriers to antenatal care during COVID-19 pandemic: a hospital-based retrospective cross-sectional study in rural South Karnataka. The New Indian Journal of Obgyn. 2021;8(1):27–32.
- 20. Hailemariam S, Agegnehu W, Misganaw D, Exploring. COVID-19 related factors of influencing antenatal care services uptake: a qualitative study among women in rural community in Southwest Ethiopia. J Prim Care Community Health. 2021:12:1–8.
- Singh DR, Sunuwar DR, Shah SK, Karki K, Sah LK, Adhikari B, et al. Impact of COVID-19 on health services utilization in Province-2 of Nepal: a qualitative study among community members and stakeholders. BMC Health Serv Res. 2021;21(1):1–14.
- Meaney S, Leitao S, Olander EK, Pope J, Matvienko-Sikar K. The impact of COVID-19 on pregnant womens' experiences and perceptions of antenatal maternity care, social support, and stress-reduction strategies. Women Birth. 2022;35(3):307–16.
- McDonald CR, Weckman AM, Wright JK, Conroy AL, Kain KC. Pregnant women in low-and middle-income countries require a special focus during the COVID-19 pandemic. Front Global Women's Health. 2020;1:564560.
- 24. Polit DF, Beck CT. Essentials of nursing research: appraising evidence for nursing practice. 9th ed. Philadelphia: Lippincott Williams & Wilkins; 2018.
- 25. Kim H, Sefcik JS, Bradway C. Characteristics of qualitative descriptive studies: a systematic review. Res Nurs Health. 2017;40(1):23–42.
- Thaddeus S, Maine D. Too far to walk: maternal mortality in context. Social science & medicine (1982). 1994;38(8):1091 – 110.
- Wulandari RD, Laksono AD, Rohmah N. Urban-rural disparities of antenatal care in South East Asia: a case study in the Philippines and Indonesia. BMC Public Health. 2021;21(1):1–9.
- Statistik BP. Profil Penduduk Indonesia Hasil SUPAS. Badan Pusat Statistik Jakarta. 2015.
- Pitoyo AJ, Triwahyudi H. Dinamika perkembangan etnis di Indonesia dalam konteks persatuan negara. Populasi. 2017;25(1):64–81.

- Suminar YD, Lutiarsi RT, Wibowo MA, Lestari ES. Profil Kesehatan Jawa Tengah. Jawa Tengah: Dinkes Jawa Tengah: Dinas Kesehatan Jawa Tengah; 2021.
- 31. Braun V, Clarke V. Using thematic analysis in psychology. Qualitative Res Psychol. 2006;3(2):77–101.
- Tong A, Sainsbury P, Craig J. Consolidated Criteria for reporting qualitative research (COREQ): a 32-Item Checklist for interviews and focus groups. Int J Qual Health Care. 2007;19:349–57.
- 33. Health IMo. Guidelines for Antenatal Care, Labor, Postpartum, and Newborn in New Normal Era. Jakarta: Indonesia Ministry of Health; 2020.
- Moyer CA, Sakyi KS, Sacks E, Compton SD, Lori JR, Williams JEO. COVID-19 is increasing ghanaian pregnant women's anxiety and reducing Healthcare seeking. Int J Gynecol Obstet. 2020;00:1–2.
- Ge Y, Shi C, Wu B. Anxiety and Adaptation of Behavior in Pregnant Zhuang Women During the COVID-19 Pandemic: A Mixed-Mode Survey. Risk Management and Healthcare Policy. 2021;14.
- WHO, Coronavirus. and pregnancy preserving maternal health across the european region 2020 [Available from: https://www.euro.who.int/en/healthtopics/Life-stages/maternal-and-newborn-health/news/news/2020/6/coronavirus-and-pregnancy-preserving-maternal-health-across-the-europeanregion.)
- Basu A, Kim HH, Basaldua R, Choi KW, Charron L, Kelsall N et al. A Cross-National Study of Factors Associated with Women's Perinatal Mental Health and Wellbeing During the COVID-19 Pandemic.Plos Ones. 2021:1–18.
- 38. Savolainen L, Oksa R, Savela N, Celuch M, Oksanen A. COVID-19 Anxiety- A longitudinal survey study of psychological ad situational risks among finnish worker. Int J Environ Res Public Health. 2021;18(2):794.
- Bakouei F, Nikpour M, Rad HA, Marzoni ZA. Exploration of the Pregnant Women's Experiences During COVID-19 Disease Crisis: A Qualitative study. 2020.
- Zhang J, Zhu L, Li S, Huang J, Ye Z, Wei Q, et al. Rural-urban Dlsparities in knowledge, Behaviors, and Mental Health during COVID-19 pandemic. Medicine. 2021;100(13):1–7.
- 41. Chen X, Chen H. Differences in preventive ehaviors of COVID-19 between urban and rural residents: lessons learned from a cross-sectional study in China. Int J Environ Res Public Health. 2020;17(4437):1–14.
- 42. Gutu B, Legese G, Fikadu N, Kumela B, Shuma F, Mosisa W, et al. Assessment of preventive behavior and associated factors towards COVID-19 in Qellam Wallaga Zone, Oromia, Ethiopia: a community-based cross-sectional study. PLoS ONE. 2021;16(4):e0251062.
- 43. Health IMo. Abolition of mudik at ied fitr 1442 and preventing corona virus disease 2019 (COVID-19) during ramadhan 1442. In: Health Mo, editor. Jakarta2021.
- 44. Karavadra B, Stockl A, Prosser-Snelling E, Simpson P, Morris E. Women's perceptions of COVID-19 and their healthcare experiences: a qualitative thematic analysis of a national survey of pregnant women in the United Kingdom. BMC Pregnancy Childbirth. 2020;20(1):1–8.
- Dunkel C. Psychological science on pregnancy: stress process, biopsychosocial models, and emerging research issues. Ann Rev Psychol. 2011;62:531–58.
- Tadesse E. Antenatal Care Service utilization of pregnant women attending Antenatal Care in Public Hospitals during the COVID-19 pandemic period. Int J Womens Health. 2020;12:1181–8.
- 47. Kimani RW, Maina R, Shumba C, Shaibu S. Maternal and newborn care during the COVID-19 pandemic in Kenya: recontextualising the community midwifery model. Hum Resour Health. 2020;75:1–5.
- Meaney S, Leitao S, Pope J, Matvienko-Sikar K. The impact of COVID-19 on pregnant women's experiences and perceptions of antenatal maternity care, social support, and stress-reduction strategies. Women Birth. 2021(21).
- Chatwin J, Butler D, Jones J, James L, Choucri L, McCarthy R. Experiences of pregnant mothers using a social media based antenatal support service during the COVID-19 lockdown in the UK: findings from a user survey. BMJ open. 2021;11(1):e040649.
- W M, Kassie GT, Asratie B, Abate M. The effects of fear and knowledge of COVID-19 on preventive practice among pregnant women who attend antenatal care in Northwest Ethiopia, 2020: institution-based cross-sectional study. Int J Women's Health. 2021;13:95–100.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

BMC by Mekar Anggraeni

Submission date: 21-Mar-2023 08:34AM (UTC+0700)

Submission ID: 2042225563

File name: Anggraeni_et_al-2023-BMC_Pregnancy_and_Childbirth.pdf (979.64K)

Word count: 8625 **Character count:** 47962