


ISSN: 1412-033X  
E-ISSN: 2085-4722

# BIODIVERSITAS

**Journal of Biological Diversity**

Volume 22 - Number 1 - January 2021



# BIODIVERSITAS

Journal of Biological Diversity  
Volume 22 – Number 1 – January 2021

ISSN/E-ISSN:  
1412-033X (printed edition), 2085-4722 (electronic)

## EDITORIAL BOARD:

Abdel Fattah N.A. Rabou (Palestine), Agnieszka B. Najda (Poland), Ajay Kumar Gautam (India), Alan J. Lymbery (Australia), Annisa (Indonesia), Bambang H. Saharjo (Indonesia), Daiane H. Nunes (Brazil), Darlina Md. Naim (Malaysia), Ghulam Hassan Dar (India), Hassan Pourbabaee (Iran), Joko R. Witono (Indonesia), Kartika Dewi (Indonesia), Katsuhiko Kondo (Japan), Kusumadewi Sri Yulita (Indonesia), Livia Wannorp (Sweden), M. Jayakara Bhandary (India), Mahdi Reyahi-Khoram (Iran), Mahendra K. Rai (India), Mahesh K. Adhikari (Nepal), Maria Panitsa (Greece), Mochamad A. Soendjoto (Indonesia), Mohib Shah (Pakistan), Mohamed M.M. Najim (Srilanka), Nurhasanah (Indonesia), Praptiwi (Indonesia), Rasool B. Tareen (Pakistan), Seyed Aliakbar Hedayati (Iran), Seyed Mehdi Talebi (Iran), Shahabuddin (Indonesia), Shahir Shamsir (Malaysia), Shri Kant Tripathi (India), Subhash C. Santra (India), Sugeng Budiharta (Indonesia), Sugiyarto (Indonesia), Taufiq Purna Nugraha (Indonesia), Yosep S. Mau (Indonesia)

## EDITOR-IN-CHIEF: Sutarno

**EDITORIAL MEMBERS:**  
English Editors: **Graham Eagleton** (grahameagleton@gmail.com), **Suranto** (surantouns@gmail.com); Technical Editor: **Solichatun** (solichatun\_s@yahoo.com), **Artini Pangastuti** (pangastuti\_tutut@yahoo.co.id); Distribution & Marketing: **Rita Rakhmawati** (oktia@yahoo.com); Webmaster: **Ari Pitoyo** (aripitoyo@yahoo.com)

**MANAGING EDITORS:**  
**Ahmad Dwi Setyawan** (unsjournals@gmail.com)


**PUBLISHER:**  
The Society for Indonesian Biodiversity

**CO-PUBLISHER:**  
Department of Biology, Faculty of Mathematics and Natural Sciences, Sebelas Maret University, Surakarta


**ADDRESS:**  
Jl. Ir. Sutami 36A Surakarta 57126. Tel. +62-271-7994097, Tel. & Fax.: +62-271-663375, email: editors@smujo.id

**ONLINE:**  
biodiversitas.mipa.uns.ac.id; smujo.id/biodiv

\*\*\*\*\*



Society for Indonesia  
Biodiversity



Sebelas Maret University  
Surakarta

Published by Smujo International for The Society for Indonesian Biodiversity and Sebelas Maret University Surakarta

## GUIDANCE FOR AUTHORS

**Aims and Scope** *Biodiversitas*, *Journal of Biological Diversity* or abbreviated as *Biodiversitas* encourages submission of manuscripts dealing with all biodiversity aspects of plants, animals and microbes at the level of the gene, species, and ecosystem as well as ethnobiology.

**Article types** The journal seeks original full-length research papers, reviews, and short communication. Manuscript of original research should be written in no more than 8,000 words (including tables and picture), or proportional with articles in this publication number. Review articles will be accommodated, while, short communication should be written at least 2,000 words, except for pre-study.

**Submission** The journal only accepts online submission, through open journal system (<https://smujo.id/biodiv/about/submissions>) or email to the editors at [unsjournals@gmail.com](mailto:unsjournals@gmail.com). Submitted manuscripts should be the original works of the author(s). The manuscript must be accompanied by a cover letter containing the article title, the first name and last name of all the authors, a paragraph describing the claimed novelty of the findings versus current knowledge. Submission of a manuscript implies that the submitted work has not been published before (except as part of a thesis or report, or abstract); and is not being considered for publication elsewhere. When a manuscript written by a group, all authors should read and approve the final version of the submitted manuscript and its revision; and agree the submission of manuscripts for this journal. All authors should have made substantial contributions to the concept and design of the research, acquisition of the data and its analysis; drafting of the manuscript and correcting of the revision. All authors must be responsible for the quality, accuracy, and ethics of the work.

**Ethics** Author(s) must obedient to the law and/or ethics in treating the object of research and pay attention to the legality of material sources and intellectual property rights.

**Copyright** If and when the manuscript is accepted for publication, the author(s) still hold the copyright and retain publishing rights without restrictions. Authors or others are allowed to multiply article as long as not for commercial purposes. For the new invention, authors are suggested to manage its patent before published.

**Open access** The journal is committed to free-open access that does not charge readers or their institutions for access. Readers are entitled to read, download, copy, distribute, print, search, or link to the full texts of articles, as long as not for commercial purposes. The license type is CC-BY-NC-SA.

**Acceptance** The only articles written in English (U.S. English) are accepted for publication. Manuscripts will be reviewed by editors and invited reviewers (double blind review) according to their disciplines. Authors will generally be notified of acceptance, rejection, or need for revision within 1 to 2 months of receipt. The manuscript is rejected if the content does not in line with the journal scope, does not meet the standard quality, inappropriate format, complicated grammar, dishonesty (i.e. plagiarism, duplicate publications, fabrication of data, citations manipulation, etc.), or ignoring correspondence in three months. The primary criteria for publication are scientific quality and biodiversity significance. **Uncorrected proofs** will be sent to the corresponding author by email as .doc or .docx files for checking and correcting of typographical errors. To avoid delay in publication, corrected proofs should be returned in 7 days. The accepted papers will be published online in a chronological order at any time, but printed in the early of each month (12 times).

**A charge** Starting on January 1, 2019, publishing costs waiver is granted to authors of graduate students from **Least Developed Countries**, who first publish the manuscript in this journal. However, other authors are charged USD 250 (IDR 3,500,000). Additional charges may be billed for language editing, USD 75-150 (IDR 1,000,000-2,000,000).

**Reprints** The sample journal reprint is only available by special request. Additional copies may be purchased when ordering by sending back the uncorrected proofs by email.

**Manuscript preparation** Manuscript is typed on A4 (210x297 mm<sup>2</sup>) paper size, in a single column, single space, 10-point (10 pt) Times New Roman font. The margin text is 3 cm from the top, 2 cm from the bottom, and 1.8 cm from the left and right. Smaller lettering size can be applied in presenting table and figure (9 pt). Word processing program or additional software can be used, however, it must be PC compatible and Microsoft Word based (.doc or .rft, not .docx). **Scientific names** of species (incl. subspecies, variety, etc.) should be written in italic, except for italic sentence. Scientific name (genera, species, author), and cultivar or strain should be mentioned completely for the first time mentioning it in the body text, especially for taxonomic manuscripts. Name of genera can be shortened after first mentioning, except generating confusion. Name of the author can be eliminated after first mentioning. For example, *Rhizopus oryzae* L. UICC 524, hereinafter can be written as *R. oryzae* UICC 524. Using trivial name should be avoided, otherwise generating confusion. **Biochemical and chemical nomenclature** should follow the order of the IUPAC - IUB. For DNA sequence, it is better used Courier New font. Symbols of standard chemical and abbreviation of chemistry name can be applied for common and clear used, for example, completely written butilic hydroxyl toluene (BHT) to be BHT hereinafter. **Metric measurement** use IS denomination, usage other system should follow the value of equivalent with the denomination of IS first mentioning. Abbreviations set of, like g, mg, mL, etc. do not follow by dot. Minus index (m<sup>-2</sup>, L<sup>-1</sup>, h<sup>-1</sup>) suggested to be used, except in things like "per-plant" or "per-plot". **Equation of mathematics** does not always can be written

down in one column with text, in that case can be written separately. **Number** one to ten are expressed with words, except if it relates to measurement, while values above them written in number, except in early sentence. The fraction should be expressed in decimal. In the text, it should be used "%" rather than "percent". Avoid expressing ideas with complicated sentence and verbiage, and used efficient and effective sentence.

**Title** of the article should be written in compact, clear, and informative sentence, preferably not more than 20 words. Name of author(s) should be completely written. **Name and institution** address should also be completely written with street name and number (location), postal code, telephone number, facsimile number, and email address. Manuscript written by a group, author for correspondence along with address is required. First page of the manuscript is used for writing above information.

**Abstract** should not be more than 200 words. **Keywords** is about five words, covering scientific and local name (if any), research theme, and special methods which used; and sorted from A to Z. All important **abbreviations** must be defined at their first mention. **Running title** is about five words. **Introduction** is about 400-600 words, covering the background and aims of the research. **Materials and Methods** should emphasize on the procedures and data analysis. **Results and Discussion** should be written as a series of connecting sentences, however, for manuscript with long discussion should be divided into subtitles. Thorough discussion represents the causal effect mainly explains for why and how the results of the research were taken place, and do not only re-express the mentioned results in the form of sentences. **Concluding** sentence should be given at the end of the discussion. **Acknowledgments** are expressed in a brief; all sources of institutional, private and corporate financial support for the work must be fully acknowledged, and any potential conflicts of interest are noted.

**Figures and Tables** of maximum of three pages should be clearly presented. Title of a picture is written down below the picture, while title of a table is written above the table. Colored figures can only be accepted if the information in the manuscript can lose without those images; chart is preferred to use black and white images. Author could consign any picture or photo for the front cover, although it does not print in the manuscript. All images property of others should be mentioned source. **There is no appendix**, all data or data analysis are incorporated into Results and Discussions. For broad data, it can be displayed on the website as a supplement.

**References** Author-year citations are required. In the text give the authors name followed by the year of publication and arrange from oldest to newest and from A to Z. In citing an article written by two authors, both of them should be mentioned, however, for three and more authors only the first author is mentioned followed by et al., for example: Saharjo and Nurhayati (2006) or (Boonkerd 2003a, b, c; Sugiyarto 2004; El-Bana and Nijs 2005; Balagadde et al. 2008; Webb et al. 2008). Extent citation as shown with word "cit" should be avoided. Reference to unpublished data and personal communication should not appear in the list but should be cited in the text only (e.g., Rifai MA 2007, pers. com. (personal communication); Setyawan AD 2007, unpublished data). In the reference list, the references should be listed in an alphabetical order (better, if only 20 for research papers). Names of journals should be abbreviated. Always use the standard abbreviation of a journal's name according to the **ISSN List of Title Word Abbreviations** ([www.issn.org/2-22661-LTWA-online.php](http://www.issn.org/2-22661-LTWA-online.php)). The following examples are for guidance.

### Journal:

Saharjo BH, Nurhayati AD. 2006. Domination and composition structure change at hemic peat natural regeneration following burning; a case study in Pelalawan, Riau Province. *Biodiversitas* 7: 154-158.

### Book:

Rai MK, Carpinella C. 2006. Naturally Occurring Bioactive Compounds. Elsevier, Amsterdam.

### Chapter in book:

Webb CO, Cannon CH, Davies SJ. 2008. Ecological organization, biogeography, and the phylogenetic structure of rainforest tree communities. In: Carson W, Schnitzer S (eds) *Tropical Forest Community Ecology*. Wiley-Blackwell, New York.

### Abstract:

Assaeed AM. 2007. Seed production and dispersal of *Rhazya stricta*. 50<sup>th</sup> annual symposium of the International Association for Vegetation Science, Swansea, UK, 23-27 July 2007.

### Proceeding:

Alikodra HS. 2000. Biodiversity for development of local autonomous government. In: Setyawan AD, Sutarno (eds.) *Toward Mount Lawu National Park; Proceeding of National Seminary and Workshop on Biodiversity Conservation to Protect and Save Germplasm in Java Island*. Universitas Sebelas Maret, Surakarta, 17-20 July 2000. [Indonesian]

### Thesis, Dissertation:

Sugiyarto. 2004. Soil Macro-invertebrates Diversity and Inter-Cropping Plants Productivity in Agroforestry System based on Sengon. [Dissertation]. Universitas Brawijaya, Malang. [Indonesian]

### Information from internet:

Balagadde FK, Song H, Ozaki J, Collins CH, Barnet M, Arnold FH, Quake SR, You L. 2008. A synthetic *Escherichia coli* predator-prey ecosystem. *Mol Syst Biol* 4: 187. [www.molecularsystemsbiology.com](http://www.molecularsystemsbiology.com)



TERESTRASI EKOSISTEM



Local snake fruit conservation in East Java, Indonesia: Community knowledge and appreciation

NOVITA K. INDAH, SERAFINAH INDRIYANI, ESTRI LARAS ARUMINGTYAS, RODIYATI AZRIANINGSIH



Effects of *Caulerpa lentillifera* added into culture media on the growth and nutritional values of *Phoronima pacifica*, a natural fish-feed crustacean

VIVI ENDAR HERAWATI, PINANDOYO, RESTIANA WISNU ARIYATI, NURMANITA RISMANINGSIH, SETO WINDARTO, SLAMET BUDI PRAYITNO, Y.S. DARMANTO, OCKY KARNA RADJASA



✓ Endophytic bacteria associated with rice roots from suboptimal land as plant growth promoters

NUR PRIHATININGSIH, HERU ADI DJATMIKO, PUJI LESTARI



Potential analysis of location, socio-culture and biodiversity as ecotourism attraction in Valentine Bay on Buano Island, West Seram, Maluku, Indonesia

Show all

# Endophytic bacteria associated with rice roots from suboptimal land as plant growth promoters

NUR PRIHATININGSIH<sup>1,✉</sup>, HERU ADI DJATMIKO<sup>1</sup>, PUJI LESTARI<sup>2</sup>

<sup>1</sup>Faculty of Agriculture, Universitas Jenderal Soedirman. Jl. Dr. Soeparno No. 61, Purwokerto Utara, Banyumas 53123, Central Java, Indonesia.

Tel./fax.: +62-281-638791, ✉email: nur.prihatiningsih@unsoed.ac.id

<sup>2</sup>Faculty of Mathematics and Natural Sciences, Universitas Jenderal Soedirman. Jl. Dr. Soeparno No. 61, Purwokerto Utara, Banyumas 53123, Central Java, Indonesia

Manuscript received: 16 November 2020. Revision accepted: 28 December 2020.

**Abstract.** Prihatiningsih N, Djatmiko HA, Lestari P. 2021. Endophytic bacteria associated with rice roots from suboptimal land as plant growth promoters. *Biodiversitas* 22: 432-437. Endophytic bacteria directly promoted plant growth and indirectly control the pathogens. This research was aimed to evaluate endophytic bacteria associated with rice root and their activity to promote plant growth and to control rice diseases. The study was conducted at the Laboratory of Plant Protection and experimental farm Faculty of Agriculture Jenderal Soedirman University, from April to August 2020. The endophytic bacterial from suboptimal land were evaluated for promoting plant growth with soaking seed before seedling and spraying them at 10, 20, and 30 days after transplanting. The experiment was arranged with six replications and four treatments namely control (untreated endophytic bacteria) SM1 (endophytic bacteria isolate from Somagede); SB1 and SB3 (from Sumbang). *Xanthomonas oryzae* pv. *oryzae* was nature inoculation because the experiment location is in the endemic category of bacterial leaf blight. The variables observed were the plant growth components i.e plant height, number of tillers, number of panicles, incubation period, disease intensity, infection rate and effectiveness control. The result of this research shows that endophytic bacteria from Somagede (SM1) is the best to enhance plant height and number of tillers, and suppress disease intensity, and delay of incubation period.

**Keywords:** Endophytic bacteria, plant growth, rice, *Xanthomonas oryzae* pv. *oryzae*

## INTRODUCTION

Endophytic bacteria living in plant tissue are non-pathogens, and produce compounds that help plant growth. Endophytic bacteria reach sufficient populations and are capable of producing plant growth-promoting compounds such as IAA (*indole acetic acid*) and gibberellins. Endophytic bacteria are associated with plants to help the absorption of nutrients because they are able to dissolve phosphate into a form available to plants (Adesemoye et al. 2009; Glick 2012; Dawwam et al. 2013). They are able to produce siderophore to chelate iron into iron-siderophore bonds available to plants. Like rhizobacteria, according to Kesaulya et al. (2015), they are able to act as biostimulants (produce phytohormones), biofertilizers (to help absorb nutrients for plants), and bioprotectants (suppress disease). Endophytic bacteria indirectly affect plants as a bioprotectant or biocontrol to several plant diseases (Berg et al. 2005; Muthukumar et al. 2017).

The major mechanism of endophytic bacteria in plants are (i) to assist nutrient availability and uptake (ii) to enhance stress tolerance (iii) to provide disease resistance (Ryan et al. 2008; Hamilton et al. 2012). Meanwhile, induced disease resistance activities are allied with the abilities to produce secondary metabolites, such as antibiotics or chitinase (Christina et al. 2013; Pieterse et al. 2014; Wang et al. 2014). Due to their beneficial functions such as plant growth promotion and disease control, endophytes can be used in the form of bio-formulations

(seed treatment, soil application, and seedling dip) in agriculture.

Bacterial leaf blight caused by *Xanthomonas oryzae* pv. *oryzae* (Xoo) is an important disease because it causes damage to leaves and can cause yield losses of up to 74% in Southeast Asia (Sarra et al. 2010). The severity of the disease depends on the virulence of the bacterial isolate which causes two primary symptoms, namely blight (seed blight) and crackle (wilt phase). During the blight phase bacteria enter through wounds or water pores on the edges and tops of the leaves causing systemic symptoms, resulting in irregular yellow wet spots, with wavy edges, irregular leaves and spots starting from the edge near the tip of the leaf (Singh et al. 2013). Xoo is a rod-shaped bacteria 0.55-0.75 x 1.35-2.17 µm, with a single polar flagellum, not spore-forming, motile, single or paired, and sometimes chained, a Gram-negative, the colony is light yellow. More than 30 races were found with sub-groups within the species (Furutani et al. 2009).

Efforts to improve plant growth and control bacterial leaf blight have been carried out by using antagonistic bacteria from the rice rhizosphere, namely *Bacillus subtilis* B1 in an organic fertilizer formula (Djatmiko et al. 2017). The leaf blight is classified as vascular disease, so it is difficult to control only with antagonistic bacteria. It is necessary to do this with endophytic bacteria that can infect the tissue, and interact directly with the Xoo. The research previously showed control of bacterial leaf blight on-screen house with the effectiveness of more than 60%

(Prihatiningsih et al. 2020a). Suboptimal land development for agriculture is generally faced with several problems, including high soil acidity and toxicity of Fe and Al as well as deficiency of nutrients N, P, K, Ca, and Mg. Therefore, improvements need to be done to the condition of soil biology and chemistry, using endophytic bacteria. Suboptimal land spread on the embankment of the river, the lower part of the river terraces, alluvial depression, plain fluvial marine, and peat (Sirappa and Titahena 2014). The specification of this research is the application of endophytic bacteria associated with rice roots as antagonistic to improve plant growth. This research aimed to evaluate endophytic bacteria associated with rice root and their activity to promote the plant growth directly and to control rice bacterial leaf blight indirectly.

## MATERIALS AND METHODS

### Preparations of endophytic bacterial associated of rice root (EBARr) from suboptimal land

Endophytic bacteria from root samples of healthy rice plants from suboptimal land around Banyumas were prepared from the results of previous research isolation. Initially, 21 isolates were obtained and then selected to produce nine potential endophytic bacterial isolates. Then the nine endophytic bacterial isolates were tested in vitro for inhibition against Xoo and in the screen house, the endophytic bacteria were tested to suppress bacterial leaf blight (Prihatiningsih et al. 2020a). The results of the two series of studies were 3 isolates that were able to suppress disease, namely SM1 (Somagede 1 isolate), SB1 (Sumbang 1), and SB3 (Sumbang 3). Furthermore, in this study, the three endophytic bacteria were tested in rice fields.

### The evaluation of endophytic bacterial associated of rice root (EBARr) to promote plant growth with observed of promoter component

#### *Hydrogen Cyanide (HCN) production test*

The nine endophytic bacterial isolates were tested as HCN producers. The paper disc was immersed in a 0.5% picric acid solution in a 2% sodium carbonate solution, placed on a NB medium containing 0.44% glycine. Then, it was affixed to the test tube wall. Endophytic bacteria producing HCN gas are shown by the change of the paper disc color from yellow to brown (Reetha et al. 2014).

#### *The siderophore production test*

The siderophore production test was carried out with SD-CASA medium, dark blue, 2% agar w/v in order to be added. As much as 10 ml of CAS-blue agar was the basis of the plate. After being solidified, it was coated with 6 ml of YPGA (*yeast extract peptone glucose agar*) medium. Then, the plate was incubated overnight at 32°C. Ten µl of supernatant endophytic bacteria was dropped on a paper disc, forming an orange or pink zone (Hu and Hu 2011).

#### *Testing IAA production*

Endophytic bacteria isolates were grown on slanted NA medium with and without L-Tryptophan (0.5%). After 2

days of incubation at room temperature then, 100 µl of Salkowski reagent A was dropped (2% 0.5 FeCl<sub>3</sub> in 35% HClO<sub>4</sub> solution). Colonies producing IAA show a change in color to pink (Mohite 2013).

#### *The phosphate solubility activity test*

The phosphate solubility activity test was assessed qualitatively using *potato-dextrose yeast extract agar* (PDYA)-CaP. Each bacterial culture was scratch inoculated in PDYA-CaP medium at 4 places, incubated at 28 ± 2°C for 10 days. The clear zone shows that the endophytic bacteria is capable of phosphate solubility (Khan et al. 2009).

#### *Resistance test of isolates to antibiotics*

The resistance test of isolates against antibiotics was carried out by growing endophytic bacteria on the NA medium by means of the pour plate method, dipping the sterile paper disc pieces in antibiotics after draining and then placing them on the bacterial medium, incubated for 24 hours. If a clear zone is formed, it means that the endophytic bacteria is sensitive to these antibiotics, while those that do not form a clear zone mean that they are resistant to these antibiotics (Yasmin et al. 2009).

### Testing of endophytic bacteria to promote plant growth

Observation of the growth components of rice plants is intended as a direct effect of endophytic bacteria on plants, including plant height and the number of tillers, number of panicles. The study was conducted with a completely randomized block design with 4 treatments and 6 replications. The treatments were controlled (untreated endophytic bacteria), B: SM1 (endophyte bacteria Somagede 1 isolate), C: SB1 (endophytic bacteria Sumbang 1 isolate), and D: SB3 (endophytic bacteria Sumbang 3 isolate). The treatment of endophytic bacteria was started by soaking the seeds before seedling for 1 night, followed by spraying the plants at 20, 30, and 40 days after planting. The observed variables were plant height, number of tillers, and number of panicle.

### The activity of EBARr as biocontrol of bacterial leaf blight

Endophytic bacteria as disease control as an indirect effect is carried out in bacterial leaf blight endemic areas and uses varieties that are susceptible-tolerant to bacterial leaf blight, namely Ciherang. Treatment and design such as testing on plant growth. The endophytic bacterial density was 10<sup>9</sup> cfu/mL. Variables observed were components of the pathosystem (incubation period, disease intensity, infection rate, control effectiveness). Observation of disease intensity using the formula  $DI = \frac{\sum (n \times v)}{Z \times N} \times 100\%$ , DI = disease intensity, n = number of plants from each scale category, v = scale category, N = number of plants observed, Z = highest category value. Xoo's attack category is: 0 = healthy, 1 = damage scale 1–5%, 3 = damage scale 6–12%, 5 = damage scale 13–25%, 7 = damage scale 26–50%, 9 = scale 51–100% damage (Djarmiko et al. 2011). The rate of infection is calculated based on the proportion of diseased tissue obtained from the value of disease intensity with the van der Plank

formula (1963),  $X_t = X_o \cdot e^{rt}$ , where  $X_t$  = proportion of diseased intensity at time  $t$  and  $X_o$  = proportion of diseased intensity at initial time,  $e$  = logarithmic number,  $r$  = infection rate and  $t$  = interval of observation time. Because of the bacterial leaf blight is a simple interest disease, calculate  $r = 2,3/t (\log 1/(1-X_t) - \log 1/(1-X_o))$  unit/day.

### Data analysis

Data at the laboratory experiment were analyzed by descriptive methods and then communicated and compared to references. Data from experimental exfarm were analyzed by ANOVA, if the effect is significant, it's continued with LSD 5%.

## RESULTS AND DISCUSSION

Based on the characterization of endophytic bacteria in previous studies and by heating treatment in an oven at 80°C for 20 minutes of the suspension of rice root endophytic bacteria in the 1st dissolution, most of the endophytic bacteria obtained showed Gram-positive, in the form of short rod shape - rod shape, forming endospores, then grouped into the genus *Bacillus* sp. (Gordon et al. 1973; Prihatiningsih et al. 2020a).

### Plant growth promoter component of EBARr

The component of plant growth promoter showed from nine endophytic bacteria that could produce HCN, siderophore, IAA, and phosphate solubility as qualitative tests (Table 1). The various respond of four assays showed the relative of qualitative. Figure 1 showed HCN production from nine endophytic bacteria before reaction to produce HCN (A) with yellow and (B) after production HCN, the paper disc become brown.

Endophytic bacteria are able to stimulate plant growth by observing plant growth-promoting components, namely bacteria as a producer of HCN, siderophore, IAA, and as a phosphate solubility (Table 1). The response of plant growth promoter components was varied from the nine isolates of endophytic bacteria that were tested as a result of the selection from the first year. Bacteria that are able to produce HCN show that these bacteria are capable of being biocontrol. This is because the role of HCN is to inhibit the proliferation or multiplication of pathogens, stimulate the formation of root hairs, thereby help to increase nutrient absorption (Olanrewaju et al. 2017).


Figure 2 showed the six endophytic bacteria produced siderophore with orange zone except for the Kr7, SR1 dan SB3, the medium still blue. The bacterial siderophore test formed an orange zone indicating that the type of siderophore was hydroxamate (Ahmed and Holmstrom 2014; Prihatiningsih et al. 2017). Siderophores are low molecular weight compounds produced by bacteria and are capable of chelating iron into a form of siderophore- $Fe^{3+}$  bonds available to plants. This siderophore is also one of the biocontrol mechanisms by binding to  $Fe^{3+}$  so the pathogen is limited in the need for  $Fe^{3+}$ . Therefore, the pathogen is inhibited from reproducing to reach the rhizosphere due to iron deficiency (Glick 2012; Prihatiningsih et al. 2017).

The endophytic bacteria were able to produce IAA which was marked with pink color in the endophytic bacteria culture which was dripped with Salkowski reagent (Figure 3). This indicates that endophytic bacteria are capable of producing the plant hormone IAA. Quantitative analysis of IAA production has conducted that SR7 and SB1 isolate were active at 79.33 and 75.88 ppm respectively (Prihatiningsih et al. 2020a). IAA functions for cell division and elongation stimulates seed and tuber germination increases the rate of xylem development and root development. IAA also functions as a vegetative growth controller, initiating the formation of adventitious and lateral roots, so that plant growth is better (Glick 2012).

**Table 1.** Plant growth components of nine rice root endophytic bacteria

Isolate code	HCN production	Siderophore production	IAA production	Phosphate solubility
Kr4	+	++	+	+
Kr5	+	+	+	+
Kr7	+	-	-	-
SR1	+	-	+	+
SR5	+	+	+	+
SR7	+	++	+	+
SM1	+	++	+	++
SB1	+	+	++	+
SB3	+	-	+++	++

Note: +: produce; ++ and +++: produce with more color (IAA) ++: more strong of produce siderophore and phosphate solubility, -: unproduced; Kr: Karangwangkal, SR: Serayu, SM: Somagede, SB: Sumbang



**Figure 1.** HCN production test of the nine endophytic bacteria (A) Before reaction and (B) After reaction as HCN production with brown color of paper



The ability of endophytic bacteria as a phosphate solubility is shown by a positive reaction in Table 2 and Figure 3, with the formation of a clear zone. Solubilization of different forms of phosphate by endophytic bacteria associated with maize can be increasing its availability for plants to promote plant growth and increase the yield (Mugiastuti et al. 2020). The ability of bacteria as a phosphate solubility helps to provide nutrients available for plants because the phosphate present in the soil in a form that is not available to plants becomes available (Olanrewaju et al. 2017).


The resistance of endophytic bacteria to antibiotics (Table 2, Figure 3) is very important as a pathogen biocontrol mechanism because it is exposed to many microorganisms that produce various antibiotics in the soil. Antibiotic-resistant endophytic bacteria will be able to survive so that they are able to compete in space and become antagonistic to pathogens (Ulloa-Ogaz et al. 2015). Isolates that are resistant to antibiotics as shown in Figure 3. SR7 isolates are resistant to Rifampicin antibiotics which are characterized by no zone formation, meaning that the growth of SR7 endophytic bacteria still occurs around the antibiotic. On the other hand, the SR5 isolate was not resistant or sensitive to the Kanamycin antibiotic which was indicated by the formation of a clear zone around the

filter paper treated with the Kanamycin antibiotic, meaning that there was no growth around the antibiotic. This is in accordance with the opinion of Yasmin et al. (2009) that the IAR test is needed to identify the endophytic bacteria resistant or sensitive to antibiotics. IAR can also be used to genotypically identify bacterial species.


**Table 2.** Intrinsic antibiotic resistance (IAR) test of rhizobacterial isolates

Isolate	Rifampicin		Kanamycin	
	Reaction	Diameter of zone (mm)	Reaction	Diameter of zone (mm)
Kr4	-	4	-	8
Kr5	-	4	-	8
Kr7	-	4	-	8
SR1	+	0	+	0
SR5	-	5	-	8
SR7	+	1	-	6
SM1	-	6	-	8
SB1	-	7	-	9
SB3	+	1	-	7

Note: +: resistance= not zone, very small (1 mm), -: sensitive: zone formed. Not zone: antibiotic resistance (+) means that be still growing in the medium with the antibiotic



**Figure 2.** Siderophore production of the endophytic bacteria



**Figure 3.** A. IAA production (a. produce of IAA and b. unproduce of IAA); B. Phosphate solubility; C. Resistance to antibiotics of the endophytic bacteria (a. SR7 resistance Rifampicin, and b. SR5 sensitive to Kanamycin)



**Table 3.** The component of plant growth with endophytic bacteria application

Isolate code	Plant height (cm)	Number of tillers	Number of panicle/clump
A (control)	96.6b	13.3b	9.875
B (SM1)	101.3a	12.6b	13.005
C (SB1)	98.5b	15.3ab	12.125
D (SB3)	98.8b	18.3a	18.875

**Table 4.** Components of plant pathosystems against bacterial leaf blight

Isolate code	Incubation period (dat)	Disease intensity (%)	Effective-ness (%)	Infection rate (unit/days)
A (control)	24	23.13a	0	0.0469
B (SM1)	30.2	10.17b	56.03	0.0435
C (SB1)	36.3	9.07b	60.79	0.0246
D (SB3)	34.2	9.27b	59.92	0.0454

### Endophytic bacteria to promote plant growth

Based on the result of the laboratory experiment than the endophytic bacteria selected to promote plant growth in the exfarm showed in Table 3, in which three endophytic bacteria can promote the increase plant height, the number of tiller, and the number of panicle. The SM1 (Somagede isolate) was able to increase plant height compared to other endophytic bacteria treatments and controls, while isolate Sumbang 3 (SB3) increased the number of tiller. This is in accordance with the opinion of Glick (2012) that endophytic bacteria are able to stimulate plant growth because they produce phytohormones such as IAA which can stimulate plant growth.

### Suppressing bacterial leaf blight by EBARr

The observation of the pathosystem components of rice bacterial leaf blight conducted at endemic areas showed that the treatment of endophytic bacteria was different from the control (Table 4). However, between endophytic bacteria was not significantly different with control, effectiveness ranging from 56.03 to 60.79%. The endophytic bacteria can delay incubation period, and suppress the infection rate. Endophytic bacteria as well as the rhizosphere bacteria are able to control bacterial disease with antibiosis mechanism as bacteriostatic which produce hydrolytic enzyme like amylase, protease, and lipase (Prihatiningsih et al. 2020b; Mugiastuti et al. 2020).

Nagendran et al. (2013) stated that endophytic bacteria were able to control bacterial leaf blight on rice with a low disease intensity of 2.80% compared to without treatment of endophytic bacteria with a disease intensity of 19.82%. Four isolates, OS52, OS40, OS23, and OS53, which were identified by nucleotide sequence analysis as *Enterobacter* sp., *B. subtilis*, *Bacillus* sp. and *Pseudomonas putida*, respectively, could increase plant growth and decrease Xoo infection under greenhouse conditions. Seed treatment with these antagonists caused disease to cease in over 60% of

plants. These species have been found as endophytes in rice (Yousefi et al. 2018). Endophytic bacteria are able to control the disease caused by bacteria and fungi because they produce hydrolytic enzyme as well as chitinase, protease similar to rhizospheric bacteria *B. subtilis* from potato rhizosphere (Lestari et al. 2017).

In conclusion, the conclusion based on this research showed that three endophytic bacteria are associated with rice root (EBARr) as prospectives as plant growth promoter, and biocontrol agents to bacterial leaf blight. The inhibition activity of EBARr similar to an exfarm test by suppressing the disease until 60.79% effectiveness. The rice growth enhanced by the application of EBARr with plant height and number of tiller so increase. Furthermore, endophytic bacteria was prospected to develop biofertilizer and biocontrol.

### ACKNOWLEDGEMENTS

Thank you for supporting this research by DRPM Kemenristek for Riset Terapan scheme 2020 with contract number 2/SP2H/AMD/LT/DRPM/2020.

### REFERENCES

- Adesemoye AO, Torbert HA, Kloepper JW. 2009. Plant growth-promoting rhizobacteria allow reduced application rates of chemical fertilizer. *Microb Ecol* 58: 921-929. DOI: 10.1007/s00248-009-9531-y.
- Ahmed E, Holmstrom SJM. 2014. Siderophores in environmental research: roles and applications. *Microb Biotechnol* 7: 196-208. DOI: 10.1111/1751-7915.12117.
- Berg G, Krechel A, Ditz M, Sikora RA, Ulrich A, Hallmann J. 2005. Endophytic and ectophytic potato-associated bacterial communities differ in structure and antagonistic function against plant pathogenic fungi. *FEMS Microbiol Ecol* 51: 215-229. DOI: 10.1016/j.femsec.2004.08.006.
- Christina A, Christopher V, Bhore SJ. 2013. Endophytic bacteria as a source of novel antibiotics: an overview. *Pharmacogn Rev* 7: 11-16. DOI: 10.4103/0973-7847.112833.
- Dawwam GE, Elbeltagy A, Emara HM, Abbas IH, Hassan MM. 2013. Beneficial effect of plant growth-promoting bacteria isolated from the roots of potato plant. *Ann Agric Sci* 58: 195-201. DOI: doi.org/10.1016/j.aos.2013.07.007.
- Djarmiko HA, Prakoso B, Prihatiningsih N. 2011. Penentuan Patotipe dan keragaman genetik *Xanthomonas oryzae* pv. *oryzae* pada tanaman padi di wilayah Karesidenan Banyumas. *Jurnal Hama Penyakit Tumbuhan Tropika* 11 (1): 35-46. [Indonesian]
- Djarmiko HA, Prihatiningsih N, Ismangil. 2017. Pupuk Organik diperkaya *Bacillus subtilis* B1. Paten sederhana dengan nomor pendaftaran paten S00201707795.
- Furutani A, Takaoka M, Sanada H, Noguchi Y, Oku T, Tsuno K, Ochiai H, Tsuge S. 2009. Identification of novel type III secretion effectors in *Xanthomonas oryzae* pv. *oryzae*. *Mol Plant-Microb Interact* 22 (1): 96-106. DOI: 10.1094/MPMI-22-1-0096.
- Glick BR. 2012. Plant growth-promoting bacteria: mechanism and applications. *Scientifica* 2012. DOI: 10.6064/2012/963401.
- Gordon RE, Haynes WC, Pang CH. 1973. *The Genus Bacillus*. Agriculture Research Service, United States Department of Agriculture, Washington DC.
- Hamilton CE, Gundel PE, Helander M, Saikkonen K. 2012. Endophytic mediation of reactive oxygen species and antioxidant activity in plants: a review. *Fungal Div* 54: 1-10. DOI: doi.org/10.1007/s13225-012-0158-9.

- Hu QP, Xu JG. 2011. A Simple double-layered chrome azurol S agar (SD-CASA) plate assay to optimize the production of siderophores by a potential biocontrol agent *Bacillus*. *Afr J Microbiol Res* 5 (25): 4321-4327. DOI: doi.org/10.5897/AJMR11.238.
- Kesaulya H, Baharuddin, Zakaria B, Syatrianty A, Syaiful. 2015. Isolation and physiological characterization of PGPR from potato plant rhizosphere in Medium Land of Buru Island. *Procedia Food Sci* 3: 190-199. DOI: doi.org/10.1016/j.profoo.2015.01.021.
- Khan AA, Jilani G, Akhtar MS, Naqvi SMS, Rasheed. 2009. Phosphorus solubilizing bacteria: Occurrence, mechanisms and their role in crop production. *J Agric Biol Sci* 1 (1): 48-58.
- Lestari P, Prihatiningsih N, Djatmiko, HA. 2017. Partial biochemical characterization of crude extract extracellular chitinase enzyme from *Bacillus subtilis* B298. *IOP Conf Ser: Mater Sci Eng* 172 (2017): 012041. DOI: 10.1088/1757-899X/172/1/012041.
- Mohite B. 2013. Isolation and characterization of indole acetic acid (IAA) producing bacteria from rhizospheric soil and its effect on plant growth. *J Soil Sci Plant Nutr* 13 (3): 638-649. DOI: 10.4067/S0718-95162013005000051.
- Mugiastuti E, Suprayogi, Prihatiningsih N, Soesanto L. 2020. Isolation and characterization of the endophytic bacteria, and their potential as maize disease control. *Biodiversitas* 21 (5): 1809-1815. DOI: 10.13057/biodiv/d210506.
- Muthukumar A, Udhayakumar R, Naveenkumar R. 2017. Role of bacterial endophytes in plant disease control. In *Endophytes: Crop Productivity and Protection* (pp. 133-161). Springer, Cham.
- Nagendran K, Karthikeyan G, Peeran MF, Raveendran M, Prabakar K, Raguchander T. 2013. Management of bacterial leaf blight disease in rice with endophytic bacteria. *World Appl Sci J* 28 (12): 2229-2241. DOI: 10.5829/idosi.wasj.2013.28.12.2009.
- Olanrewaju OS, Glick BR, Babalola OO. 2017. Mechanisms of action of plant growth promoting bacteria. *World J Microbiol Biotechnol* 33: 197. DOI: 10.1007/s11274-017-2364-9.
- Pieterse CM, Zamioudis C, Berendsen RL, Weller DM, Van Wees SC, Bakker PA. 2014. Induced systemic resistance by beneficial microbes. *Annu Rev Phytopathol* 52: 347-375. DOI: 10.1146/annurev-phyto-082712-102340.
- Prihatiningsih N, Djatmiko HA, Lestari P. 2017. Aktivitas siderofor *Bacillus subtilis* sebagai pemacu pertumbuhan dan pengendali patogen tanaman terung. *Jurnal Hama dan Penyakit Tumbuhan Tropika* 17 (2): 170-178. [Indonesian]
- Prihatiningsih N, Djatmiko HA, Lestari P. 2020a. Screening of competent rice root endophytic bacteria to promote rice growth and bacterial leaf blight disease control. *Jurnal Hama dan Penyakit Tumbuhan Tropika* 20 (1): 78-84. [Indonesian]
- Prihatiningsih N, Arwiyanto T, Hadisutrisno B, Widada J. 2020. Characterization of *Bacillus* spp. from rhizosphere of potato Granola variety as an antibacterial against *Ralstonia solanacearum*. *Biodiversitas* 21 (9): 4199-4204. DOI: 10.13057/biodiv/d210934.
- Reetha AK, Pavani SL, Mohan S. 2014. Hydrogen cyanide production ability by bacterial antagonist and their antibiotics inhibition potential on *Macrophomina phaseolina* (Tassi.) Goid. *Int J Curr Microbiol Appl Sci* 3 (5): 172-178.
- Ryan RP, Germaine K, Franks A, Ryan DJ, Dowling DN. 2008. Bacterial endophytes: recent developments and applications. *FEMS Microbiol Lett* 278: 1-9. DOI: 10.1111/j.1574-6968.2007.00918.x.
- Sarra S, Diarra L, Dembele M, Coulibaly MM, Sese Y. 2010. Characterization of bacterial leaf blight epidemic in the Office du Niger (Mali) and search for a sustainable resistance against the pathogen. *Second Africa Rice Congress, Bamako Mali; Innovation and Partnerships to Realize Africa's Rice Potential*.
- Singh AK, Sarma Bk, Singh PK, Nandan R. 2013. Screening of rice (*Oryza sativa* L.) germplasms against *Xanthomonas oryzae* pv. *oryzae*. *J Ecofriendly Agric* 8 (1): 86-88.
- Sirappa MP, Titahena MLJ. 2014. Improvement of suboptimal land productivity approach by land and plant management. *J Trop Soils* 19 (2): 1-11. DOI: http://dx.doi.org/10.5400/jts.2014.v19i2.99-109.
- Ulloa-Ogaz AL, Muñoz-Castellanos LN, Nevárez-Moorillón GV. 2015. Biocontrol of phytopathogens: Antibiotic production as mechanism of control. *Formatex* 2015: 305-309.
- van der Plank, JE. 1963. *Plant Disease: Epidemic and Control*. Academic Press, New York.
- Wang M, Xing Y, Wang J, Xu Y, Wang G. 2014. The role of the chi1 gene from the endophytic bacteria *Serratia proteamaculans* 336x in the biological control of wheat take-all. *Can J Microbiol* 60: 533-540. DOI: 10.1139/cjm-2014-0212.
- Yasmin F, Othman R, Sijam K, Saad MS. 2009. Characterization of beneficial properties of plant growth-promoting rhizobacteria isolated from sweet potato rhizosphere. *Afr J Microbiol Res* 3 (11): 815-821. http://www.academicjournals.org/ajmr.
- Yousefi H, Hassanzadeh N, Behboudi K, Firouzjahi FB. 2018. Identification and determination of characteristics of endophytes from rice plants and their role in biocontrol of bacterial blight caused by *Xanthomonas oryzae* pv. *oryzae*. *Hell Plant Prot J* 11: 19-33. DOI: 10.2478/hppj-2018-0003.