

 Jurnal Kimia
Sains & Aplikasi
Journal of Scientific & Applied Chemistry

**Jurnal Kimia
Sains & Aplikasi**
Journal of Scientific & Applied Chemistry

ISSN: 1410-8917
e-ISSN: 2597-9914

Accredited by the Ministries of Research,
Technology and Higher Education
No: 3/E/KPT/2019

 **Current issue:** [Vol 26, No 3 \(2023\): Volume 26 Issue 3 Year 2023 \(https://ejournal.undip.ac.id/index.php/ksa/issue/current\)](https://ejournal.undip.ac.id/index.php/ksa/issue/current) | [Archives](#)
(<https://ejournal.undip.ac.id/index.php/ksa/issue/archive>). | [Start Submission](#)
(<https://ejournal.undip.ac.id/index.php/ksa/about/submissions>).

 **Starting in 2020, JKSA is published every month** (<https://ejournal.undip.ac.id/index.php/ksa/announcement/view/253>)

 **JKSA was awarded the Sinta-2 accreditation** (<https://ejournal.undip.ac.id/index.php/ksa/announcement/view/225>)

 **Change in publishing frequency** (<https://ejournal.undip.ac.id/index.php/ksa/announcement/view/224>)

Currently, JKSA is accredited with Sinta 2 with a decree no. 3/E/KP/2019 (<https://www.dropbox.com/s/dpa9orzqutotdtx/2019-I-Elektronik.pdf?dl=0>) of the Directorate General of Strengthening for Research and Development, Ministry of Research, Technology and Higher Education, Republic of Indonesia



Jurnal Kimia Sains dan Aplikasi (p-ISSN: [1410-8917](http://issn2.lipi.go.id/issn.cgi?daftar&1180430817&1&&) (<http://issn2.lipi.go.id/issn.cgi?daftar&1180430817&1&&>)) and e-ISSN: [2597-9914](http://issn2.lipi.go.id/issn.cgi?daftar&1505120918&1&&) (<http://issn2.lipi.go.id/issn.cgi?daftar&1505120918&1&&>)) is an open access and peer reviewed journal published by Department of Chemistry, Diponegoro University. This journal is published every month, twelve times per year in the end of month. JKSA publishes research papers, review and short communication in field of Chemistry.

Jurnal Kimia Sains dan Aplikasi accept articles written in Bahasa Indonesia and English (preferable). Although we accept the manuscript in Indonesian, however we encourage the authors to write manuscripts in English.

[illegible]

Last update: 23rd March 2023

Number of documents: 612

Cites in Google Scholar: 2069

Google Scholar IF: 2069/612 = 3.381

h-Index: 16, i10-Index: 60

Scopus: 338 citations form 194 articles

Google Scholar citation [here](https://scholar.google.co.id/citations?user=fzjW0kgAAAAJ&hl=en)
(<https://scholar.google.co.id/citations?user=fzjW0kgAAAAJ&hl=en>).

Scopus citation [here](https://www.scopus.com/results/results.uri?)
(<https://www.scopus.com/results/results.uri?>

```
src=dm&sort=plf-  
f&sid=80b5bb6cd79162345cd08f711c8181a2&sot=b&sdt=b&sl=74&s=  
f&ws=r-f&ps=r-f&cs=r-  
f&origin=resultslist&zone=queryBar).
```

Journal Content

Search


Search Scope

All 


Search

Browse

- **By Issue**
(<https://ejournal.undip.ac.id/index.php/ksa/issue/archive>)
- **By Author**
(<https://ejournal.undip.ac.id/index.php/ksa/search/authors>)
- **By Title**
(<https://ejournal.undip.ac.id/index.php/ksa/search/titles>)
- **Other Journals**
(<https://ejournal.undip.ac.id/index.php/index/search>)
- **Categories**
(<https://ejournal.undip.ac.id/index.php/index/search/categories>)




(<https://ejournal.undip.ac.id/index.php/ksa/stats>)



Visitor map

Article abstract views (total: 1446017)



(<https://ejournal.undip.ac.id/index.php/ksa/stats>)

Recent articles (#tabRecent)

Most cited articles (#tabCited).

Contact (#tabContact).

Vol 26, No 3 (2023): Volume 26 Issue 3 Year 2023

Extraction of Silica (SiO₂) from Rare-Earth Metal Zircon (ZrSiO₄) as Lithium-Ion Battery Anode Material

(<https://ejournal.undip.ac.id/index.php/ksa/article/view/49011>)

Citations

?

(<https://badge.dimensions.ai/details/doi/10.14710/jksa.26.3.79-84?domain=https://ejournal.undip.ac.id>)

| Language: **EN (#)** | DOI: **[10.14710/jksa.26.3.79-84](https://doi.org/10.14710/jksa.26.3.79-84)**

(<https://doi.org/10.14710/jksa.26.3.79-84>)

PDF

(<https://ejournal.undip.ac.id/index.php/ksa/article/view/49011/23877>)

79-84

Putri Lestari, Rhiska Aria Berliani, Erlangga Aria Pratama, Ridwan Assabila Al-Jannata, Desy Vita Pratiwi, Hendri Widiyandari

Tyrosinase Inhibitory of Silver Nanoparticles Synthesized using Morus Nigra Leaves Extract

(<https://ejournal.undip.ac.id/index.php/ksa/article/view/51074>)

Citations

?

(<https://badge.dimensions.ai/details/doi/10.14710/jksa.26.3.85-90?domain=https://ejournal.undip.ac.id>)

| Language: **EN (#)** | DOI: **[10.14710/jksa.26.3.85-90](https://doi.org/10.14710/jksa.26.3.85-90)**

(<https://doi.org/10.14710/jksa.26.3.85-90>)

PDF

(<https://ejournal.undip.ac.id/index.php/ksa/article/view/51074/23878>)

85-90

Tunas Alam, Ricky Ardiansyah, Sharfina Maulidayanti, Deva Azvara, Frida Octavia Purnomo, Dicky Annas

Thermal Properties of Acetylated Betung Bamboo (Dendrocalamus asper) Pulp – Polypropylene Biocomposites

(<https://ejournal.undip.ac.id/index.php/ksa/article/view/52082>)

Citations

?

(<https://badge.dimensions.ai/details/doi/10.14710/jksa.26.3.91-100?domain=https://ejournal.undip.ac.id>)

| Language: **EN (#)** | DOI: **[10.14710/jksa.26.3.91-100](https://doi.org/10.14710/jksa.26.3.91-100)**

(<https://doi.org/10.14710/jksa.26.3.91-100>)

PDF

(<https://ejournal.undip.ac.id/index.php/ksa/article/view/52082/23879>)

91-100

Wida Banar Kusumaningrum, Sukma Surya Kusumah, Ismadi Ismadi, Rochmadi Rochmadi, Subyakto Subyakto

An Eco-Friendly Synthesis of Silver Nanoparticles Using Extract of Capsicum chinense Jacq. as A Bio-reducing Agent

(<https://ejournal.undip.ac.id/index.php/ksa/article/view/53326>)

Citations

?

(<https://badge.dimensions.ai/details/doi/10.14710/jksa.26.3.101-108?domain=https://ejournal.undip.ac.id>)

| Language: **EN (#)** | DOI: **[10.14710/jksa.26.3.101-108](https://doi.org/10.14710/jksa.26.3.101-108)**

(<https://doi.org/10.14710/jksa.26.3.101-108>)

PDF

(<https://ejournal.undip.ac.id/index.php/ksa/article/view/53326/23880>)

101-108

Boyfannie Ivan Putra, Hermin Sulistyarti, Rurini Retnowati, Mohamad Rafi

Quantitative Structure-Activity Relationship of 3-Thiocyanate-1H-Indoles Derived Compounds as Antileukemia by AM1, PM3, and RM1 Methods

(<https://ejournal.undip.ac.id/index.php/ksa/article/view/53504>)

Citations

?

(<https://badge.dimensions.ai/details/doi/10.14710/jksa.26.3.109-117?domain=https://ejournal.undip.ac.id>)

| Language: **EN (#)** | DOI: **[10.14710/jksa.26.3.109-117](https://doi.org/10.14710/jksa.26.3.109-117)**

(<https://doi.org/10.14710/jksa.26.3.109-117>)

PDF

(<https://ejournal.undip.ac.id/index.php/ksa/article/view/53504/23881>)

109-117

Ponco Iswanto, Irvan Maulana Firdaus, Ahmad Fawwaz Dafaulhaq, Ahmad Ghifari Ramadhani, Maylani Permata Saputri, Heny Ekowati

Editorial Policies

- Focus and Scope
- Section Policies
- Peer Review Process / Policy
- Article Processing Charges
- Publication Ethics and Malpractice Statement
- Content Licence
- Indexing
- Crossmark
- Article Withdrawal
- Open Access Policy
- Privacy Statement

About the Journal

- Editorial Team
- Peer Reviewers
- Publisher
- Journal History
- Article Metrics
- Contact
- Sitemap
- About this Publishing System

Announcements

- See complete announcements

Information

- For Readers
- For Authors
- For Librarians
- [UNDIP journal portal](#)
- [UNDIP journal portal 2](#)
- [UNDIP journal portal 3](#)

Visitor: 390900 [View My Stats](#)Jurnal Kimia Sains dan Aplikasi is indexed in:

This work is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](#).

Copyright ©2023 [Universitas Diponegoro](#). Powered by [Public Knowledge Project OJS](#) and [Mason Publishing OJS theme](#).

<https://ejournal.undip.ac.id/index.php/ksa>

2/2



Jurnal Kimia

Sains & Aplikasi

Jurnal Kimia

Sains & Aplikasi

Journal of Scientific & Applied Chemistry

OPEN ACCESS

PEER REVIEW

ISSN: 1410-8917

e-ISSN: 2597-9914

Accredited by the Ministries of Research, Technology and Higher Education

No: 3/E/KPT/2019

[Home \(https://ejournal.undip.ac.id/index.php/ksa/index/\)](https://ejournal.undip.ac.id/index.php/ksa/index/) / [About the Journal \(https://ejournal.undip.ac.id/index.php/ksa/about/\)](https://ejournal.undip.ac.id/index.php/ksa/about/) / [Editorial Team \(https://ejournal.undip.ac.id/index.php/ksa/about/editorialTeam/\)](https://ejournal.undip.ac.id/index.php/ksa/about/editorialTeam/)

Editorial Team

People > [Editorial Team \(https://ejournal.undip.ac.id/index.php/ksa/about/editorialTeam/\)](https://ejournal.undip.ac.id/index.php/ksa/about/editorialTeam/) | [Peer Reviewers \(https://ejournal.undip.ac.id/index.php/ksa/about/displayMembership/422/1\)](https://ejournal.undip.ac.id/index.php/ksa/about/displayMembership/422/1)

Editor in Chief



Adi Darmawan (ScopusID: [55953897600 \(http://www.scopus.com/authid/detail.uri?authorId=55953897600\)](http://www.scopus.com/authid/detail.uri?authorId=55953897600))
[id \(http://orcid.org/0000-0001-5744-5789\)](http://orcid.org/0000-0001-5744-5789) [P \(https://publons.com/researcher/3141033\)](https://publons.com/researcher/3141033) Chemistry Department, Faculty of Sciences and Mathematics, Diponegoro University Jl. Prof. Soedarto, SH., Tembalang, Semarang, Indonesia

Associate editors

- 
- Dr. Choiril Azmiyawati (ScopusID: [55543514300 \(http://www.scopus.com/authid/detail.uri?authorId=55543514300\)](http://www.scopus.com/authid/detail.uri?authorId=55543514300))
[id \(http://orcid.org/0000-0002-4143-9832\)](http://orcid.org/0000-0002-4143-9832) Chemistry Department, Faculty of Sciences and Mathematics, Diponegoro University Jl. Prof. Soedarto, SH., Tembalang, Semarang, Indonesia
- 
- Didik Setiyo Widodo (ScopusID: [57195404137 \(http://www.scopus.com/authid/detail.uri?authorId=57195404137\)](http://www.scopus.com/authid/detail.uri?authorId=57195404137))
[id \(https://orcid.org/0000-0001-8411-9700\)](https://orcid.org/0000-0001-8411-9700) Chemistry Department, Faculty of Sciences and Mathematics, Diponegoro University Jl. Prof. Soedarto, SH., Tembalang, Semarang, Indonesia
- 
- Mr Ismiyarto Ismiyarto (ScopusID: [57217110370 \(http://www.scopus.com/authid/detail.uri?authorId=57217110370\)](http://www.scopus.com/authid/detail.uri?authorId=57217110370))
[id \(http://orcid.org/0000-0002-3939-3433\)](http://orcid.org/0000-0002-3939-3433) Chemistry Department, Faculty of Sciences and Mathematics, Diponegoro University Jl. Prof. Soedarto, SH., Tembalang, Semarang, Indonesia
- 
- Dr. Mukhammad Asy'ari (ScopusID: [56117266100 \(http://www.scopus.com/authid/detail.uri?authorId=56117266100\)](http://www.scopus.com/authid/detail.uri?authorId=56117266100))
[id \(http://orcid.org/0000-0002-3489-1644\)](http://orcid.org/0000-0002-3489-1644) Chemistry Department, Faculty of Sciences and Mathematics, Diponegoro University Jl. Prof. Soedarto, SH., Tembalang, Semarang, Indonesia
- 
- Dr. Nor Basid Adiwibawa Prasetya (ScopusID: [56574376400 \(http://www.scopus.com/authid/detail.uri?authorId=56574376400\)](http://www.scopus.com/authid/detail.uri?authorId=56574376400))
[id \(http://orcid.org/0000-0002-6956-3667\)](http://orcid.org/0000-0002-6956-3667) Chemistry Department, Faculty of Sciences and Mathematics, Diponegoro University Jl. Prof. Soedarto, SH., Tembalang, Semarang, Indonesia
- 
- Dr. Yayuk Astuti (ScopusID: [57100033100 \(http://www.scopus.com/authid/detail.uri?authorId=57100033100\)](http://www.scopus.com/authid/detail.uri?authorId=57100033100))
[id \(http://orcid.org/0000-0002-2107-3829\)](http://orcid.org/0000-0002-2107-3829) Chemistry Department, Faculty of Sciences and Mathematics, Diponegoro University Jl. Prof. Soedarto, SH., Tembalang, Semarang, Indonesia

International Advisory Editorial Boards

- 
- Dr. Amin Fatoni (ScopusID: [55488648900 \(http://www.scopus.com/authid/detail.uri?authorId=55488648900\)](http://www.scopus.com/authid/detail.uri?authorId=55488648900))
[id \(http://orcid.org/0000-0002-6550-2461\)](http://orcid.org/0000-0002-6550-2461) Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia
- 
- Prof. Dr Fitria Rahmawati (ScopusID: [36053591500 \(http://www.scopus.com/authid/detail.uri?authorId=36053591500\)](http://www.scopus.com/authid/detail.uri?authorId=36053591500))
[id \(http://orcid.org/0000-0002-3145-9063\)](http://orcid.org/0000-0002-3145-9063) Department of Chemistry, Faculty of Mathematics and Natural Sciences, Sebelas Maret University, Indonesia
- 
- Dr Gaurav A Bhaduri (ScopusID: [28367493600 \(http://www.scopus.com/authid/detail.uri?authorId=28367493600\)](http://www.scopus.com/authid/detail.uri?authorId=28367493600))
[id \(http://orcid.org/0000-0002-7714-8877\)](http://orcid.org/0000-0002-7714-8877) Indian Institute of Technology Jammu (IIT JMU), India
- 
- Dr. Guozhao Ji (ScopusID: [55262553900 \(http://www.scopus.com/authid/detail.uri?authorId=55262553900\)](http://www.scopus.com/authid/detail.uri?authorId=55262553900))
[id \(https://orcid.org/0000-0003-4556-2675\)](https://orcid.org/0000-0003-4556-2675) School of Environmental Science and Technology, Dalian University of Technology Dalian, Liaoning, China
- 
- Dr. Ibrahim A. I. Hassan (ScopusID: [55652057500 \(http://www.scopus.com/authid/detail.uri?authorId=55652057500\)](http://www.scopus.com/authid/detail.uri?authorId=55652057500))
Department of Chemistry, South Valley University Qena, Egypt, Egypt
- 
- Dr. Mus'ab Abdul Razak (ScopusID: [38961852200 \(http://www.scopus.com/authid/detail.uri?authorId=38961852200\)](http://www.scopus.com/authid/detail.uri?authorId=38961852200))
[id \(http://orcid.org/0000-0001-5120-1345\)](http://orcid.org/0000-0001-5120-1345) Department of Chemical and Environmental Engineering, Faculty of Engineering, Universiti Putra Malaysia, Malaysia

Journal Administrator



Aulia Ekadenti
Chemistry Department, Faculty of Sciences and Mathematics, Diponegoro University Jl. Prof. Soedarto, SH., Tembalang, Semarang, Indonesia

Currently, JKSA is accredited with Sinta 2 with a decree [no. 3/E/KP/2019 \(https://www.dropbox.com/s/dpa9orzqutotdtx/2019-I-Elektronik.pdf?dl=0\)](https://www.dropbox.com/s/dpa9orzqutotdtx/2019-I-Elektronik.pdf?dl=0) of the Directorate General of Strengthening for Research and Development, Ministry of Research, Technology and Higher Education, Republic of Indonesia



Last update: 23rd March 2023
Number of documents: 612
Cites in Google Scholar: 2069
Google Scholar IF: 2069/612 = 3.381
h-Index: 16, i10-Index: 60
Scopus: 338 citations form 194 articles
Google Scholar citation [here \(https://scholar.google.co.id/citations?user=fzjW0kgAAAAJ&hl=en\)](https://scholar.google.co.id/citations?user=fzjW0kgAAAAJ&hl=en)
Scopus citation [here \(https://www.scopus.com/results/results.uri?src=dm&sort=plf-f&sid=80b5bb6cd79162345cd08f711c8181a2&sot=b&sdt=b&sl=74&sf&ws=r-f&ps=r-f&cs=r-f&origin=resultslist&zone=queryBar\)](https://www.scopus.com/results/results.uri?src=dm&sort=plf-f&sid=80b5bb6cd79162345cd08f711c8181a2&sot=b&sdt=b&sl=74&sf&ws=r-f&ps=r-f&cs=r-f&origin=resultslist&zone=queryBar)

Journal Content

Search

Search Scope

All

Search


- Browse
- [By Issue \(https://ejournal.undip.ac.id/index.php/ksa/issue/archive\)](https://ejournal.undip.ac.id/index.php/ksa/issue/archive)
 - [By Author \(https://ejournal.undip.ac.id/index.php/ksa/search/authors\)](https://ejournal.undip.ac.id/index.php/ksa/search/authors)
 - [By Title \(https://ejournal.undip.ac.id/index.php/ksa/search/titles\)](https://ejournal.undip.ac.id/index.php/ksa/search/titles)
 - [Other Journals \(https://ejournal.undip.ac.id/index.php/index/search\)](https://ejournal.undip.ac.id/index.php/index/search)
 - [Categories \(https://ejournal.undip.ac.id/index.php/index/search/categories\)](https://ejournal.undip.ac.id/index.php/index/search/categories)

Visitor: 390903 [View My Stats](#)Jurnal Kimia Sains dan Aplikasi is indexed in:



 This work is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-sa/4.0/).

Copyright ©2023 [Universitas Diponegoro](#). Powered by [Public Knowledge Project OJS](#) and [Mason Publishing OJS theme](#).



Quantitative Structure–Activity Relationship of 3–Thiocyanate–1H–Indoles Derived Compounds as Antileukemia by AM1, PM3, and RM1 Methods

Ponco Iswanto ^{a,*}, Irvan Maulana Firdaus ^b, Ahmad Fawwaz Dafaulhaq ^b,
Ahmad Ghifari Ramadhani ^b, Maylani Permata Saputri ^a, Heny Ekowati ^c

^a Department of Chemistry, Faculty of Mathematics and Natural Sciences, Jenderal Soedirman University, Jl. dr. Soeparno No 61, Karangwangkal, Purwokerto 53122, Indonesia

^b Al Irsyad Al Islamiyyah Boarding School Senior High School, Jl. Raya Kebumen–Baturraden KM. 10, Banyumas, Central Java 53151, Indonesia

^c Department of Pharmacy, Faculty of Health Sciences, Jenderal Soedirman University, Jl. dr. Soeparno, Karangwangkal, Purwokerto 53122, Indonesia

* Corresponding author: ponco.iswanto@unsoed.ac.id

<https://doi.org/10.14710/jksa.26.3.109-117>

Article Info

Article history:

Received: 04th April 2023

Revised: 04th May 2023

Accepted: 22nd May 2023

Online: 31st May 2023

Keywords:

anticancer; QSAR; 3–thiocyanato–1H–indoles; semi-empirical

Abstract

Cancer is a disease with fatal consequences; thus, searching for innovative compounds with anticancer properties remains an active pursuit. One of the highly promising candidates is a compound derived from 3–thiocyanato–1H–indoles. However, the number of derivative compounds is currently limited. A quantitative structure and activity relationship (QSAR) study was conducted on derivative compounds 3–thiocyanato–1H–indoles to establish equations that predict the anticancer activity of more effective derivatives. This study aims to compare the effectiveness of the AM1 (Austin Model 1), PM3 (Parameterized Model 3), and RM1 (Recife Model 1) semi-empirical methods, which are new techniques implemented in the Hyperchem version 8.0. Twenty experimental data were used, 16 derivatives of 3–thiocyanato–1H–indoles as regression compounds (fitting) and four derivatives as test compounds. QSAR analysis was performed based on multiple linear regression calculations on 3–thiocyanato–1H–indoles derivative compounds by plotting IC_{50} (μM) as the dependent variable and descriptors as the independent variable. The best QSAR equation was obtained from the AM1 semiempirical calculation method with the following equation: $IC_{50} = -1.705 + 0.511(\Delta) + 0.346(Dipol) + 18.287(qC_9) - 0.645(Log P) + 13.952(qC_6)$, with $n = 20$; $r = 0.814$; $r^2 = 0.662$; The standard error (SE) = 1.044; $F_{count}/F_{table} = 1.851$; PRESS = 15.219.

1. Introduction

Cancer remains a significant challenge in the healthcare sector. Based on the Ministry of Health in 2020, cancer ranks as the third leading cause of death in Indonesia, following behind heart disease and stroke. Among the various types of cancer, leukemia—commonly known as blood cancer—is the prevalent type in children [1]. Leukemia arises from the excessive and uncontrolled production of white blood cells, disrupting the normal activity of other blood cells [2]. Unfortunately, due to delayed diagnosis and treatment, around 60% of children

with leukemia enter medical treatment in advanced stages [3].

Current cancer treatment primarily involves chemotherapy, radiotherapy, and surgery. Researchers are actively pursuing new anticancer drug development since current anticancer agents have limitations, such as their potential to affect normal cells [4]. Conventional new drug development in the laboratory entails multiple steps such as design, synthesis, purification, and identification, which can consume time, energy, and material resources. This process can be wasteful if the

results are less efficacious than existing drugs [5, 6]. Therefore, discovering potent, safe, and selective anticancer compounds is an important aspect of developing anticancer drugs [4].

Computational chemistry has a significant role in medicinal chemistry, especially in drug design, the theoretical prediction of chemical properties, and molecule activity [7]. One of the applications of computational chemistry is the quantitative structure-activity relationship (QSAR) study [8]. This approach is beneficial as it quantitatively investigates the correlation between molecular structure and experimentally measured biological activity values [9]. Several QSAR studies have been conducted in the search for anticancer compounds, such as those utilizing the Austin model 1 (AM1) and parameterized model 3 (PM3) semi-empirical methods for the QSAR application of estradiol-derived compounds [5, 10], as well as the PM3 method for the QSAR of calanone-derived compounds [11]. Modified AM1 has also been used for calculating alkane/water partition coefficients [12], while PM3 has been used in molecular simulations [13]. Furthermore, the RM1 (Recife Model 1) method, a method used in Monte Carlo simulation [14] and available in Hyperchem version 8, is widely used for calculating the enthalpy of formation, dipole moment, net charge, ionization potential, and structure geometry [15].

Despite their use as synthesis intermediates for complex bioactive chemicals, 3-thiocyanate-1H-indoles are rarely studied [16]. 3-thiocyanate-1H-indoles are a combination of heterocyclic indoles and thiocyanate compounds, where indoles are abundant natural products with biologically active properties [17], while thiocyanate has the ability to inhibit cancer cells [18]. The 3-thiocyanate-1H-indole compound and its derivatives have been tested against human cancer HL-60, HEP-2, NCI-H292, and MCF-7 cell lines and have demonstrated good activity [4]. However, only a few derivatives of 3-thiocyanate-1H-indoles have been produced, and some of these are no anticancer activity [19]. Therefore, further studies are needed to develop better anticancer compound derivatives for use as potential drugs [20].

This study compared the semi-empirical methods of AM1, PM3, and RM1 [21]. The aim was to analyze the QSAR of 3-thiocyanate-1H-indoles derivative compounds as anticancer agents using the three methods to obtain the best mathematical equation model [22]. Semi-empirical methods offer several advantages over *Ab initio*, such as faster calculations, small storage space, and the ability to accurately model compounds through electronic structure calculations and experiment parameterizations [23]. Furthermore, the best QSAR equation is expected to be used for the prediction of modification of new 3-thiocyanate-1H-indoles derivatives with more potent anticancer activity.

2. Materials and Methods

2.1. Materials

The computer system used in this study was equipped with an Intel Core i7-9700 processor, 16 GB

RAM, and Windows 10 operating system. The software included Hyperchem software version 8.0 [21] and SPSS version 25 [24]. Twenty compounds derived from 3-thiocyanato-1H-indoles and their biological activity as anticancer compounds were utilized (Figure 1) [4].




Figure 1. Structure of the parent model compound of 3-thiocyanate-1H-indoles derivatives [4]

Table 1. Fitting and test compound substituents, the IC_{50} (μM) experimental of 3-thiocyanato 1H-indoles derived compounds and the activity against HL-60 cell line [4]

Compound	R1	R2	R3	IC_{50} (μM) experiment
Fitting compound				
2	H	H	4-Me-C ₆ H ₄	2.53
3	H	H	OMe	5.63
5	H	H	CN	5.62
6	Ph	H	H	1.43
7	Me	H	4-Me-C ₆ H ₄	4.05
8	4-MeO-C ₆ H ₄	H	H	1.42
9	Me	H	H	1.54
10	4-Cl-C ₆ H ₄	H	H	3.58
11	H	C ₆ H ₅	H	0.63
12	H	4-Me-C ₆ H ₄	H	2.27
13	H	4-MeO-C ₆ H ₄	H	4.45
14	H	3CF ₃ -C ₆ H ₄	H	0.69
15	H	4-Cl-C ₆ H ₄	H	1.08
16	Me	C ₆ H ₅	H	1.39
17	Me	4-Me-C ₆ H ₄	H	1.50
20	Me	4-Cl-C ₆ H ₄	H	1.10
Test compound				
1	H	H	H	3.21
4	H	H	Br	3.00
18	Me	4-MeO-C ₆ H ₄	H	1.35
19	Me	3CF ₃ -C ₆ H ₄	H	2.56

The selection of the test compounds was based on: the smallest molecule (compound 1), the largest molecule (compound 19), the electron-withdrawing compound (compound 4), and the electron-donating compound (compound 18) (Table 1). These test compounds were expected to represent fitting compounds, and they can aid in selecting the best equation model through backward analysis, which was further tested by calculating the PRESS value [23]. The QSAR method was the Hansch method which consisted of hydrophobicity (π), electronic (σ), and steric (Es).

2.2. Structure optimization

Each 3-thiocyanate-1H-indoles derivate compound was created as a 2D structural model using the Hyperchem software. The model was then transformed into a 3D structure through the “Add H and Model Build”. The structure geometry was optimized by minimizing molecular energy to attain the most stable molecular conformation. This was achieved using RHF spin pairing, lowest state, Polak-Ribiere optimization algorithm, and Root Mean Square (RMS) Gradient 0.001 kcal/Å.mol [25].

2.3. Correlation analysis

Bivariate correlation analysis was conducted using the two-tailed method and Pearson coefficient through SPSS 25 software. During this step, the correlation level between each descriptor and anticancer activity (IC_{50}) was investigated [11].

2.4. Descriptor determination

The QSAR method was Hansch method. The descriptor was composed of electrons, lipophilicity, and steric hindrance. After the optimum structure, the electrons, in this case, the charge and the dipole moment, the lipophilicity, and steric hindrance; the last two were in the QSAR properties. Determining descriptors is essential in identifying the best QSAR equation. The Hanch analysis uses electronic, steric, and hydrophobic parameters [10]. This study utilized the following descriptors as electronic parameters: atomic charge of the parent compound, hydration energy, dipole moment, HOMO, LUMO, and HOMO-LUMO energy difference (ΔE). Steric parameters included van der Waals volume (Vvdw), molecular weight (BM), and polarizability (α), while the hydrophobic parameter was the partition coefficient (Log P).

2.5. Determination of the QSAR equation

The best QSAR equation for predicting anticancer activity was determined using multilinear regression analysis. The analysis was performed by applying the backward method on 16 fitting compounds, with IC_{50} as the dependent variable and the descriptors as the independent variables. Statistical analysis was carried out on the output data of the backward analysis to select the equation model. The analysis included the correlation coefficient (r), partition coefficient (r^2), standard deviation (SD), F_{count}/F_{table} , and PRESS on four test compounds. The chosen model was then analyzed using the Enter method for all 20 compounds derived from 3-thiocyanate-1H-indoles to obtain the best QSAR equation [26]. This analysis was performed using the SPSS software.

3. Results and Discussion

3.1. Structure geometry optimization

Geometry optimization obtains the most stable molecular structure with the lowest energy and minimal atomic force [27]. This study employed three semi-empirical methods, namely AM1, PM3, and RM1 [21], to optimize the geometry of the compounds. The semi-empirical method is a type of electronic structure calculation developed based on molecular orbital theory and mathematical models. These methods simplify and approximate the computational procedure. The method involves only valence electrons in solving the Schrödinger equation, making optimization time shorter than the ab initio methods. After optimization, the compound's conformational shape differs from its original state, resulting in a broader structure due to the induction effect of electron clouds on specific atoms and groups, causing them to be further apart.

A derivative of 3-thiocyanate-1H-indoles was optimized before and after using the AM1, PM3, and RM1 methods. The total energy obtained from the AM1 method was -44.065.7702998 kcal/mol, PM3 gave a total energy of -40247.4055128 kcal/mol, and RM1 resulted in a total energy of -43665.4491644 kcal/mol. This result suggests that the AM1 method gave the smallest total energy. These optimized compound structures were used to obtain data on physical and chemical properties.

3.2. Descriptor calculation results

The optimized derivative compound of 3-thiocyanate-1H-indoles was used to calculate its physicochemical characteristics. The measured values included atomic charges for qN1, qC2, qC3, qC4, qC5, qC6, qC7, qC8, qC9, qS10, qC11, qN12, hydration energy, dipole moment, HOMO, LUMO, HOMO and LUMO energy difference (ΔE) as electronic parameters. Steric parameters included van der Waals volume (Vvdw), molecular weight (BM), and polarizability (α). At the same time, the partition coefficient (Log P) was measured as a hydrophobic parameter [28].

The calculation results for the AM1, PM3, and RM1 methods are shown in Tables 2, 3, and 4, respectively. Notably, the dipole moment, partition coefficient (Log P), and net atomic charge of the C2 (qC2) show the highest trend in all three methods. For the AM1 method, the dipole moment, Log P, and qC2 were 6.265 Debye, -0.89, and 0.059 C, respectively. For the PM3 method, the dipole moment, Log P, and qC2 were 6.633 Debye, -0.89, and -0.107 C, respectively. The data for the RM1 method was 6.863 Debye for the dipole moment, -0.89 for Log P, and 0.121 C for qC2.

Table 2. Calculation data of AM1 method descriptors

Compound	Dipole	Vvdw	Hidration	log P	α	BM	qN1	qC2	qC3	qC4	qC5	qC6	qC7	qC8	qC9	qS10	qC11	qN12	HOMO	LUMO	ΔE
Fitting compound																					
2	4.462	772.49	-9.15	0.17	30.84	264.34	-0.198	-0.05	-0.383	-0.05	-0.079	-0.039	-0.1	-0.143	0.003	0.577	-0.375	-0.006	-8.144	-0.258	7.886
3	5.673	594.63	-11.29	-1.58	21.81	204.25	-0.2	-0.041	-0.392	-0.019	-0.121	0.062	-0.175	-0.109	-0.029	0.573	-0.377	0	-8.151	-0.14	8.011
5	6.977	587.99	-14.36	-0.87	21.19	199.23	-0.197	0.025	-0.42	-0.043	-0.044	-0.026	-0.082	-0.137	0.002	0.554	-0.384	-0.009	-9.278	-0.807	8.471
6	4.468	720.71	-8.36	0.12	29	250.32	-0.103	-0.05	-0.376	-0.052	-0.09	-0.142	-0.113	-0.136	0.027	0.58	-0.377	-0.001	-8.195	-0.521	7.674
7	4.883	825.15	-5.87	0.42	32.67	278.37	-0.153	-0.049	-0.383	-0.054	-0.08	-0.04	-0.101	-0.142	0.009	0.576	-0.375	-0.007	-8.066	-0.226	7.84
8	4.911	569.37	-6.68	-0.34	21.18	188.25	-0.097	-0.052	-0.377	-0.052	-0.09	-0.143	-0.113	-0.136	0.026	0.578	-0.377	-0.001	-8.18	-0.059	8.121
9	5.147	796.77	-10.05	-0.87	31.47	280.34	-0.155	-0.049	-0.383	-0.053	-0.086	-0.148	-0.11	-0.142	0.008	0.574	-0.375	-0.006	-7.969	-0.497	7.472
10	2.921	764.45	-8.01	-0.1	30.93	284.76	-0.105	-0.052	-0.371	-0.052	-0.088	-0.141	-0.111	-0.137	0.028	0.582	-0.379	0.002	-8.143	-0.75	7.393
11	4.906	715.99	-9.24	0.11	29	250.32	-0.199	0.048	-0.361	-0.064	-0.084	-0.151	-0.104	-0.148	0.016	0.582	-0.384	-0.006	-7.957	-0.766	7.191
12	5.206	767.13	-8.06	0.26	30.84	264.34	-0.2	0.051	-0.363	-0.063	-0.084	-0.151	-0.105	-0.148	0.015	0.582	-0.384	-0.007	-7.914	-0.754	7.16
13	6.265	792.7	-10.93	-0.89	31.47	280.34	-0.202	0.059	-0.368	-0.061	-0.085	-0.151	-0.105	-0.148	0.014	0.579	-0.383	-0.007	-7.874	-0.715	7.159
14	1.254	796.61	-8.56	0.68	30.56	318.32	-0.197	0.034	-0.35	-0.068	-0.082	-0.15	-0.1	-0.147	0.02	0.58	-0.385	0.003	-8.209	-1.147	7.062
15	4.148	759.41	-8.91	-0.12	30.93	284.76	-0.199	0.042	-0.356	-0.066	-0.083	-0.151	-0.102	-0.148	0.018	0.582	-0.385	-0.001	-8.056	-0.982	7.074
16	5.309	751.56	-6.89	0.35	30.84	264.34	-0.151	0.06	-0.366	-0.064	-0.085	-0.15	-0.107	-0.146	0.013	0.582	-0.388	-0.009	-7.888	-0.738	7.15
17	5.505	805.25	-5.72	0.51	32.67	278.37	-0.151	0.063	-0.368	-0.063	-0.086	-0.15	-0.107	-0.146	0.013	0.581	-0.388	-0.01	-7.846	-0.728	7.118
20	4.724	795.04	-6.57	0.13	32.76	298.79	-0.15	0.054	-0.361	-0.065	-0.084	-0.149	-0.105	-0.146	0.015	0.582	-0.388	-0.005	-7.987	-0.95	7.037
Test compound																					
1	4.751	518.25	-9.99	-0.59	19.34	174.22	-0.2	-0.05	-0.384	-0.054	-0.086	-0.148	-0.109	-0.143	0.001	0.575	-0.375	-0.005	-8.266	-0.085	8.181
4	5.833	580.79	-9.52	-0.54	21.97	253.12	-0.197	-0.05	-0.378	-0.058	-0.056	-0.187	-0.085	-0.142	0.011	0.58	-0.381	0.005	-8.441	-0.327	8.114
18	4.542	829.47	-8.6	-0.64	33.31	294.37	-0.152	0.072	-0.374	-0.06	-0.087	-0.149	-0.108	-0.145	0.012	0.577	-0.386	-0.01	-7.826	-0.688	7.138
19	7.191	830.98	-6.24	0.92	32.4	332.34	-0.15	0.045	-0.354	-0.068	-0.082	-0.149	-0.102	-0.147	0.018	0.588	-0.388	0.002	-8.099	-1.103	6.996

The PM3 method calculates a -0.107 C charge, which was negative, while the other two methods both produce positive values for qC2. This indicates that PM3 is less feasible for use, as PM3 is better suited for compounds containing transition metal atoms. The dipole moment values for all three methods were large, indicating substituents containing electronegative atoms [29]. The Log P data for all three methods are the same, indicating that the compound contains many functional groups contributing to its polarity [23].

The net atomic charge is a crucial descriptor in determining various chemical reactions and physicochemical properties of compounds. It measures intermolecular interactions and can be positive or negative, depending on the group attached to the atom. The net positive atomic charge resulted from electron-withdrawing groups, such as methoxy, which reduce the electron density. The net negative atomic charge arises from electron-contributing groups, such as methyl or alkyl groups, which increase the electron density [11]. The PM3 method calculated the atomic charge descriptors, which showed similar positive and negative charge values as those calculated by the AM1 and RM1 methods, except for qN1, which was positively charged, and qC9, which was negatively charged. The difference may be due to the

PM3 calculation method predicting that all N atoms are pyramidal, which is a weakness of the method [30].

The log P value determines the distribution of molecules in the body. A more positive log P value suggests that the compound tends to distribute in the nonpolar phase rather than the polar phase. Since the human body consists of lipid and water phases, the log P value significantly affects the diffusion process of active substances [31]. Polarizability measures a molecule's ability to form an instantaneous dipole or induce one in another molecule. Dipole attraction arises from an unsymmetrical charge distribution in the molecule, resulting in two ends with different charges. Dipole moments are crucial in drug activity, particularly in the interaction between compounds and receptor targets. If there is sufficient contact over a large area, a significant inter-dipole interaction energy can be formed. The polarizabilities obtained from the AM1, PM3, and RM1 calculations have almost the same value, indicating that the three methods are equivalent in calculating polarizability. However, the dipole moment calculation results show that the AM1 method has a relatively smaller value than PM3 and RM1.

Table 3. Calculation data of PM3 method descriptors

Compound	Dipole	Vvdw	Hydration	log P	α	BM	qN1	qC2	qC3	qC4	qC5	qC6	qC7	qC8	qC9	qS10	qC11	qN12	HOMO	LUMO	ΔE
Fitting compound																					
2	5.089	770.25	-9.12	0.17	30.84	264.34	0.323	-0.198	-0.28	-0.044	-0.065	-0.032	-0.086	-0.093	-0.181	0.394	-0.286	-0.026	-8.179	-0.268	7.911
3	6.236	596.16	-11.26	-1.58	21.81	204.25	0.321	-0.189	-0.291	-0.012	-0.112	0.078	-0.154	-0.059	-0.212	0.391	-0.289	-0.019	-8.129	-0.165	7.964
5	7.569	588.65	-14.34	-0.87	21.19	199.23	0.33	-0.137	-0.322	-0.035	-0.015	-0.021	-0.067	-0.085	-0.182	0.386	-0.303	-0.023	-9.241	-0.735	8.506
6	5.003	711.42	-8.18	0.12	29	250.32	-0.248	0.051	-0.358	-0.067	-0.052	-0.117	-0.076	-0.118	0.082	0.486	-0.152	-0.167	-8.128	-0.482	7.646
7	5.441	823.17	-5.84	0.42	32.67	278.37	0.383	-0.178	-0.28	-0.046	-0.065	-0.034	-0.086	-0.095	-0.155	0.393	-0.285	-0.027	-8.099	-0.227	7.872
8	5.823	797.09	-10.03	-0.87	31.47	280.34	0.385	-0.203	-0.269	-0.053	-0.062	-0.111	-0.087	-0.1	-0.151	0.394	-0.289	-0.021	-8.107	-0.435	7.672
9	5.732	571.74	-6.71	-0.34	21.18	188.25	0.28	-0.178	-0.28	-0.052	-0.059	-0.115	-0.084	-0.099	-0.153	0.392	-0.287	-0.023	-8.14	-0.065	8.075
10	4.356	764.6	-8.04	-0.1	30.93	284.76	0.37	-0.199	-0.263	-0.053	-0.061	-0.109	-0.084	-0.101	-0.146	0.397	-0.29	-0.018	-8.275	-0.684	7.591
11	5.423	715.8	-9.24	0.11	29	250.32	0.321	-0.12	-0.263	-0.053	-0.059	-0.115	-0.082	-0.1	-0.171	0.402	-0.305	-0.026	-8.116	-0.738	7.378
12	5.649	766.32	-8.05	0.26	30.84	264.34	0.32	-0.115	-0.265	-0.052	-0.059	-0.115	-0.082	-0.1	-0.172	0.401	-0.304	-0.027	-8.072	-0.726	7.346
13	6.633	791.81	-10.88	-0.89	31.47	280.34	0.316	-0.107	-0.27	-0.05	-0.06	-0.115	-0.083	-0.1	-0.172	0.401	-0.304	-0.027	-8.028	-0.688	7.34
14	2.172	794.38	-8.57	0.68	30.56	318.32	0.325	-0.135	-0.253	-0.056	-0.058	-0.113	-0.077	-0.1	-0.167	0.402	-0.307	-0.016	-8.368	-1.117	7.251
15	4.998	758.55	-8.9	-0.12	30.93	284.76	0.322	-0.124	-0.26	-0.054	-0.058	-0.115	-0.08	-0.1	-0.17	0.402	-0.306	-0.023	-8.21	-0.953	7.257
16	5.795	751.81	-6.89	0.35	30.84	264.34	0.29	-0.088	-0.266	-0.055	-0.059	-0.116	-0.08	-0.104	-0.157	0.4	-0.305	-0.028	-8.048	-0.702	7.346
17	5.94	804.1	-5.72	0.51	32.67	278.37	0.289	-0.084	-0.269	-0.054	-0.06	-0.116	-0.081	-0.103	-0.157	0.4	-0.305	-0.029	-8.006	-0.691	7.315
20	5.443	794.82	-6.57	0.13	32.76	298.79	0.291	-0.092	-0.263	-0.056	-0.059	-0.116	-0.079	-0.104	-0.156	0.401	-0.306	-0.025	-8.142	-0.914	7.228
Test compound																					
1	5.393	520.17	-10.02	-0.59	19.34	174.22	0.321	-0.198	-0.279	-0.05	-0.059	-0.114	-0.084	-0.097	-0.179	-0.393	-0.288	-0.023	-8.222	-0.107	8.115
4	5.948	591.7	-9.83	-0.54	21.97	253.12	0.33	-0.139	-0.326	-0.023	-0.03	-0.118	-0.075	-0.079	-0.194	0.383	-0.3	-0.028	-9.133	-0.547	8.586
18	5.055	829.02	-8.56	-0.64	33.31	294.37	0.286	-0.074	-0.275	-0.052	-0.061	-0.116	-0.082	-0.103	-0.158	0.397	-0.304	-0.029	-7.983	-0.654	7.329
19	7.759	829.09	-6.22	0.92	32.4	332.34	0.291	-0.103	-0.256	-0.059	-0.056	-0.115	-0.075	-0.105	-0.153	0.405	-0.305	-0.017	-8.256	-1.063	7.193

The HOMO-LUMO gap (ΔE) is a descriptor that measures the ease a molecular system can excite to a higher electronic state. The HOMO energy is directly linked to the ionization potential and the molecule's susceptibility to attacking electrophiles. In contrast, the LUMO energy relates to electron affinity and the molecule's vulnerability to nucleophile attack. The energy difference between the HOMO and LUMO is crucial in determining molecular stability. A large HOMO-LUMO gap (ΔE) suggests high stability and low reactivity in chemical reactions. This gap is also used to estimate the molecule's lowest excitation energy. The AM1 method produced relatively smaller ΔE values than the PM3 and RM1 methods (Table 2), indicating low stability and increased compound reactivity. Hydration energy and van der Waals volume are also considered significant in determining the biological activity of anticancer compounds, as their values vary across different compounds [21].

3.3. Correlation calculation results of descriptors with IC₅₀

The correlation analysis was performed using the two-tailed and Pearson correlation coefficients. The results were used to select the most suitable descriptors as independent variables for multiple linear regression (MLR) analysis. This analysis helped to determine the extent to which the independent variables (descriptors) influenced the IC₅₀ value to construct a reliable QSAR equation. The correlation values between the descriptors and anticancer activity are presented in Table 5.

The chosen descriptor should have a significant correlation value and a significance level below 0.05 for a 95% confidence interval. Thus, the descriptors with substantial correlations for the AM1 method were ΔE , dipole, hydration, qC9, qC8, HOMO, log P, qC3, qC6, and qS10. For the PM3 method, the descriptors selected were ΔE , qC7, dipole, hydration, α , qC8, log P, qC6, BM, and qC4. Finally, the chosen descriptors for the RM1 method were ΔE , hydration, qC6, dipole, volume, log P, qC8, BM, α , and qC4.

Table 4. Calculation data of RM1 method descriptors

Compound	Dipole	Vvdw	Hydration	log P	α	BM	qN1	qC2	qC3	qC4	qC5	qC6	qC7	qC8	qC9	qS10	qC11	qN12	HOMO	LUMO	ΔE
Fitting compound																					
2	5.015	761.58	-8.88	0.17	30.84	264.34	-0.311	0.053	-0.368	-0.068	0.04	-0.059	-0.063	-0.121	0.062	0.483	-0.149	-0.173	-8.238	-0.203	8.035
3	6.092	588.77	-10.95	-1.58	21.81	204.25	-0.313	0.064	-0.379	-0.033	-0.085	0.068	-0.139	-0.087	0.028	0.479	-0.151	-0.166	-8.224	-0.108	8.116
5	7.388	581.55	-14.1	-0.87	21.19	199.23	-0.313	0.126	-0.413	-0.064	0.011	-0.084	0.039	-0.116	0.067	0.481	-0.164	-0.173	-9.365	-0.734	8.631
6	5.003	711.42	-8.18	0.12	29	250.32	-0.248	0.051	-0.358	-0.067	-0.052	-0.117	-0.076	-0.118	0.082	0.486	-0.152	-0.167	-8.291	-0.462	7.829
7	5.444	813.43	-5.64	0.42	32.67	278.37	-0.294	0.054	-0.367	-0.068	-0.04	-0.06	-0.064	-0.12	0.071	0.482	-0.149	-0.174	-8.158	-0.169	7.989
8	5.555	785.79	-9.79	-0.87	31.47	280.34	-0.242	0.049	-0.359	-0.068	-0.052	-0.118	-0.076	-0.118	0.081	0.485	-0.151	-0.168	-8.184	-0.422	7.762
9	5.718	564.24	-6.54	-0.34	21.18	188.25	-0.295	0.054	-0.369	-0.068	-0.049	-0.122	-0.075	-0.119	0.068	0.48	-0.15	-0.172	-8.267	-0.02	8.247
10	2.629	755.79	-7.84	-0.1	30.93	284.76	-0.252	0.048	-0.352	-0.067	-0.05	-0.116	-0.073	-0.119	0.083	0.49	-0.154	-0.164	-8.445	-0.754	7.691
11	5.568	708.28	-9.07	0.11	29	250.32	-0.311	0.11	-0.336	-0.08	-0.047	-0.126	-0.069	-0.125	0.073	0.489	-0.156	-0.171	-8.217	-0.698	7.519
12	5.882	760.84	-7.89	0.26	30.84	264.34	-0.311	0.113	-0.338	-0.079	-0.048	-0.126	-0.069	-0.125	0.073	0.488	-0.155	-0.172	-8.173	-0.688	7.485
13	6.863	784.62	-10.67	-0.89	31.47	280.34	-0.313	0.121	-0.342	-0.077	-0.048	-0.126	-0.07	-0.125	0.072	0.486	-0.155	-0.172	-8.122	-0.647	7.475
14	2.056	785.32	-8.37	0.68	30.56	318.32	-0.309	0.096	-0.324	-0.084	-0.046	-0.124	-0.065	-0.125	0.078	0.486	-0.155	-0.164	-8.463	-1.092	7.371
15	4.323	752.81	-8.73	-0.12	30.93	284.76	-0.31	0.1	-0.327	-0.083	-0.046	-0.125	-0.066	-0.126	0.076	0.488	-0.156	-0.166	-8.373	-0.996	7.377
16	5.477	769.77	-6.72	0.35	30.84	264.34	-0.284	0.18	-0.405	-0.049	-0.047	-0.118	-0.083	-0.11	0.05	0.479	-0.159	-0.191	-8.707	-0.418	8.289
17	5.591	820.46	-5.59	0.51	32.67	278.37	-0.284	0.183	-0.406	-0.049	-0.048	-0.118	-0.083	-0.11	0.05	0.478	-0.159	-0.192	-8.676	-0.413	8.263
20	5.217	819.42	-6.4	0.13	32.76	298.79	-0.283	0.17	-0.399	-0.051	-0.046	-0.117	-0.08	-0.11	0.051	0.48	-0.161	-0.187	-8.844	-0.709	8.135
Test compound																					
1	5.314	513.44	-9.82	-0.59	19.34	174.22	-0.313	0.054	-0.369	-0.068	-0.048	-0.122	-0.073	-0.121	0.059	0.481	-0.15	-0.171	-8.355	-0.048	8.307
4	6.7	576.8	-9.32	-0.54	21.97	253.12	-0.309	0.056	-0.365	-0.07	-0.022	-0.1	-0.053	-0.116	0.064	0.488	-0.156	-0.161	-8.566	-0.351	8.215
18	5.407	818.56	-8.34	-0.64	33.31	294.37	-0.287	0.131	-0.348	-0.072	-0.049	-0.124	-0.073	-0.123	0.068	0.483	-0.162	-0.178	-8.058	-0.628	7.43
19	7.7	819.85	-6.06	0.92	32.4	332.34	-0.286	0.106	-0.329	-0.08	-0.045	-0.124	-0.066	-0.126	0.074	0.493	-0.161	-0.167	-8.844	-1.054	7.79

Table 5. The correlation values between the anticancer activity and descriptors

		AM1 method						
	IC ₅₀	Dipole	qN1	qC2	qC3	qC4	qC5	qC6
		0.457	-0.264	-0.313	-0.697	0.395	0.093	0.682
		qC7	qC8	qC9	qS10	qC11	qN12	Log P
		-0.245	0.570	-0.601	-0.590	0.338	-0.001	-0.552
		Vvdw	Hydration	HOMO	LUMO	ΔE	α	BM
		-0.379	-0.560	-0.519	0.352	0.590	-0.439	-0.405
		PM3 method						
	IC ₅₀	Dipole	qN1	qC2	qC3	qC4	qC5	qC6
		0.494	0.225	-0.373	-0.314	0.681	-0.061	0.705
		qC7	qC8	qC9	qS10	qC11	qN12	Log P
		-0.413	0.666	-0.292	-0.149	-0.060	0.170	-0.548
		Vvdw	Hidrasi	HOMO	LUMO	ΔE	α	BM
		-0.387	-0.555	-0.400	0.357	0.570	-0.448	-0.414
		RM1 method						
	IC ₅₀	Dipole	qN1	qC2	qC3	qC4	qC5	qC6
		0.410	-0.289	-0.258	-0.279	0.312	0.133	0.629
		qC7	qC8	qC9	qS10	qC11	qN12	Log P
		0.206	0.384	-0.293	-0.142	0.099	0.244	-0.548
		Vvdw	Hydration	HOMO	LUMO	ΔE	α	BM
		-0.411	-0.547	-0.173	0.280	0.390	-0.448	-0.414

The chosen descriptor should have a significant correlation value and a significance level below 0.05 for a 95% confidence interval. Thus, the descriptors with substantial correlations for the AM1 method were ΔE , dipole, hydration, qC9, qC8, HOMO, log P, qC3, qC6, and qS10. For the PM3 method, the descriptors selected were ΔE , qC7, dipole, hydration, α , qC8, log P, qC6, BM, and qC4. Finally, the chosen descriptors for the RM1 method were ΔE , hydration, qC6, dipole, volume, log P, qC8, BM, α , and qC4.

3.4. Determination of the best equation model with multiple linear regression

Analysis using the backward method of fitting compounds resulted in developing 6, 6, and 7 QSAR equation models for AM1, PM3, and RM1, respectively (Table 6, 7, 8). The selection of the best QSAR equation requires that the correlation coefficient (r) be greater than 0.8, the Fcount value should exceed the Ftable ($F_{\text{count}}/F_{\text{table}} > 1$) for a 95% confidence level, and the model should have a small standard error (SE) and a small predict the residual sum of square (PRESS) value [11].

Table 6. QSAR AM1 equation model from multiple linear regression analysis using a backward method

Model	Variable	R	r ²	SE	F _{count} /F _{table}	PRESS
1	ΔE, Dipole, Hidration, qC ₉ , qC ₈ , HOMO, logP, qC ₃ , qC ₆ , qS ₁₀	0.947	0.898	0.9395	4.382/4.74=0.924	163.637
2	ΔE, Dipole, Hidration, qC ₉ , qC ₈ , HOMO, logP, qC ₆ , qS ₁₀	0.947	0.898	0.8577	5.841/4.1=1.424	161.876
3	ΔE, Dipole, qC ₉ , qC ₈ , HOMO, logP, qC ₆ , qS ₁₀	0.946	0.895	0.8026	7.487/3.73=2.221	154.696
4	ΔE, Dipole, qC ₉ , HOMO, logP, qC ₆ , qS ₁₀	0.941	0.886	0.7831	8.895/3.5=2.541	125.836
5	ΔE, Dipole, qC ₉ , logP, qC ₆ , qS ₁₀	0.929	0.863	0.8113	9.411/3.37=2.792	76.165
6	ΔE, Dipole, qC ₉ , logP, qC ₆	0.908	0.825	0.8696	9.397/3.33=2.821	54.746

Table 7. QSAR PM3 equation model from multiple linear regression analysis using a backward method

Model	Variable	R	r ²	SE	F _{count} /F _{table}	PRESS
1	ΔE, qC ₇ , Dipole, Hidration, α, qC ₈ , logP, qC ₆ , BM, qC ₄	0.961	0.923	0.8150	5.988/4.74=1.255	307.388
2	ΔE, qC ₇ , Dipole, Hidration, polarizability, qC ₈ , logP, qC ₆ , qC ₄	0.961	0.923	0.7440	7.984/4.1=1.947	305.539
3	qC ₇ , Dipole, Hidration, α, qC ₈ , logP, qC ₆ , qC ₄	0.961	0.923	0.6898	10.447/3.73=2.8	309.889
4	qC ₇ , Dipole, Hidration, qC ₈ , logP, qC ₆ , qC ₄	0.958	0.917	0.6691	12.606/3.5=3.601	323.525
5	qC ₇ , Dipole, Hidration, qC ₈ , logP, qC ₄	0.945	0.894	0.7128	12.636/3.37=3.749	167.775
6	qC ₇ , Dipole, qC ₈ , logP, qC ₄	0.932	0.869	0.7504	13.306/3.33=3.995	114.348

Table 8. QSAR RM1 equation model from multiple linear regression analysis using a backward method

Model	Variable	R	r ²	SE	F _{count} /F _{table}	PRESS
1	ΔE, Hidration, qC ₆ , Dipole, Volume, logP, qC ₈ , BM, α, qC ₄	0.879	0.773	1.3981	1.704/4.74=0.359	15.730
2	ΔE, Hidration, qC ₆ , Volume, logP, qC ₈ , BM, α, qC ₄	0.879	0.773	1.2765	2.272/4.1=0.554	17.850
3	ΔE, Hidration, qC ₆ , Volume, logP, qC ₈ , α, qC ₄	0.879	0.773	1.1819	2.981/3.73=0.799	17.943
4	ΔE, Hidration, qC ₆ , Volume, qC ₈ , α, qC ₄	0.872	0.760	1.1370	3.620/3.5=1.034	21.163
5	ΔE, Hidration, qC ₆ , qC ₈ , α, qC ₄	0.834	0.695	1.2088	3.415/3.37=1.013	10.166
6	Hidration, qC ₆ , qC ₈ , α, qC ₄	0.823	0.678	1.1783	4.208/3.33=1.263	4.135
7	Hidration, qC ₆ , qC ₈ , qC ₄	0.821	0.674	1.1298	5.691/3.36=1.693	4.719

These QSAR equation models underwent statistical testing based on the values of r , r^2 , SE, and F [32], and the PRESS value was calculated for the four test compounds [33]. Model 6 was identified as the best for the AM1, PM3, and RM1 methods because it had an r -value in the range of 0.9, the smallest PRESS value, and the highest $F_{\text{count}}/F_{\text{table}}$ value [34]. In selecting the QSAR equation model, the PRESS value was given significant consideration as a smaller PRESS value indicates that the calculation of compound activity prediction is closer to the experimental data. The final QSAR equation for all 3-thiocyanate-1H-indole derivative compounds was obtained using the Enter method, which analyzed the selected models from the AM1, PM3, and RM1 methods.

The results of the Enter method analysis are shown in Table 10.

Table 9. Output of AM1, PM3, and RM1 with Enter method

Model	Variable	R	r ²	SE	F _{count} /F _{table}	PRESS
AM1	ΔE, Dipole, qC ₉ , logP, qC ₆	0.814	0.662	1.04484	5.481/2.96=1.851	15.219
PM3	qC ₇ , Dipole, qC ₈ , logP, qC ₄	0.761	0.578	1.16667	3.852/2.96=1.301	19.055
RM1	Hidration, qC ₆ , qC ₈ , α, qC ₄	0.788	0.622	1.10531	4.6/2.96=1.554	17.103

By comparing the r , r^2 , SE, $F_{\text{count}}/F_{\text{table}}$, and PRESS values, AM1 suggests the best method. Therefore, the best QSAR equation for 3-thiocyanate-1H-indoles was $IC_{50} = -1.705 + 0.511(\Delta E) + 0.346(\text{Dipol}) + 18.287(qC_9) - 0.645(\text{Log P}) + 13.952(qC_6)$, with $n = 20$; $r = 0.814$; $r^2 = 0.662$; $SE = 1.044$; $F_{\text{count}}/F_{\text{table}} = 1.851$; $PRESS = 15.219$. The correlation graph between predicted and experimental activities (IC_{50}) shows a slope close to 1 (Table 10, Figure 2) [26], meaning that the resulting equation can provide a good level of prediction.

Table 10. PRESS value of the final QSAR equation with Enter method

Compound	IC ₅₀ prediction	IC ₅₀ experiment	PRESS
1	2.45	3.21	0.5776
2	3.27	2.53	0.5476
3	5.70	5.63	0.0049
4	2.40	3	0.3600
5	5.27	5.62	0.1225
6	2.20	1.43	0.5929
7	3.33	4.05	0.5184
8	2.84	1.42	2.0164
9	2.53	1.54	0.9801
10	1.70	3.58	3.5344
11	1.78	0.63	1.3225
12	1.75	2.27	0.2704
13	2.84	4.45	2.5921
14	0.17	0.69	0.2704
15	1.64	1.08	0.3136
16	1.70	1.39	0.0961
17	1.65	1.5	0.0225
18	2.06	1.35	0.5041
19	2.02	2.56	0.2916
20	1.63	1.1	0.2809
Σ PRESS			15.219




Figure 2. Correlation graph between predicted and experimental activity (IC_{50})

4. Conclusion

Based on the results, the RM1 semi-empirical method was the latest semi-empirical calculation method in the Hyperchem Program version 8 and unsuitable for QSAR 3-thiocyanate-1H-indoles derivative compounds. The AM1 semi-empirical method was the best for QSAR of 3-thiocyanate-1H-indoles derivative compounds compared to PM3 and RM1 methods. The AM1 semi-empirical method, obtained $r^2 = 0.662$, was a measure of the goodness of fit of a model. The standard error (SE) was 1.044.

Acknowledgment

We would like to thank Intan Tia Nurrochmah for her help in checking the bibliography.

References

- [1] Kementerian Kesehatan Republik Indonesia, *Profil Kesehatan Indonesia Tahun 2019*, Kementerian Kesehatan Republik Indonesia, Jakarta, 2020,
- [2] M. Fahrudin Ghazali, Ade Eviyanti, Sistem Pakar Diagnosa Dini Penyakit Leukimia Dengan Metode Certainty Factor, *Kinetik: Game Technology, Information System, Computer Network, Computing, Electronics, and Control*, 1, 3, (2016), 135-146 <https://doi.org/10.22219/kinetik.v1i3.122>
- [3] Mukhlis Ramadhan, Sistem Pakar Dalam Mengidentifikasi Penyakit Kanker Pada Anak Sejak Dini dan Cara Penanggulangannya, *Jurnal SAINTIKOM*, 10, 2, (2011), 125-135
- [4] Richa Monika, Dhingra Neelima, 3-Thiocyanato-1H-indoles as potential anticancer agents: Two dimensional quantitative structure activity relationship study, *International Journal of Pharmaceutical Chemistry and Analysis*, 3, 4, (2016), 198-204
- [5] Syarifah Nugraheni, Ponco Iswanto, Iqmal Tahir, Hubungan Kuantitatif Struktur dan Aktivitas Antikanker Senyawa Turunan Estradiol Hasil Perhitungan Metode Semiempiris AM1, *Prosiding Seminar Nasional Kimia XV ISSN NO*, 2004
- [6] Asril Burhan, Andi Nur Aisyah, Akbar Awaluddin, Zulham Zulham, Burhanuddin Taebe, Abdul Gafur, Efek antioksidan dan antikanker ekstrak batang murbei (*Morus alba* L.) secara *in vitro*, *Kartika: Jurnal Ilmiah Farmasi*, 7, 1, (2019), 17-21
- [7] Petko Alov, Ivanka Tsakovska, Ilza Pajeva, Computational studies of free radical-scavenging properties of phenolic compounds, *Current Topics in Medicinal Chemistry*, 15, 2, (2015), 85-104 <http://dx.doi.org/10.2174/1568026615666141209143702>
- [8] Tao Huang, Guohui Sun, Lijiao Zhao, Na Zhang, Rugang Zhong, Yongzhen Peng, Quantitative structure-activity relationship (QSAR) studies on the toxic effects of nitroaromatic compounds (NACs): A systematic review, *International Journal of Molecular Sciences*, 22, 16, (2021), 8557 <https://doi.org/10.3390/ijms22168557>
- [9] Amit Kumar Halder, M Natália Dias Soeiro Cordeiro, Development of multi-target chemometric models for the inhibition of class I PI3K enzyme isoforms: A case study using QSAR-Co tool, *International Journal of Molecular Sciences*, 20, 17, (2019), 4191 <https://doi.org/10.3390/ijms20174191>
- [10] Ponco Iswanto, Isnaeni Tatik Rosdiyana, Iqmal Tahir, Hubungan kuantitatif struktur dan aktivitas antikanker senyawa turunan estradiol hasil perhitungan metode semiempiris PM3, *Berkala MIPA*, 17, 1, (2007), 12-20
- [11] Eva Vulina Y. D., Mochammad Chasani, Mokhammad Abdulghani, Hubungan kuantitatif struktur-aktivitas (HKSA) antikanker senyawa turunan kalanon dengan metode semi empiris PM3 (Parameterized Model 3), *Molekul*, 7, 2, (2012), 130-142 <http://dx.doi.org/10.20884/1.jm.2012.7.2.115>
- [12] Panagiotis C. Petris, Paul Becherer, Johannes G. E. M Fraaije, Alkane/water partition coefficient calculation based on the modified AM1 method and internal hydrogen bonding sampling using COSMO-RS, *Journal of Chemical Information and Modeling*, 61, 7, (2021), 3453-3462 <https://doi.org/10.1021/acs.jcim.0c01478>
- [13] Xiaoliang Pan, Pengfei Li, Junming Ho, Jingzhi Pu, Ye Mei, Yihan Shao, Accelerated computation of free energy profile at ab initio quantum mechanical/molecular mechanical accuracy via a semi-empirical reference potential. II. Recalibrating semi-empirical parameters with force matching, *Physical Chemistry Chemical Physics*, 21, 37, (2019), 20595-20605 <https://doi.org/10.1039/C9CP02593F>
- [14] Elton A. S. de Castro, Daniel A. B. de Oliveira, Sergio A. S. Farias, Ricardo Gargano, João B. L. Martins, Structure and electronic properties of azadirachtin, *Journal of Molecular Modeling*, 20, (2014), 2084 <https://doi.org/10.1007/s00894-014-2084-0>
- [15] Nathalia B. D. Lima, Gerd B. Rocha, Ricardo O. Freire, Alfredo M. Simas, RM1 semiempirical model: chemistry, pharmaceutical research, molecular biology and materials science, *Journal of the Brazilian Chemical Society*, 30, 4, (2019), 683-716 <https://doi.org/10.21577/0103-5053.20180239>
- [16] Qi Guan, Chunming Han, Daiying Zuo, Zengqiang Li, Qian Zhang, Yanpeng Zhai, Xuewei Jiang, Kai Bao, Yingliang Wu, Weige Zhang, Synthesis and evaluation of benzimidazole carbamates bearing indole moieties for antiproliferative and antitubulin activities, *European Journal of Medicinal Chemistry*,

- 87, (2014), 306–315
<https://doi.org/10.1016/j.ejmech.2014.09.071>
- [17] Martyn Inman, Christopher J. Moody, Indole synthesis—something old, something new, *Chemical Science*, 4, 1, (2013), 29–41
<https://doi.org/10.1039/C2SC21185H>
- [18] Robert J. Capon, Colin Skene, Edward Hsiang-Te Liu, Ernest Lacey, Jennifer H. Gill, Kirstin Heiland, Thomas Friedel, The isolation and synthesis of novel nematocidal dithiocyanates from an Australian marine sponge, *Oceanapia* sp, *The Journal of Organic Chemistry*, 66, 23, (2001), 7765–7769
<https://doi.org/10.1021/jo0106750>
- [19] Saw Simeon, Nathjanaan Jongkon, Construction of quantitative structure activity relationship (QSAR) Models to predict potency of structurally diversified janus kinase 2 inhibitors, *Molecules*, 24, 23, (2019), 4393 <https://doi.org/10.3390/molecules24234393>
- [20] Samina Kausar, Andre O. Falcao, Analysis and comparison of vector space and metric space representations in QSAR modeling, *Molecules*, 24, 9, (2019), 1698
<https://doi.org/10.3390/molecules24091698>
- [21] Hypercube, in, Hypercube Inc., Florida, USA, 2007,
- [22] José Diogo L. Dutra, Manoel A. M. Filho, Gerd B. Rocha, Alfredo M. Simas, Ricardo O. Freire, RM1 semiempirical quantum chemistry: Parameters for trivalent Lanthanum, Cerium and Praseodymium, *PLoS ONE*, 10, 7, (2015), e0124372
<https://doi.org/10.1371/journal.pone.0124372>
- [23] Ponco Iswanto, Maylani Permata Saputri, Vaulina Y. D. Eva, Sari Paramita, Quantitative Structure Activity Relationship Analysis of 1, 3, 4-Thiadiazole Derivatives as Anti-Inflammatory using Parameterized Model 3 Method, *Journal of Physics: Conference Series*, 2020
<https://doi.org/10.1088/1742-6596/1494/1/012031>
- [24] IBM, in, USA, 2016,
- [25] Eva Vaulina, Ponco Iswanto, Model QSAR Senyawa Fluorokuinolon Baru Sebagai Zat Antibakteri *Salmonella thypimurium*, *Molekul*, 1, 1, (2006), 10–18
<http://dx.doi.org/10.20884/1.jm.2006.1.1.17>
- [26] Ponco Iswanto, Eva Vaulina Yulistia Delsy, Ely Setiawan, Fiandy Aminullah Putra, Quantitative Structure-Property Relationship Analysis Against Critical Micelle Concentration of Sulfonate-Based Surfactant Based on Semiempirical Zindo/1 Calculation, *Molekul*, 14, 2, (2019), 78–83
<http://dx.doi.org/10.20884/1.jm.2019.14.2.467>
- [27] Sari Paramita, Maylani Permata, Eva Vaulina Y.D., Nasrokhah Nasrokhah, Ponco Iswanto, Pemilihan Metode Perhitungan Kimia Komputasi Semi-Empiris Untuk Pengembangan 1, 3, 4-Thiadiazole, *Indonesian Journal of Chemical Research*, 8, 1, (2020), 51–56
<https://doi.org/10.30598/10.30598/ijcr.2020.8-pon>
- [28] Adina Zharifah, Eysa Kusumowardani, Akbar Saputro, Devvi Sarwinda, Efficient dynamic molecular simulation using QSAR model to know inhibition activity in breast cancer medicine, *AIP Conference Proceedings*, 2017
<https://doi.org/10.1063/1.4991192>
- [29] Piotr Kawczak, Leszek Bober, Tomasz Bączek, The comparison of semiempirical and ab initio molecular modeling methods in activity and property evaluation of selected antimicrobial sulfonamides, *Medicinal Chemistry Research*, 28, (2019), 778–787
<https://doi.org/10.1007/s00044-019-02334-4>
- [30] Harno Dwi Pranowo, Abdul Kadir Rukmana Hetadi, *Pengantar Kimia Komputasi*, Universitas Gadjah Mada, Yogyakarta, 2011,
- [31] Nursalam Hamzah, Nur Syamsi Dhuha, Hasma Nur Putrianti, Hubungan Kuantitatif Skruktur Aktivitas Senyawa Turunan 1-Benzene Acyl-2-(Methylindol-3-Yl)-Bensimidazole Sebagai Inhibitor Pertumbuhan MCF-7, *Jurnal farmasi*, UIN Alauddin Makassar, Makassar, 2017
- [32] Soekardjo Siswandono, *Prinsip-prinsip rancangan obat*, Universitas Airlangga, Surabaya, 1995,
- [33] Emmy Yuanita, Ni Komang Tri Dharmayani, Maria Ulfa, Jufrizal Syahri, Quantitative structure–activity relationship (QSAR) and molecular docking of xanthone derivatives as anti-tuberculosis agents, *Journal of Clinical Tuberculosis and Other Mycobacterial Diseases*, 21, (2020), 100203
<https://doi.org/10.1016/j.jctube.2020.100203>
- [34] Renske P. J. Hoondert, Rik Oldenkamp, Dick de Zwart, Dik van de Meent, Leo Posthuma, QSAR - based estimation of species sensitivity distribution parameters: An exploratory investigation, *Environmental Toxicology and Chemistry*, 38, 12, (2019), 2764–2770 <https://doi.org/10.1002/etc.4601>



Ponco Iswanto 1 <ponco.iswanto@unsoed.ac.id>

[JKSA] Editor Decision - your submission needs to be revised

1 message

Dr. Mukhammad Asy'ari <asyari@live.undip.ac.id>
Reply-To: "Dr. Mukhammad Asy'ari" <asyari@live.undip.ac.id>
To: "Dr. Ponco Iswanto" <ponco.iswanto@unsoed.ac.id>

Thu, May 4, 2023 at 11:57 AM

Dear Dr. Ponco Iswanto

We have reached a decision regarding your submission to Jurnal Kimia Sains dan Aplikasi, "Quantitative Structure-Activity Relationship of 3-Thiocyanate-1H-Indoles Derived Compounds as Antileukemia by AM1, PM3 and RM1 Methods".

Our decision is that you need to revise this manuscript according to the feedback from the review.

How to prepare a revised manuscript.

- (1) Carefully read reviewer(s)' comments and make appropriate revisions.
- (2) Revised parts should be highlighted by changing the background color or the color of the words. The revised manuscript is in .docx format. Make sure that the revised manuscripts are cleared from the reviewer comments (none of the reviewers' comments are in the revised manuscript)

* How to prepare a response letter (a reply letter to the reviewer's question)*

- (1) Copy all the comments given by the reviewers.
- (2) Respond each comment, question, or suggestion one by one. The answers to the reviewer questions is in docx format. An example of the answers to the reviewer questions is attached.
- (3) Indicate changes you make in the manuscript.

* How to upload your revised manuscript. *

- (1) The revised manuscript and the response letter should be put together by compressing it into a zip file
- (2) Login JKSA through your OJS account
- (3) Find your active manuscript listed under "review" and click it
- (4) Click "In Review" listed under "Status"
- (5) Upload your file through "Upload Author Version" under "Editor Decision"

Please revise this manuscript no more than 2 (two) weeks from now. If within a month you do not revise this manuscript, then your manuscript will be treat as a new submission.

Regards

Dr. Mukhammad Asy'ari
Chemistry Department, Faculty of Sciences and Mathematics, Diponegoro University
Jl. Prof. Soedarto, SH., Tembalang, Semarang
Phone 081572250245
Fax -
asyari@live.undip.ac.id

Editor comment:
Please cite articles published by JKSA

Reviewer B:

In the manuscript "Quantitative Structure-Activity Relationship of 3-Thiocyanate-1H-Indoles Derived Compounds as Antileukemia by AM1, PM3 and RM1 Methods" Ponco et al. presented an interesting study pertaining to the drug development for Antileukemia. Whereas the work seems to be carefully done, a few points need attention before publication.

1. In Table 1, Author listed the IC₅₀ (μM) value. But no protein target was mentioned.

What's the targeted protein of 3-thiocyanato 1H-indoles derived compounds has been performed by experimental analysis to obtain those the IC₅₀ (μM) data.

2. Subsection 2.4. Structure optimization "The structure geometry was optimized by minimizing molecular energy to attain the most stable molecular conformation." How the author determined that the molecular structure of a compound reaches the stable conformation?

3. Subsection 2.5.Determination QSAR equation "The analysis was performed by applying the backward method on 16 fitting compounds, with IC₅₀ as the dependent variable and the descriptors as the independent variables". Authors should mention the systematic protocols in more detail to obtain those descriptors.

4. Authors concluded that AM1 semiempirical method is the best QSAR model for tested compounds. But, from those 20-compound derivatives, which compounds have promising inhibitory activity against leukemia?

Jurnal Kimia Sains dan Aplikasi
<http://ejournal.undip.ac.id/index.php/ksa>



Contoh Catatan Revisi oleh Author_English version.docx

405K



Ponco Iswanto 1 <ponco.iswanto@unsoed.ac.id>

[JKSA] Proofreading article [ID-53504]

5 messages

Jurnal Kimia Sains dan Aplikasi <jksa@live.undip.ac.id>
To: Ponco Iswanto 1 <ponco.iswanto@unsoed.ac.id>

Mon, Jun 5, 2023 at 2:35 PM

Yang kami hormati: Bapak Dr. Ponco Iswanto

Dengan email ini kami lampirkan artikel Bapak yang telah kami edit. Mohon untuk melakukan pengecekan kembali artikel tersebut. Apabila terdapat hal yang kurang sesuai, mohon untuk highlight dan beri komentar pada dokumen PDF dan kirim kembali kepada kami. Terima kasih.

Salam

Adi Darmawan
Editor in Chief
Jurnal Kimia Sains dan Aplikasi
email: jksa@live.undip.ac.id



 **Volume 26 No. 3 - 5 - Ponco Iswanto.pdf**
1546K

Ponco Iswanto 1 <ponco.iswanto@unsoed.ac.id>
To: Jurnal Kimia Sains dan Aplikasi <jksa@live.undip.ac.id>

Mon, Jun 5, 2023 at 6:29 PM

Baik, terima kasih.

ponco
[Quoted text hidden]

Ponco Iswanto 1 <ponco.iswanto@unsoed.ac.id>
To: Jurnal Kimia Sains dan Aplikasi <jksa@live.undip.ac.id>

Mon, Jun 5, 2023 at 8:35 PM

b Al Irsyad Al Islamiyyah Integrated Islamic Senior High School

seharusnya:

b Al Irsyad Al Islamiyyah Boarding School Senior High School

Dr. Ponco Iswanto
[Quoted text hidden]

 **Ponco_perbaiki_nama_sekolah_Volume 26 No. 3 - 5 - Ponco Iswanto.pdf**
1338K

Jurnal Kimia Sains dan Aplikasi <jksa@live.undip.ac.id>
To: Ponco Iswanto 1 <ponco.iswanto@unsoed.ac.id>

Tue, Jun 6, 2023 at 8:28 AM

Yth. Bapak Ponco Iswanto

Terima kasih atas koreksinya. Kami telah mengganti afiliasi b menjadi Al Irsyad Al Islamiyyah Boarding School Senior High School.

Bersamaan dengan email ini, kami lampirkan manuskrip yang telah kami edit.

Salam

Editorial Team

Jurnal Kimia Sains dan Aplikasi

email: jksa@live.undip.ac.id



[Quoted text hidden]



Volume 26 No. 3 - 5 - Ponco Iswanto.pdf
1546K

Ponco Iswanto 1 <ponco.iswanto@unsoed.ac.id>

To: Jurnal Kimia Sains dan Aplikasi <jksa@live.undip.ac.id>

Tue, Jun 6, 2023 at 5:46 PM

Terima kasih.

Ponco Iswanto

[Quoted text hidden]



Ponco Iswanto 1 <ponco.iswanto@unsoed.ac.id>

[JKSA] Submission Acknowledgement

1 message

Adi Darmawan <adidarmawan@live.undip.ac.id>
Reply-To: "Dr. Ponco Iswanto" <ponco.iswanto@unsoed.ac.id>
To: "Dr. Ponco Iswanto" <ponco.iswanto@unsoed.ac.id>

Tue, Apr 4, 2023 at 4:40 PM

Dear Dr. Ponco Iswanto

Thank you for submitting the manuscript, "Quantitative Structure-Activity Relationship of 3-Thiocyanate-1H-Indoles Derived Compounds as Antileukemia by AM1, PM3 and RM1 Methods" to Jurnal Kimia Sains dan Aplikasi. With the online journal management system that we are using, you will be able to track its progress through the editorial process by logging in to the journal web site:

Manuscript URL: <https://ejournal.undip.ac.id/index.php/ksa/author/submission/53504>

Username: dr_poncoiswanto1

Jurnal Kimia Sains dan Aplikasi (JKSA) follows the Open Access model. The publication cost should be covered by the author's institution or research funds. These Open Access charges replace subscription charges and allow the Jurnal Kimia Sains dan Aplikasi (JKSA) to make the valuable published materials freely accessible to all interested online visitors, especially the researchers and scholars.

The article processing charge is for professional copyediting; for efficient and thorough peer review; for immediate publication quickly upon acceptance; for the article to be freely and for universally accessible in various formats online.

Payment is made after successful review of the manuscript; acceptance email is sent to the corresponding author with payment details. No charge is made for articles that are rejected after peer review.

The article processing charge is Rp. 1.000.000, however JKSA waives the Article Processing Charges (APCs) for Non-Indonesian authors.

For papers submitted in Indonesian, if the paper is then accepted, it is requested that the paper be translated into English. We recommend that the paper be translated by a professional translator and for that there is a translation fee of Rp. 300,000

All authors are requested to submit a Copyright Transfer Agreement after the manuscript is confirmed to be accepted for publication. Copyright transfer agreement form can be downloaded from the Copyright notice section under the Submission tab.

Please don't hesitate to ask any question you have. You can contact me or send an email to jksa@live.undip.ac.id. Your emails will be respond in a very short time.

Thank you for considering this journal as a venue for your work.

Regards

Adi Darmawan
Jurnal Kimia Sains dan Aplikasi

Jurnal Kimia Sains dan Aplikasi
<http://ejournal.undip.ac.id/index.php/ksa>



Ponco Iswanto 1 <ponco.iswanto@unsoed.ac.id>

[JKSA] [ID-53504] - Editor Decision - your submission is accepted

4 messages

Dr. Mukhammad Asy'ari <asyari@live.undip.ac.id>
Reply-To: "Dr. Mukhammad Asy'ari" <asyari@live.undip.ac.id>
To: "Dr. Ponco Iswanto" <ponco.iswanto@unsoed.ac.id>

Mon, May 22, 2023 at 7:58 AM

Dear Dr. Ponco Iswanto

We have reached a decision regarding your submission to Jurnal Kimia Sains dan Aplikasi, "Quantitative Structure-Activity Relationship of 3-Thiocyanate-1H-Indoles Derived Compounds as Antileukemia by AM1, PM3 and RM1 Methods".

Our decision is to accept your submission. Congratulations

We are pleased to inform you that Jurnal Kimia Sains dan Aplikasi would like to publish your manuscript in our next issue. Before this manuscript proceeds to the layouting process, please pay attention to the following:

- Send a signed Copyright transfer agreement. You can download the Copyright transfer agreement in the Copyright notice section. Please fill in handwriting and sign and then scan and email us (email: jksa@live.undip.ac.id).
- Send a graphical abstract
- Optionally, send the supplementary files (if requested), including research instrument/materials, data analysis, data sets, and source text.
- Complete the metadata including ALL authors' names, affiliations, ORCID URLs, Google scholar URLs, and countries, article title, abstract, keywords, and references.

In accordance to the Journal policy, you are required to immediately pay the publication fee of Rp 1.000,000 by transfer to the following bank account:

Bank name: PT. BANK NEGARA INDONESIA (PERSERO) TBK
SWIFT Code: BNINIDJAUDS
Account number: 423759081
Account holder: Adi Darmawan
Bank Address: BNI Undip Branch, Semarang, Indonesia

Please send the proof of remittance by email to the editorial office of Jurnal Kimia Sains dan Aplikasi (email: jksa@live.undip.ac.id). JKSA waives the article processing charge for non-Indonesian author. After payment, in few days you will receipt email for the further process, i.e. copy-editing, layout and proofreading. Once your manuscript is moved to publishing, our production editor will keep you informed of your article's progress in the production process. You may also receive a proof of your manuscript for final review.

Thank you for your valuable contribution to the journal.

Regards

JKSA's Editor,

Dr. Mukhammad Asy'ari
Chemistry Department, Faculty of Sciences and Mathematics, Diponegoro University
Jl. Prof. Soedarto, SH., Tembalang, Semarang
Phone: 081572250245
Email: asyari@live.undip.ac.id

Jurnal Kimia Sains dan Aplikasi
<http://ejournal.undip.ac.id/index.php/ksa>

Ponco Iswanto 1 <ponco.iswanto@unsoed.ac.id>
To: "Dr. Mukhammad Asy'ari" <asyari@live.undip.ac.id>

Thu, May 25, 2023 at 5:39 AM

Dear Dr. Mukhammad Asy'ari,

I want to ask something...

- Complete the metadata including ALL authors' names, affiliations, ORCID URLs, Google scholar URLs, and countries, article title, abstract, keywords, and references.

Could I make in .docx format?

Thank you very much.

Best wishes

Dr. Ponco Iswanto

[Quoted text hidden]

Mukhammad Asy'ari <asyari@live.undip.ac.id>
To: Ponco Iswanto 1 <ponco.iswanto@unsoed.ac.id>

Thu, May 25, 2023 at 6:28 AM

Dear Dr. Iswanto Poncho,

Previously, I would like to congratulate you on the acceptance of your article at JKSA, regarding your question, the steps are as follows:

Login to the JKSA website using your username and password.

Go to: Review (Active) -- click the title of the article -- click edit metadata -- complete: ORCID ID SINTA ID... etc. and add the author and complete their metadata fields.

I hope that this answer will help you, thank you.

Best Regards,

Dr. Mukhammad Asy'ari, M.Si

Biochemistry Lab. Chemistry Department,

Faculty of Science and Mathematics

Diponegoro University, Semarang, Central Java, Indonesia

HP: 081572250245

Email: asyari@live.undip.ac.id or asharychem@gmail.com

Editor Staff

Jurnal Kimia Sains dan Aplikasi (JKSA)

Email: jksa@live.undip.ac.id



[Quoted text hidden]

Ponco Iswanto 1 <ponco.iswanto@unsoed.ac.id>
To: Mukhammad Asy'ari <asyari@live.undip.ac.id>

Thu, May 25, 2023 at 9:29 AM

Alhamdulillah...

Thank you for your email..

Best Regards..

Dr. Ponco Iswanto

[Quoted text hidden]



Ponco Iswanto 1 <ponco.iswanto@unsoed.ac.id>

[JKSA] [ID-53504] Revised Version Acknowledgement

1 message

Adi Darmawan <adidarmawan@live.undip.ac.id>
Reply-To: "Dr. Ponco Iswanto" <ponco.iswanto@unsoed.ac.id>
To: "Dr. Ponco Iswanto" <ponco.iswanto@unsoed.ac.id>

Tue, May 9, 2023 at 6:22 AM

Dr. Ponco Iswanto:

Thank you for submitting the revision of manuscript, "Quantitative Structure-Activity Relationship of 3-Thiocyanate-1H-Indoles Derived Compounds as Antileukemia by AM1, PM3 and RM1 Methods" to Jurnal Kimia Sains dan Aplikasi. With the online journal management system that we are using, you will be able to track its progress through the editorial process by logging in to the journal web site:

Manuscript URL: <https://ejournal.undip.ac.id/index.php/ksa/author/submission/53504>

Username: dr_poncoiswanto1

Editor: Mukhammad Asy'ari

If you have any questions, please contact me. Thank you for considering this journal as a venue for your work.

Adi Darmawan
Jurnal Kimia Sains dan Aplikasi

Jurnal Kimia Sains dan Aplikasi
<http://ejournal.undip.ac.id/index.php/ksa>