


**ISSN: 2332-2071**  
**Volume 10 Number 4 2022**



# ***Mathematics and Statistics***

*<http://www.hrpublishing.org>*

**Horizon Research Publishing, USA**



## Mathematics and Statistics

Mathematics and Statistics is an international peer-reviewed journal that publishes original and high-quality research papers in all areas of mathematics and statistics. As an important academic exchange platform, scientists and researchers can know the most up-to-date academic trends and seek valuable primary sources for reference. The subject areas include, but are not limited to the following fields: Algebra, Applied Mathematics, Approximation Theory, Combinatorics, Computational Statistics, Computing in Mathematics, Operations Research Methodology, Discrete Mathematics, Mathematical Physics, Geometry and Topology, Logic and Foundations of Mathematics, Number Theory, Numerical Analysis, Probability Theory, Central Limit Theorem Computation, Sample Survey, Statistical Modelling, Statistical Theory.

## General Inquires

Publish with HRPUB, learn about our policies, submission guidelines etc.

Email: [editor@hrpub.org](mailto:editor@hrpub.org)

Tel: +1-626-626-7940

## Subscriptions (Electronic Version Only)

**Journal Title:** Mathematics and Statistics

**Journal's Homepage:** [http://www.hrpub.org/journals/jour\\_info.php?id=34](http://www.hrpub.org/journals/jour_info.php?id=34)

**Publisher:** Horizon Research Publishing Co.,Ltd

**Publication Frequency:** bimonthly

**Electronic Version:** freely online available at [http://www.hrpub.org/journals/jour\\_info.php?id=34](http://www.hrpub.org/journals/jour_info.php?id=34)

## Online Submission

Manuscripts should be submitted by Online Manuscript Tracking System (<http://www.hrpub.org/submission.php>).

If you are experiencing difficulties during the submission process, please feel free to contact the editor at [submission@hrpub.org](mailto:submission@hrpub.org).

## Copyright

Authors retains all copyright interest or it is retained by other copyright holder, as appropriate and agrees that the manuscript remains permanently open access in HRPUB 's site under the terms of the Creative Commons Attribution International License (CC BY). HRPUB shall have the right to use and archive the content for the purpose of creating a record and may reformat or paraphrase to benefit the display of the record.

## Creative Commons Attribution License (CC-BY)

All articles published by HRPUB will be distributed under the terms and conditions of the Creative Commons Attribution License(CC-BY). So anyone is allowed to copy, distribute, and transmit the article on condition that the original article and source is correctly cited.

## Open Access

Open access is the practice of providing unrestricted access to peer-reviewed academic journal articles via the internet. It is also increasingly being provided to scholarly monographs and book chapters. All original research papers published by HRPUB are available freely and permanently accessible online immediately after publication. Readers are free to copy and distribute the contribution under creative commons attribution-non commercial licence. Authors can benefit from the open access publication model a lot from the following aspects:

- High Availability and High Visibility-free and unlimited accessibility of the publication over the internet without any restrictions;
- Rigorous peer review of research papers----Fast, high-quality double blind peer review;
- Faster publication with less cost----Papers published on the internet without any subscription charge;
- Higher Citation----open access publications are more frequently cited.

# Mathematics and Statistics

---

## Editor-in-Chief

Prof. Dshalalow Jewgeni

Florida Inst. of Technology, USA

## Honorary Editor-in-Chief

Prof. Ravi P. AGARWAL

Texas A&M University, USA

## Deputy Editor-in-Chief

Prof. Sergey Sudoplatov

Sobolev Institute of Mathematics, Russia

Prof. Francisco Gallego Lupianez

Universidad Complutense de Madrid, Spain

Prof. Irina Burova

St. Petersburg State University, Russia

## Members of Editorial Board

Prof. Rachid El Harti

Hassan First University, Morocco

Prof. H. M. Srivastava

University of Victoria, Canada

Prof. Veli Shakhmurov

Okan University, Turkey

Prof. Liang-yun Zhang

Nanjing Agricultural University, China

Prof. Birol Altın

Gazi University, Turkey

Prof. Emanuele Galligani

University of Modena and Reggio Emilia, Italy

Prof. Ramalingam Shanmugam

Texas State University, USA

Prof. Manoj Sahnı

Pandit Deendayal Petroleum University, India

Prof. Ayhan Esl

Malatya Turgut Ozal University, Turkey

Prof. Mahmoud Mohammed Mostafa El-Borai

Alexandria University, Egypt

Prof. Serguei Pergamenchtchikov

Rouen University, France

Prof. Allaberen Ashyralyev

Near East University, Turkey

Prof. Semra Erpolat Taşabat

Mimar Sinan Fine Arts University, Turkey

Prof. Olivier Olela Otafudu

University of the Witwatersrand, South Africa

Prof. Leonid N. Yasnitsky

Perm State University, Russia

Prof. Andriy Zagorodnyuk

Vasyl Stefanyk Precarpathian National University, Ukraine

Dr. Debaraj Sen

Concordia University, Canada

Dr. Antonio Maria Scarfone

National Research Council, Italy

Dr. Vadim Kryakvin

Southern Federal University, Russia

Dr. Azam Imomov

Karshi State University, Uzbekistan

Dr. Evgeny Maleko

Magnitogorsk State Technical University, Russia

Dr. İmdat İçsan

Giresun University, Turkey

Dr. Pradeep Kumar Pandey

Jaypee University of Information Technology, India

Dr. N. Gunasekaran

Shibaura Institute of Technology, Japan

## Mathematics and Statistics

Volume 10 Number 4 2022

**Markowitz Random Set and Its Application to the Paris Stock Market Prices***(<https://www.doi.org/10.13189/ms.2022.100401>)*

Ahssaine Bourakadi, Naima Soukher, Baraka Achraf Chakir, Driss Mentagui.....701

**Solving Lorenz System by Using Lower Order Symmetrized Runge-Kutta Methods***(<https://www.doi.org/10.13189/ms.2022.100402>)*

N. Adan, N. Razali, N. A. Zainuri, N. A. Ismail, A. Gorgey, N. I. Hamdan .....713

**Advancement of Generalized Method of Moment Estimation (GMM) For Spatial Dynamic Panel Simultaneous Equations Models with Fixed Time Effect***(<https://www.doi.org/10.13189/ms.2022.100403>)*

Dwi Endah Kusriani, Setiawan, Heri Kuswanto, Budi Nurani Ruchjana .....729

**Analysis of IBFS for Transportation Problem by Using Various Methods***(<https://www.doi.org/10.13189/ms.2022.100404>)*

S. K. Sharma, Keshav Goel .....741

**Cluster Analysis on Various Cluster Validity Indexes with Average Linkage Method and Euclidean Distance (Study on Compliant Paying Behavior of Bank X Customers in Indonesia 2021)***(<https://www.doi.org/10.13189/ms.2022.100405>)*

Solimun Solimun, Adji Achmad Rinaldo Fernades .....747

**Perfect Codes in the Spanning and Induced Subgraphs of the Unity Product Graph***(<https://www.doi.org/10.13189/ms.2022.100406>)*

Mohammad Hassan Mudaber, Nor Haniza Sarmin, Ibrahim Gambo.....754

**Uncertainty Optimization-Based Rough Set for Incomplete Information Systems***(<https://www.doi.org/10.13189/ms.2022.100407>)*

Arvind Kumar Sinha, Pradeep Shende .....759

**Double Duplication of Special Classes of Cycle Related Graphs***(<https://www.doi.org/10.13189/ms.2022.100408>)*

R.Kuppan, L.Shobana.....773

**Bayesian Model Averaging in Modeling of State Specific Failure Rates in HIV/AIDS Progression***(<https://www.doi.org/10.13189/ms.2022.100409>)*

Nahashon Mwirigi, Prof. Richard Simwa, Dr. Mary Wainaina, Dr. Stanley Sewe .....782

<b>Construction and Selection of Double Inspection Single Sampling Plan for an Independent Process Using Bivariate Poisson Distribution</b> ( <a href="https://www.doi.org/10.13189/ms.2022.100410">https://www.doi.org/10.13189/ms.2022.100410</a> ) D. Senthilkumar, P. Sabarish .....	799
<b>Subset Intersection Group Testing Strategy</b> ( <a href="https://www.doi.org/10.13189/ms.2022.100411">https://www.doi.org/10.13189/ms.2022.100411</a> ) Maheswaran Srinivasan .....	808
<b>A New Ranking Approach for Solving Fuzzy Transportation Problem with Pentagonal Fuzzy Number</b> ( <a href="https://www.doi.org/10.13189/ms.2022.100412">https://www.doi.org/10.13189/ms.2022.100412</a> ) V. Vidhya, K. Ganesan .....	816
<b>Analysis of Limiting Ratios of Special Sequences</b> ( <a href="https://www.doi.org/10.13189/ms.2022.100413">https://www.doi.org/10.13189/ms.2022.100413</a> ) A. Dinesh Kumar, R. Sivaraman .....	825
<b>Generalized Biased Estimator for Beta Regression Model: Simulation and Application</b> ( <a href="https://www.doi.org/10.13189/ms.2022.100414">https://www.doi.org/10.13189/ms.2022.100414</a> ) Rasha A. Farghali, Samah M. Abo-El-Hadid .....	833
<b>The Power and Its Graph Simulations on Discrete and Continuous Distributions</b> ( <a href="https://www.doi.org/10.13189/ms.2022.100415">https://www.doi.org/10.13189/ms.2022.100415</a> ) Budi Pratikno, Nailatul Azizah, Avita Nur Azizah .....	844
<b>A Study on Sylow Theorems for Finding out Possible Subgroups of a Group in Different Types of Order</b> ( <a href="https://www.doi.org/10.13189/ms.2022.100416">https://www.doi.org/10.13189/ms.2022.100416</a> ) Md. Abdul Mannan, Md. Amanat Ullah, Uttam Kumar Dey, Mohammad Alauddin .....	851
<b>Dynamics of Nonlinear Operator Generated by Lebesgue Quadratic Stochastic Operator with Exponential Measure</b> ( <a href="https://www.doi.org/10.13189/ms.2022.100417">https://www.doi.org/10.13189/ms.2022.100417</a> ) Nur Zatul Akmar Hamzah, Siti Nurlaili Karim, Mathuri Selvarajoo, Noor Azida Sahabudin .....	861
<b>Numerical Solution of Nonlinear Fredholm Integral Equations Using Half-Sweep Newton-PKSOR Iteration</b> ( <a href="https://www.doi.org/10.13189/ms.2022.100418">https://www.doi.org/10.13189/ms.2022.100418</a> ) Labiyyana Hanif Ali, Jumat Sulaiman, Azali Saudi, Xu Ming Ming .....	868
<b>2-Odd Labeling of Graphs Using Certain Number Theoretic Concepts and Graph Operations</b> ( <a href="https://www.doi.org/10.13189/ms.2022.100419">https://www.doi.org/10.13189/ms.2022.100419</a> ) Ajaz Ahmad Pir, Tabasum Mushtaq, A. Parthiban .....	875
<b>Evolution Equations of Pseudo Spherical Images for Timelike Curves in Minkowski 3-Space</b> ( <a href="https://www.doi.org/10.13189/ms.2022.100420">https://www.doi.org/10.13189/ms.2022.100420</a> ) H. S. Abdel-Aziz, H. Serry, M. Khalifa Saad .....	884

# Mathematics and Statistics

## Call for Papers

Mathematics and Statistics is an international peer-reviewed journal that publishes original and high-quality research papers in all areas of mathematics and statistics. As an important academic exchange platform, scientists and researchers can know the most up-to-date academic trends and seek valuable primary sources for reference.

### Aims & Scope


- Algebra
- Applied Mathematics
- Approximation Theory
- Combinatorics
- Computational Statistics
- Computing in Mathematics
- Operations Research Methodology
- Discrete Mathematics
- Mathematical Physics
- Geometry and Topology
- Logic and Foundations of Mathematics
- Number Theory
- Numerical Analysis
- Probability Theory
- Central Limit Theorem Computation
- Sample Survey
- Statistical Modelling
- Statistical Theory

### Editorial Board

- |   |   |
|---|---|
| Prof. Dshalalow Jewgeni                 | Florida Inst. of Technology, USA                          |
| Prof. Ravi P. AGARWAL                   | Texas A&M University, USA                                 |
| Prof. Sergey Sudoplatov                 | Sobolev Institute of Mathematics, Russia                  |
| Prof. Francisco Gallego Lupianez        | Universidad Complutense de Madrid, Spain                  |
| Prof. Irina Burova                      | St. Petersburg State University, Russia                   |
| Prof. Rachid El Harti                   | Hassan First University, Morocco                          |
| Prof. H. M. Srivastava                  | University of Victoria, Canada                            |
| Prof. Veli Shakhmurov                   | Okan University, Turkey                                   |
| Prof. Liang-yun Zhang                   | Nanjing Agricultural University, China                    |
| Prof. Birol Altin                       | Gazi University, Turkey                                   |
| Prof. Emanuele Galligani                | University of Modena and Reggio Emilia, Italy             |
| Prof. Ramalingam Shanmugam              | Texas State University, USA                               |
| Prof. Manoj Sahni                       | Pandit Deendayal Petroleum University, India              |
| Prof. Ayhan Esil                        | Malatya Turgut Ozal University, Turkey                    |
| Prof. Mahmoud Mohammed Mostafa El-Borai | Alexandria University, Egypt                              |
| Prof. Serguei Pergamenchikov            | Rouen University, France                                  |
| Prof. Allaberen Ashyralyev              | Near East University, Turkey                              |
| Prof. Semra Erpolat Tasabat             | Mimar Sinan Fine Arts University, Turkey                  |
| Prof. Olivier Olela Otafudu             | University of the Witwatersrand, South Africa             |
| Prof. Leonid N. Yasnitsky               | Perm State University, Russia                             |
| Prof. Andriy Zagorodnyuk                | Vasyl Stefanyk Precarpathian National University, Ukraine |
| Dr. Debaraj Sen                         | Concordia University, Canada                              |
| Dr. Antonio Maria Scarfone              | National Research Council, Italy                          |
| Dr. Vadim Kryakvin                      | Southern Federal University, Russia                       |

### Manuscripts Submission

Manuscripts to be considered for publication have to be submitted by Online Manuscript Tracking System(<http://www.hrpub.org/submission.php>). If you are experiencing difficulties during the submission process, please feel free to contact the editor at [submission@hrpub.org](mailto:submission@hrpub.org).



### Contact Us

Horizon Research Publishing  
2880 ZANKER RD STE 203  
SAN JOSE, CA 95134  
USA  
Email: [editor@hrpub.org](mailto:editor@hrpub.org)

Submit your paper at [http://www.hrpub.org/journals/jour\\_submitmanuscript.php?id=34](http://www.hrpub.org/journals/jour_submitmanuscript.php?id=34)

Available Online [http://www.hrpub.org/journals/jour\\_info.php?id=34](http://www.hrpub.org/journals/jour_info.php?id=34)



# The Power and Its Graph Simulations on Discrete and Continuous Distributions

Budi Pratikno\*, Nailatul Azizah, Avita Nur Azizah

Department of Mathematics, Faculty of Mathematics and Natural Science, Jenderal Soedirman University, Purwokerto, Indonesia

Received May 17, 2022; Revised June 17, 2022; Accepted July 22, 2022

## Cite This Paper in the Following Citation Styles

(a): [1] Budi Pratikno, Nailatul Azizah, Avita Nur Azizah, "The Power and Its Graph Simulations on Discrete and Continuous Distributions," *Mathematics and Statistics*, Vol. 10, No. 4, pp. 844 - 850, 2022. DOI: 10.13189/ms.2022.100415.

(b): Budi Pratikno, Nailatul Azizah, Avita Nur Azizah (2022). *The Power and Its Graph Simulations on Discrete and Continuous Distributions. Mathematics and Statistics*, 10(4), 844 - 850. DOI: 10.13189/ms.2022.100415.

Copyright©2022 by authors, all rights reserved. Authors agree that this article remains permanently open access under the terms of the Creative Commons Attribution License 4.0 International License

**Abstract** We determined the power and its graph simulations on the discrete Poisson and Chi-square distributions. There are four important steps of the research methodology summarized as follow: (1) determine the sufficient statistics (if possible), (2) create the rejection area (UMPT test is sometime used), (3) derive the formula of the power, and (4) determine the graphs using the data (in simulation). The formula of the power and their curves are then created using *R* code. The result showed that the power of the illustration of the discrete (Binomial distribution) depended on the number of trials  $n$  and bound of the rejection area. The curve of the power is sigmoid (*S*-curve) and tends to be zero when parameter shape ( $\theta$ ) is greater than 0.4. It decreases (started from  $\theta = 0.2$ ) as the parameter theta increases. In the Poisson context, the curve of the power of the Poisson distribution is not *S*-curve, and it only depends on the parameter shape  $\lambda$ . We note that the curve of the power of the Poisson is quickly to be one for  $n$  greater than 2 and  $\lambda$  less than 10. In this case, the size of the Poisson distribution is greater than 0.05, so it is not a reasonable thing even the power is close to be one. In this context, we have to choose the maximum power and minimum size. In the context of Chi-square distribution, the graph of the power and size functions depend on rejection region boundary ( $k$ ). Here, we note that skewness of the *S*-curve is positive as the  $k$  increases. Similarly, the size also depends on the  $k$  (and constant), and it decrease as the  $k$  increases. We here also noted that the power is quickly to be one for large degree of freedom ( $r$ ).

**Keywords** Poisson Discrete Distribution, Chi-Square

Continuous Distribution, Parameter Shape, R-code

## 1. Introduction

In the theory of statistics, there are three important concepts of the hypothesis testing in rejecting or accepting null hypothesis ( $H_0$ ), namely (1) a probability error type I ( $\alpha$ ), (2) a probability error type II ( $\beta$ ) and (3) power of the test ( $\pi(\theta)$ ) (Wackerly, et al. [5]). Here, the power is a significant method to test the hypothesis on parameter shape. We then study more details about the power of the hypothesis testing on some distributions. Furthermore, the power is defined as a probability to reject  $H_0$  under  $H_1$  on  $H_0 : \theta = \theta_0$  versus  $H_1 : \theta \neq \theta_0$ , for parameter shape  $\theta$  (Wackerly, et al. [5]).

Following to the previous research, many authors, such as Pratikno [2], Khan and Pratikno [22] and Khan [12], used the power in testing intercept with non-sample prior information (NSPI). They used the probability integral of the cumulative distribution function (cdf) of the continuous distributions. Moreover, Pratikno [2] and Khan et al. [11] used the power and size to compute the cdf of the bivariate noncentral *F* (BNCF) distribution in multivariate and multiple regression models. Here, many authors, such as Khan [12, 13, 14], Khan and Saleh [15, 16, 17, 20, 21], Khan and Hoque [19], Saleh [1], Yunus [6], and Yunus and

Khan [7, 8, 9, 10], have contributed to the research of the power in the context of the hypothesis area. In the context of the hypothesis testing with NSPI on multivariate and multiple regression models, Pratikno [2] and Khan et al. [11] used the BNCF distribution to compute the power using *R-code*. This is due to the computation of the probability integral of the probability distribution function (pdf) and cdf of the BNCF distribution are very complicated and hard (see Pratikno [2] and Khan et al. [18]), so the *R* code is used.

Unlike previous research analyze focusing on continuous distribution, we only consider two distributions (Poisson and Chi-square). To illustrate the simple power, we present the power of the Binomial distribution. Furthermore, the steps to compute the power of the Binomial, Poisson and Chi-square distributions are similar to the previous theory are: (1) we have to determine the sufficient statistics (if possible), (2) we create the rejection area using uniformly the most powerful test (UMPT, if needed), (3) we derive the formula of the power of the discrete and continuous distributions, and (4) finally, we graphically analysed the power. A simulation is then conducted using the generate data.

The concept of power and size (as initiate, Binomial distribution) of the testing hypothesis is presented in Section 2. The derived formula and the analysis of the power of the power and size of the Binomial, Poisson and Chi-square distributions are then given in Section 3. The conclusion is in Section 4.

## 2. The Power and Size of One-Side Hypothesis Testing

Following Pratikno [2], Khan [12,13,14], Khan and Saleh [15,16,17,20,21], Khan and Hoque [19], Saleh [1], Yunus [6], and Yunus and Khan [7, 8, 9, 10], we noted that the power and size of the tests provide a significant method to find the conclusion of the hypothesis testing parameter shape. Here, we must choose the maximum power and minimum size as an indicator. Here, the power and size are defined as a probability to reject  $H_0$  under  $H_1$  in testing hypothesis, and probability to reject  $H_0$  under  $H_0$ , respectively (Wackerly, et al. [5]). Following Pratikno [2], we then write the power and size in testing hypothesis,  $H_0 : \theta = \theta_0$  versus  $H_1 : \theta > \theta_0$  (or  $H_1 : \theta = \theta_1$ ) as, respectively,

$$\begin{aligned}\pi(\theta_1) &= P(\text{reject } H_0 \mid \text{under } H_1) \\ &= P(\text{reject } H_0 \mid \theta = \theta_1)\end{aligned}\quad (1)$$

$$\begin{aligned}\pi(\theta_0) &= P(\text{reject } H_0 \mid \text{under } H_0) \\ &= P(\text{reject } H_0 \mid \theta = \theta_0)\end{aligned}\quad (2)$$

where  $\alpha$  is probability of type error I and  $\beta$  is probability of type error II. The details of the power and size in testing coefficient parameters on the regression models are found on Pratikno [2], and the power and size on several continuous distributions are also found Pratikno et al. [3,4].

## 3. The Power and Size of Discrete and Continuous Distributions

### 3.1. The Power and Size of the Binomial Distribution


To illustrate the simple power, we firstly derived the formula of the power and size of the discrete Binomial distribution. The power and size of this distribution are computed in one-side hypothesis testing on several  $n$  and bound of the rejection areas. Let,  $X_i$  follows Bernoulli distribution with parameter  $\theta$ . Take a trial  $n=12$ , then

$$Y = \sum_{j=1}^{n=12} X_j \text{ follows Binomial distribution with } n=12$$

and  $p = \theta$  and is written as  $Y : B(n, \theta)$ . Here, we decide (an example  $\theta = 0.7$ ) to test  $H_0 : \theta = 0.7$  versus  $H_1 : \theta > 0.7$  (as  $\theta_1$ ), with rejection area  $\{(x_1, \dots, x_{12}) : Y \leq 5\}$ , therefore the power function on the binomial distribution is then given as

$$\begin{aligned}\pi(\theta) &= P(\text{reject } H_0 \mid \text{under } H_1 : \theta) \\ &= \sum_{y=0}^5 \binom{12}{y} \theta^y (1-\theta)^{12-y} \\ &= \binom{12}{0} \theta^0 (1-\theta)^{12} + \binom{12}{1} \theta^1 (1-\theta)^{11} + \dots + \binom{12}{5} \theta^5 (1-\theta)^7 \\ &= \binom{12}{0} \theta^0 (1-\theta)^{12} + \binom{12}{1} \theta^1 (1-\theta)^{11} + \dots + \binom{12}{5} \theta^5 (1-\theta)^7 \\ &= (1-\theta)^7 (1 + 7\theta + 28\theta^2 + 84\theta^3 + 210\theta^4 + 462\theta^5)\end{aligned}\quad (3)$$

Using the equation (3) and *R-code*, we then produced the graphs (curves) of the power in figure 1.



**Figure 1.** The power of the Binomial Distribution at several  $n$  and  $Y=5$  and 9

Figure 1 showed that the curve of the power of the Binomial distribution (figure 1 (a) and (b)) are sigmoid ( $S$  curve) and depends on the number of trials ( $n$ ) and the bound of the rejection area ( $Y$ ). They are going to be zero for  $\theta > 0.4$ . The curve decreases (started from  $\theta = 0.2$ ) as the parameter increases. From figure 1 (a) and (b), it is clear that both  $n$  and  $Y$  have significant effect on the shape of the curve (see figure 1 (a) and (b), they move to the right). Here, the maximum power is one and the minimum power is zero. The size is then produced using the equation (3) under  $H_0$ , as

$$\begin{aligned}\alpha &= P(\text{reject } H_0 | \text{under } H_0) \\ &= P(Y \leq 5 | \theta = 0.7) \\ &= 0.04.\end{aligned}$$

It is clear that the size is constant and is less than 0.05, as expected

### 3.2. The Power and Size of the Poisson Distribution

Let,  $X_1, K, X_n$  follow Poisson distribution, the

probability distribution function (pdf) of random variable  $X$  is then given by

$$f(x, \lambda) = \frac{e^{-\lambda} \lambda^{x_i}}{x_i!} \quad (4)$$

with  $x = 0, 1, 2, \dots$ , and  $\lambda > 0$ . The pdf curve of the Poisson distribution (positive skew) tends to be normal for large values  $\lambda$ , where the center of the pdf curve always moves to the right when  $\lambda$  increases.

To find the power, we then derive sufficient statistics and a rejection area using factorization theorem and UMP test. In other words, let  $S$  be sufficient statistics. The join distribution of the Poisson distribution is then expressed as

$$f(x_1, \dots, x_n; \lambda) = g(s, \lambda) h(x_1, \dots, x_n) \quad (5)$$

where the pdf of the join distribution of the Poisson distribution is

$$\begin{aligned}f(x_1, \dots, x_n; \lambda) &= \prod_{i=1}^n f(x_i, \lambda) \\ &= \prod_{i=1}^n \frac{e^{-\lambda} \lambda^{x_i}}{x_i!} = \frac{e^{-n\lambda} \lambda^{\sum x_i}}{\prod_{i=1}^n x_i!}.\end{aligned} \quad (6)$$

We therefore conclude that  $s = \sum_{i=1}^n x_i$  sufficient statistics, this is due to the equation (6) can be expressed as

$$\begin{aligned}f(x_1, \dots, x_n; \lambda) &= \frac{e^{-n\lambda} \lambda^{\sum x_i}}{\prod_{i=1}^n x_i!} \\ &= \left( \frac{e^{-n\lambda}}{\prod_{i=1}^n x_i!} \right) (\lambda^s) \\ &= h(x_1, \dots, x_n) g(s, \lambda)\end{aligned} \quad (7)$$

The rejection area is then derived using *uniformly the most powerful* (UMP) test as follow. Using the properties of *maximum likelihood ratio* (MLR) of the

$f(x_1, \dots, x_n; \lambda)$  on  $S = \sum_{i=1}^n X_i$   $\left( \sum_{i=1}^n X_i > k \right)$  and


UMP-test, we then get the probability to reject  $H_0$  under  $H_0$  (the size or  $\alpha$ ) and the probability to reject  $H_0$  under  $H_1$  (the power) in testing  $H_0: \lambda = \lambda_0$  versus

$H_1 : \lambda > \lambda_0$ , are, respectively,

$$\begin{aligned}\alpha &= \alpha(\lambda) = \alpha^* = P\left(\sum_{i=1}^n X_i > k \mid \lambda_0\right) \\ &= P\left(\sum_{i=1}^n X_i > \text{Pois}(n\lambda) \mid \lambda_0\right) \\ &= \sum_{x=0}^n \frac{e^{-(n\lambda_0)} (n\lambda_0)^x}{x!}\end{aligned}\quad (8)$$

$$\begin{aligned}\pi(\lambda) &= (\text{Probability reject } H_0 \text{ under } H_1) \\ &= P\left(\sum_{i=1}^n X_i > \text{Poi}(n\lambda) \mid \lambda\right) \\ &= 1 - P\left(\sum_{i=1}^n X_i \leq \text{Pois}(n\lambda) \mid \lambda\right) \\ &= 1 - \left(\sum_{x=0}^n \frac{e^{-n\lambda} (n\lambda)^x}{x!} \mid \lambda\right)\end{aligned}\quad (9)$$

Using the equation (8) and (9), we presented the graph of the power of the Poisson distribution (figure 2), and the value of the size and power for  $n=3$  in testing  $H_0 : \lambda = 1$  versus  $H_1 : \lambda > 3$ , respectively, as



**Figure 2.** The power of the Poisson Distribution at several  $n$

$$\alpha = 1 - \left(\sum_{x=0}^n \frac{e^{-n\lambda_0} (n\lambda_0)^x}{x!}\right) = 0.40779 > 0.05,$$

$$\pi(\lambda) = 1 - \left(\sum_{x=0}^n \frac{e^{-n\lambda} (n\lambda)^x}{x!}\right) = 0.99985; 1.$$

From figure 2, we see that the power of the Poisson distribution tends to be 1 when  $\lambda < 10$ . Here, the simulation of the  $n$  has not influenced to the curve of the

power yet. Thus, we conclude value of  $n$  whether small or large does not affect to change the shape of the curve of the power. Similarly, for large  $\lambda$ , the shape of the curve of the power does not change. Here, the size 0.408 is greater than 0.05 (too high), and it is not as expected.

### 3.3. The Power and Size of the Chi-square Distribution

Let,  $X$  be a random variable that follows Chi-square distribution. The probability distribution function (pdf) of random variable  $X$  is then given by

$$f(x) = \frac{1}{2^{\frac{r}{2}} \Gamma\left(\frac{r}{2}\right)} x^{\frac{r}{2}-1} e^{-x/2}, x \geq 0 \quad (10)$$

with  $r$  is the degree of freedom (as parameter). The cdf of this distribution is the written as

$$F_x(x) = \int_0^x f(x) dx = \int_0^x \frac{1}{2^{\frac{r}{2}} \Gamma\left(\frac{r}{2}\right)} x^{\frac{r}{2}-1} e^{-x/2} dx$$


The power of this distribution in testing parameter shape  $H_0 : r = r_0$  versus  $H_1 : r > r_0$  ( $r_0$  is determined as 1), is then obtained as

$$\begin{aligned}\pi(r) &= P(\text{reject } H_0 \mid \text{under } H_1, r_0 = r) \\ &= P(S > k \mid r) \\ &= 1 - P(S \leq k \mid r) \\ &= 1 - \int_0^k \frac{1}{2^{\frac{r}{2}} \Gamma\left(\frac{r}{2}\right)} s^{\frac{r}{2}-1} e^{-s/2} ds \\ &= 1 - \left[ -\frac{1}{\Gamma\left(\frac{r}{2}\right)} \left[ \tau\left(\frac{r}{2}, v^{2/r}\right) \right]_0^k \right] \\ &= 1 + \frac{\left[ \tau\left(\frac{r}{2}, \frac{s}{2}\right) \right]_0^k}{\Gamma\left(\frac{r}{2}\right)}\end{aligned}\quad (11)$$

Here,  $s = \sum_{i=1}^n x_i$  is a sufficient statistics and


$v = \left(\frac{s}{2}\right)^{r/2}$ . Using the equation (11), we then produced

the graphs of the power and size as presented in figure 3 and figure 4.



**Figure 3.** The power of the Chi-square Distribution at several  $k$

From figure 3, we see that the curves of the power depend on of the values  $k$ . They skew to the right (S-curve positive) as  $k$  increases.



**Figure 4.** The Size of the Chi-square Distribution at several  $k$

Similarly, we also see from figure 4 that the sizes are constant and depended on the  $k$ , and they decrease as the  $k$  increases. To illustrate the values of the size of the Chi-square distribution, we present a simulation  $k=5$  and  $k=10$ , on  $r=1$ , as below

For  $k=5$ ,

$$\begin{aligned}\alpha = \pi(1) &= 1 + \frac{\left[ \tau\left(\frac{r}{2}, \frac{s}{2}\right) \right]_0^5}{\tau\left(\frac{r}{2}\right)} \\ &= 1 + \frac{\left[ \tau\left(\frac{r}{2}, \frac{5}{2}\right) - \tau\left(\frac{r}{2}, 0\right) \right]}{\tau\left(\frac{r}{2}\right)} \\ &\approx 0.025.\end{aligned}$$

For  $k=10$ ,

$$\begin{aligned}\alpha = \pi(1) &= 1 + \frac{\left[ \tau\left(\frac{r}{2}, \frac{s}{2}\right) \right]_0^{10}}{\tau\left(\frac{r}{2}\right)} \\ &= 1 + \frac{\left[ \tau\left(\frac{r}{2}, 5\right) - \tau\left(\frac{r}{2}, 0\right) \right]}{\tau\left(\frac{r}{2}\right)} \\ &\approx 0.002.\end{aligned}$$

## 4. Conclusions

To find the power of the Poisson distribution, we

consider sufficient statistics and *UMP test* for getting the rejection area. In the Binomial distribution context, the curve of the power depends on the number of trials  $n$  and the bound of the rejection area. The curves tend to be zero when  $\theta > 0.4$ , and it decreases (started from  $\theta = 0.2$ ) as the parameter increases. We also note that, the curve is sigmoid (*S* curve). In the Poisson distribution context, the result showed that the power of the Poisson (not sigmoid, *S* curve) tends to be 1 on several simulation  $n(n \geq 2)$  and  $\lambda(\lambda < 10)$ . In the context of chi-square distribution, we note that the curves of the power depend on the  $k$  and the skewness of the *S*-curve is positive as  $k$  increases. On the size context, we note that the size is constant. The size also depends on the  $k$  and decreases as  $k$  increases.

## Acknowledgements

I am thankful to the LPPM UNSOED for providing me with a research grant.

## REFERENCES

- [1] K. Md. E. Saleh. Theory of preliminary test and Stein-type estimation with applications. John Wiley and Sons, Inc., New Jersey, 2006.
- [2] Pratikno. Test of Hypothesis for Linear Models with Non-Sample Prior Information. Unpublished PhD Thesis, University of Southern Queensland, Australia, 2012.
- [3] Pratikno. The noncentral  $t$  distribution and its application on the power of the tests. *Far East Journal of Mathematical Science (FJMS)*, 106 (2), 463-474, 2018.
- [4] B. Pratikno, Power of hypothesis testing parameters shape of the distributions. *Far East Journal of Mathematical Science (FJMS)*, 110 (1), 15-22, 2019
- [5] D. Wackerly, W.Mendenhall III, and R. L.Scheaffer. Mathematical statistics with application, 7th Ed. Thomson Learning, Inc., Belmont, CA, USA, 2008.
- [6] R. M. Yunus. Increasing power of M-test through pre-testing. Unpublished PhD Thesis, University of Southern Queensland, Australia, 2010.
- [7] R. M. Yunus and S. Khan. Test for intercept after pre-testing on slope a robust method. In: 9th Islamic Countries Conference on Statistical Sciences (ICCS-IX): Statistics in the Contemporary World - Theories, Methods and Applications, 2007.
- [8] R. M. Yunus and S. Khan. Increasing power of the test through pre-test a robust method. *Communications in Statistics-Theory and Methods*, 40, 581-597, 2011a.
- [9] R. M. Yunus and S. Khan. M-tests for multivariate regression model. *Journal of Nonparametric Statistics*, 23, 201-218, 2011b.
- [10] R. M. Yunus and S. Khan. The bivariate noncentral chi-square distribution - Acompound distribution approach. *Applied Mathematics and Computation*, 217, 6237-6247, 2011c.
- [11] S. Khan, B. Pratikno, A.I.N. Ibrahim and R.M Yunus, The correlated bivariate noncentral F distribution and Its application. *Communications in Statistics—Simulation and Computation*, 45 3491–3507, 2016.
- [12] S. Khan. Estimation of the Parameters of two Parallel Regression Lines Under Uncertain Prior Information. *Biometrical Journal*, 44, 73-90, 2003.
- [13] S. Khan. Estimation of parameters of the multivariate regression model with uncertain prior information and Student-t errors. *Journal of Statistical Research*, 39(2) 2005, 79-94.
- [14] S. Khan. Shrinkage estimators of intercept parameters of two simple regression models with suspected equal slopes. *Communications in Statistics - Theory and Methods*, 37, 247-260, 2008.
- [15] S. Khan. and A. K. Md. E. Saleh. Preliminary test estimators of the mean based on p-samples from multivariate Student-t populations. *Bulletin of the International Statistical Institute. 50th Session of ISI, Beijing*, 599-600, 1995.
- [16] S. Khan. and A. K. Md. E. Saleh. Shrinkage pre-test estimator of the intercept parameter for a regression model with multivariate Student-t errors. *Biometrical Journal*, 39, 1-17, 1997.
- [17] S. Khan. and A. K. Md. E. Saleh. On the comparison of the pre-test and shrinkage estimators for the univariate normal mean. *Statistical Papers*, 42(4), 451-473, 2001.
- [18] S. Khan., Z. Hoque and A. K. Md. E. Saleh. Improved estimation of the slope parameter for linear regression model with normal errors and uncertain prior information. *Journal of Statistical Research*, 31 (1), 51-72, 2002.
- [19] S. Khan. and Z. Hoque. Preliminary test estimators for the multivariate normal mean based on the modified W, LR and LM tests. *Journal of Statistical Research*, Vol 37, 43-55, 2003.
- [20] S. Khan. and A. K. Md. E. Saleh. Estimation of intercept parameter for linear regression with uncertain non-sample prior information. *Statistical Papers*. 46, 379-394, 2005.
- [21] S. Khan. and A. K. Md. E. Saleh. Estimation of slope for linear regression model with uncertain prior information and Student-t error. *Communications in Statistics - Theory and Methods*, 37(16), 2564-258, 2008.
- [22] S. Khan and B. Pratikno, Testing Base Load with Non-Sample Prior Information on Process Load. *Statistical Papers*, 54 (3), 605-617, 2013.