

PEER REVIEW

A2

LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW
KARYA ILMIAH : JURNAL ILMIAH

Judul Jurnal Ilmiah (Artikel) : Impact of Distance from the Forest Edge on The Wild Bee Diversity on The Northern Slope of Mount Slamet.

Penulis Jurnal Ilmiah *) : 1 Imam Widhiono (*nama pengusul dicetak tebal)
 2 **Eming Sudiana**

Jumlah Penulis : 2

Status Penulis : Penulis Ke-2

Identitas Jurnal Ilmiah :

- a. Nama Jurnal : Journal of Biology & Biology Education "Biosaintifika"
- b. Nomor ISSN : p-ISSN 2085-191X e-ISSN 2338-7610
- c. Edisi/Volume, Nomor : September 2016/Volume 8 Nomor 2
- d. Penerbit : Fakultas MIPA Universitas Negeri Semarang.
- e. DOI artikel : <http://dx.doi.org/10.15294/biosaintifika.v8i2.5058>
- f. Alamat Web : <https://journal.unnes.ac.id/nju/index.php/biosaintifika/article/view/5058>
- g. Terindeks di : DOAJ, SINTA 2

Kategori Publikasi Jurnal Ilmiah : ☐ Jurnal Ilmiah Internasional /Internasional Bereputasi
 (beri v pada kategori yang tepat) ☒ Jurnal Ilmiah Nasional Terakreditasi
☐ Jurnal Nasional/Nasional terindeks di..... *

Hasil Penilaian Peer Review :

| Komponen Yang Dinilai | Nilai Maksimal Jurnal Ilmiah | | | Nilai Akhir Yang Diperoleh |
|---|--|-----------------------------|--------------------------|----------------------------|
| | Internasional/Internasional bereputasi | Nasional Terakreditasi | Nasional Terindeks *) | |
| | <input type="checkbox"/> | <input type="checkbox"/> 25 | <input type="checkbox"/> | |
| a Kelengkapan unsur isi artikel (10%) | | 10 % X 25= | 2.5 | 2.5 |
| b Ruang lingkup dan kedalaman pembahasan (30%) | | 30 % X25= | 7.5 | 7.4 |
| c Kecukupan dan kemutahiran data/informasi dan metodologi (30%) | | 30 % X25= | 7.5 | 7.5 |
| d Kelengkapan unsur dan kualitas terbitan/jurnal (30%) | | 30 % X25= | 7.5 | 7.4 |
| Total = (100%) | | 25 | | 24.7 |
| Nilai Pengusul (40 % x Total) | | 10 | | 9.86 |
| Catatan Penilaian artikel oleh Reviewer: | | | | |
| 1. Tentang kelengkapan dan kesesuaian unsur | : Lengkap dan Sesuai | | | |
| 2. Tentang ruang lingkup dan kedalaman pembahasan | : Ruang lingkup dan kedalaman baik | | | |
| 3. Kecukupan dan kemutahiran data serta metodologi | : Cukup, Mutahir | | | |
| 4. Kelengkapan unsur kualitas penerbit | : Lengkap | | | |
| 5. Indikasi plagiasi | : Tidak ada | | | |
| 6. Kesesuaian bidang ilmu | : Sesuai | | | |

Purwokerto,

*) Wajib diisi

Reviewer 1



Dr. Dwi Nugroho Wibowo, M.S.
 NIP. 196111251986011001
 Jabatan/Gol. : Lektor Kepala/(Gol. IV/c)
 Bidang Ilmu : Ekologi
 Unit Kerja : Fakultas Biologi Unsoed

Reviewer 2



Drs. Edy Yani, M.S.
 NIP. 195811301984031001
 Jabatan/Gol. : Lektor Kepala/(Gol. IV/c)
 Bidang Ilmu : Ekologi Tumbuhan
 Unit Kerja : Fakultas Biologi Unsoed

A2

LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW
KARYA ILMIAH : JURNAL ILMIAH

Judul Jurnal Ilmiah (Artikel) : Impact of Distance from the Forest Edge on The Wild Bee Diversity on The Northern Slope of Mount Slamet.

Penulis Jurnal Ilmiah *) : 1 Imam Widhiono (*nama pengusul dicetak tebal)
 2 Eming Sudiana

Jumlah Penulis : 2

Status Penulis : Penulis Ke-2

Identitas Jurnal Ilmiah :

- a. Nama Jurnal : Journal of Biology & Biology Education "Biosaintifika"
- b. Nomor ISSN : p-ISSN 2085-191X e-ISSN 2338-7610
- c. Edisi/Volume, Nomor : September 2016/Volume 8 Nomor 2
- d. Penerbit : Fakultas MIPA Universitas Negeri Semarang.
- e. DOI artikel : <http://dx.doi.org/10.15294/biosaintifika.v8i2.5058>
- f. Alamat Web : <https://journal.unnes.ac.id/nju/index.php/biosaintifika/article/view/5058>
- g. Terindeks di : DOAJ, SINTA 2

Kategori Publikasi Jurnal Ilmiah : ☐ Jurnal Ilmiah Internasional /Internasional Bereputasi
 (beri v pada kategori yang tepat) ☐ Jurnal Ilmiah Nasional Terakreditasi
☐ Jurnal Nasional/Nasional terindeks di..... *

Hasil Penilaian Peer Review :

| Komponen Yang Dinilai | Nilai Maksimal Jurnal Ilmiah | | | Nilai Akhir Yang Diperoleh |
|--|---|-------------------------------------|--------------------------|----------------------------|
| | Internasional/ Internasional bereputasi | Nasional Terakreditasi | Nasional *) | |
| | 25 | <input checked="" type="checkbox"/> | <input type="checkbox"/> | |
| a Kelengkapan unsur isi artikel (10%) | 10 % X 25= 2.5 | | | 2.5 |
| b Ruang lingkup dan kedalaman pembahasan (30%) | 30 % X25= 7.5 | | | 7.2 |
| c Kecukupan dan kemutakhiran data/informasi dan metodologi (30%) | 30 % X25= 7.5 | | | 7.5 |
| d Kelengkapan unsur dan kualitas terbitan/jurnal (30%) | 30 % X25= 7.5 | | | 7.2 |
| Total = (100%) | 25 | | | 26.4 |
| Nilai Pengusul (40 % x Total) | 10 | | | 9.76 |

Catatan Penilaian artikel oleh Reviewer:

1. Tentang kelengkapan dan kesesuaian unsur : LENGKAP & SESUAI
2. Tentang ruang lingkup dan kedalaman pembahasan : RUANG LINGKUP BERSIFAT & BAKI KEDALAMAN
3. Kecukupan dan kemutakhiran data serta metodologi : CUKUP - MUTAKHIR
4. Kelengkapan unsur kualitas penerbit : LENGKAP
5. Indikasi plagiasi : TIDAK ADA
6. Kesesuaian bidang ilmu : SESUAI

Purwokerto, 10 Oktober 2020*) wajib diisi

Reviewer 1



Dr. Dwi Nugroho Wibowo, M.S.
 NIP. 196111251986011001
 Jabatan/Gol. : Lektor Kepala/(Gol. IV/c)
 Bidang Ilmu : Ekologi
 Unit Kerja : Fakultas Biologi Unsoed



Mengesahkan :
 Dekan

Dr. Imam Widhiono M.Z., M.S.
 NIP. 198904201985031002
 Unit Kerja : Fakultas Biologi Unsoed

A2

LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW
KARYA ILMIAH : JURNAL ILMIAH

Judul Jurnal Ilmiah (Artikel) : Impact of Distance from the Forest Edge on The Wild Bee Diversity on The Northern Slope of Mount Slamet.

Penulis Jurnal Ilmiah *) : 1 Imam Widhiono (*nama pengusul dicetak tebal)
 2 **Eming Sudiana**

Jumlah Penulis : 2

Status Penulis : Penulis Ke-2

Identitas Jurnal Ilmiah :

- a. Nama Jurnal : Journal of Biology & Biology Education "Biosaintifika"
- b. Nomor ISSN : p-ISSN 2085-191X e-ISSN 2338-7610
- c. Edisi/Volume, Nomor : September 2016/Volume 8 Nomor 2
- d. Penerbit : Fakultas MIPA Universitas Negeri Semarang.
- e. DOI artikel : <http://dx.doi.org/10.15294/biosaintifika.v8i2.5058>
- f. Alamat Web : <https://journal.unnes.ac.id/nju/index.php/biosaintifika/article/view/5058>
- g. Terindeks di : DOAJ, SINTA 2

Kategori Publikasi Jurnal Ilmiah : ☐ Jurnal Ilmiah Internasional /Internasional Bereputasi
 (beri v pada kategori yang tepat) ☐ Jurnal Ilmiah Nasional Terakreditasi
☐ Jurnal Nasional/Nasional terindeks di..... *

Hasil Penilaian Peer Review :


| Komponen Yang Dinilai | Nilai Maksimal Jurnal Ilmiah | | | Nilai Akhir Yang Diperoleh |
|---|---|---------------------------|-------------|----------------------------|
| | Internasional/ Internasional bereputasi | Nasional Terakreditasi | Nasional *) | |
| | 25 | | | |
| a Kelengkapan unsur isi artikel (10%) | 10 % X 25= 2.5 | | | 4,5 |
| b Ruang lingkup dan kedalaman pembahasan (30%) | 30 % X25= 7.5 | | | 7,5 |
| c Kecukupan dan kemutahiran data/informasi dan metodologi (30%) | 30 % X25= 7.5 | | | 7,4 |
| d Kelengkapan unsur dan kualitas terbitan/jurnal (30%) | 30 % X25= 7.5 | | | 7,8 |
| Total = (100%) | 25 | | | 24,9 |
| Nilai Pengusul (40 % x Total) | 10 | | | 9,96 |

Catatan Penilaian artikel oleh Reviewer:

- Tentang kelengkapan dan kesesuaian unsur : lengkap & sesuai
- Tentang ruang lingkup dan kedalaman pembahasan : sangat baik
- Kecukupan dan kemutahiran data serta metodologi : mutakhir
- Kelengkapan unsur kualitas penerbit : lengkap
- Indikasi plagiasi : tidak ada
- Kesesuaian bidang ilmu : sesuai

Purwokerto, 5-10-2020 *) wajib diisi

Reviewer 2


 Drs. Edy Yani, M.S.
 NIP. 195811301984031001
 Jabatan/Gol. : Lektor Kepala/(Gol. IV/c)
 Bidang Ilmu : Ekologi Tumbuhan
 Unit Kerja : Fakultas Biologi Unsoed



Mengetahui :
 Dekan

Prof. Dr. rer. nat. Imam Widhiono M.Z., M.S.
 NIP. 195902201985031002
 Unit Kerja : Fakultas Biologi Unsoed

ARTIKEL

DB₂
p-ISSN 2085-191X (print)
e-ISSN 2338-7610 (online)

Biosaintifika

Journal of Biology & Biology Education



Sumber foto: Widhiono & Sudiana (2016)

Nationally Accredited by Ministry of Research, Technology and Higher Education 2015

| | | | | | |
|---------------|----------|----------|---------------|----------------------------|---|
| Biosaintifika | Volume 8 | Number 2 | Pages 135-247 | Semarang September 2016 | p-ISSN 2085-191X (print) e-ISSN 2338-7610 (online) |
|---------------|----------|----------|---------------|----------------------------|---|


[Home](#) > [Archives](#) > **Vol 8, No 2 (2016)**

Vol 8, No 2 (2016)

September 2016

DOI: <https://doi.org/10.15294/biosaintifika.v8i2>

Available online since 18th September 2016

Table of Contents

Articles

Growth Pattern and Copper Accumulation in Callus of *Datura metel*

Yulita Nurchayati, Santosa Santosa, Laurentius Hartanto Nugroho, Ari Indrianto

[DOI](#) [10.15294/biosaintifika.v8i2.5177](https://doi.org/10.15294/biosaintifika.v8i2.5177) Views of Abstract: 758 | PDF: 415

PDF
135-140

The Effects of Dose *Rhizoctonia Binucleat* (BNR) and Phosphorus to Nitrate Reductase Activity (NRA) and Chlorophyll of *Vanilla* Seedling (*Vanilla planifolia* Andrews)

Haryuni Haryuni, Tyas Soemarah Kurnia Dewi

[DOI](#) [10.15294/biosaintifika.v8i2.6328](https://doi.org/10.15294/biosaintifika.v8i2.6328) Views of Abstract: 669 | PDF: 441

PDF
141-147

Impact of Distance from the Forest Edge on The Wild Bee Diversity on the Northern Slope of Mount Slamet

Imam Widhiono, Eming Sudiana

[DOI](#) [10.15294/biosaintifika.v8i2.5058](https://doi.org/10.15294/biosaintifika.v8i2.5058) Views of Abstract: 539 | PDF: 356

PDF
148-154

Protein Profile and Hematological Parameters of Mice post Injected with Irradiated *Plasmodium berghei*

Imam Rosadi, Mukh Syaifudin, Dewi Elfidasari

[DOI](#) [10.15294/biosaintifika.v8i2.6341](https://doi.org/10.15294/biosaintifika.v8i2.6341) Views of Abstract: 577 | PDF: 320

PDF
155-164

The Effect of Organic Nutrient and Growth Regulators on Seed Germination, Embryo and Shoots Development of *Dendrobium antennatum* Lindl. Orchid by In Vitro

Edy Setiti Wida Utami, Sucipto Haryanto

[DOI](#) [10.15294/biosaintifika.v8i2.5165](https://doi.org/10.15294/biosaintifika.v8i2.5165) Views of Abstract: 792 | PDF: 532

PDF
165-171

The Effectiveness of Experimental Diet with Varying Levels of Papain on The Growth Performance, Survival Rate and Feed Utilization of *Keureling* Fish (*Tor tambra*)

Zainal Abidin Muchlisin, Fardin Afrido, Tanzil Murda, Nur Fadli, Abdullah A. Muhammadar, Zulkarnain Jalil, Cut Yulvizar

[DOI](#) [10.15294/biosaintifika.v8i2.5777](https://doi.org/10.15294/biosaintifika.v8i2.5777) Views of Abstract: 1004 | PDF: 559

PDF
172-177

The Correlation Between Dietary Habits and Dental Hygiene Practice with Dental Caries Among School Children at Urban Area in Semarang

Omaran Ibrahim Mohammed Ali, Oedijani Oedijani, Fatimah Muis

[DOI](#) [10.15294/biosaintifika.v8i2.6489](https://doi.org/10.15294/biosaintifika.v8i2.6489) Views of Abstract: 500 | PDF: 336

PDF
178-184

Characterization of Three Species of Thrips on Weeping Fig, Nutmeg, and Marine Seruni Plants Based on Mtcoi DNA Sequences

Nia Kurniawaty, Purnama Hidayat, Aunu Rauf

[DOI](#) [10.15294/biosaintifika.v8i2.5448](https://doi.org/10.15294/biosaintifika.v8i2.5448) Views of Abstract: 445 | PDF: 326

PDF
185-192

The Effectiveness of Local Plants from Lom and Sawang Ethnic as Antimalarial Medicine

Henny Helmi, Budi Afriyansyah, Wivied Ekasari

[DOI](#) [10.15294/biosaintifika.v8i2.5437](https://doi.org/10.15294/biosaintifika.v8i2.5437) Views of Abstract: 491 | PDF: 401

PDF
193-200

A Preliminary Study of Bryophytes in Enggano Island, Bengkulu, Indonesia

Ainun Nadhifah, Muhammad Imam Surya

[DOI](#) [10.15294/biosaintifika.v8i2.5239](https://doi.org/10.15294/biosaintifika.v8i2.5239) Views of Abstract: 659 | PDF: 462

PDF
201-205

The Potential Fruit Crop of Cibodas Botanical Garden

Suluh Normasiwi, Muhammad Imam Surya

[DOI](#) [10.15294/biosaintifika.v8i2.5235](https://doi.org/10.15294/biosaintifika.v8i2.5235) Views of Abstract: 553 | PDF: 475

PDF
206-213

Flavonoid Production in Callus Cultures from Mesocarp of *Stelechocarpus burahol*

PDF



ABOUT THE JOURNAL

[Aims and Scope](#)

[Publication Ethics](#)

[Indexing & Abstracting](#)

[Editorial Team](#)

[Reviewer Team](#)

[Contact](#)

COLLABORATE WITH



FOR AUTHORS

[Guidelines for Author](#)

[Peer Review Process](#)

[Author Fees](#)

[Online Submission](#)

[Google Citation Analysis](#)

[Scopus Citation Analysis](#)



Readers

| | |
|------------|--------|
| ID 141,701 | TR 443 |
| US 15,118 | AU 388 |
| IN 1,901 | IR 343 |
| MY 1,622 | NL 307 |
| PH 908 | PK 284 |
| TH 768 | MX 248 |
| CN 753 | DE 238 |
| BR 630 | VN 237 |
| JP 521 | HK 236 |
| GB 475 | ZA 234 |
| SG 475 | CA 234 |

Pageviews: 513,339

Flags Collected: 166



[08455725](#) View My Stats

USER

Username

Password

☐ Remember me

Ethnotaxonomical Study of Mole Crab (Crustacea: Hippoidea) on Coastal Community of Cilacap*Dian Bhagawati, Sutrisno Anggoro, Mohammad Zainuri, Lachmudin Sya'rani* 10.15294/biosaintifika.v8i2.6491  Views of Abstract: 711 | PDF: 511PDF
222-231**Harmful Algal in Banyuasin Coastal Waters, South Sumatera***Riris Aryawati, Dietrich Geoffrey Bengen, Tri Prartono, Hilda Zulkifli* 10.15294/biosaintifika.v8i2.6356  Views of Abstract: 824 | PDF: 566PDF
232-239**The Effects of Terminalia catappa L. Leaves Extract on the Water Quality Properties, Survival and Blood Profile of Ornamental fish (Betta sp) Cultured***Rudy Agung Nugroho, Hetty Manurung, Dewi Saraswati, Deasy Ladyescha, Firman Muhammad Nur* 10.15294/biosaintifika.v8i2.6519  Views of Abstract: 1682 | PDF: 736PDF
240-247This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Search Scope

All

Search

Browse

- » By Issue
- » By Author
- » By Title
- » Other Journals



Impact of Distance from the Forest Edge on The Wild Bee Diversity on the Northern Slope of Mount Slamet

✉ Imam Widhiono, Eming Sudiana

DOI: 10.15294/biosaintifika.v8i2.5058

Faculty of Biology, Jenderal Soedirman University, Indonesia

History Article

Received 13 February 2016
Approved 16 June 2016
Published 18 September 2016

Keywords:

abundance; diversity; distance;
forest edges; species; wild bees

Abstract

In agricultural landscape in northern slope of Mount Slamet, diversity of wild bee species as pollinator depend on forested habitats. This study aimed to assess the effects of distance from the forest edge on the diversity of wild bees on strawberry and tomato crops. This study was conducted from July 2014 to October 2014. The experimental fields contained tomato and strawberry with a total area of 4 ha (2 ha each) and divided into five plots based on distance from the forest edge (0, 50, 100, 150, and 200 m). Wild bee was caught with kite netting in 7.00 -9.00 in ten consecutive days. Wild bee diversity differed according to distance from the forest edge, the highest value was at 0 m for strawberry plots ($H' = 2.008$, $E = 0.72$ and $Chao1 = 16$) and for tomato plots, the highest diversity was at 50 m from the forest edge ($H' = 2.298$, $E = 0.95$ and $Chao1 = 11$) and the lowest was at 200 m in both plots. Wild bee species richness and abundance decreased with distance, resulting in the minimum diversity and abundance of wild bee at 200 m from forest edge in both crops.

How to Cite

Widhiono, I., & Sudiana, E. (2016). Impact of Distance from the Forest Edge on The Wild Bee Diversity on the Northern Slope of Mount Slamet. *Biosaintifika: Journal of Biology & Biology Education*, 8(2), 148-154.

© 2016 Semarang State University

✉ Correspondence Author:

Jl. Dr. Soeparno No 63 Purwokerto 53122
E-mail: imamwidhiono@yahoo.com

p-ISSN 2085-191X
e-ISSN 2338-7610

INTRODUCTION

Since 1998, the forest areas on the northern slopes of Mount Slamet have been developed as agricultural areas (Perhutani, 2000), which have resulted in landscape fragmentation by reducing the amount of natural and semi-natural habitats. Natural and semi-natural habitats provide essential resources for wild bees, which act as crop pollinators in agricultural areas. The major agricultural commodities in this region are strawberries and tomatoes; pollination of both crops is dependent on the presence of wild bees in forested habitats. Wild bees (Hymenoptera: Apoidea) are considered the most important group of insect pollinators in agricultural landscapes. Habitat loss and fragmentation leads to changes in wild bee abundance and species richness, which may affect crop pollination.

Habitat loss and isolation due to agricultural intensification represent major threats to insect pollinator diversity because ecologically valuable wild bee habitats remaining in agricultural landscapes tend to be confined to a relatively small proportion of semi-natural habitats. Forest edges, as the remaining semi-natural habitat in these areas, provide essential resources for pollinators within agricultural landscapes and may help to maintain pollination services in agroecosystems. Recent research by Widhiono & Sudiana (2015) found that forest edges have abundant flowering wild plant species, which are food sources for wild bees. Therefore, the forest areas on the northern slopes of Mount Slamet play an important role by harboring wild bees as pollinators for adjacent farmland. Forest edge areas within agricultural landscapes often provide habitat for wild bee species, from which they forage on flowering crops in agricultural fields. Several studies have shown the importance of natural or semi-natural habitats, such as forest edges, in sustaining pollinator populations close to fruit crops (Carvalho et al., 2010).

The optimal foraging theory predicts that the mean richness level of wild bee pollinators decreases with distance from the forest edge (Cresswell et al., 2000), because a majority of wild bee pollinators are central place foragers with fixed nest sites (such as in the soil, plant stems, or trees) within the forest edges. In tropical ecosystems in Asia, wild bee abundance and species richness are affected by distance from the forest edge, and the abundance of insect pollinators often declines with distance from the forest edge (Klein et al., 2003). Wild bee abundance and richness decreases with distance from the fo-

rest edge, resulting in a decline in the mean levels of flower-visitor richness and visitation rate in croplands around forest edges. In the study area, strawberries and tomatoes are planted around forest edges, so distance from the forest edge affects the diversity and abundance of wild bee pollinators. This study aimed to assess the effect of distance from the forest edge on the diversity and abundance of wild bees on strawberry and tomato crops. The results will facilitate the development of a strategy for insect pollinator conservation in agricultural areas.

METHODS

Study site

This study was conducted from July 2014 to October 2014 on the northern slope of Mount Slamet, Central Java, in Serang village (altitude ± 1100 m asl (above sea levels), which is located at $7^{\circ}14'21''$ S and $109^{\circ}17'37.42''$ E. The experimental fields contained tomato (*Lycopersicon esculentum*) and strawberry (*Fragaria x annanasa*) crops adjacent to the forest edge, with a total area of 4 ha (2 ha each). Each agricultural field was divided into five plots based on distance from the forest edge (0, 50, 100, 150, and 200 m).

Sampling Methods

Wild bee sampling was performed in every 40 plants in each plot using the *scan method* i.e., observing wild bees visiting and pollinating flowers. Sampling was conducted in the morning (7.00-9.00 AM) every week and replicate five times periods. For identification, wild bees were collected using kite netting. Furthermore, all wild bee specimens sent to LIPI Bogor to species identification.

Data Analysis

The species richness and abundances of wild bees recorded in the five plots were analyzed. A general linear model was applied using the SPSS ver. 18.0 software. The dependent variables were wild bee species number and abundance, and the categorical variable was the distance from the forest edge (0, 50, 100, 150, and 200 m). Raw data from the field were used to reveal species richness with estimators Chao 1, species diversity (Shannon-Weiner index), Evenness (E), and relative abundance of different species in a sampling site (Magurran, 2003). Comparisons of species composition according to the distance from the forest edge were performed using single linkage cluster analysis based on Bray-Curtis similarity (McAleece et al., 1997). All diversity parameters

were analyzed using the Biodiversity Pro software (McAleece et al., 1997).

RESULTS AND DISCUSSION

We recorded 680 wild bees representing 15 species or species groups in 12 genera in strawberry plots, and 379 individuals representing 11 species in tomato plots. The most abundant and widespread bee species in the strawberry plots were *Trigona laeviceps* (36.3%), *Apis cerana* (28.6%), and *Rophaledia romandi* (12.4%) while the most abundant species in tomato plots were *A.cerana* (18.7%), *Megachille relativa* (12.95%), and *Ame-gilla cingulata* (12.1%). (Tabel 1). The differences of species dominance between two crops were due to differences of flower characters. Flower visitors of strawberries dominated by small bees, while tomatoes flower dominated by buzzing bees. One of the features of the tomatoes flowers is the sporical opening of its anthers which requires the agitation of the flowers by the presence of pollinators that vibrate their indirect flight muscles for the release of pollen grains. Teppner (2005), observed that *Bombus* and *Lasioglossum*, can be good pollinators of the flowers by vibrating their anthers easily. Harter et al., (2002) note that some families of bees from that perform buzz pollination and dominate at tomatoes fields are Andrenidae, Apidae, Colletidae, Halictidae, and Megachilidae.

Wild bee population and wild bee species richness according to the distance from the forest edge in strawberry and tomato plots, showed that the highest species richness was at 0 m and the lowest at 200 m from the forest edge for both crops (Figure 2).

The species diversity indices (H' , Evenness E) and species estimator (Chao 1) differed according to the distance from the forest edge. The highest value was at 0 m in the strawberry plots ($H' = 2.008$, $E = 0.72$ and Chao1= 16) and the lowest at 200 m ($H' = 1.708$, $E = 0.77$ and Chao1= 9). In tomato plots, the highest diversity was at 50 m from the forest edge ($H' = 2.298$, $E = 0.95$ and Chao1= 11) and the lowest was at 200 m ($H' = 2.156$, $E = 0.98$ and Chao1= 9) (Table 2). From data analysis showed that Chao1, (the simplest nonparametric species richness estimator), not significantly different with the result. This result indicated that the sample in both plantations has the full assemblage of species, including those species not detected in the set of samples (Nicholas and Chao, 2013).

The data suggest a relationship between the distance from the forest edge and the likelihood of the diversity of wild bees in strawberry and tomato fields. A Kruskal-Wallis test showed that between the distance from the forest has significant differences ($p < 0.05$) on wild bee populations and wild bee species richness in both strawberry and tomato plots. Bee species richness was sig-




Figure 1. Wild bee species found in sampling area ie. A. *Amegilla cingulata*, B. *Apis cerana*, C. *Rophaledia romandi*, D. *Megachille relativa*




Figure 2. Wild bee population (A) and wild bee species richness (B) according to the distance from the forest edge in strawberry and tomato plots.

Table 1. Wild bee abundance in strawberry and tomato plots according to the distance from the forest edge.

| Insect species | Strawberry | | | | | Tomato | | | | |
|-----------------------|----------------------------|-------|-------|-------|-------|--------|------|-------|-------|-------|
| | Diatance from forest edges | | | | | | | | | |
| | 0 m | 50 m | 100 m | 150 m | 200 m | 0 m | 50 m | 100 m | 150 m | 200 m |
| <i>A. cerana</i> | 46 | 48 | 29 | 41 | 31 | 22 | 7 | 30 | 5 | 7 |
| <i>A.dorsata</i> | 6 | 6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| <i>T. laeviceps</i> | 59 | 53 | 50 | 45 | 40 | 0 | 0 | 0 | 0 | 0 |
| <i>R.romandi</i> | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 |
| <i>R. fasciata</i> | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 |
| <i>M.relativa</i> | 10 | 7 | 8 | 4 | 6 | 11 | 11 | 11 | 10 | 6 |
| <i>A.cingulata</i> | 6 | 6 | 6 | 6 | 3 | 12 | 14 | 6 | 6 | 6 |
| <i>A.zonata</i> | 7 | 7 | 7 | 5 | 6 | 13 | 13 | 7 | 7 | 6 |
| <i>X.latipes</i> | 1 | 0 | 0 | 0 | 0 | 13 | 5 | 1 | 1 | 0 |
| <i>X. virginica</i> | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| <i>Polistes sp</i> | 3 | 3 | 0 | 0 | 0 | 5 | 3 | 3 | 3 | 3 |
| <i>Nomia sp.</i> | 4 | 0 | 4 | 4 | 0 | 4 | 4 | 4 | 4 | 4 |
| <i>L.malachurum</i> | 3 | 3 | 3 | 3 | 3 | 7 | 5 | 3 | 3 | 0 |
| <i>D.campaniforme</i> | 9 | 6 | 7 | 4 | 4 | 1 | 9 | 6 | 9 | 9 |
| <i>P.politus</i> | 3 | 0 | 3 | 1 | 0 | 0 | 0 | 0 | 0 | 0 |
| Insect abundance | 170 | 151 | 129 | 125 | 105 | 100 | 83 | 83 | 60 | 53 |
| Mean | 11.33 | 10.07 | 8.60 | 8.33 | 7.00 | 9.09 | 7.55 | 7.55 | 5.45 | 4.82 |
| STDev | 17.09 | 16.67 | 13.46 | 14.29 | 11.97 | 5.86 | 3.72 | 7.90 | 2.70 | 2.86 |

nificantly affected by distance to the main habitat. Distance from the forest edge was associated with decreased wild bee abundance and richness. Distance also greatly affected assemblage composition. Our results suggest that distance strongly determines the spatial distribution of bees in the study area. Distance from the forest edge had a significant effect; (Spearman's correlation; $r^2 =$

0.96, $p < 0.05$ in strawberry plots with equations $Y = 167.200 - 0.312 x$ and $r^2 = 0.75$, $p < 0.05$ in tomato plots with equations $Y = 99.200 - 0.234 x$), the same effects has also showed on species richness $r^2 = 0.96$, $p < 0.05$ with equations $Y = 13.800 - 0.024 x$ in strawberry plots and $r^2 = 0.52$, $p < 0.05$ with equations $Y = 11.400 - 0.008 x$ in tomatoes plots), this mean that bee abundance declined

Table 2. Species richness and diversity parameters in strawberry and tomato plots according to the distance from the forest edge

| Parameters | Strawberry | | | | | Tomato | | | | |
|------------------|----------------------------|-------|-------|-------|-------|--------|-------|-------|-------|-------|
| | Diatance from forest edges | | | | | | | | | |
| | 0 m | 50 m | 100 m | 150 m | 200 m | 0 m | 50 m | 100 m | 150 m | 200 m |
| Species richness | 15 | 11 | 11 | 11 | 9 | 11 | 11 | 11 | 11 | 9 |
| STDev species | 0.00 | 0.46 | 0.46 | 0.46 | 0.51 | 0.00 | 0.00 | 0.00 | 0.00 | 0.40 |
| Shannon H' | 2.008 | 1.812 | 1.901 | 1.757 | 1.708 | 2.207 | 2.290 | 2.041 | 2.27 | 2.156 |
| Evenness | 0.742 | 0.756 | 0.793 | 0.733 | 0.777 | 0.921 | 0.955 | 0.851 | 0.95 | 0.981 |
| Chao-1 | 16 | 11 | 11 | 11 | 9 | 11 | 11 | 11 | 11 | 9 |




Figure 3. Species composition similarity between the distance from the forest in Strawberry (A) and Tomato (B) using Bray-Curtis similarity index.

with increasing distance from forest edge. This result is in agreement with that of Ricketts et al. (2008), who reported that the native pollinator visitation rate drops to 50% of the maximum at a location 668 m from natural habitat, and is consistent with previous reports of the effect of forest on bee visits and pollination services (De Marco & Coelho 2004; Chacoff & Aizen 2006).

These results suggest that forest edges are important sources of pollinators, likely because they provide “partial habitats” (Holland & Fahrig, 2000), such as mating, foraging, nesting, and nesting materials sites, which bees need to complete their life cycle (Bailey et al., 2014). Klein et al. (2003), in their study of coffee pollination in agroforestry systems at Lore Lindu (Central Sulawesi), verified that the number of social bee species diminished with the distance between fragments. De Marco & Coelho (2004) verified that cultivation near forest fragments (distance <1km) results in 14.6% greater production when compared to distant systems. Ricketts et al., (2008) reported a greater increase, 20%, in Costa Rica. This effect of distance from semi-natural habitats suggests that variables related to landscape ecology, such as the permeability of the matrix to the dispersion of pollinators, may be essential

(Jauker et al., 2009). Forest edges could provide one or more important partial habitats for diverse bee species in agricultural landscapes, in particular when associated with flowering agricultural crops (Le Feon et al., 2011). Species composition according to the distance from the forest edge was evaluated using a single linkage cluster analysis based on Bray–Curtis similarity. The species composition at forest edges exhibited the highest similarity (92.8% in strawberry plots and 79.7% in tomato plots, respectively) at the closest distance (50 m) and lowest similarity (76.3% in strawberry plots and 58.8% in tomato plots) at the farthest distance (200 m) from forest edges in plots of both crops. (Figure 3)

These research findings may be due to differences in foraging distance among wild bee taxa. Foraging distance has been shown to increase the body size of bees (Greenleaf et al., 2007). Foraging distance, therefore, determines the spatial scale at which wild bees can provide pollination services to crops (Greenleaf & Kremen 2006). Many wild bees that pollinate crops nest in natural habitats and forage on crops within their daily travel distance (Schulke & Waser, 2001). Foraging bees are likely to fly short distances, recent investigations predicted maximum for-

ging distances of 100–200 m for small bee species and up to 1100 m for very large species based on mainly indirect methods (Zurbuchen et al., 2010) and change directions between successive visits to high-reward patches and fly longer distances in the same direction to low-reward patches (Ne'man et al., 2006). These studies suggest that forest edges are likely to be a source of pollinators of various crops. Indeed, forest edges exhibit a complex vegetation structure and undisturbed soil, offering shelter for bees and a wide range of nesting sites for both cavity and ground-nesting bees (Artz & Waddington, 2006). In addition, they provide a diversity of flowering wild plants throughout the bees' activity period (Margrath et al., 2013). Widhiono & Sudiana (2015) found that forest edges in this study area harbor diverse flowering wild plant species. Several studies have shown the importance of natural or semi-natural habitats for sustaining pollinator populations or pollination services close to fruit crops (Chacoff & Aizen, 2006; Ricketts et al., 2008; Carvalheiro et al., 2010). Other studies have reported a negative impact of distance on forests on pollination services or bee abundance and richness in tropical ecosystems (De Marco & Coelho 2004; Chacoff et al., 2008). Finally, these studies also suggest that the pollination of strawberry and tomato crops could be negatively affected by being situated too far from the forest edge.

CONCLUSION

Forest edge habitats are important for enhancing pollinator diversity in agricultural landscapes. They harbor diverse wild bee communities and can be considered classical source habitats. Dispersal from these forest edge habitats into the agricultural area, however, is strongly affected by the distance from the forest edge.

ACKNOWLEDGEMENTS

We are thankful to Yulia Arnitasari, and Farda Komarudin who have helped data collection and encouraged this work from the beginning. This research was a part of main research that founded by Jenderal Soedirman University. We are also thankful to authorities of Institute of Research and Community Services, Jenderal Soedirman University for support.

REFERENCES

Artz, D. R. & Waddington, K. D. (2006). The effects of neighbouring tree islands on pollinator density

- and diversity, and on pollination of a wet prairie species, *Asclepias lanceolata* (Apocynaceae). *Journal of Ecology*, 94(3), 597–608.
- Bailey, S., Requier, F., Nusillard, B., Roberts, S. P., Potts, S. G., & Bouget, C. (2014). Distance from forest edge affects bee pollinators in oil-seed rape fields. *Ecology and evolution*, 4(4), 370–380.
- Chacoff, N. P., & Aizen, M. A. (2006). Edge effects on flower-visiting insects in grapefruit plantations bordering premontane subtropical forest. *Journal of Applied Ecology*, 43(1), 18–27.
- Chacoff, N. P., Aizen, M. A., & Aschero, V. (2008). Proximity to forest edge does not affect crop production despite pollen limitation. *Proceedings of the Royal Society of London B: Biological Sciences*, 275(1637), 907–913.
- Carvalheiro, L. G., Seymour, C. L., Veldtman, R., & Nicolson, S. W. (2010). Pollination services decline with distance from natural habitat even in biodiversity-rich areas. *Journal of Applied Ecology*, 47(4), 810–820.
- Cresswell, J. E. (1999). The influence of nectar and pollen availability on pollen transfer by individual flowers of oil-seed rape (*Brassica napus*) when pollinated by bumblebees (*Bombus lapidarius*). *Journal of Ecology*, 87(4), 670–677.
- De Marco Jr, P., & Coelho, F. M. (2004). Services performed by the ecosystem: forest remnants influence agricultural cultures' pollination and production. *Biodiversity & Conservation*, 13(7), 1245–1255.
- Greenleaf, S. S., & Kremen, C. (2006). Wild bee species increase tomato production and respond differently to surrounding land use in Northern California. *Biological Conservation*, 133(1), 81–87.
- Greenleaf, S. S., Williams, N. M., Winfree, R., & Kremen, C. (2007). Bee foraging ranges and their relationship to body size. *Oecologia*, 153(3), 589–596.
- Harter, B., Leistikow, C., Wilms, W., Truylio, B., & Engels, W. (2002). Bees collecting pollen from flowers with poricidal anthers in a south Brazilian Araucaria forest: a community study. *Journal of Apicultural Research*, 41(1–2), 9–16.
- Holland, J., & Fahrig, L. (2000). Effect of woody borders on insect density and diversity in crop fields: a landscape-scale analysis. *Agriculture, ecosystems & environment*, 78(2), 115–122.
- Jauker, F., Diekötter, T., Schwarzbach, F., & Wolters, V. (2009). Pollinator dispersal in an agricultural matrix: opposing responses of wild bees and hoverflies to landscape structure and distance from main habitat. *Landscape Ecology*, 24(4), 547–555.
- Klein, A. M., Steffan-Dewenter, I., & Tscharnkte, T. (2003). Fruit set of highland coffee increases with the diversity of pollinating bees. *Proceedings of the Royal Society of London B: Biological Sciences*, 270(1518), 955–961.
- Le Féon, V., Burel, F., Chifflet, R., Henry, M., Ricroch,

- A., Vaissière, B. E., & Baudry, J. (2013). Solitary bee abundance and species richness in dynamic agricultural landscapes. *Agriculture, Ecosystems & Environment*, 166, 94-101.
- Magrach, A., Santamaría, L., & Larrinaga, A. R. (2013). Forest edges show contrasting effects on an austral mistletoe due to differences in pollination and seed dispersal. *Journal of Ecology*, 101(3), 713-721.
- Magurran, A. E. (2013). *Measuring biological diversity*. New Jersey: John Wiley & Sons.
- McAleece, N., Lambhead, P. J. D., Paterson, G. L. J., & Gage, J. D. (1997). *Biodiversity Pro*. The Natural History Museum, London.
- Gotelli, N. J., Chao, A., & Levin, S. (2013). Measuring and estimating species richness, species diversity, and biotic similarity from sampling data. *Encyclopedia of biodiversity*, 5, 195-211.
- Ne'eman, G., Shavit, O., Shaltiel, L., & Shmida, A. (2006). Foraging by male and female solitary bees with implications for pollination. *Journal of Insect Behavior*, 19(3), 383-401.
- Perhutani. 2000. *Rencana Pengelolaan Pelestarian Hutan, Kelas hutan Damar*. Perencanaan Hutan II, Yogyakarta.
- Ricketts, T. H., Regetz, J., Steffan-Dewenter, I., Cunningham, S. A., Kremen, C., Bogdanski, A., ... & Morandin, L. A. (2008). Landscape effects on crop pollination services: are there general patterns?. *Ecology letters*, 11(5), 499-515.
- Schulke, B., & Waser, N. M. (2001). Long-distance pollinator flights and pollen dispersal between populations of *Delphinium nuttallianum*. *Oecologia*, 127(2), 239-245.
- Teppner, H. (2005). Pollinators of tomato, *Solanum lycopersicum* (Solanaceae), in central Europe. *Phyton*, 45(2), 217-235.
- Widhiono, I. & Sudiana, E. (2015). Peran tumbuhan liar dalam konservasi keragaman serangga penyerbuk Ordo Hymenoptera. *Pros Sem Nas Masy Biodiv Indon*. Vol 1, No 7, 1586-1590.
- Zurbuchen, A., Landert, L., Klaiber, J., Müller, A., Hein, S., & Dorn, S. (2010). Maximum foraging ranges in solitary bees: only few individuals have the capability to cover long foraging distances. *Biological Conservation*, 143(3), 669-676.

TURNITIN

Impact of Distance from the Forest Edge on The Wild Bee Diversity on the Northern Slope of Mount Slamet

by Eming Sudiana

Submission date: 23-Aug-2020 11:15PM (UTC+0700)

Submission ID: 1372913008

File name: BIOSAINTEFIKA_VOL_8_NO_2_2016__ARTIKEL.pdf (377.82K)

Word count: 3738

Character count: 18793



Impact of Distance from the Forest Edge on The Wild Bee Diversity on the Northern Slope of Mount Slamet

Imam Widhiono, Eming Sudiana

DOI: 10.15294/biosaintifika.v8i2.5058

Faculty of Biology, Jenderal Soedirman University, Indonesia

History Article

Received 13 February 2016
Approved 16 June 2016
Published 18 September 2016

Keywords:

abundance; diversity; distance;
forest edges; species; wild bees

Abstract

In agricultural landscape in northern slope of Mount Slamet, diversity of wild bee species as pollinator depend on forested habitats. This study aimed to assess the effects of distance from the forest edge on the diversity of wild bees on strawberry and tomato crops. This study was conducted from July 2014 to October 2014. The experimental fields contained tomato and strawberry with a total area of 4 ha (2 ha each) and divided into five plots based on distance from the forest edge (0, 50, 100, 150, and 200 m). Wild bee was caught with kite netting in 7.00 -9.00 in ten consecutive days. Wild bee diversity differed according to distance from the forest edge, the highest value was at 0 m for strawberry plots ($H' = 2.008$, $E = 0.72$ and $Chao1 = 16$) and for tomato plots, the highest diversity was at 50 m from the forest edge ($H' = 2.298$, $E = 0.95$ and $Chao1 = 11$) and the lowest was at 200 m in both plots. Wild bee species richness and abundance decreased with distance, resulting in the minimum diversity and abundance of wild bee at 200 m from forest edge in both crops.

How to Cite

Widhiono, I., & Sudiana, E. (2016). Impact of Distance from the Forest Edge on The Wild Bee Diversity on the Northern Slope of Mount Slamet. *Biosaintifika: Journal of Biology & Biology Education*, 8(2), 148-154.

© 2016 Semarang State University

✉ Correspondence Author:
Jl. Dr. Soeparno No 63 Purwokerto 53122
E-mail: imamwidhiono@yahoo.com

p-ISSN 2085-191X
e-ISSN 2338-7610

INTRODUCTION

Since 1998, the forest areas on the northern slopes of Mount Slamet have been developed as agricultural areas (Perhutani, 2000), which have resulted in landscape fragmentation by reducing the amount of natural and semi-natural habitats. Natural and semi-natural habitats provide essential resources for wild bees, which act as crop pollinators in agricultural areas. The major agricultural commodities in this region are strawberries and tomatoes; pollination of both crops is dependent on the presence of wild bees in forested habitats. Wild bees (Hymenoptera: Apoidea) are considered the most important group of insect pollinators in agricultural landscapes. Habitat loss and fragmentation leads to changes in wild bee abundance and species richness, which may affect crop pollination.

Habitat loss and isolation due to agricultural intensification represent major threats to insect pollinator diversity because ecologically valuable wild bee habitats remaining in agricultural landscapes tend to be confined to a relatively small proportion of semi-natural habitats. Forest edges, as the remaining semi-natural habitat in these areas, provide essential resources for pollinators within agricultural landscapes and may help to maintain pollination services in agroecosystems. Recent research by Widhiono & Sudiana (2015) found that forest edges have abundant flowering wild plant species, which are food sources for wild bees. Therefore, the forest areas on the northern slopes of Mount Slamet play an important role by harboring wild bees as pollinators for adjacent farmland. Forest edge areas within agricultural landscapes often provide habitat for wild bee species, from which they forage on flowering crops in agricultural fields. Several studies have shown the importance of natural or semi-natural habitats, such as forest edges, in sustaining pollinator populations close to fruit crops (Carvalho et al., 2010).

The optimal foraging theory predicts that the mean richness level of wild bee pollinators decreases with distance from the forest edge (Cresswell et al., 2000), because a majority of wild bee pollinators are central place foragers with fixed nest sites (such as in the soil, plant stems, or trees) within the forest edges. In tropical ecosystems in Asia, wild bee abundance and species richness are affected by distance from the forest edge, and the abundance of insect pollinators often declines with distance from the forest edge (Klein et al., 2003). Wild bee abundance and richness decreases with distance from the fo-

rest edge, resulting in a decline in the mean levels of flower-visitor richness and visitation rate in croplands around forest edges. In the study area, strawberries and tomatoes are planted around forest edges, so distance from the forest edge affects the diversity and abundance of wild bee pollinators. This study aimed to assess the effect of distance from the forest edge on the diversity and abundance of wild bees on strawberry and tomato crops. The results will facilitate the development of a strategy for insect pollinator conservation in agricultural areas.

METHODS

Study site

This study was conducted from July 2014 to October 2014 on the northern slope of Mount Slamet, Central Java, in Serang village (altitude ± 1100 m asl (above sea levels), which is located at $7^{\circ}14'21''$ S and $109^{\circ}17'37.42''$ E. The experimental fields contained tomato (*Lycopersicon esculentum*) and strawberry (*Fragaria x ammanasa*) crops adjacent to the forest edge, with a total area of 4 ha (2 ha each). Each agricultural field was divided into five plots based on distance from the forest edge (0, 50, 100, 150, and 200 m).

Sampling Methods

Wild bee sampling was performed in every 40 plants in each plot using the scan method i.e., observing wild bees visiting and pollinating flowers. Sampling was conducted in the morning (7.00-9.00 AM) every week and replicate five times periods. For identification, wild bees were collected using kite netting. Furthermore, all wild bee specimens sent to LIPI Bogor to species identification.

Data Analysis

The species richness and abundances of wild bees recorded in the five plots were analyzed. A general linear model was applied using the SPSS ver. 18.0 software. The dependent variables were wild bee species number and abundance, and the categorical variable was the distance from the forest edge (0, 50, 100, 150, and 200 m). Raw data from the field were used to reveal species richness with estimators Chao 1, species diversity (Shannon-Weiner index), Evenness (E), and relative abundance of different species in a sampling site (Magurran, 2003). Comparisons of species composition according to the distance from the forest edge were performed using single linkage cluster analysis based on Bray-Curtis similarity (McAleece et al., 1997). All diversity parameters

were analyzed using the Biodiversity Pro software (McAleece et al., 1997).

RESULTS AND DISCUSSION

We recorded 680 wild bees representing 15 species or species groups in 12 genera in strawberry plots, and 379 individuals representing 11 species in tomato plots. The most abundant and widespread bee species in the strawberry plots were *Trigona laeviceps* (36.3%), *Apis cerana* (28.6%), and *Ropalidia romandi* (12.4%) while the most abundant species in tomato plots were *A.cerana* (18.7%), *Megachille relativa* (12.95%), and *Ame-gilla cingulata* (12.1%). (Tabel 1). The differences of species dominance between two crops were due to differences of flower characters. Flower visitors of strawberries dominated by small bees, while tomatoes flower dominated by buzzing bees. One of the features of the tomatoes flowers is the sporical opening of its anthers which requires the agitation of the flowers by the presence of pollinators that vibrate their indirect flight muscles for the release of pollen grains. Teppner (2005), observed that *Bombus* and *Lasioglossum*, can be good pollinators of the flowers by vibrating their anthers easily. Harter et al., (2002) note that some families of bees from that perform buzz pollination and dominate at tomatoes fields are Andrenidae, Apidae, Colletidae, Halictidae, and Megachilidae.

Wild bee population and wild bee species richness according to the distance from the forest edge in strawberry and tomato plots, showed that the highest species richness was at 0 m and the lowest at 200 m from the forest edge for both crops (Figure 2).

The species diversity indices (H' , Evenness E) and species estimator (Chao 1) differed according to the distance from the forest edge. The highest value was at 0 m in the strawberry plots ($H' = 2.008$, $E = 0.72$ and $Chao1 = 16$) and the lowest at 200 m ($H' = 1.708$, $E = 0.77$ and $Chao1 = 9$). In tomato plots, the highest diversity was at 50 m from the forest edge ($H' = 2.298$, $E = 0.95$ and $Chao1 = 11$) and the lowest was at 200 m ($H' = 2.156$, $E = 0.98$ and $Chao1 = 9$) (Table 2). From data analysis showed that $Chao1$, (the simplest nonparametric species richness estimator), not significantly different with the result. This result indicated that the sample in both plantations has the full assemblage of species, including those species not detected in the set of samples (Nicholas and Chao, 2013).

The data suggest a relationship between the distance from the forest edge and the likelihood of the diversity of wild bees in strawberry and tomato fields. A Kruskal-Wallis test showed that between the distance from the forest has significant differences ($p < 0.05$) on wild bee populations and wild bee species richness in both strawberry and tomato plots. Bee species richness was sig-



Figure 1. Wild bee species found in sampling area ie. A. *Amegilla cingulata*, B. *Apis cerana*, C. *Ropalidia romandi*, D. *Megachille relativa*




Figure 2. Wild bee population (A) and wild bee species richness (B) according to the distance from the forest edge in strawberry and tomato plots.

Table 1. Wild bee abundance in strawberry and tomato plots according to the distance from the forest edge.

| Insect species | Strawberry | | | | | Tomato | | | | |
|------------------------|----------------------------|-------|-------|-------|-------|----------------------------|------|-------|-------|-------|
| | Distance from forest edges | | | | | Distance from forest edges | | | | |
| | 0 m | 50 m | 100 m | 150 m | 200 m | 0 m | 50 m | 100 m | 150 m | 200 m |
| <i>A. cerana</i> | 46 | 48 | 29 | 41 | 31 | 22 | 7 | 30 | 5 | 7 |
| <i>A. dorsata</i> | 6 | 6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| <i>T. laeviceps</i> | 59 | 53 | 50 | 45 | 40 | 0 | 0 | 0 | 0 | 0 |
| <i>R. romandi</i> | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 |
| <i>R. fasciata</i> | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 |
| <i>M. relativa</i> | 10 | 7 | 8 | 4 | 6 | 11 | 11 | 11 | 10 | 6 |
| <i>A. cingulata</i> | 6 | 6 | 6 | 6 | 3 | 12 | 14 | 6 | 6 | 6 |
| <i>A. zonata</i> | 7 | 7 | 7 | 5 | 6 | 13 | 13 | 7 | 7 | 6 |
| <i>X. latipes</i> | 1 | 0 | 0 | 0 | 0 | 13 | 5 | 1 | 1 | 0 |
| <i>X. virginica</i> | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| <i>Polistes sp.</i> | 3 | 3 | 0 | 0 | 0 | 5 | 3 | 3 | 3 | 3 |
| <i>Nomia sp.</i> | 4 | 0 | 4 | 4 | 0 | 4 | 4 | 4 | 4 | 4 |
| <i>L. malachurum</i> | 3 | 3 | 3 | 3 | 3 | 7 | 5 | 3 | 3 | 0 |
| <i>D. campaniforme</i> | 9 | 6 | 7 | 4 | 4 | 1 | 9 | 6 | 9 | 9 |
| <i>P. politus</i> | 3 | 0 | 3 | 1 | 0 | 0 | 0 | 0 | 0 | 0 |
| Insect abundance | 170 | 151 | 129 | 125 | 105 | 100 | 83 | 83 | 60 | 53 |
| Mean | 11.33 | 10.07 | 8.60 | 8.33 | 7.00 | 9.09 | 7.55 | 7.55 | 5.45 | 4.82 |
| STDev | 17.09 | 16.67 | 13.46 | 14.29 | 11.97 | 5.86 | 3.72 | 7.90 | 2.70 | 2.86 |

nificantly affected by distance to the main habitat. Distance from the forest edge was associated with decreased wild bee abundance and richness. Distance also greatly affected assemblage composition. Our results suggest that distance strongly determines the spatial distribution of bees in the study area. Distance from the forest edge had a significant effect; (Spearman's correlation; $r^2 =$

0.96, $p < 0.05$ in strawberry plots with equations $Y = 167.200 - 0.312 x$ and $r^2 = 0.75$, $p < 0.05$ in tomato plots with equations $Y = 99.200 - 0.234 x$), the same effects has also showed on species richness $r^2 = 0.96$, $p < 0.05$ with equations $Y = 13.800 - 0.024 x$ in strawberry plots and $r^2 = 0.52$, $p < 0.05$ with equations $Y = 11.400 - 0.008 x$ in tomatoes plots), this mean that bee abundance declined

Table 2. Species richness and diversity parameters in strawberry and tomato plots according to the distance from the forest edge

| Parameters | Strawberry | | | | | Tomato | | | | |
|------------------|----------------------------|-------|-------|-------|-------|--------|-------|-------|-------|-------|
| | Distance from forest edges | | | | | | | | | |
| | 0 m | 50 m | 100 m | 150 m | 200 m | 0 m | 50 m | 100 m | 150 m | 200 m |
| Species richness | 15 | 11 | 11 | 11 | 9 | 11 | 11 | 11 | 11 | 9 |
| STDev species | 0.00 | 0.46 | 0.46 | 0.46 | 0.51 | 0.00 | 0.00 | 0.00 | 0.00 | 0.40 |
| Shannon H' | 2.008 | 1.812 | 1.901 | 1.757 | 1.708 | 2.207 | 2.290 | 2.041 | 2.27 | 2.156 |
| Evenness | 0.742 | 0.756 | 0.793 | 0.733 | 0.777 | 0.921 | 0.955 | 0.851 | 0.95 | 0.981 |
| Chao-1 | 16 | 11 | 11 | 11 | 9 | 11 | 11 | 11 | 11 | 9 |




Figure 3. Species composition similarity between the distance from the forest in Strawberry (A) and Tomato (B) using Bray-Curtis similarity index.

with increasing distance from forest edge. This result is in agreement with that of Ricketts et al. (2008), who reported that the native pollinator visitation rate drops to 50% of the maximum at a location 668 m from natural habitat and is consistent with previous reports of the effect of forest on bee visits and pollination services (De Marco & Coelho 2004; Chacoff & Aizen 2006).

These results suggest that forest edges are important sources of pollinators, likely because they provide “partial habitats” (Holland & Fahrig, 2000), such as mating, foraging, nesting, and nesting materials sites, which bees need to complete their life cycle (Bailey et al., 2014). Klein et al. (2003), in their study of coffee pollination in agroforestry systems at Lore Lindu (Central Sulawesi), verified that the number of social bee species diminished with the distance between fragments. De Marco & Coelho (2004) verified that cultivation near forest fragments (distance <1km) results in 14.6% greater production when compared to distant systems. Ricketts et al., (2008) reported a greater increase, 20%, in Costa Rica. This effect of distance from semi-natural habitats suggests that variables related to landscape ecology, such as the permeability of the matrix to the dispersion of pollinators, may be essential

(Jauker et al., 2009). Forest edges could provide one or more important partial habitats for diverse bee species in agricultural landscapes, in particular when associated with flowering agricultural crops (Le Feon et al., 2011). Species composition according to the distance from the forest edge was evaluated using a single linkage cluster analysis based on Bray-Curtis similarity. The species composition at forest edges exhibited the highest similarity (92.8% in strawberry plots and 79.7% in tomato plots, respectively) at the closest distance (50 m) and lowest similarity (76.3% in strawberry plots and 58.8% in tomato plots) at the farthest distance (200 m) from forest edges in plots of both crops. (Figure 3)

These research findings may be due to differences in foraging distance among wild bee taxa. Foraging distance has been shown to increase the body size of bees (Greenleaf et al., 2007). Foraging distance, therefore, determines the spatial scale at which wild bees can provide pollination services to crops (Greenleaf & Kremen 2006). Many wild bees that pollinate crops nest in natural habitats and forage on crops within their daily travel distance (Schulke & Waser, 2001). Foraging bees are likely to fly short distances, recent investigations predicted maximum foraging

ging distances of 100–200 m for small bee species and up to 1100 m for very large species based on 23 only indirect methods (Zurbuchen et al., 2010) and change directions between successive visits to high-reward patches and fly longer distances in the same direction to low-reward patches (Ne'man et al., 2006). These studies suggest that forest edges are likely to be a source of pollinators of various crops. Indeed, forest edges exhibit a complex vegetation structure and undisturbed 35, offering shelter for bees and a wide range of nesting sites for both cavity and ground-nesting bees (Artz & Waddington, 2006). In addition, they provide a diversity of flowering wild plants throughout the bees' activity period (Margrath et al., 2013). Widhiono & Sudiana (2015) found that forest edges in this study area harbor diverse flowering wild plant species. Several studies have shown the importance of natural or semi-natural habitats for sustaining pollinator populations or pollination serv 34 close to fruit crops (Chacoff & Aizen, 2006; Ricketts et al., 2008; Carvalheiro et al., 2010). Other studies have reported a negative impact of distance on forests on pollination services or bee abundance and richness in tropical ecosystems (De Marco & Coelho 2004; Chacoff et al., 2008). Finally, these studies also suggest that the pollination of strawberry and tomato crops could be negatively affected by being situated too far from the forest edge.

CONCLUSION

10 Forest edge habitats are important for enhancing pollinator diversity in agricultural landscapes. They harbor diverse wild bee communities and can be considered classical source habitats. Dispersal from these forest edge habitats into the agricultural area, however, is strongly affected by the distance from the forest edge.

ACKNOWLEDGEMENTS

We are thankful to Yulia Arnitasari, and Farda Komarudin who have helped data collection and encouraged this work from the beginning. This research was a part of main research that founded by Jenderal Soedirman University. We are also 1 thankful to authorities of Institute of Research and Community Services, Jenderal Soedirman University for support.

REFERENCES

- Artz, D. R. & Waddington, K. D. (2006). The effects of neighbouring tree islands on pollinator density

and diversity, and on pollination of a wet prairie species, *Asclepias lanceolata* (Apocynaceae). *Journal of Ecology*, 94(3), 597–608

- 9 Bailey, S., Requier, F., Nusillard, B., Roberts, S. P., Potts, S. G., & Bouget, C. (2014). Distance from forest edge affects bee pollinators in oil-seed rape fields. *Ecology and evolution*, 4(4), 370–380.

- 17 Chacoff, N. P., & Aizen, M. A. (2006). 5 Edge effects on flower-visiting insects in grapefruit plantations bordering premontane subtropical forest. *Journal of Applied Ecology*, 43(1), 18–27. Chacoff, N. P., Aizen, M. A., & Aschero, V. (2008). Proximity to forest edge does not affect crop production despite pollen limitation. *Proceedings of the Royal Society of London B: Biological Sciences*, 275(1637), 907–913.

- 14 12 Carvalheiro, L. G., Seymour, C. L., Veldtman, R., & Nicolson, S. W. (2010). Pollination services decline with distance from natural habitat even in biodiversity-rich areas. *Journal of Applied Ecology*, 47(4), 810–820.

- Cresswell, J. E. (1999). The influence of nectar and pollen availability on pollen transfer by individual flowers of oil-seed rape (*Brassica napus*) when pollinated by bumblebees (*Bombus lapidarius*). *Journal of Ecology*, 87(4), 670–677.

- De Marco Jr, P., & Coelho, F. M. (2004). Services performed by the ecosystem: forest remnants influence agricultural cultures' pollination and production. *Biodiversity & Conservation*, 13(7), 1245–1255.

- 40 Greenleaf, S. S., & Kremen, C. (2006). Wild bee species increase tomato production and respond differently to surrounding land use in Northern California. *Biological Conservation*, 133(1), 81–87.

- Greenleaf, S. S., Williams, N. M., Winfree, R., & Kremen, C. (2007). Bee foraging ranges and their relationship to body size. *Oecologia*, 153(3), 589–596.

- 6 Harter, B., Leistikow, C., Wilms, W., Truyllo, B., & Engels, W. (2002). Bees collecting pollen from flowers with poricidal anthers in a south Brazilian Araucaria forest: a community study. *Journal of Apicultural Research*, 41(1–2), 9–16.

- 15 Holland, J., & Fahrig, L. (2000). Effect of woody borders on insect density and diversity in crop fields: a landscape-scale analysis. *Agriculture, ecosystems & environment*, 78(2), 115–122.

- 4 Jauker, F., Diekötter, T., Schwarzbach, F., & Wolters, V. (2009). Pollinator dispersal in an agricultural matrix: opposing responses of wild bees and hoverflies to landscape structure and distance from main habitat. *Landscape Ecology*, 24(4), 547–555.

- Klein, A. M., Steffan-Dewer, I., & Tscharntke, T. (2003). Fruit set of highland coffee increases with the diversity of pollinating bees. *Proceedings of the Royal Society of London B: Biological Sciences*, 270(1518), 955–961.

- 8 Le Féon, V., Burel, F., Chifflet, R., Henry, M., Ricroch,

- A., Vaissière, B. E., & Baudry, J. (2013). Solitary bee abundance and species richness in dynamic agricultural landscapes. *Agriculture, Ecosystems & Environment*, 166, 94-101.
- 11 Magrath, A., Santamaría, L., & Larrinaga, A. R. (2013). Forest edges show contrasting effects on an austral mistletoe due to differences in pollination and seed dispersal. *Journal of Ecology*, 101(3), 713-721.
- 29 Magurran, A. E. (2013). *Measuring biological diversity*. New Jersey: John Wiley & Sons.
- 24 McAleece, N., Lamshead, P. J. D., Paterson, G. L. J., & Gage, J. D. (1997). *Biodiversity Pro*. The Natural History Museum, London.
- 16 Gotelli, N. J., Chao, A., & Levin, S. (2013). Measuring and estimating species richness, species diversity, and biotic similarity from sampling data. *Encyclopedia of biodiversity*, 5, 195-211.
- 13 Ne'eman, G., Shavit, O., Shaltiel, L., & Shmida, A. (2006). Foraging by male and female solitary bees with implications for pollination. *Journal of Insect Behavior*, 19(3), 383-401.
- Perhutani. 2000. *Rencana Pengelolaan Pelestarian Hutan, Kelas hutan Damar*. Perencanaan Hutan II, Yogyakarta.
- 7 Ricketts, T. H., Regetz, J., Steffan-Dewenter, I., Cunningham, S. A., Kremen, C., Bogdanski, A., ... & Morandin, L. A. (2008). Landscape effects on crop pollination services: are there general patterns?. *Ecology letters*, 11(5), 499-515.
- 22 Schulke, B., & Waser, N. M. (2001). Long-distance pollinator flights and pollen dispersal between populations of *Delphinium nuttallianum*. *Oecologia*, 127(2), 239-245.
- Teppner, H. (2005). Pollinators of tomato, *Solanum lycopersicum* (Solanaceae), in central Europe. *Phyton*, 45(2), 217-235.
- 19 Widhiono, I. & Sudiana, E. (2015). Peran tumbuhan liar dalam konservasi keragaman serangga penyerbuk Ordo Hymenoptera. *Pros Sem Nas Masy Biodiv Indon*. Vol 1, No 7, 1586-1590.
- Zurbuchen, A., Landert, L., Klaiber, J., Müller, A., Hein, S., & Dorn, S. (2010). Maximum foraging ranges in solitary bees: only few individuals have the capability to cover long foraging distances. *Biological Conservation*, 143(3), 669-676.

Impact of Distance from the Forest Edge on The Wild Bee Diversity on the Northern Slope of Mount Slamet

ORIGINALITY REPORT

25%

SIMILARITY INDEX

19%

INTERNET SOURCES

22%

PUBLICATIONS

15%

STUDENT PAPERS

PRIMARY SOURCES

1

downloads.hindawi.com

Internet Source

2%

2

Frank Jauker. "Pollinator dispersal in an agricultural matrix: opposing responses of wild bees and hoverflies to landscape structure and distance from main habitat", Landscape Ecology, 04/2009

Publication

1%

3

Submitted to UH, Manoa

Student Paper

1%

4

Submitted to Royal Holloway and Bedford New College

Student Paper

1%

5

Ashworth, L.. "Pollinator-dependent food production in Mexico", Biological Conservation, 200905

Publication

1%

6

www.scielo.br

Internet Source

1%

| | | |
|----|---|-----|
| 7 | paduaresearch.cab.unipd.it Internet Source | 1 % |
| 8 | Annika Louise Hass, Lara Brachmann, Péter Batáry, Yann Clough, Hermann Behling, Teja Tscharntke. "Maize-dominated landscapes reduce bumblebee colony growth through pollen diversity loss", <i>Journal of Applied Ecology</i> , 2018 Publication | 1 % |
| 9 | cdn.foe.co.uk Internet Source | 1 % |
| 10 | Frank Jauker, Tim Diekötter, Franziska Schwarzbach, Volkmar Wolters. "Pollinator dispersal in an agricultural matrix: opposing responses of wild bees and hoverflies to landscape structure and distance from main habitat", <i>Landscape Ecology</i> , 2009 Publication | 1 % |
| 11 | parasiticplants.org Internet Source | 1 % |
| 12 | Ivey, C. T., P. Martinez, and R. Wyatt. "Variation in pollinator effectiveness in swamp milkweed, <i>Asclepias incarnata</i> (Apocynaceae)", <i>American Journal of Botany</i> , 2003. Publication | 1 % |
| 13 | Submitted to Drexel University Student Paper | 1 % |

| | | |
|----|---|-----|
| 14 | Submitted to Oxford Brookes University Student Paper | 1 % |
| 15 | www.biobio-indicator.org Internet Source | 1 % |
| 16 | Submitted to University of Wales, Bangor Student Paper | 1 % |
| 17 | Natacha P. Chacoff. "Proximity to forest edge does not affect crop production despite pollen limitation", Proceedings of The Royal Society B Biological Sciences, 04/22/2008 Publication | 1 % |
| 18 | Submitted to Associatie K.U.Leuven Student Paper | 1 % |
| 19 | docobook.com Internet Source | 1 % |
| 20 | Submitted to Warren Wilson College Student Paper | 1 % |
| 21 | e-citations.ethbib.ethz.ch Internet Source | 1 % |
| 22 | hss.ulb.uni-bonn.de Internet Source | 1 % |
| 23 | Gidi Ne'eman, Ofrit Shavit, Liora Shaltiel, Avi Shmida. "Foraging by Male and Female Solitary Bees with Implications for Pollination", Journal | 1 % |

of Insect Behavior, 2006

Publication

| | | |
|----|---|------|
| 24 | orca.cf.ac.uk Internet Source | <1 % |
| 25 | complete.bioone.org Internet Source | <1 % |
| 26 | agrivita.ub.ac.id Internet Source | <1 % |
| 27 | Submitted to UC, Boulder Student Paper | <1 % |
| 28 | tel.archives-ouvertes.fr Internet Source | <1 % |
| 29 | Zawar Hussain, Akbar Ali Khan. "A new index for measuring evenness", Communications in Statistics - Theory and Methods, 2017 Publication | <1 % |
| 30 | Submitted to Universitas Jenderal Soedirman Student Paper | <1 % |
| 31 | boris.unibe.ch Internet Source | <1 % |
| 32 | www.carpathianconvention.org Internet Source | <1 % |
| 33 | drji.org Internet Source | <1 % |

34

Internet Source

<1 %

35

H.T. Ngo, J. Gibbs, T. Griswold, L. Packer.
"Evaluating bee (Hymenoptera: Apoidea)
diversity using Malaise traps in coffee
landscapes of Costa Rica", The Canadian
Entomologist, 2013

Publication

<1 %

36

Rosana Halinski, Lucas Alejandro Garibaldi,
Charles Fernando dos Santos, André Luis
Acosta et al. "Forest fragments and natural
vegetation patches within crop fields contribute
to higher oilseed rape yields in Brazil",
Agricultural Systems, 2020

Publication

<1 %

37

Kasina, Muo, Manfred Kraemer, Christopher
Martius, and Dieter Wittmann. "Diversity and
activity density of bees visiting crop flowers in
Kakamega, Western Kenya", Journal of
Apicultural Research, 2009.

Publication

<1 %

38

www.cienciaanimal.ufpa.br

Internet Source

<1 %

39

rosenheim.faculty.ucdavis.edu

Internet Source

<1 %

40

Arthur Carlos de Oliveira, Camila Nonato

Junqueira, Solange Cristina Augusto. "Pesticides affect pollinator abundance and productivity of sunflower (L.) ", Journal of Apicultural Research, 2018

<1 %

Publication

41 Willem Proesmans, Dries Bonte, Guy Smagghe, Ivan Meeus et al. "Small forest patches as pollinator habitat: oases in an agricultural desert?", Landscape Ecology, 2019

<1 %

Publication

42 Imam Widhiono, Eming Sudiana, Darsono Darsono. "Diversity of Wild Bees along Elevational Gradient in an Agricultural Area in Central Java, Indonesia", Psyche: A Journal of Entomology, 2017

<1 %

Publication

43 Vincent Bretagnolle, Sabrina Gaba. "Weeds for bees? A review", Agronomy for Sustainable Development, 2015

<1 %

Publication

44 Marini, Lorenzo, Marino Quaranta, Paolo Fontana, Jacobus C. Biesmeijer, and Riccardo Bommarco. "Landscape context and elevation affect pollinator communities in intensive apple orchards", Basic and Applied Ecology, 2012.

<1 %

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off