


PAPER • OPEN ACCESS

## Preface

To cite this article: 2020 *J. Phys.: Conf. Ser.* **1494** 011001

View the [article online](#) for updates and enhancements.



**IOP | ebooks™**

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.



# Soedirman's International Conference on Mathematics and Applied Sciences 2019

Java Heritage Hotel Purwokerto  
October 23<sup>rd</sup> - 24<sup>th</sup>, 2019





# Committees

## Steering Committee/ Editorial Board

1. Prof. Abdellah Salhi, Ph.D. (University of Essex, UK)
2. Prof. Guiseppe Di Fazio, Ph.D. (University of Catania, Italy)
3. Prof. Hendra Gunawan, Ph.D. (Bandung Institute of Technology, Indonesia)
4. Prof. Dr.-Ing. R. Wahyu Widanarto (Jenderal Soedirman University, Indonesia)
5. Prof. Dr. Hadi Nur (University Teknologi Malaysia, Malaysia)
6. Dr. Hirokazu Saito (Tokyo University of Science, Japan)
7. Assoc. Prof. Dr. Imran Parvez (Hajee Mohammad Danesh Science and Technology University (HSTU), Bangladesh)
8. Dr. Jas Raj Subba (Royal University of Bhutan, Bhutan)
9. Dr. Saluma Samanman (Faculty of Science and Technology, Princess of Naradhiwas University, Narathiwat, Thailand)
10. Amin Fatoni, Ph. D. (Jenderal Soedirman University, Indonesia)
11. Dr. Eng. Mukhtar Effendi (Jenderal Soedirman University, Indonesia)
12. Drs. Wuryatmo A. Sidik, Dipl.Sc., M.Sc., Ph.D. (Jenderal Soedirman University, Indonesia)
13. Sri Maryani, Ph. D. (Jenderal Soedirman University, Indonesia)
14. Dadan Hermawan, Ph. D. (Jenderal Soedirman University, Indonesia)
15. Wahyu Tri Cahyanto, Ph. D. (Jenderal Soedirman University, Indonesia)
16. Dr. Abdullah Nur Aziz (Jenderal Soedirman University, Indonesia)
17. Dr. Jajang (Jenderal Soedirman University, Indonesia)
18. Dr. Hartiwi Diastuti (Jenderal Soedirman University, Indonesia)
19. Dr. Santi Nur Handayani (Jenderal Soedirman University, Indonesia)
20. Uyi Sulaeman, Ph. D. (Jenderal Soedirman University, Indonesia)
21. Bambang Hendriya G, Ph. D. (Jenderal Soedirman University, Indonesia)
22. R. Farzand Abdullatif, Ph. D. (Jenderal Soedirman University, Indonesia)

**Conference Chair** : Dr. Idha Sihwaningrum

**Conference Co-chair** :

Scientific : Amin Fatoni, Ph.D.  
Programs : Dr. Eng. Mukhtar Effendi



### Organizing Committee:

1. Dr. Nunung Nurhayati
2. Ari Asnani, Ph. D.
3. Dr. Zaroh Irayani
4. Rina Reorita, M. Si.
5. Dwi Kartika, M. Sc.
6. Wihantoro, M. Sc.
7. Abdul Naseh, A. Md.
8. Dr. Abdullah Nur Aziz, M. Si.
9. Frenky Oktavinanda
10. Kharis, S. Si.
11. Drs. Budi Pratikno, Ph. D.
12. Siti Muniroh, S. Si.
13. Andi Setiawan, A. Md.
14. Dena Martini, SE
15. Dr. Puji Lestari
16. Latief Nur Muhammad, ST
17. Listiyani Tri Hastuti, SE
18. Dian Riana Ningsih, M. Si.
19. Senny Widyaningsih, M. Si.
20. Siti Rahmah Nurshiami, M. Si.
21. Renny, M. Si.
22. Dr. Hartiwi Diastuti
23. Sukirlan ETR, MM
24. Rini Susanti, S. Si.
25. Sherly Arya Iskandar, SE
26. Eko Jatmiko, SH



## Report from The Organizing Committee

It is indeed my great pleasure and honor to welcome you all to Soedirman's International Conference on Mathematics and Applied Sciences (SICoMAS) 2019. The conference running this year is the first SICoMAS series hosted by Faculty of Mathematics and Natural Sciences Jenderal Soedirman University. As the development of technology and management of world resources for our future based on the innovation in Mathematics and Sciences, this conference takes issue "Innovation in Mathematics and Applied Sciences for better future".

SICoMAS 2019 aims to provide a platform for researchers, lecturers, teachers, students, practitioners, and industrial professionals to share knowledge, exchange ideas, collaborate, and present research results in the fields of Mathematics, Chemistry, Physics, and their applications. Hence, my sincere gratitude goes to our four keynote speakers (Prof. Dr. Hadi Nur from University Teknologi Malaysia, Prof. Dr. Hirokazu Saito from Tokyo University of Science, Dr. Devi Putra, ST, M.Sc. from Pertamina Research and Technology, and Uyi Sulaeman, Ph.D. from Jenderal Soedirman University), and our six invited speakers (Prof. Dr. Youtoh Imai from Nishogakusha University, Prof. Riyanto, Ph.D. from Universitas Islam Indonesia, Dr. Moh. Adhib Ulil Absor from Gadjah Mada University, Bambang Hendriya Guswanto, Ph.D, Dadan Hermawan, Ph.D. and Dr. Eng. Mukhtar Effendi, M. Eng. from Jenderal Soedirman University) for sharing their expertise in this conference. My deepest appreciation also goes to our 80 presenters and 7 non presenters for their commitment to participate in this conference.

As the output of this conference, some selected papers in the field of chemistry will be published in Jurnal Molekul which is accredited Sinta 1; and other selected papers in the fields of Mathematics, Physics, Physical Chemistry, and Innovative Chemistry Education will be published in IOP Conference Series Journal. So, I greatly thank Jenderal Soedirman University, all our contributors, and all the members of the committee for the invaluable support that makes this conference a reality.

Finally, I would like to apologize for any short comings found in this conference; and hopefully this two-day conference will be engraved in your memory.


The chair of SICoMAS 2019

PAPER • OPEN ACCESS

## Peer review statement

To cite this article: 2020 *J. Phys.: Conf. Ser.* **1494** 011002

View the [article online](#) for updates and enhancements.



**IOP | ebooks™**

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

## Peer review statement

All papers published in this volume of *Journal of Physics: Conference Series* have been peer reviewed through processes administered by the proceedings Editors. Reviews were conducted by expert referees to the professional and scientific standards expected of a proceedings journal published by IOP Publishing.



PAPER • OPEN ACCESS

## The validity of the properties of real numbers set to hyperreal numbers Set

To cite this article: R Masithoh *et al* 2020 *J. Phys.: Conf. Ser.* **1494** 012021

View the [article online](#) for updates and enhancements.



**IOP | ebooks™**

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.



# The validity of the properties of real numbers set to hyperreal numbers Set

**R Masithoh, B H Guswanto, and S Maryani**

Analysis and Algebra Laboratory, Department of Mathematics, Faculty of Mathematics and Natural Sciences, Jenderal Soedirman University, Jl. Dr. Soeparno 61, Karangwangkal, Purwokerto 53123, INDONESIA

E-mail: rahmasithoh@gmail.com

**Abstract.** This research discusses the validity of the properties of real numbers set to hyperreal numbers set, i.e. algebraic, ordered, and completeness properties, by using a finitely additive measure. This finitely additive measure is a map from the power set of natural numbers set onto set  $\{0,1\}$ . The subset of natural numbers set has measure zero if it's finite and one if it's infinite. The set of hyperreal numbers is constructed from equivalence classes of the set of all sequences of real numbers by using a relation involving the finitely additive measure, that is, two sequences of real numbers are said to be related if and only if those two sequences are the same almost everywhere. In the hyperreal numbers set, there exist infinitesimal numbers besides 0. Infinitesimal number is a number which is less than any positive real number and greater than any negative real number. So, in hyperreal number set, there are some smallest positive numbers. The results show that the hyperreal numbers set is an ordered and complete field.

## 1. Introduction

The set  $\mathbb{R}$  of real numbers can be constructed by using the concept of equivalence classes of the set of all rational Cauchy-sequence (see [3]). Abraham Robinson [2] had expanded the real numbers set. This expansion is called hyperreal numbers set  ${}^*\mathbb{R}$ . This numbers set is constructed by using the concept of equivalence classes of real numbers sequence. For example, both  $\{1/n\}$  and  $\{1/n^2\}$  are the sequences that converge to zero, but have different convergence rates. In  ${}^*\mathbb{R}$ , convergence rate is also considered. Then, in  ${}^*\mathbb{R}$ , both sequences are in different classes.

Another interesting property of hyperreal numbers set  ${}^*\mathbb{R}$  is there exist more than one “small” or infinitesimal number. Infinitesimal number is a number which is less than any positive real numbers and greater than any negative real numbers. Differently from hyperreal numbers set  ${}^*\mathbb{R}$ , real numbers set  $\mathbb{R}$  has one infinitesimal number, that is zero. Moreover, in  ${}^*\mathbb{R}$ , there is a “quantification” concept for “large” or infinite numbers. In  ${}^*\mathbb{R}$ , infinite property can be manifested as a number. Therefore, this paper investigates whether or not algebraic, ordered, and completeness properties, which are valid in real numbers set  $\mathbb{R}$  (see [1]), are also valid in hyperreal numbers set  ${}^*\mathbb{R}$ .

## 2. Methods

The study of the validity of the properties of real numbers set  $\mathbb{R}$  to hyperreal numbers set  ${}^*\mathbb{R}$  is performed by the following steps.



1. Constructing hyperreal numbers set using the concept of equivalence classes and measure  $m: P(\mathbb{N}) \rightarrow \{0,1\}$  with  $P(\mathbb{N})$  is the power set of natural numbers set  $\mathbb{N}$ .
2. Studying the properties of real numbers set which are valid in hyperreal numbers set, such as algebraic, ordered, and completeness properties by using the measure  $m$ .

### 3. Results and Discussion

Hyperreal numbers set  ${}^*\mathbb{R}$  is constructed by using the concept of equivalence classes in the set of real numbers sequences (see [2]).

#### 3.1 Construction of hyperreal numbers set ${}^*\mathbb{R}$

Hyperreal numbers set is constructed of using a measure  $m: P(\mathbb{N}) \rightarrow \{0,1\}$  with  $P(\mathbb{N})$  is the power set of natural numbers set  $\mathbb{N}$  which is defined by the following definition.

##### Definition 3.1

The measure  $m$  is finitely additive measure on  $P(\mathbb{N})$ , if

- i. for each  $A \subseteq \mathbb{N}$ ,  $m(A) \in \{0,1\}$ ;
- ii. measure  $m(\mathbb{N}) = 1$  and  $m(A) = 0$  for all finite set  $A \subseteq \mathbb{N}$ .

The measure  $m$  is finitely additive measure which means  $m(A \cup B) = m(A) + m(B)$ , for all disjoint sets  $A$  and  $B$ . The measure  $m$  divides the subset of  $\mathbb{N}$  into two parts, those are, a “large” or infinite set with a measure one and a “small” or finite set with a measure zero.

Any subset  $A \subset \mathbb{N}$  satisfies one of the conditions  $m(A) = 1$  or  $m(A^c) = 1$ . For any  $A, B \subseteq \mathbb{N}$  with  $m(A) = 1$  and  $m(B) = 1$ , then

$$m((A \cap B)^c) = m(A^c \cup B^c) \leq m(A^c) + m(B^c) = 0 + 0 = 0$$

which implies  $m(A \cap B) = 1$ . Next discussion is the construction of  ${}^*\mathbb{R}$  by using the concept of equivalence classes on the set of all real numbers sequences.

##### Definition 3.2

Let  $\mathfrak{R}$  is the set of all real numbers sequences and  $\sim$  is an equivalence relation on  $\mathfrak{R}$  which is defined by the following. For all  $\{a_n\}, \{b_n\} \in \mathfrak{R}$

$$\{a_n\} \sim \{b_n\} \text{ if and only if } m(\{n : a_n = b_n\}) = 1. \quad (1)$$

In other words,  $\{a_n\}$  is the same as  $\{b_n\}$  almost everywhere. This following is the definition of hyperreal numbers set by using equivalence relation  $\sim$ .

##### Definition 3.3

Let  $\{a_n\} \in \mathfrak{R}$ . The equivalence classes  $a$  with respect to the relation  $\sim$  is defined by

$${}^*a = \langle a_n \rangle = \{\{x_n\} \in \mathfrak{R} : \{x_n\} \sim \{a_n\}\}. \quad (2)$$

Next, hyperreal numbers set  ${}^*\mathbb{R}$  is a set of all equivalence classes in set  $\mathfrak{R}$  and is denoted by

$${}^*\mathbb{R} = \{\langle a_n \rangle : \{a_n\} \in \mathfrak{R}\}. \quad (3)$$

To study the properties of hyperreal numbers set which is analogous to real numbers set, the addition, multiplication, and order operation on  ${}^*\mathbb{R}$  are defined.

### Definition 3.4

For each  $\langle a_n \rangle, \langle b_n \rangle \in {}^*\mathbb{R}$ , the addition operation “+”, multiplication operation “ $\cdot$ ”, and order operation “ $<$ ” on  ${}^*\mathbb{R}$  are defined by

$$\begin{aligned}\langle a_n \rangle + \langle b_n \rangle &= \langle a_n + b_n \rangle \\ \langle a_n \rangle \cdot \langle b_n \rangle &= \langle a_n \cdot b_n \rangle \\ \langle a_n \rangle < \langle b_n \rangle &\text{ if and only if } m(\{n : a_n < b_n\}) = 1.\end{aligned}\quad (4)$$

Next, each  $a \in \mathbb{R}$  can be written as  ${}^*a = \langle a_n \rangle \in {}^*\mathbb{R}$  which means  $m(\{n : a_n = a\}) = 1$ . The set  ${}^*\mathbb{R}$  is divided into three parts, those are infinitesimal numbers, finite numbers, and infinite numbers.

### Definition 3.5

- a) An element  ${}^*x \in {}^*\mathbb{R}$  is said to be infinitesimal if  $-a < {}^*x < a$  for each  $a \in \mathbb{R}^+$ .
- b) An element  ${}^*x \in {}^*\mathbb{R}$  is said to be finite if  $-a < {}^*x < a$  for some  $a \in \mathbb{R}^+$ .
- c) An element  ${}^*x \in {}^*\mathbb{R}$  is said to be infinite if  ${}^*x \leq -a$  or  ${}^*x \geq a$  for each  $a \in \mathbb{R}^+$ .

### Example 3.1

The numbers  ${}^*0$ ,  ${}^*\delta_1 = \langle 1/n \rangle$ , and  ${}^*\delta_2 = \langle 1/n^2 \rangle$  in  ${}^*\mathbb{R}$  are infinitesimal. Since  $m(\{n : -a < 0_n < a\}) = 1$  then

$$m(\{n : -a < 1/n < a\}) = 1$$

and

$$m(\{n : -a < 1/n^2 < a\}) = 1$$

for each  $a \in \mathbb{R}^+$ , respectively. In addition,  ${}^*\delta_1 = \langle 1/n \rangle$  and  ${}^*\delta_2 = \langle 1/n^2 \rangle$  are clearly two different numbers. The number  $\langle n \rangle$  is infinite positive and number  $\langle -n^2 \rangle$  is infinite negative.

### Theorem 3.1

Every  ${}^*x \in {}^*\mathbb{R}$  which is finite can be written uniquely as  ${}^*x = a + \varepsilon$  with  $a \in \mathbb{R}$  and  $\varepsilon$  is infinitesimal.

Next,  ${}^*x$  and  ${}^*y$  are said to be infinitely close if  ${}^*x - {}^*y$  infinitesimal. This relationship between  ${}^*x$  and  ${}^*y$  is denoted by  ${}^*x \approx {}^*y$ .

### Definition 3.6

For each finite number  ${}^*x \in {}^*\mathbb{R}$ , an unique real number  $a$  that satisfies  ${}^*x \approx a$  is called the part of standard  ${}^*x$  and denoted by  $\text{st}({}^*x)$ . Conversely, for each  $a \in \mathbb{R}$ , the set of all  ${}^*x \in {}^*\mathbb{R}$  with  $a = \text{st}({}^*x)$  is called monad  $a$  and denoted by  $\text{mon}(a)$ .

### 3.2 Internal set

A subset of  ${}^*\mathbb{R}$  can be constructed by using measure  $m$  [2]. It is used to discuss the completeness properties of  ${}^*\mathbb{R}$ .

### Definition 3.7

Let  $A_n \in \mathbb{R}$  where  $n = 1, 2, 3, \dots$ . The sequence  $\{A_n\}$  defines  $\langle A_n \rangle \in {}^*\mathbb{R}$  where

$$\langle x_n \rangle \in \langle A_n \rangle \text{ if and only if } m(\{n: x_n \in A_n\}) = 1 \quad (5)$$

and the subset of  ${}^*\mathbb{R}$  which is obtained in this way is called an internal set.

### 3.3 The properties of hyperreal numbers set

**3.3.1 Algebraic properties.** In hyperreal numbers set  ${}^*\mathbb{R}$ , there are two binary operations, those are hyperreal addition and hyperreal multiplication operation. Hyperreal numbers set  ${}^*\mathbb{R}$  is a field with an identity element

$${}^*0 = \{\langle 0_n \rangle \in \mathfrak{R} : m(\{n: 0_n = 0\}) = 1\}$$

under the addition operation and the identity element

$${}^*1 = \{\langle 1_n \rangle \in \mathfrak{R} : m(\{n: 1_n = 1\}) = 1\}.$$

under the multiplication operation. Both identities under appropriate operations are unique. In addition, the multiplication of each element in  ${}^*\mathbb{R}$  and  ${}^*0$  is  ${}^*0$ .

#### Theorem 3.2

- a) If  ${}^*z, {}^*a \in {}^*\mathbb{R}$  and  ${}^*z + {}^*a = {}^*a$ , then  ${}^*z = {}^*0$
- b) If  ${}^*u, {}^*b \in {}^*\mathbb{R}$ ,  ${}^*b \neq {}^*0$ , and  ${}^*u \cdot {}^*b = {}^*b$ , then  ${}^*u = {}^*1$
- c) If  ${}^*a \in {}^*\mathbb{R}$ , then  ${}^*a \cdot {}^*0 = {}^*0$

#### Prove 3.2.a)

Let  ${}^*z, {}^*a \in {}^*\mathbb{R}$  where  ${}^*z = \langle z_n \rangle$  and  ${}^*a = \langle a_n \rangle$ . Observe that  ${}^*z + {}^*a = {}^*a$  which means  $\langle z_n \rangle + \langle a_n \rangle = \langle a_n \rangle$  and based on addition definition in  ${}^*\mathbb{R}$ ,  $\langle z_n + a_n \rangle = \langle a_n \rangle$  or in other words  $m(\{n: z_n + a_n = a_n\}) = 1$ . For each  $n \in \{n: z_n + a_n = a_n\}$ , if  $z_n + a_n = a_n$  then  $z_n = 0$ . Since  $m(\{n: z_n + a_n = a_n\}) = 1$  then  $m(\{n: z_n = 0\}) = 1$ . So, it can be proven that  ${}^*z = {}^*0$ . ■

Furthermore, under multiplication operation, every element in  ${}^*\mathbb{R} \setminus \{ {}^*0 \}$  has a unique inverse element and the multiplication of two numbers in  ${}^*\mathbb{R}$  that produces  ${}^*0$  is fulfilled when one or both of these numbers is  ${}^*0$ .

#### Theorem 3.3

- a) If  ${}^*a \neq {}^*0$  and  ${}^*a, {}^*b \in {}^*\mathbb{R}$  such that  ${}^*a \cdot {}^*b = {}^*1$ , then  ${}^*b = 1 / {}^*a$ .
- b) If  ${}^*a, {}^*b \in {}^*\mathbb{R}$  and  ${}^*a \cdot {}^*b = {}^*0$  then  ${}^*a = {}^*0$  or  ${}^*b = {}^*0$ .

#### Prove 3.3.b)

Let  ${}^*a, {}^*b \in {}^*\mathbb{R}$  where  ${}^*a = \langle a_n \rangle$  and  ${}^*b = \langle b_n \rangle$ . Observe that

$$\langle a_n \rangle \cdot \langle b_n \rangle = \langle a_n \cdot b_n \rangle,$$

which means

$$m(\{n: a_n \cdot b_n = 0\}) = 1.$$

Also, observe that

$$\{n: a_n \cdot b_n = 0\} = \{n: a_n = 0\} \cup \{n: b_n = 0\}.$$

If  $m(\{n: a_n = 0\}) = 0$  and  $m(\{n: b_n = 0\}) = 0$ , then  $\{n: a_n = 0\}$  and  $\{n: b_n = 0\}$  are finite sets. Consequently,  $\{n: a_n \cdot b_n = 0\}$  is finite or  $m(\{n: a_n \cdot b_n = 0\}) = 0$ . In this case, there is a contradiction. So, it must be  $m(\{n: a_n = 0\}) = 1$  or  $m(\{n: b_n = 0\}) = 1$ . In other words,  ${}^*a = {}^*0$  or  ${}^*b = {}^*0$  ■

**3.3.2 Ordered Properties.** Ordered properties in hyperreal numbers set is analogous to the ordered properties in real numbers set.

#### Theorem 3.4

- a) If  ${}^*a, {}^*b, {}^*c \in {}^*\mathbb{R}$  such that  ${}^*a > {}^*b$  and  ${}^*b > {}^*c$  then  ${}^*a > {}^*c$ ;  
 b) If  ${}^*a, {}^*b, {}^*c \in {}^*\mathbb{R}$  and  ${}^*a > {}^*b$  then  ${}^*a + {}^*c > {}^*b + {}^*c$ ;  
 c) If  ${}^*a, {}^*b, {}^*c \in {}^*\mathbb{R}$  such that  ${}^*a > {}^*0$  and  ${}^*b < {}^*c$ , then  ${}^*a \cdot {}^*b < {}^*a \cdot {}^*c$ , also if  ${}^*a, {}^*b, {}^*c \in {}^*\mathbb{R}$  such that  ${}^*a < {}^*0$  and  ${}^*b < {}^*c$  then  ${}^*a \cdot {}^*b > {}^*a \cdot {}^*c$ ;  
 d) If  ${}^*a, {}^*b \in {}^*\mathbb{R}$  and  ${}^*0 = \langle 0_n \rangle \in {}^*\mathbb{R}$  such that  ${}^*a \cdot {}^*b > {}^*0$ , then  ${}^*a > {}^*0$  and  ${}^*b > {}^*0$ , or  ${}^*a < {}^*0$  and  ${}^*b < {}^*0$ ;  
 e) If  ${}^*a, {}^*b \in {}^*\mathbb{R}$  and  ${}^*0 = \langle 0_n \rangle \in {}^*\mathbb{R}$  such that  ${}^*a \cdot {}^*b < {}^*0$ , then  ${}^*a > {}^*0$  and  ${}^*b < {}^*0$ , or  ${}^*a < {}^*0$  and  ${}^*b > {}^*0$ .

#### Prove

- a) We must prove that  ${}^*a > {}^*c$  with  ${}^*a = \langle a_n \rangle$ ,  ${}^*b = \langle b_n \rangle$ , and  ${}^*c = \langle c_n \rangle$ . Observe that  ${}^*a > {}^*b$  and  ${}^*b > {}^*c$  which mean  $\langle a_n \rangle > \langle b_n \rangle$  and  $\langle b_n \rangle > \langle c_n \rangle$ . This means that  $m(\{n: a_n > b_n\}) = 1$  and  $m(\{n: b_n > c_n\}) = 1$ . We must show that  $m(\{n: a_n > c_n\}) = 1$ . For every  $n \in \{n: a_n > b_n\} \cap \{n: b_n > c_n\}$ , if  $a_n > b_n$  and  $b_n > c_n$  then  $a_n > c_n$ . Next, since  $m(\{n: a_n > b_n\}) = 1$  and  $m(\{n: b_n > c_n\}) = 1$ , then

$$m(\{n: a_n > b_n\} \cap \{n: b_n > c_n\}) = 1.$$

Consequently,  $m(\{n: a_n > c_n\}) = 1$ . So, it can be proven that  ${}^*a > {}^*c$ .

- b) We must prove that  ${}^*a + {}^*c > {}^*b + {}^*c$  with  ${}^*a = \langle a_n \rangle$ ,  ${}^*b = \langle b_n \rangle$ , and  ${}^*c = \langle c_n \rangle$ . Observe that  ${}^*a > {}^*b$  or  $\langle a_n \rangle > \langle b_n \rangle$ . This means  $m(\{n: a_n > b_n\}) = 1$ . We must show that  $m(\{n: a_n + c_n > b_n + c_n\}) = 1$ . For every  $n \in \{n: a_n > b_n\}$ , if  $a_n > b_n$  then  $a_n + c_n > b_n + c_n$ . Next, since  $m(\{n: a_n > b_n\}) = 1$ , then

$$m(\{n: a_n + c_n > b_n + c_n\}) = 1.$$

So, we can prove that  ${}^*a + {}^*c > {}^*b + {}^*c$ .

- c) We must prove that  ${}^*a \cdot {}^*b < {}^*a \cdot {}^*c$  with  ${}^*a = \langle a_n \rangle$ ,  ${}^*b = \langle b_n \rangle$ , and  ${}^*c = \langle c_n \rangle$ . Observe that  ${}^*a > {}^*0$  and  ${}^*b < {}^*c$  which mean  $\langle a_n \rangle > \langle 0_n \rangle$  and  $\langle b_n \rangle < \langle c_n \rangle$ . This means  $m(\{n: a_n > 0\}) = 1$  and  $m(\{n: b_n < c_n\}) = 1$ . We must show that  $m(\{n: a_n b_n < a_n c_n\}) = 1$ . For every  $n \in \{n: a_n > 0\} \cap \{n: b_n < c_n\}$ , if  $a_n > 0$  and  $b_n < c_n$  then  $a_n b_n < a_n c_n$ . Next, since

$$m(\{n: a_n > 0\}) = 1$$

and

$$m(\{n: b_n < c_n\}) = 1$$

then  $m(\{n: a_n > 0\} \cap \{n: b_n < c_n\}) = 1$ . Therefore,  $m(\{n: a_n b_n < a_n c_n\}) = 1$ . So, we proved that  ${}^*a \cdot {}^*b < {}^*a \cdot {}^*c$ . Using the same way, it can be proven that if  ${}^*a, {}^*b, {}^*c \in {}^*\mathbb{R}$  such that  ${}^*a < {}^*0$  and  ${}^*b < {}^*c$ , then  ${}^*a \cdot {}^*b > {}^*a \cdot {}^*c$ .

d) We must prove that if  ${}^*a \cdot {}^*b > {}^*0$  then  ${}^*a > {}^*0$  and  ${}^*b > {}^*0$ , or  ${}^*a < {}^*0$  and  ${}^*b < {}^*0$  with  ${}^*a = \langle a_n \rangle$ ,  ${}^*b = \langle b_n \rangle$ , and  ${}^*0 = \langle 0_n \rangle \in {}^*\mathbb{R}$ . In order to do that, We must show that  $\langle a_n \rangle > \langle 0_n \rangle$  and  $\langle b_n \rangle > \langle 0_n \rangle$ , or  $\langle a_n \rangle < \langle 0_n \rangle$  and  $\langle b_n \rangle < \langle 0_n \rangle$ . In other words, we must show that  $m(\{n: a_n > 0\}) = 1$  and  $m(\{n: b_n > 0\}) = 1$ , or  $m(\{n: a_n < 0\}) = 1$  and  $m(\{n: b_n < 0\}) = 1$ . Observe that  ${}^*a \cdot {}^*b > {}^*0$  which means  $m(\{n: a_n b_n > 0\}) = 1$  and

$$\{n: a_n b_n > 0\} = \{n: a_n > 0 \text{ and } b_n > 0\} \cup \{n: a_n < 0 \text{ and } b_n < 0\}.$$

Since  $m(\{n: a_n b_n > 0\}) = 1$ , then

$$m(\{n: a_n > 0 \text{ and } b_n > 0\} \cup \{n: a_n < 0 \text{ and } b_n < 0\}) = 1.$$

Suppose

$$m(\{n: a_n > 0 \text{ and } b_n > 0\}) = 0$$

and

$$m(\{n: a_n < 0 \text{ and } b_n < 0\}) = 0.$$

Consequently,

$$\begin{aligned} m(\{n: a_n > 0 \text{ and } b_n > 0\} \cup \{n: a_n < 0 \text{ and } b_n < 0\}) &= m(\{n: a_n > 0 \text{ and } b_n > 0\}) + m(\{n: a_n < 0 \text{ and } b_n < 0\}) \\ &= 0 + 0 \\ &= 0. \end{aligned}$$

This contradicts to

$$m(\{n: a_n b_n > 0\}) = m(\{n: a_n > 0 \text{ and } b_n > 0\} \cup \{n: a_n < 0 \text{ and } b_n < 0\}) = 1.$$

So, must

$$m(\{n: a_n > 0 \text{ and } b_n > 0\}) = 1$$

or

$$m(\{n: a_n < 0 \text{ and } b_n < 0\}) = 1.$$

e) Using the same way as in proof of d), we can prove that if  ${}^*a \cdot {}^*b < {}^*0$ , then  ${}^*a > {}^*0$  and  ${}^*b < {}^*0$ , or  ${}^*a < {}^*0$  and  ${}^*b > {}^*0$ . ■

In real numbers set, natural numbers are positive real numbers[1]. Also, natural numbers in hyperreal numbers set are positive hyperreal numbers. Moreover, in real numbers set, there is no smallest positive real numbers[1]. In hyperreal numbers set, there exist infinitesimal numbers, for example  $\langle 1/n \rangle$ , since  $m(\{n: -a < 1/n < a\}) = 1$  for each  $a \in \mathbb{R}^+$ . Therefore, in hyperreal numbers set, there are smallest positive hyperreal numbers.

**3.3.3 Completeness Properties.** This following is a discussion of completeness properties of  ${}^*\mathbb{R}$ , that is the completeness properties which is related to the supremum and infimum concepts in  ${}^*\mathbb{R}$ .

### Definition 3.8

Let  ${}^*A$  is a non-empty subset of  ${}^*\mathbb{R}$ .

- a) The set  ${}^*A$  is called bounded above if there exists  ${}^*u \in {}^*\mathbb{R}$  such that  ${}^*a \leq {}^*u$  for each  ${}^*a \in {}^*A$ . The hyperreal number such  ${}^*u$  is called as upper bound of  ${}^*A$ .
- b) The set  ${}^*A$  is called bounded below if there exists  ${}^*v \in {}^*\mathbb{R}$  such that  ${}^*v \leq {}^*a$  for each  ${}^*a \in {}^*A$ . The hyperreal number such  ${}^*v$  is called lower bound of  ${}^*A$ .
- c) The set  ${}^*A$  is called bounded if  ${}^*A$  is both bounded above and bounded below. Conversely, a set  ${}^*A$  is called unbounded, if  ${}^*A$  is not bounded above or not bounded below.

### Definition 3.9

Let  ${}^*A$  is a non-empty subset of  ${}^*\mathbb{R}$ .

- a) If  ${}^*A$  bounded above, then a number  ${}^*u$  is called a supremum (least upper bound) of  ${}^*A$  if it satisfies these conditions:
- ${}^*u$  is an upper bound of  ${}^*A$ ;
  - if  ${}^*s$  is any upper bound of  ${}^*A$ , then  ${}^*u \leq {}^*s$ .
- b) If  ${}^*A$  bounded below, then  ${}^*v$  is called an infimum (greatest lower bound) of  ${}^*A$  if it satisfies these conditions:
- ${}^*v$  is a lower bound of  ${}^*A$ ;
  - if  ${}^*t$  is any lower bound of  ${}^*A$ , then  ${}^*t \leq {}^*v$ .

### Theorem 3.5

Let  ${}^*u, {}^*v \in {}^*\mathbb{R}$ , and  ${}^*A$  is a non-empty subset of  ${}^*\mathbb{R}$ .

- a) An element  ${}^*u$  is a supremum  ${}^*A$  if and only if  ${}^*z < {}^*u$  implies  ${}^*a^z > {}^*z$  for some  ${}^*a^z \in {}^*A$ .
- b) An element  ${}^*v$  is an infimum  ${}^*A$  if and only if  ${}^*z > {}^*v$  implies  ${}^*a^z < {}^*z$  for some  ${}^*a^z \in {}^*A$ .

#### Prove 3.5.a)

Since  ${}^*A = \langle A_i \rangle$  is a non-empty internal set of  ${}^*\mathbb{R}$ , then there exist  ${}^*a \in {}^*A$  such that  $m(\{i: a_i \in A_i\}) = 1$ . Let  ${}^*u = \sup {}^*A$  is supremum  ${}^*A$ , which means for every  ${}^*a \in {}^*A$ ,  $m(\{i: a_i \leq u_i = \sup A_i\}) = 1$ . Based on supremum theorem in real numbers set,  $u_i$  is a supremum  $A_i$  if and only if, for each  $i \in \{i: a_i \leq u_i\}$ , if  $z_i < u_i$  then there exists  $a_i^z \in A_i$  such that  $z_i < a_i^z$  [1]. Since  $m(\{i: a_i \leq u_i\}) = 1$ , then

$$m(\{i: z_i < a_i^z\}) = 1.$$

Consequently, if  ${}^*z = \langle z_i \rangle < {}^*u$  then there exists  ${}^*a^z = \langle a_i^z \rangle$  such that  ${}^*z < {}^*a^z$ . ■

### Theorem 3.6

Let  ${}^*u, {}^*v \in {}^*\mathbb{R}$ ,  ${}^*A$  is a non-empty internal set of  ${}^*\mathbb{R}$ .

- a) An element  ${}^*u$  is a supremum of  ${}^*A$ , if and only if for every  ${}^*\varepsilon > {}^*0$  there exists  ${}^*a^\varepsilon \in {}^*A$  such that  ${}^*a^\varepsilon > {}^*u - {}^*\varepsilon$ .
- b) An element  ${}^*v$  is an infimum of  ${}^*A$ , if and only if for each  ${}^*\varepsilon > {}^*0$  there exist  ${}^*a^\varepsilon \in {}^*A$  such that  ${}^*a^\varepsilon < {}^*v + {}^*\varepsilon$ .

#### Prove 3.6.a)

Since  ${}^*A = \langle A_i \rangle$  is a non-empty internal set of  ${}^*\mathbb{R}$ , then there exist  ${}^*a \in {}^*A$  such that  $m(\{i: a_i \in A_i\}) = 1$ . Let  ${}^*u = \sup {}^*A$  is a supremum of  ${}^*A$ , which means for each  ${}^*a \in {}^*A$ ,  $m(\{i: a_i \leq u_i = \sup A_i\}) = 1$ . Based on supremum theorem of real numbers set,  $u_i$  is a supremum  $A_i$  if and only if, for each  $i \in \{i: a_i \leq u_i\}$ , every  $\varepsilon_i > 0$ , there exist  $a_i^\varepsilon \in A_i$  such that  $a_i^\varepsilon > u_i - \varepsilon_i$  [1]. Since  $m(\{i: a_i \leq u_i\}) = 1$ , then

$m(\{i: a_i^\varepsilon > u_i - \varepsilon_i\}) = 1$ . Consequently, for each  ${}^*\varepsilon = \langle \varepsilon_i \rangle > {}^*0$  there exists  ${}^*a^\varepsilon = \langle a_i^\varepsilon \rangle \in {}^*A$  such that  ${}^*a^\varepsilon > {}^*u - {}^*\varepsilon$ . ■

### Completeness Properties

- a) Every non-empty internal set of  ${}^*\mathbb{R}$  that has an upper bound also has a least upper bound.  
 b) Every non-empty internal set of  ${}^*\mathbb{R}$  that has a lower bound also has a greatest lower bound.

#### Prove a)

Let  ${}^*A = \langle A_i \rangle$  is an internal set in  ${}^*\mathbb{R}$  which is bounded above  ${}^*a = \langle a_i \rangle$ . For each  ${}^*x \in {}^*A$ ,  ${}^*x \leq {}^*a$  which means if  $\langle x_i \rangle \in \langle A_i \rangle$  then  $\langle x_i \rangle \leq \langle a_i \rangle$ . This means  $m(\{i: x_i \in A_i\}) = 1$  and  $m(\{i: x_i \leq a_i\}) = 1$ . consequently, almost every  $A_i$  bounded above by  $a_i$ . Next, we must prove that  ${}^*b = \langle \sup A_i \rangle$  is a sup  ${}^*A$  by showing:

- i. for each  $\langle x_i \rangle \in \langle A_i \rangle$ ,  $m(\{i: \sup A_i \geq x_i\}) = 1$ .  
 ii. for each upper bound  ${}^*a$  of  ${}^*A$ ,  $m(\{i: \sup A_i \leq a_i\}) = 1$ .

Since  $\sup A_i$  is a least upper bound of  $A_i$ , then for every  ${}^*x \in {}^*A$ ,  $m(\{i: \sup A_i \geq x_i\}) = 1$  and for every upper bound  ${}^*a$  of  ${}^*A$ ,  $m(\{i: \sup A_i \leq a_i\}) = 1$ . So, it can be proven that  ${}^*b = \langle \sup A_i \rangle$  is least upper bound of  ${}^*A$ . ■

Hyperreal numbers set can be described as follows.




Figure 3.1 Hyperreal Numbers Set

Based on Figure 3.1, hyperreal numbers set is divided into three parts, those are negative infinite, finite, and positive infinite numbers. Based on Theorem 3.1, standard part of hyperreal numbers which are finite, are real numbers. In other words, the set of all standard part of hyperreal numbers which is finite is real numbers set.

### 4. Conclusions

Based on result and discussion above, it can be concluded that hyperreal numbers set  ${}^*\mathbb{R}$  is the set of all equivalence classes in set  $\mathfrak{R}$  and denoted by

$${}^*\mathbb{R} = \{\langle a_n \rangle: \{a_n\} \in \mathfrak{R}\}.$$

Hyperreal numbers set is a field with the identity element under the addition operation, that is



$${}^*0 = \left\{ \{0_n\} \in \mathfrak{R} : m(\{n : 0_n = 0\}) = 1 \right\}$$

and the identity element under the multiplication operation, that is

$${}^*1 = \left\{ \{1_n\} \in \mathfrak{R} : m(\{n : 1_n = 1\}) = 1 \right\}.$$

















The ordered properties of real numbers set also valid to hyperreal numbers set, except the ordered properties of  $\mathbb{R}$  which says there is no a smallest positive real number. In the hyperreal numbers set there are infinitesimal numbers other than zero, for example  $\langle 1/n \rangle$ . All completeness properties in real numbers set is also valid in hyperreal numbers set  ${}^*\mathbb{R}$ . Thus, it can be concluded that the hyperreal numbers set is an ordered and complete field.

















## References

- [1] Bartle R G and Sherbert D R. 2010 *Introduction to real analysis* (New York: John Wiley & Sons, Inc.)
- [2] Cutland N (Ed) 1988 *Nonstandard analysis and its application* (New York: Cambridge University Press)
- [3] Strichartz R S 2000 *The way of analysis* ( Burlington: Jones and Bartlett Publishers, Inc.)

## DAFTAR ISI PROCEEDING SICOMAS 2019

<b>OPEN ACCESS</b> <b>The proof of properties of dihedral group and its commutative elements</b> H Muhammad, L Susilowati and Fatmawati <a href="#">+ Open abstract</a> <a href="#">View article</a> <a href="#">PDF</a>	012003
<b>OPEN ACCESS</b> <b>Petri Net Model and Max-Plus Algebra on Queue in Clinic UNS Medical Center</b> T. Sulistyaningsih, Siswanto and Pangadi <a href="#">+ Open abstract</a> <a href="#">View article</a> <a href="#">PDF</a>	012004
<b>OPEN ACCESS</b> <b>Cat swarm optimization and Levenberg-Marquardt for model identification and prediction tuberculosis disease spreading</b> A Nuari, A Damayanti and A B Pratiwi <a href="#">+ Open abstract</a> <a href="#">View article</a> <a href="#">PDF</a>	012005
<b>OPEN ACCESS</b> <b>Numerical comparisons based on new NCP-functions</b> I Nurhidayat and L C Han <a href="#">+ Open abstract</a> <a href="#">View article</a> <a href="#">PDF</a>	012006
<b>OPEN ACCESS</b> <b>Ant System Algorithm in Renewing Pheromone for Completing Quadratic Assignment Problem</b> F M Utami, Renny and S R Nurshiami <a href="#">+ Open abstract</a> <a href="#">View article</a> <a href="#">PDF</a>	012007
<b>OPEN ACCESS</b> <b>Clustering of fish freshness using discrete wavelet transform and Kohonen self organizing map</b> J Anvy, A Damayanti and A B Pratiwi <a href="#">+ Open abstract</a> <a href="#">View article</a> <a href="#">PDF</a>	012008
<b>OPEN ACCESS</b> <b>Annihilator graph of semiring of matrices over Boolean semiring</b> R Yudatama, V Y Kurniawan and S B Wiyono <a href="#">+ Open abstract</a> <a href="#">View article</a> <a href="#">PDF</a>	012009
<b>OPEN ACCESS</b> <b>Interval min-plus algebraic structure and matrices over interval min-plus algebra</b> A R Awallia, Siswanto and V Y Kurniawan <a href="#">+ Open abstract</a> <a href="#">View article</a> <a href="#">PDF</a>	012010

<p><b>OPEN ACCESS</b></p> <p><b>On Cartesian product of intuitionistic Q-fuzzy ideal on ordered semigroup</b></p> <p>N Hidayat, I Yanti, Z Fitriah, D Miftah, S Anam, S I Rahman and F Hidayat</p> <p>+ Open abstract    View article    PDF</p>	012011
<p><b>OPEN ACCESS</b></p> <p><b>Implementation of Bayesian Mixture Models in identifying subpopulation of breast cancer patients based on blood test measurements</b></p> <p>N Dwimantara, S Abdullah, A Bustamam and A Rachman</p> <p>+ Open abstract    View article    PDF</p>	012012
<p><b>OPEN ACCESS</b></p> <p><b>Determining soil fertility using principal componen regression analysis of oil palm plantation in West Sulawesi, Indonesia</b></p> <p>U S Pasaribu, N Nurhayati, N F F Ilmi and K N Sari</p> <p>+ Open abstract    View article    PDF</p>	012013
<p><b>OPEN ACCESS</b></p> <p><b>Half-space model problem for a compressible fluid model of Korteweg type with slip boundary condition</b></p> <p>Suma Inna, Sri Maryani and Hirokazu Saito</p> <p>+ Open abstract    View article    PDF</p>	012014
<p><b>OPEN ACCESS</b></p> <p><b>The metric dimension of <math>k</math>-subdivision graphs</b></p> <p>L Susilowati, S Zahidah, R D Nastiti and M I Utoyo</p> <p>+ Open abstract    View article    PDF</p>	012015
<p><b>OPEN ACCESS</b></p> <p><b>Jacobson graph construction of ring <math>\mathbb{Z}_3^n</math>, for <math>n &gt; 1</math></b></p> <p>A Novictor, L Susilowati and Fatmawati</p> <p>+ Open abstract    View article    PDF</p>	012016
<p><b>OPEN ACCESS</b></p> <p><b>Betweenness centrality in corona product of <math>K_n</math> and <math>C_m</math> graph</b></p> <p>T I Haryadi and L Susilowati</p> <p>+ Open abstract    View article    PDF</p>	012017
<p><b>OPEN ACCESS</b></p> <p><b>Algorithm and computer program to determine metric dimension of graph</b></p> <p>F Muhammad and L Susilowati</p> <p>+ Open abstract    View article    PDF</p>	012018

<b>OPEN ACCESS</b> <b>An Adams type inequality of fractional integral operator on hypergroups</b> I Sihwaningrum, A Wardayani and Y Dasril + Open abstract  View article  PDF	012019
<b>OPEN ACCESS</b> <b>Regular rings and their properties</b> A Wardayani, I Kharismawati and I Sihwaningrum + Open abstract  View article  PDF	012020
<b>OPEN ACCESS</b> <b>The validity of the properties of real numbers set to hyperreal numbers Set</b> R Masithoh, B H Guswanto and S Maryani + Open abstract  View article  PDF	012021
<b>OPEN ACCESS</b> <b>Removal of rhodamine b using 4-hydroxy-3-methoxyphenylcalix[4]resorcinarene</b> S N Handayani, Irmanto and N N Indriyani + Open abstract  View article  PDF	012022
<b>OPEN ACCESS</b> <b>Adsorption of Gold(III), Copper(II), Nickel(II) on Amino Silica Hybrid Coated Magnetite</b> M T Thahir, N Nuryono and S J Santosa + Open abstract  View article  PDF	012023
<b>OPEN ACCESS</b> <b>Kinetics Adsorption of Strontium(II) by Silica Xerogel from Fly Ash</b> N A Kundari, K Megasari and C Yusephin + Open abstract  View article  PDF	012024
<b>OPEN ACCESS</b> <b>Triterpenoid and Steroid from the Rind of <i>Chisocheton macrophyllus</i> (Meliaceae)</b> V M Sari, Nurlelasari, D Harneti, R maharani, N Indrayati, M N Azmi and U Supratman + Open abstract  View article  PDF	012025
<b>OPEN ACCESS</b> <b>Introducing Colorimetric Analysis with Document Scanner for High School Students</b> A Fatoni, Supiani, D W Dwiasi and M D Anggraeni + Open abstract  View article  PDF	012026

<b>OPEN ACCESS</b>	012027
<b>The pH dependence of Ag<sub>3</sub>PO<sub>4</sub> synthesis on visible light photocatalytic activities</b>	
I Futihah, A Riapanitra, S Yin and U Sulaeman	
<a href="#">+ Open abstract</a> <a href="#">View article</a> <a href="#">PDF</a>	
<b>OPEN ACCESS</b>	012028
<b>Improved reuse and affinity of enzyme using immobilized amylase on alginate matrix</b>	
Zusfahair, D R Ningsih, D Kartika, M Kurniasih, R Nofiani and A Fatoni	
<a href="#">+ Open abstract</a> <a href="#">View article</a> <a href="#">PDF</a>	
<b>OPEN ACCESS</b>	012029
<b>Effect of ultraviolet and visible lights on degradation of congo red dye using Fe<sup>2+</sup>/H<sub>2</sub>O<sub>2</sub></b>	
J Tama, K Riyani and T Setyaningtyas	
<a href="#">+ Open abstract</a> <a href="#">View article</a> <a href="#">PDF</a>	
<b>OPEN ACCESS</b>	012030
<b>Introducing Iron Analysis with Smartphone Camera for High School Students</b>	
A Fatoni, Zusfahair and M D Anggraeni	
<a href="#">+ Open abstract</a> <a href="#">View article</a> <a href="#">PDF</a>	
<b>OPEN ACCESS</b>	012031
<b>Quantitative Structure Activity Relationship Analysis of 1,3,4-Thiadiazole Derivatives as Anti-Inflammatory using Parameterized Model 3 Method</b>	
Ponco Iswanto, Maylani Permata Saputri, Vaulina Y.D. Eva and Sari Paramita	
<a href="#">+ Open abstract</a> <a href="#">View article</a> <a href="#">PDF</a>	
<b>OPEN ACCESS</b>	012032
<b>Application of High Performance Liquid Chromatography Method for Triadimenol Analysis in Water Sample</b>	
D Hermawan, Cacu, Suwandri, A Fatoni, Mudasir and H Y Aboul-Enein	
<a href="#">+ Open abstract</a> <a href="#">View article</a> <a href="#">PDF</a>	
<b>OPEN ACCESS</b>	012033
<b>The Effect of Multiplanes Graphene Flakes Coating on Ag Nanowire Transparent Conductive</b>	
Nur Fiqalbi, Ratih Dwi Cahyaningrum, Bambang Prihandoko and Harsojo	
<a href="#">+ Open abstract</a> <a href="#">View article</a> <a href="#">PDF</a>	
<b>OPEN ACCESS</b>	012034
<b>Mapping of gold mineralization using 3D inversion magnetic data at Zone X, West Java</b>	
Muzzammil Al Macky, Mohammad Syamsu Rosid and Agus Pajrin Jaman	
<a href="#">+ Open abstract</a> <a href="#">View article</a> <a href="#">PDF</a>	

- 
- OPEN ACCESS** 012035  
**Hydrocarbon Reservoir Characterization of 'Y' Field in Kutai Basin East Kalimantan Using Artificial Neural Network Method**  
Y R Hendryan, M S Rosid and Haryono  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012036  
**Study of Thin Film Cu Nanowire and Reduced Graphene Oxide as A Gas Sensor**  
Noviana Candra Wijayanti, Ratih Dwi Cahyaningrum and Harsojo  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012037  
**Study of Transparent Conductive of Ag Nanowire Under Ambient Environment**  
Ratih Dwi Cahyaningrum and Harsojo  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012038  
**Estimation of Potential of Iron Sand in The Eastern Coastal Area of Cilacap Regency Based on The Local Magnetic Anomalies Data**  
Sehah, M A Kurniawan, I Andriyanto, B Arismawan, A Risyad, A M Wibowo, S A Raharjo and Sugito  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012039  
**Estimation of Ideal Configuration and Dimension of Pico Hydropower using Euler-Lagrange Equation and Runge-Kutta Method**  
J. Aminuddin, Nurhayati, A. Widiyani, P. Razi, Wihantoro, A.N. Aziz, R.F. Abdullatif and A. Arifin  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012040  
**CO oxidation through the formation of carboxyl intermediate on Pt(111) surface: A first principles study**  
S Zulaehah, W T Cahyanto, I N Fitriani, F Abdullatif, W Widanarto and M Effendi  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012041  
**Designing of Muffler Part for Car Exhaust System with Low Emission and Noise using Conjugate Gradient Method**  
J. Aminuddin, Wihantoro, Bilalodin, Sunardi and N. Rauf  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012042  
**The effects of milling time on structure, magnetic properties and microwave absorption capability of strontium lanthanum ferrite compounds**  
Mukhtar Effendi, Endra Nugraha, Wahyu Tri Cahyanto and Wahyu Widanarto

---

**OPEN ACCESS**

012043

**Magnetic and microwave absorption properties of  $\text{Mn}^{4+}$  doped barium-natural ferrites prepared by the modified solid-state reaction method**

H. R. Luthfianti, W. Widanarto, S. K. Ghoshal, M. Effendi and W. T. Cahyanto

[+ Open abstract](#)[View article](#)[PDF](#)

---

**OPEN ACCESS**

012044

**Cimandiri Fault Identification Using Earthquake Tomography Double-Difference Method**

R G Simanjorang, M S Rosid, A S Sembiring, Daryono and N Heryandoko

[+ Open abstract](#)[View article](#)[PDF](#)

---

**OPEN ACCESS**

012045

**Effect of Carbon Active Addition to Electrochemical Performance of  $\text{Li}_4\text{Ti}_5\text{O}_{12}/\text{SnO}_2$  Composite Anode on Lithiumion Battery**

Ebsan Simamora, Jeffrey Riady, Bambang Priyono and Anne Zulfia Syahril

[+ Open abstract](#)[View article](#)[PDF](#)



**SICoMAS 2019**  
**Soedirman's International Conference**  
**on Mathematics and Applied Sciences**  
Purwokerto INDONESIA, October 23-24, 2019  
<https://conference.unsoed.ac.id/index.php/SICoMAS/>



No. : T/1710/UN23.11.SICOMAS/DL.17/2019  
Subject : **Invited Speaker Invitation**

July 12th, 2019

Bambang Hendriya Guswanto, Ph.D.  
Jenderal Soedirman University, Indonesia

Dear Bambang Hendriya Guswanto, Ph.D.

We are pleased to invite you as an **Invited Speaker** on the "**Soedirman's International Conference on Mathematics and Applied Sciences (SICoMAS)**" which will be held on **October 23-24, 2019** at the Java Heritage Hotel Purwokerto, Central Java, Indonesia. The themes of the conference will be on the following topics: "**Innovation in Mathematics and Applied Sciences for Better Future**"

Should you require any information, please do not hesitate to contact us. We are looking forward to meeting you in Purwokerto.

Sincerely yours,

Dr. Idha Sihwaningrum, M.Sc.St.  
Chair of the Organizing Committee

**Faculty of Mathematics and Natural Sciences,**  
**Universitas Jenderal Soedirman**  
Jl. Dr. Soeparno 61, Karangwangkal, Purwokerto  
Jawa Tengah, INDONESIA 53125  
Phone: +62 281 637793 | Fax: +62 281 637793

**Contact:**  
Idha Sihwaningrum (+62 812-2759-820)  
Wahyu Widanarto (+62 857-22699-6007)  
Amin Fatoni (+62 812-2771-7444)  
Email: [sicomas@unsoed.ac.id](mailto:sicomas@unsoed.ac.id)