

Jurnal Rekayasa Kimia & Lingkungan

(Journal of Chemical Engineering & Environment)

Accredited By:
Director General of Higher Education, Research & Technology
No: 158/E/KPT/2021

ISSN 1412-5064
E-ISSN 2356-1651

(Sinta 2)

HOME ABOUT LOGIN REGISTER CATEGORIES SEARCH CURRENT ARCHIVES ANNOUNCEMENTS EDITORIAL TEAM RE

Vol. 9 No. 4 Desember 2013

ISSN 1412-5064

Jurnal
Rekayasa Kimia
&
Lingkungan
Journal of Chemical Engineering and Environment

Home - Vol 17, No 1 (2022)

JURNAL REKAYASA KIMIA & LINGKUNGAN

Jurnal Rekayasa Kimia dan Lingkungan (ISSN: 1412-5064 / E-ISSN: 2356-1651) with the variant title *Journal of Chemical Engineering and Environment* is an open-access journal published twice a year (June and December) by the Chemical Engineering Department, Faculty of Engineering, Universitas Syiah Kuala, Indonesia.

The printed version of JRKL has been published since 2004. The online version has been published since 2006. The journal is indexed by the Indonesian Publication Index (IPI), Google Scholar, and the Indonesian ministry of research, technology, and higher education (Sinta 2).

JRKL has been accredited to SINTA 2 since 2018 based on the Decree of the Ministry of Research, Technology and Higher Education No: 32A/E/KPT/2018 and extended by the Decree of the Director General of Higher Education, Research and Technology No: 158/E/KPT/2021.

Beginning with issue Volume 17 Number 1 in 2022, JRKL will transition to a continuous publishing model. This means that as soon as an article is finished, it will be published online, resulting in a steady stream of topical, high-quality content. When an issue cycle ends, the articles go to print, and the next issue begins online. Because the publication of articles is no longer dictated by our print schedule, your readers, students, and colleagues can access and cite articles much more quickly.

The scope of journal include food and biochemical engineering, catalytic reaction engineering, clean energy technology, environmental and safety technology, fundamentals of chemical engineering and applied industrial engineering, industrial engineering, material science engineering, process and control engineering, polymer and petrochemical technology, membrane technology, agro-industrial technology, separation and purification technology, environmental modelling, environmental and information sciences, and water and waste water treatment and management.

The editors welcome manuscripts in the form of research articles, literature review, or case reports on the above topics. To be published in the Jurnal Rekayasa Kimia dan Lingkungan, the manuscripts may not be submitted to other journals or are being considered for publication in other journals.



9 772356 165006

VISITOR STATS

Viewers

	140,819		468		293
	11,109		452		243
	813		418		242
	774		346		229

Pageviews: 389,697



TOOLS



INFORMATION

- For Readers
- For Authors
- For Librarians

ANNOUNCEMENTS

Article Language Requirement

STARTING IN DECEMBER 2019 EDITION, VOLUME 14, NUMBER 2, THE RKL JOURNAL WILL PUBLISH A FULL VERSION IN ENGLISH. THEREFORE, THE AUTHOR IS REQUIRED TO PREPARE THE MANUSCRIPT IN THE ENGLISH VERSION.

Posted: 2019-07-10

More...

Call for Paper: Call For Paper

JRKL invites researchers to participate in the publication of manuscripts of laboratory research papers or review papers. JRKL is published twice a year, June and December editions.

Posted: 2017-06-28

More...

More Announcements...

VOL 17, NO 1 (2022): JURNAL REKAYASA KIMIA DAN LINGKUNGAN (JANUARY-JUNE, 2022)

TABLE OF CONTENTS

Articles

Effect of Acid Concentration on the Properties of Microcrystalline Cellulose from Pineapple Crown Leaf
Fitriani Fitriani, Sri Aprilia, Nasrul Arahman, Muhammad Roil Bilad
10.23955/rkl.v17i1.21010

Synthesis of Renewable Diesel from Pure Plant Oil of Kapuk (Ceiba pentandra) Seed Using NiMo γ -Al₂O₃ Catalysts

PDF
1-7

PDF
8-17



USER

Username
Password
☐ Remember me

JOURNAL

Search

Search Scope
All

Browse

- By Issue
- By Author
- By Title
- Other Journals
- Categories

- PUBLIC
- AUTHC
- MANUS
- PUBLIS

INDEXED I



Edwin Permana, Excel Emerlan Samiar Wasistha, Septina Is Heriyanti, Sri Djangkung Sumbogo Murti,
Lenny Marlinda, Indra Lasmana Tarigan
[10.23955/rkl.v17i1.21673](#)



[Distribution pattern of Pb and Zn contamination in rivers near industrial zone in Aceh, Indonesia, revealed by principal component analysis \(PCA\)](#)

PDF
18-26

Lelifajri Lelifajri, Rahmadi Rahmadi, Rinaldi Idroes, Supriatno Supriatno, Eka Safitri, Nazaruddin
Nazaruddin, Ilham Maulana, Khairi Suhud
[10.23955/rkl.v17i1.22672](#)



[Synthesis and characterization of nanoemulsion of citronella oil \(Cymbopogon Nardus L. Rendle\) and its Sunscreen Potential Evaluation](#)

PDF
27-34

Undri Rastuti, Uyi Sulaeman, Senny Widyaningsih, Sity Khalidah Zia, Ryan Aditya Mahendra
[10.23955/rkl.v17i1.23077](#)

[Adsorption Of Cr\(VI\) Ions Using Sugar Palm Fruit Peel \(Arenga Pinnata\) Immobilized in Ca-Alginate Bead](#)

PDF
35-43

Intan Lestari, Widy Prasetyo, Indra Lasmana Tarigan, Edwin Permana
[10.23955/rkl.v17i1.22873](#)

[Optimizing Gold Recovery of Artisanal Mining: A Lesson Learned from Kenya](#)

PDF
44-53

Izzan Nur Aslam, Nestor Orcon, Bern Klein, Pocut Nurul Alam
[10.23955/rkl.v17i1.23223](#)

[Ethanol Dehydration in Fixed Bed Column Using Pellet Adsorbent From Natural Zeolite Lampung](#)

PDF
54-64

Jonathan Kristian Adisaputro, Simparmin Br. Ginting
[10.23955/rkl.v17i1.24866](#)

JURNAL REKAYASA KIMIA & LINGKUNGAN

Jurusan Teknik Kimia Universitas Syiah Kuala, Jl. Tgk. Syech Abdur Rauf No.7, Kopelma Darussalam, Banda Aceh, INDONESIA

PRINCIPAL CONTACT

Nasrul Arahman, Prof. Dr. S.T., M.T.
Phone: +62813-6092-7917
E-mail: rkl@che.unsyiah.ac.id, nasrular@unsyiah.ac.id

SUPPORT CONTACT

Mirna Rahmah Lubis
E-mail: mirna@che.unsyiah.ac.id

Wahyu Rinaldi, ST, M.Sc.
E-mail: wahyu.rinaldi@che.unsyiah.ac.id

00185009

VISITORS

Jurnal Rekayasa Kimia & Lingkungan

(Journal of Chemical Engineering & Environment)

Accredited By:
Director General of Higher Education, Research & Technology
No: 158/E/KPT/2021

ISSN 1412-5064
E-ISSN 2356-1412

(Sinta 2)

HOME ABOUT LOGIN REGISTER CATEGORIES SEARCH CURRENT ARCHIVES ANNOUNCEMENTS EDITORIAL TEAM RE

Vol. 9 No. 4 Desember 2013

ISSN 1412-5064

Jurnal
**Rekayasa Kimia
&
Lingkungan**
Journal of Chemical Engineering and Environment



Jurnal Rekayasa Kimia & Lingkungan Vol. 9 No. 4 Desember 2013 ISSN 1412-5064



9 772356 166006

VISITOR STATS

Viewers

	140,819		468		293
	11,109		452		243
	813		418		242
	774		346		229

Pageviews: 389,697



TOOLS



INFORMATION

- For Readers
- For Authors
- For Librarians

Home > Editorial Team

EDITORIAL TEAM

EDITOR IN CHIEF

PROF. DR. NASRUL ARAHMAN, S.T., M.T.

Chemical Engineering Department, Syiah Kuala University, Banda Aceh 23211, Indonesia

TRACK EDITOR

PROF. DR. M. FAISAL, S.T., M. ENG.

Chemical Engineering Department, Syiah Kuala University, Banda Aceh 23211, Indonesia

PROF. DR. MUHAMMAD DANI SUPARDAN, S.T., M.T.

Chemical Engineering Department, Syiah Kuala University, Banda Aceh 23211, Indonesia

MIRNA RAHMAH LUBIS, S.T., M.S.

Chemical Engineering Department, Syiah Kuala University, Banda Aceh 23211, Indonesia

WAHYU RINALDI, S.T., M.SC.

Chemical Engineering Department, Syiah Kuala University, Banda Aceh 23211, Indonesia

DR. SYAWALIAH, S.T.

Chemical Engineering Department, Syiah Kuala University, Banda Aceh 23211, Indonesia

DR. MUKRAMAH, S.T.

Chemical Engineering Department, Syiah Kuala University, Banda Aceh 23211, Indonesia

EDITORIAL BOARD

PROF. DR. ZUCHRA HELWANI

Chemical Engineering Department, Riau University, Pekanbaru, Indonesia

PROF. DR. IR. HUSNI HUSIN, MT.

Chemical Engineering Department, Syiah Kuala University, Banda Aceh 23211, Indonesia

DR. MUHAMMAD ROIL BILAD

Faculty of Integrated Technologies, Universiti Brunei Darussalam, Bandar Seri Begawan, Brunei Darussalam

ASSOCIATE PROF. SAEID RAJABZADEH

Research Center for Membrane and Film Technology, Kobe University, Japan



USER

Username

Password

☐ Remember Me

JOURNAL

Search

Search Scope

All

Browse

- By Issue
- By Author
- By Title
- Other Journals
- Categories

PUBLIC

AUTHOR

MANUSCRIPT

PUBLISHED

INDEXED



ASSOCIATE PROFESSOR DR. FAISAL ABNISA

Department of Chemical and Materials, Faculty of Engineering, King Abdulaziz University, Saudi Arabia





DR. DEWI AGUSTINA IRYANI

Chemical Engineering Department, Lampung University, Indonesia





ASSOC. PROF. DR. ERNESTO HERNANDEZ

School of Engineering, Technology and Design, Canterbury Christ Church (CCC) University, United Kingdom



ASSOC. PROF. JOHANN FELLNER

Institute for Water Quality, Resource and Waste Management, Vienna University of Technology, Austria



ASSOCIATE PROF. DR. AGUS SAPTORO

Department of Chemical Engineering, Curtin University Malaysia, Miri, Malaysia



ASSOC. PROF. HAMED KARKHANECHI

Department of Chemical Engineering, Ferdowsi University of Mashhad, Mashhad, Iran



DR. POERNOMO GUNAWAN

School of Chemical & Biomedical Engineering, Nanyang Technological, University Singapore, Singapore



DR. NIK ABDUL HADI MD NORDIN

Chemical Engineering Department, Universiti Teknologi PETRONAS, Seri Iskandar, Malaysia



PROF. T. M. INDRA MAHLIA

School of Information, Systems and Modelling, University of Technology Sydney, Australia



JURNAL REKAYASA KIMIA & LINGKUNGAN

Jurusan Teknik Kimia Universitas Syiah Kuala, Jl. Tgk. Syech Abdur Rauf No.7, Kopelma Darussalam, Banda Aceh, INDONESIA

PRINCIPAL CONTACT

Nasrul Arahman, Prof. Dr. S.T., M.T.

Phone: +62813-6092-7917

E-mail: rkl@che.unsyiah.ac.id, nasrular@unsyiah.ac.id

SUPPORT CONTACT

Mirna Rahmah Lubis

E-mail: mirna@che.unsyiah.ac.id

Wahyu Rinaldi, ST, M.Sc.

E-mail: wahyu.rinaldi@che.unsyiah.ac.id

00185009

VISITORS

Jurnal Rekayasa Kimia & Lingkungan

(Journal of Chemical Engineering & Environment)

Accredited By:
Director General of Higher Education, Research & Technology
No: 158/E/KPT/2021



ISSN 1412-5064
E-ISSN 2356-1024

HOME ABOUT LOGIN REGISTER CATEGORIES SEARCH CURRENT ARCHIVES ANNOUNCEMENTS EDITORIAL TEAM RE

Vol. 9 No. 4 Desember 2013

ISSN 1412-5064

Jurnal
**Rekayasa Kimia
&
Lingkungan**
Journal of Chemical Engineering and Environment




Jurnal Rekayasa Kimia & Lingkungan Vol. 9 No. 4 Desember 2013 ISSN 1412-5064



9 772356 166006

CURRENT ISSUE



VISITOR STATS

Viewers See more ▶

140,819	468	293
11,109	452	243
813	418	242
774	346	229

Pageviews: 389,697

FLAG counter

TOOLS



INFORMATION

- For Readers
- For Authors
- For Librarians

Home > Archives > Vol 17, No 1 (2022)

VOL 17, NO 1 (2022)

Jurnal Rekayasa Kimia dan Lingkungan (January-June, 2022)

TABLE OF CONTENTS

Articles

Effect of Acid Concentration on the Properties of Microcrystalline Cellulose from Pineapple Crown Leaf
Fitriani Fitriani, Sri Aprilia, Nasrul Arahman, Muhammad Roil Bilad
10.23955/rkl.v17i1.21010

Synthesis of Renewable Diesel from Pure Plant Oil of Kapuk (Ceiba pentandra) Seed Using NiMo/γ-Al₂O₃ Catalysts
Edwin Permana, Excel Emerlan Samiar Wasistha, Septina Is Heriyanti, Sri Djangkung Sumbogo Murti, Lenny Marlinda, Indra Lasmana Tarigan
10.23955/rkl.v17i1.21673

Distribution pattern of Pb and Zn contamination in rivers near industrial zone in Aceh, Indonesia, revealed by principal component analysis (PCA)
Lelifajri Lelifajri, Rahmadi Rahmadi, Rinaldi Idroes, Supriatno Supriatno, Eka Safitri, Nazaruddin, Nazaruddin, Ilham Maulana, Khairi Suhud
10.23955/rkl.v17i1.22672

Synthesis and characterization of nanoemulsion of citronella oil (Cymbopogon Nardus L. Rendle) and its Sunscreen Potential Evaluation
Undri Rastuti, Uyi Sulaeman, Senny Widyaningsih, Sity Khalidah Zia, Ryan Aditya Mahendra
10.23955/rkl.v17i1.23077

Adsorption Of Cr(VI) Ions Using Sugar Palm Fruit Peel (Arenga Pinnata) Immobilized in Ca-Alginate Bead
Intan Lestari, Widy Prasetyo, Indra Lasmana Tarigan, Edwin Permana
10.23955/rkl.v17i1.22873

Optimizing Gold Recovery of Artisanal Mining: A Lesson Learned from Kenya
Izzan Nur Aslam, Nestor Orcon, Bern Klein, Pocut Nurul Alam
10.23955/rkl.v17i1.23223

Ethanol Dehydration in Fixed Bed Column Using Pellet Adsorbent From Natural Zeolite Lampung
Jonathan Kristian Adisaputro, Simpamin Br. Ginting
10.23955/rkl.v17i1.24866



PDF
1-7

PDF
8-17

USER

Username
Password
☐ Remember Me

JOURNAL

Search

Search Scope

Browse

• By Issue
• By Author
• By Title
• Other Journals
• Categories

• PUBLIC
• AUTHENTIC
• MANUSCRIPT
• PUBLISHED

INDEXED IN



JURNAL REKAYASA KIMIA & LINGKUNGAN

Jurusan Teknik Kimia Universitas Syiah Kuala, Jl. Tgk. Syech Abdur Rauf No.7, Kopelma Darussalam, Banda Aceh, INDONESIA

PRINCIPAL CONTACT

Nasrul Arahman, Prof. Dr. S.T., M.T.
Phone: +62813-6092-7917
E-mail: rkl@che.unsyiah.ac.id, nasrular@unsyiah.ac.id

SUPPORT CONTACT

Mirna Rahmah Lubis
E-mail: mirna@che.unsyiah.ac.id

Wahyu Rinaldi, ST, M.Sc.
E-mail: wahyu.rinaldi@che.unsyiah.ac.id

00185009

VISITORS



SERTIFIKAT

Direktorat Jenderal Penguatan Riset dan Pengembangan,
Kementerian Riset, Teknologi, dan Pendidikan Tinggi



Kutipan dari Keputusan Direktur Jenderal Penguatan Riset dan Pengembangan,
Kementerian Riset, Teknologi, dan Pendidikan Tinggi Republik Indonesia
Nomor: 30/E/KPT/2018
Tentang Hasil Akreditasi Jurnal Ilmiah Periode 2 Tahun 2018

Jurnal Rekayasa Kimia dan Lingkungan (Journal of chemical engineering and environment)

E-ISSN: 23561661

Penerbit: Jurusan Teknik Kimia, Universitas Syiah Kuala, Banda Aceh

Ditetapkan sebagai Jurnal Ilmiah

TERAKREDITASI PERINGKAT 2

Akreditasi berlaku selama 5 (lima) tahun, yaitu
Volume 12 Nomor 1 Tahun 2017 sampai Volume 16 Nomor 1 Tahun 2021

Jakarta, 24 oktober 2018

Direktur Jenderal Penguatan Riset dan Pengembangan



Dr. Muhammad Dimiyati
NIP. 195912171984021001



← → ↺ sinta.kemdikbud.go.id/journals/detail?id=957



HOME

ABOUT

AUTHORS

SUBJECTS

AFFILIATIONS

SOURCES

REGISTRATION

FAQ

AUTHOR LOGIN



New! - Science And Technology Index (SINTA) Version 3.0

[Click Here](#)

Journal Profile



Jurnal Rekayasa Kimia dan Lingkungan (RKL)

eISSN : 23561661 | pISSN : 14125064

Science | Social | Engineering
Universitas Syiah Kuala



S2
Sinta Score

14
H-Index

1147
Citations

13
H5-Index

925
5 Year Citations



Synthesis, characterization, and sunscreen potential evaluation of citronella oil nanoemulsion (*Cymbopogon Nardus L. Rendle*)

Undri Rastuti^{1*}, Uyi Sulaeman¹, Senny Widyaningsih¹, Sity Khalidah Zia¹,
Ryan Aditya Mahendra¹

¹ Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Jenderal Soedirman.
Jl. dr. Soeparno Karangwangkal, Purwokerto 53123, Indonesia.

*E-mail: undrirastuti@yahoo.co.id

Article History

Received: 11 October 2021; Received in Revision: 30 December 2021; Accepted: 16 February 2022

Abstract

Citronella (*Cymbopogon nardus L. Rendle*) essential oil contains active compounds such as antioxidants and has the potential to be developed as a sunscreen. Nanoemulsion is a water and oil dispersion stabilized by a surfactant layer and has a particle size below 100 nm. This study described the formulation, characterization of citronella oil nanoemulsions, and determination of their activity as a sunscreen through the Sun Protection Factor (SPF) values. The nanoemulsion formulation was made with 3 variations of oil concentration: F1 (1%), F2 (3%), and F3 (5%). The results of the organoleptic test were following the standards which were clear and no phase separation occurred. In addition, the pH value of the 1%, 3%, and 5% prepared nanoemulsions were categorized as safe to be used on the skin. The prepared formulas were categorized as O/W nanoemulsion types. The viscosity values were found to be 12.76, 25.13, and 84.05 cP respectively. Furthermore, the percent transmittance values were 98.4, 98.9, and 99.9%. The particle sizes, respectively, were 9.4 (100%), 10.07 (97%), and 8.96 nm (82.5%). The centrifugation and freeze-thaw cycle tests showed good physical stability. The results of the sunscreen potential test of the 1%, 3%, and 5% citronella oil nanoemulsion showed SPF values of 1.03, 1.13, and 1.14 with %Te values of 77.42, 70.21, and 67.86% respectively, and %Tp values of 95.94, 94.04, and 93.61% respectively. From this research, the citronella oil nanoemulsion were found out to have a higher sunscreen activity than the pure citronella oil.

Keywords: Citronella essential oil (*Cymbopogon nardus L. Rendle*), nanoemulsion, antioxidant, sunscreen, SPF.

1. Introduction

Research on nanoemulsion has been developed over the last few decades (Koroleva and Yurtov, 2012). Nanoemulsion is a dispersion of water and oil stabilized by a surfactant and has a particle size below 100 nm (Paquin et al., 2015). The surfactants plays a major role to increase the benefit of preparing the active substances into nanoemulsion. The surfactant supports the mass transfer resistance of the oil release into continuous phase from the dispersed phase and also supports the particle size reduction leading to a higher activity of the active substances (Agrawal, et al., 2016). This topic of research was highly developed because nanoemulsion has various advantages over emulsion. Good physical stability, large surface area, and high bioavailability can be obtained in nanoemulsions. These advantages allow nanoemulsions to be applied in various fields such as pharmacology, food, cosmetics, etc. (Nirmala and Nagarajan, 2017).

The application of nanoemulsions could be applied in essential oils. The essential oils have secondary metabolites and can be obtained from plants (Seow et al., 2014). Essential oil has a high economic value because it has many pharmacological effects (Eden et al., 2018). Citronella essential oil is one of the various essential oils obtained in Indonesia. It can be produced by the distillation of fragrant citronella leaves (*Cymbopogon Nardus L. Rendle*). The Citronella essential oil has good pharmacological effects such as anticancer (Ong et al., 2006), antimicrobial (Wei and Wee, 2013), and antioxidants (Mileva et al., 2014). Citronella oil contains an antioxidant compound that could be utilized as a natural sunscreen. Sunscreen is a substance that can protect the skin from exposure to harmful UV rays (free radicals). The most important component in sunscreen is antioxidant compounds that can reduce free radicals (Gause and Chauhan, 2016).

Although citronella essential oil has been proven to have many pharmacological effects, it still has some drawbacks, such as low physical stability and poor water solubility. These limitations can be handled by modifying citronella oil into nanoemulsions (Nirmala and Nagarajan, 2017). From the research by Sakulu et al. (2009), it was proven that preparing citronella oil into nanoemulsion help to increase the repellent activity of the substance. From the research by Harni et al (2020), it also proven that preparing citronella oil into nanoemulsion was effective to increase the inhibition towards the pathogen *P. Palmivora*.

In this research, the synthesis and characterization of citronella oil nanoemulsion were carried out to evaluate its sunscreen potency compared to the pure citronella oil's sunscreen activity. The nano emulsification method used is the low-energy phase inversion method, Emulsion Inversion Point (EIP). In the EIP method, the W/O macroemulsion is prepared at a constant temperature and then slowly diluted with water. During the dilution process, the system passes through an inversion point where a W/O to O/W phase transformation occurs. At this inversion point, the oil-water interfacial tension is so low that it leads to the formation of a small droplet (Forgiarini et al., 2001).

2. Methodology

2.1. Materials

The materials used in this research were citronella oil (*C. nardus* L. Rendle), surfactant tween 80 (MERCK, Germany), propylene glycol, methylene blue (MERCK, Germany), distilled water, and ethanol (MERCK, Germany). The research started with the synthesis of nanoemulsion, followed by the characterization of nanoemulsion, and ended with the sunscreen evaluation of the prepared citronella oil nanoemulsion

2.2. Synthesis of Nanoemulsion

The synthesis of citronella oil nanoemulsion was carried out in three steps, following the formulation in Table 1. The first step was the preparation of the oil phase. The oil phase was a mixture of tween 80 as a surfactant, propylene glycol as a cosurfactant, and citronella essential oil. This oil phase was homogenized using a magnetic stirrer at a speed of 750 rpm at 50 °C for 30 minutes. The second step was mixing the oil phase and water phase (distilled water). The distilled water was added slowly into the oil phase. The

third step was the homogenization process of both the oil phase and water phase for 30 minutes at 50 °C with a speed of 1200 rpm using a magnetic stirrer.

Table 1. Citronella oil nanoemulsion formulation

Formula	Material (%)			
	CO	Tween 80	PG	Distilled water
F1	1	25	10	Add 100%
F2	3	25	10	
F3	5	25	10	

2.2. Characterization of citronella oil nanoemulsion

The characterization of citronella oil nanoemulsion was carried out by conducting an organoleptic test, a pH test (Seibert et al., 2019), a nanoemulsion type test, viscosity value measurement, transmittance percentage test, particle size measurement, centrifugation, and freeze-thaw cycle test (Gurpreet and Singh, 2018).

Organoleptic test was conducted to identify the characteristic of nanoemulsion by observing the color, odor, phases separation, and clarity of the prepared formula. The pH value was measured to identify if the nanoemulsion prepared is safe to be used on the skin. The nanoemulsion type was evaluated by adding a few drops of methylene blue into nanoemulsion to determine the dispersed and the dispersing phase (oil in water or water in oil). The viscosity of nanoemulsion was measured using ostwald viscometer with 3 times repetitions. Transmittance percentage test was conducted to measure the clarity level of the nanoemulsion as the early stage to determine the droplet size of nanoemulsion. The transmittance percentage test was conducted by using Spectrophotometer UV-Vis (λ 650 nm). The transmittance percentage test was followed by particle size measurement using Particle Size Analyzer to ensure the real droplets size formed in the nanoemulsion. The Centrifugation and freeze-thaw cycle test was conducted to determine the stability of the formula. The formula was centrifuged 10.000 rpm for 30 minutes and its organoleptic was observed. The formula was then stored in under the temperature of 4 °C for 24 hours, moved to 25 °C for 24 hours and then moved to 40 °C for another 24 hours for the freeze-thaw cycle test. The freeze-thaw cycle was repeated for 3 times.

2.3. Nanoemulsion Sunscreen Potency Test

a) Determination of SPF Value

A total of 0.2 mL of citronella oil and each formula were diluted with ethanol solvent until the volume reached 10 mL (20,000 ppm) (Widyawati et al., 2019). Each formula was measured for its SPF value using a UV-Vis Spectrophotometer (Shimadzu 1800) at a wavelength range of 290-400 nm at 5 nm intervals using ethanol as a blank. The SPF value was calculated by the equation, while the AUC value can be determined by the equation (1) (Mansur et al., 1986):

$$\text{Log SPF} = \frac{\sum AUC}{\lambda_n - \lambda_1} [AUC]_{\lambda_p - A}^{\lambda_p} = \frac{A(p-a) + A(p)}{2} \{\lambda(p) - \lambda(p-a)\} \quad (1)$$

- b) Determination of the Erythema transmission percentage (%Te) value

The percentage of erythema transmission was determined by measuring the transmittance of each diluted sample in the wavelength range of 292.5-317.5 nm. The value of %Te can be calculated by the equation (2) (Schalka et al., 2019):

$$\%Te = \frac{E_e}{\sum Fe} = \frac{\sum T.Fe}{\sum Fe} \quad (2)$$

- c) Determination of the Pigmentation transmission percentage (%Tp) value

The percentage of pigmentation transmission can be determined by measuring the transmittance of each sample that has been diluted in the wavelength range of 292.5-372.5 nm. The %Tp value can be calculated using the equation (3) (Schalka et al., 2019):

$$\%Tp = \frac{E_p}{\sum Fp} = \frac{\sum T.Fp}{\sum Fp} \quad (3)$$

3. Results and Discussion

3.1. Citronella Oil Nanoemulsion

The synthesis of citronella oil nanoemulsion was carried out using a low-energy method. The low-energy method is based on phase inversion in the emulsion upon changes in composition or temperature (Koroleva and Yurtov, 2012). The low energy method used in this study is the Emulsion Inversion Point (EIP) method or the composition phase reversal method. The principle of the EIP method is that the W/O macroemulsion is prepared at a constant temperature and then slowly diluted with water. During this dilution process, the system passes through an inversion point where a W/O to O/W phase transformation occurs. At this point of inversion, the oil-water interfacial tension is so

low that small droplet sizes can be formed (Forgiarini et al., 2001).

The nanoemulsion obtained by the EIP method was light yellow and clear. In the first synthesis step, the oil phase was made first so that the oil droplets are wrapped by a layer of surfactant and cosurfactant to form micelles. When the water phase was added in the second stage, the surface tension will decrease. This allows the emulsification process to occur spontaneously. In addition, the homogenization process using a stirrer allows the process of forming small oil droplet sizes (Forgiarini et al., 2001).

3.2. Organoleptic Results

The prepared nanoemulsions were obtained in slightly yellow to yellow, slightly fragrant to fragrant. The phases did not separate, and the clarity was very high (Table 2). There was an alteration of the scent characteristic between citronella oil nanoemulsions that were stored for 0 week and 4 weeks.

This means that the prepared citronella oil nanoemulsion was relatively less stable, especially in terms of aroma. Changes in the aroma are expected due to the merging of small droplets into larger droplets, causing a change in the density that likely decreases the aroma. In addition, the high clarity of the formulation was found to be unchanged. In this research, tween 80 was used as a surfactant. The use of surfactants provides high clarity and inhibits the phase separation that occurs in the emulsion (Costa et al., 2012).

Table 2. Organoleptic test results on citronella oil nanoemulsion preparation

Parameters	0 week			4 weeks		
	F1	F2	F3	F1	F2	F3
Color	LY	LY	LY	LY	Y	Y
Scent	LF	LF	F	F	SF	SF
Phase Separation	VI	VI	VI	VI	VI	VI
Clarity	VC	VC	VC	VC	VC	VC

Remarks:

LY : Less Yellow
 SF : Slightly Fragrant
 VC : Very clear
 VI : Very Inseparable
 F : Fragrant
 Y : Yellow

LF : Less Fragrant

3.3. pH

The pH results obtained from each nanoemulsion varied and decreased until the 3rd week (Table 3). In the citronella oil nanoemulsion F1 (1%), the pH value decreased from 6.5 to 6.3 at weeks 0 to 3. Citronella oil nanoemulsion F2 (3%) has a pH value of 6.3 at week 0 and 6.1 at week 3. Furthermore, the pH value of Citronella oil nanoemulsion F3 (5%) has changed from 6.2 at week 0 to 6.0 at week 3. The obtained pH data was then analyzed using two-way ANOVA. The concentrations (1, 3, and 5%) showed a significant value of 0.001 (p -value < 0.05), indicating that the concentration affected the pH obtained by the nanoemulsion.

Table 3. pH measurement results of citronella oil nanoemulsion preparation

Time (Week)	pH		
	F1	F2	F3
0	6.5	6.3	6.2
1	6.4	6.3	6.1
2	6.3	6.2	6.0

Time (weeks) showed a significant value, namely 4.55×10^{-5} (p -value < 0.05).

3.4. Nanoemulsion Type

The nanoemulsion type test was conducted by adding methylene blue to each prepared nanoemulsion. Figure 1 displayed that the methylene blue solution completely dissolved in 1%, 3%, and 5% citronella oil nanoemulsion. This result indicates that the type of nanoemulsion formed was oil in water (O/W) type. The O/W type is formed because the nanoemulsion system is made up of water so that methylene blue can dissolve completely (Gurpreet and Singh, 2018). The mechanism of dissolution of methylene blue in the water is solvation. Methylene blue ionizes in water into anions and cations, then the ions are surrounded by water molecules according to the induced dipole force. The type of oil in water (O/W) nanoemulsion is as expected because it is safe when used topically (Koroleva and Yurtov, 2012).



Figure 1. Result of examination of nanoemulsion type in each sample. (A) before dripping with methylene blue, (B) after dripping with methylene blue

3.5. Viscosity

Based on Table 4, it shows that the greater concentration of citronella oil in the prepared nanoemulsion will reduce the volume of water, thereby increasing their viscosity value. An increase in the amount of oil causes more dispersed droplets so that the friction between the droplets is higher and increases the viscosity. An increase in the amount of oil also causes an increase in the density of a particle. When the density of the particles is large, the mass will be heavier.

Table 4. The viscosity test results of citronella oil nanoemulsion preparation

No	Sample	Viscosity (cP)
1.	F1	12.76
2.	F2	25.13
3.	F2	84.05

The greater the particle density, the more the fluid flow is hampered. This increases the viscosity value. In addition, viscosity is influenced by the cohesive force between molecules, namely the force of the attraction between similar molecules. The increase in the amount of oil causes the cohesive force in the system to be large. This is what causes the viscosity to increase (Giancoli, 2009).

3.6. Transmittance Percentage

A good nanoemulsion formula will have clear visuals with transmittance $> 90\%$ (Costa et al., 2012). The highest transmittance value was found in 1% citronella oil nanoemulsion (Table 5) with more water composition. Therefore, it is expected to have a higher clarity level.

Table 5. %T measurement results of citronella oil nanoemulsion

No.	Sample	% T
1.	F1	99.9
2.	F2	98.9
3.	F3	98.4

3.7. Particle Size

Table 6 indicates that the three nanoemulsion formulations have been included in the size range below 100 nm (Gupta et al., 2016). The highest level of homogeneity was owned by 1% citronella oil nanoemulsion, followed by 3% and 5% citronella oil nanoemulsions. A higher oil concentration in the prepared nanoemulsions results in a less homogeneous formulation because it might need a longer time and faster stirring to homogenize the mixtures. Meanwhile, in this study, the homogenization time and stirrer speed were made the same for the three formulas.

F1	9.4 nm (100%)	0.0815
F2	10.07 nm (97%)	0.0122
	5530 nm (3%)	
F3	8.96 nm (82.5%)	2.5030
	216.8 nm (17.5%)	

The optimum formula was found in citronella oil nanoemulsion with a particle size of 9.4 nm. The particle size of less than 90 nm is known to increase the stability of nanoemulsion preparations against sedimentation (Gupta et al., 2016).

Table 6. Size distribution of citronella oil nano-emulsion

Formula	Particle Size	PDI
---------	---------------	-----




Figure 2. Nanoemulsion particle size distribution

The polydispersity index value serves as an indicator of a homogeneous particle size distribution. A low polydispersity index value indicates that the dispersion system is more stable in the long term (Gao et al., 2008). A polydispersity index of <0.5 indicates a fairly uniform particle size (monodisperse). The results obtained in this study indicate that 1% and 3% citronella oil nanoemulsions are included in the monodisperse category because they have a polydispersity index value of <0.5 . Meanwhile, 5% citronella oil nanoemulsion has a very high polydispersity index value of 2.503. This shows that 5% citronella oil nanoemulsion is still heterodisperse and unstable.

3.8. Centrifugation

The results of the centrifugation test (Table 7) showed that after the centrifugation test on each nanoemulsion preparation, there was no change in terms of turbidity, phase separation, and precipitation. The synthesized citronella oil nanoemulsion preparation has a good level of physical stability (Gurpreet and Singh, 2018).

Table 7. Centrifugation test results for citronella oil nanoemulsion

After centrifugation	Sample		
	F1	F2	F3
Turbidity	No	No	No
Phase separation	No	No	No
Precipitation	No	No	No

3.8. Freeze-thaw Cycle

The results of the freeze-thaw cycle test (Table 8) were observed by physical observation whether there was a change in terms of organoleptic, clarity, phase separation, and precipitation. The synthesized citronella oil nanoemulsion showed a good level of stability because there were no changes in terms of organoleptic, clarity, phase separation, and precipitation after the freeze-thaw cycle test. This indicates that the prepared nanoemulsions are stable to temperature changes. This stability is proposed due to the oil in water being protected by a surfactant layer that forms micelles. Micelle will prevent droplets from combining with the dispersing medium (Gurpreet & Singh, 2018).

Table 8. Freeze-thaw cycle test results for citronella oil nanoemulsion

Parameters	Before			After		
	F1	F2	F2	F1	F2	F3
Color	LY	LY	LY	LY	LY	LY
Clarity	C	C	C	C	C	C
Scent	DS	DS	DS	DS	DS	DS
Phases Separation	N	N	N	N	N	N
Precipitation	N	N	N	N	N	N

Remarks:

LY = Light Yellow

DS = Distinctive smell

N = None

C = Clear

3.9. Sunscreen Potential Evaluation

a. SPF value

The SPF value was tested by dissolving the citronella oil nanoemulsion into ethanol and measuring its absorbance using a UV-Vis spectrophotometer with a wavelength range of 290-400 nm. The measurement of the SPF value was also carried out on pure citronella oil dissolved in ethanol with the same concentration, namely 1%, 3%, and 5%, to compare the SPF value of the nanoemulsion of citronella oil and the SPF value of pure citronella oil. The results of the measurement of the SPF value can be seen in Figure 3.

Table 9. SPF, %Te, and %Tp values of citronella oil nanoemulsion and citronella oil

Sample	CO			CO Nanoemulsion		
	SPF	%Te	%Tp	SPF	%Te	%Tp
F1	1.01	93.79	98.41	1.03	77.42	95.94
F2	1.02	89.73	96.29	1.13	70.21	94.04
F3	1.03	83.58	94.84	1.14	67.86	93.61




Figure 3. SPF value measurement results of citronella oil (CO) and citronella oil nanoemulsion (CON)

Based on the results of the measurement of the SPF value, the SPF values of the citronella oil nanoemulsion with an oil concentration of 1%, 3%, and 5% were obtained to be 1.03, 1.13, and 1.14 respectively. The SPF value of pure citronella oil with the same concentration was 1.01, 1.02, and 1.03. The results showed that the SPF values of the prepared nanoemulsion were higher than the citronella oil. The SPF value increased along with the increase in the concentration of citronella oil so that the SPF value could be increased by adding the concentration of citronella oil into the nanoemulsion.

b. %Te value

The test of the erythema transmission value (%Te) was tested at a wavelength of 292.5–317.5 nm. The results of the measurement of the %Te value can be seen in Figure 3.




Figure 4. %Te values of citronella oil (CO) and citronella oil nanoemulsion (CON)

According to the tests, the %Te values of the citronella oil nanoemulsion formula with a concentration of 1, 3, and 5% were 77.42, 70.21, and 67.86% respectively. The %Te values of the pure citronella oil with the same oil concentration were obtained to be 93.79, 89.73, and 83.58%.

Based on Figure 4, it can be seen that with the same oil content, the %Te value of citronella oil in nanoemulsion form is always lower than that of pure citronella oil. This result indicates that the amount of UV light transmitted after hitting the pure citronella oil is higher than the amount of UV light transmitted after hitting the nanoemulsion. Based on the measurement of the %Te value, it is also proven that by making citronella oil in the nanoemulsion form, it will increase the effectiveness of sun protection.

c. %Tp value determination

The pigmentation transmission percentage (%Tp) assay was carried out at a wavelength of 322–317.5 nm. The measurement of the %Tp values is displayed in Figure 4.

The %Tp values obtained from 1, 3, and 5% citronella oil nanoemulsion were 95.94, 94.04, and 93.61% respectively. The %Tp values of the pure citronella oil with the same concentration were calculated to be 98.41, 96.29, and 94.84%. **Figure 5** indicates that with the same oil content, the %Tp values of citronella oil in nanoemulsion are always lower than that of pure citronella oil. Therefore, based on the measurement of the %Tp value, it is also confirmed that citronella oil in the form of nanoemulsion will increase the effectiveness of sun protection.




Figure 5. %Tp values of citronella oil (CO) and citronella oil nanoemulsion (CON)

In general, based on the measurement results of SPF, %Te, and %Tp values, citronella oil nanoemulsions with an oil concentration of 1, 3, and 5% do not show good sunscreen activity. However, their sunscreen properties are better, with higher SPF, %Te, and %Tp values, than the pure citronella oil with the same oil concentration. This finding is because the nanoemulsion has a smaller globule size that creates the wider UV absorption area.

4. Conclusion

Citronella oil nanoemulsions were formulated into 3 formulations with different concentrations, namely 1%, 3%, and 5%. The ratio of tween 80 and propylene glycol was 25:10. The characteristics of the prepared citronella oil nanoemulsion have a distinctive organoleptic visualization, an acceptable pH value to the skin (6.0-6.5) with nanoemulsions type of O/W. The nanoemulsion viscosity for the formulations of 1%, 3%, and 5% were 12.76, 25.13, and 84.05 cP, while the percent transmittance of the nanoemulsion with the respective concentrations were 98.4, 98.9, and 99.9%. The particle size distribution of 1%, 3%, and 5% nanoemulsions were 9.4 nm (100%), 10.07 nm (97%), and 8.96 nm (82.5%) respectively. Furthermore, the citronella oil nanoemulsions showed good physical stability when tested by centrifugation and freeze-thaw cycle tests. Analysis of the potential as a sunscreen of the 1%, 3%, and 5% citronella oil nanoemulsion showed the SPF values of 1.03, 1.13, and 1.14 respectively, the %Te value of 77.42, 70.21, and 67.86%, while the %Tp value of 95.94, 94.04, and 93.61%. The prepared nanoemulsions was proven to have the higher sunscreen activity compared to the pure citronella oil's sunscreen activity with the same concentration. The more effective ratio of the oil, surfactant, and water could be explored in the next research to get formula with the best stability and activity.

Acknowledgment

This research was supported by Research and Community Service (LPPM) of Universitas Jenderal Soedirman through 'Riset Peningkatan Kompetensi'

References

- Agrawal, N., Ganesh, L., Maddikeri, M., Aniruddha, B., Pandit, P. 2016. Sustained release for formulation of citronella oil nanoemulsion using cavitation techniques. *Ultrasonics Sonochemistry*

- 36, 367-374. <http://doi.org/10.1016/j.ultsonch.2016.11.037>
- Costa, J.A., Lucas, E.F., Queirós, Y.G.C., Mansur, C.R.E., 2012. Evaluation of nanoemulsions in the cleaning of polymeric resins. *Colloids Surfaces A Physicochem. Eng. Asp.* 415, 112–118. <https://doi.org/10.1016/j.colsurfa.2012.10.011>
- Eden, W.T., Alighiri, D., Cahyono, E., Supardi, K.I., Wijayati, N., 2018. Fractionation of Java Citronella Oil and Citronellal Purification by Batch Vacuum Fractional Distillation, in: *IOP Conference Series: Materials Science and Engineering*. p. 394. <https://doi.org/10.1088/1757-899X/349/1/012067>
- Forgiarini, A., Esquena, J., González, C., Solans, C., 2001. Formation of nano-emulsions by low-energy emulsification methods at constant temperature. *Langmuir* 17, 2076–2083. <https://doi.org/10.1021/la001362n>
- Gao, L., Zhang, D., Chen, M., 2008. Drug nanocrystals for the formulation of poorly soluble drugs and its application as a potential drug delivery system. *J. Nanoparticle Res.* 10, 845–862. <https://doi.org/10.1007/s1105100893574>
- Gause, S., Chauhan, A., 2016. UV-blocking potential of oils and juices. *Int. J. Cosmet. Sci.* 38, 354–363. <https://doi.org/10.1111/ics.12296>
- Giancoli, D.C., 2009. *Physics for Scientists & Engineers with Modern Physics*, in: *Physics for Scientists & Engineers with Modern Physics*. Pearson Prentice Hall.
- Gupta, A., Eral, H.B., Hatton, T.A., Doyle, P.S., 2016. Controlling and predicting droplet size of nanoemulsions: Scaling relations with experimental validation. *Soft Matter* 12, 1425–1458. <https://doi.org/10.1039/c5sm02051d>
- Gurpreet, K., Singh, S.K., 2018. Review of nanoemulsion formulation and characterization techniques. *Indian J. Pharm. Sci.* 80, 781–789.
- Harni, R., Kherati, K., Noveriza, R., Yuliani, S., 2020. Effectiveness of citronella oil nanoemulsion on *Phytophthora palmivora* Butl. black pod pathogen. *Journal of Industrial and Beverage Crops*. 7(3), 127-135.
- Koroleva, M.Y., Yurtov, E. V., 2012. Nanoemulsions: the properties, methods of preparation and promising applications. *Russ. Chem. Rev.* 81, 21–43. <https://doi.org/10.1070/rc2012v08-1n01abeh004219>
- Mansur, J. de S., Breder, M.N.R., Mansur, M.C. d'Ascensão, Azulay, R.D., 1986. Determination of SUN Protection Factor by Spectrophotometry. *An. bras. dermatol* 61, 121–124.
- Mileva, M., Kusovaski, V., Krastev, D., Dobрева, A., Galabov, A., 2014. Chemical composition, in vitro antiradical and antimicrobial activities of Bulgarian Rosa alba L. essential oil against some oral pathogens. *Int. J. Curr. Microbiol. App. Sci.* 3, 11–20.
- Nirmala, M., Nagarajan, R., 2017. Recent Research Trends in Fabrication and Applications of Plant Essential Oil Based Nanoemulsions. *J. Nanomed. Nanotechnol.* 8, 434–444. <https://doi.org/10.4172/2157-7439.1000434>
- Ong, T.P., Heidor, R., de Conti, A., Dagli, M.L.Z., Moreno, F.S., 2006. Farnesol and geraniol chemopreventive activities during the initial phases of hepatocarcinogenesis involve similar actions on cell proliferation and DNA damage, but distinct actions on apoptosis, plasma cholesterol and HMGCoA reductase. *Carcinogenesis* 27, 1194–1203. <https://doi.org/10.1093/carcin/bgi291>
- Paquin, F., Rivnay, J., Salleo, A., Stingelin, N., Silva, C., 2015. Multi-phase semicrystalline microstructures drive exciton dissociation in neat plastic semiconductors. *J. Mater. Chem. C* 3, 10715–10722. <https://doi.org/10.1039/b000000x>
- Sakulku, U., Nuchuchua, O., Uawongyart, N., Puttipipatkachorn, S., Sootitawant, A., Ruktanonchai, U., 2009. Characterization and mosquito repellent activity of citronella oil nanoemulsion. *National Journal of Pharmaceutichs.* 372, 105-111.
- Schalka, S., Corrêa, M. de P., Sawada, L.Y.,

- Canale, C.C., de Andrade, T.N., 2019. A novel method for evaluating sun visible light protection factor and pigmentation protection factor of sunscreens. *Clin. Cosmet. Investig. Dermatol.* 12, 605–616. <https://doi.org/10.2147/CCID.S207256>
- Seibert, J.B., Rodrigues, I.V., Carneiro, S.P., Amparo, T.R., Lanza, J.S., Frézard, F.J.G., de Souza, G.H.B., Santos, O.D.H. dos, 2019. Seasonality study of essential oil from leaves of *Cymbopogon densiflorus* and nanoemulsion development with antioxidant activity. *Flavour Fragr. J.* 34, 5–14. <https://doi.org/10.1002/ffj.3472>
- Seow, Y.X., Yeo, C.R., Chung, H.L., Yuk, H.G., 2014. Plant Essential Oils as Active Antimicrobial Agents. *Crit. Rev. Food Sci. Nutr.* 54, 625–644. <https://doi.org/10.1080/10408398.2011.599504>
- Wei, L.S., Wee, W., 2013. Chemical composition and antimicrobial activity of *Cymbopogon nardus* citronella essential oil against systemic bacteria of aquatic animals. *Iran. J. Microbiol.* 5, 147–152.