



Determinant Factors Stroke Prevention Behavior among Hypertension Patient in Indonesia

Arif Setyo Upoyo*, Atyanti Isworo, Yunita Sari, Agis Taufik, Annas Sumeru, Akhyarul Anam

Department of Nursing, Faculty of Health Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia

Abstract

Edited by: Sasho Stoleski
Citation: Upoyo AS, Isworo A, Sari Y, Taufik A, Sumeru A, Anam A. Determinant Factors Stroke Prevention Behavior among Hypertension Patient in Indonesia. Open Access Maced J Med Sci. 2021 May 07; 9(E):336-339. <https://doi.org/10.3889/oamjms.2021.6040>
Keywords: Behavior; Stroke prevention; Hypertension; Self-awareness
***Correspondence:** Arif Setyo Upoyo, Department of Nursing, Faculty of Health Sciences, Universitas Jenderal Soedirman. E-mail: afkarfadhoh@gmail.com
Received: 17-Mar-2021
Revised: 24-Apr-2021
Accepted: 28-Apr-2021
Copyright: © 2021 Arif Setyo Upoyo, Atyanti Isworo, Yunita Sari, Agis Taufik, Annas Sumeru, Akhyarul Anam
Funding: This study was supported by the Research and Community Service Institute of Universitas Jenderal Soedirman, Indonesia
Competing Interest: The authors have declared that no competing interest exists
Open Access: This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

Stroke is the leading cause of death in the world and in Indonesia. Hypertension is a major risk of stroke. Modified stroke risk factors are behavioral factors. The purpose of this study is to identify the factors that influence stroke prevention behavior in hypertensive patients. The research method uses cross sectional. Total participants were 461 hypertensive patients who came to Community Health Center. Sampling technique used consecutive sampling. The questionnaire used contains demographics, knowledge, self-efficacy, awareness of stroke risk, and stroke prevention behaviors which include diet, exercise, smoking, and drinking alcohol. Data analysis used Chi-square and logistic regression. The results showed that 65.7% of respondents showed poor stroke prevention behavior. Factors related to stroke prevention behavior include self-efficacy ($p = 0.043$; 95%CI = 1.01–2.22; OR = 1.5), perception ($p = 0.038$; 95%CI = 0.45–0.98; OR = 0.663), and awareness of stroke risk ($p \leq 0.001$; 95%CI = 2.255–6.333; OR = 3.779). The dominant factor influencing stroke prevention behavior in hypertensive patients is awareness of stroke risk.

Introduction

Stroke is a major health-care problem in South, East, and Southeast Asia. With a majority of the world's population living in the developing countries of these regions, the global burden of stroke will have the largest contribution from Asia [1]. The prevalence of stroke in Indonesia increases every year. Based on the 2013 Basic Health Research, the prevalence of stroke was 7 per thousand population, while in 2018, the prevalence of stroke was 10.9 per thousand population [2].

Risk factors for stroke include factors that cannot be modified and factors that can be modified. Factors that cannot be modified include age, sex, race/ethnicity, and genetics, while factors that can be modified include hypertension, diabetes, atrial fibrillation, dyslipidemia, diet, physical activity, obesity, metabolic syndrome, alcohol consumption, and smoking [3], [4], [5]. Risk factors for stroke that can be modified are influenced by patient behavior.

Hypertension is one of the main factors of stroke. Primary hypertension is a major cause of cardiovascular disease morbidity which has a significant influence on the incidence of stroke [6], [7]. Goldstein states that the prevalence of stroke in hypertensive patients aged 50 years is 20% with a risk ratio of 4 and the prevalence

continues to increase with increasing age [8], whereas according to Ghani *et al.*, hypertension sufferers have a risk of 2.87 times getting a stroke [7].

The purpose of this study was to determine the factors that influence stroke prevention behavior in hypertensive patients.

Methodology

Design

The research method used cross-sectional study. The research was conducted in April–August 2019.

Sample

The subjects of the study were 461 participants of hypertension. Sampling technique used consecutive sampling from five public health centers in Banyumas Regency, Central of Java. Inclusion criteria of sample were hypertensive patient, >35 years old and visit to public health center. Exclusion criteria were hypertensive patients with complications and refused as a respondent.

Instrument

The questionnaire used contained demographics, knowledge, self-efficacy, awareness of stroke risk, and stroke prevention behaviors. Knowledge instrument modified from Hypertension Knowledge-Level Scale by Erkoc *et al.* [9] that divided into three categories: Low, middle, and high. Self-efficacy and awareness of stroke risk used numeric score 1–10, high score if above average, and low score if below or equal to average. Stroke prevention behaviors which included low salt and cholesterol diet, exercise, smoking, and drinking alcohol used Likert score 1–4, total score categorized good if the total score is more than average and bad if the total score is below or equal to average.

Data analysis

Data analysis used Chi-square for bivariate analysis and logistic regression for multivariate analysis with SPSS 16.

Results

Based on Table 1, it is known that the respondents are mostly female (80.9%), over 60 years old (56.8%), still married (71.4%), living together with family (49.9%), with income below the regional minimum wage (65.7%), and low education (70.9%). Respondent characteristics that significantly influence behavior include gender ($p=0.000$; 95%CI=7.12–22.28; OR=12.59), age ($p=0.000$; 95%CI = 0.24–0.56; OR = 0.369), marital status ($p=0.008$; 95%CI = 1.17–2.89; OR = 1.83), and income ($p=0.008$; 95%CI = 0.39–0.87; OR = 0.58). Female, old age, married, and low income have a tendency to behave badly in stroke prevention behavior.

Based on Table 2, it is known that the majority showed poor stroke prevention behavior (65.7%), high

knowledge of hypertension (89.4%), low self-efficacy (56.2%), good perception (57.3%), information about low stroke (41.4%), and low stroke risk awareness (60.1%).

Factors related to stroke prevention behavior include self-efficacy ($p=0.043$; 95%CI = 1.01–2.22; OR = 1.50), perception ($p=0.038$; 95%CI = 0.45–0.98; OR = 0.663), and awareness of stroke risk ($p=0.000$; 95%CI = 0.24–0.53; OR = 0.354).

The results of multivariate analysis showed that factors related to stroke prevention behavior were age, sex, efficacy, and awareness of stroke risk. The dominant factor influencing behavior is awareness of stroke risk with OR 3779 (Table 3).

Discussion

Most respondents are female, the elderly, have low education, and earn below the regional minimum wage standard. The risk of hypertension increases with age. This is related to vascular changes that cause an increased peripheral resistance. Postmenopausal women increase the risk of developing hypertension due to hormonal changes. Low education and income can affect health behavior.

The result shows that the majority (65.7%) of hypertension patients had a poor stroke prevention behavior. The poor behavior is high consumption of sodium and fat and also lack of regular exercise. The factors that influence stroke prevention behavior in this study include perception, self-efficacy, and stroke risk awareness.

The majority of the respondents have poor perception and low self-efficacy in stroke prevention. Perception and self-efficacy can influence behavior. Imprecise perception in stroke prevention and low self-efficacy will inhibit stroke prevention behavior [10].

One of the factors that influence perception is information. In this study, we obtained bad perception

Table 1: Characteristics of respondents

Characteristics of respondents	n	%	OR	95%CI	p
Sex					
Male	88	19.1	12.59	7.12–22.28	<0.001
Female	373	80.9			
Age					
<60 years old	199	43.2	0.369	0.24–0.56	<0.001
>60 years old	262	56.8			
Marital status					
Married	329	71.4	1.83	1.17–2.89	0.008
Not yet	132	28.6			
Living together					
Living alone	12	2.6	1.27	0.86–1.87	0.227
With couple	230	49.9			
With family	201	43.6			
Other	18	3.9			
Income					
Under minimum standard payment	303	65.7	0.58	0.39–0.87	0.008
Standard payment	24	5.2			
Upper minimum standard payment	134	29.1			
Level education					
No	40	8.7	0.807	0.39–1.63	0.551
Elementary	327	70.9			
Junior high school	57	12.4			
Senior high school	34	7.4			
High education	3	0.7			

Table 2: Stroke prevention behavior and associated factors

Variable	n	%	OR	95% CI	p
Stroke prevention behavior					
Good	158	34.3	-	-	-
Bad	303	65.7			
Knowledge					
Low	0	0	0.830	0.44–1.58	0.568
Middle	49	10.6			
High	412	89.4			
Self-efficacy					
Low	259	56.2	1.50	1.01–2.22	0.043
High	202	43.8			
Perception					
Bad	264	57.3	0.663	0.45–0.98	0.038
Good	197	42.7			
Information about stroke					
Low	191	41.4	1.19	0.811–1.766	0.366
Middle	126	27.3			
Good	144	31.2			
Stroke risk awareness					
Low	277	60.1	0.354	0.24–0.53	<0.001
High	184	39.9			

Table 3: Multivariate analysis results (n = 461)

Variable	SE	Sig.	OR	CI 95%	
				Min.	Max.
Sex	0.339	<0.001	0.072	0.037	0.139
Age	0.261	<0.001	2.598	1.557	4.335
Marital status	0.327	0.218	0.669	0.352	1.269
Income	0.260	0.832	0.946	0.569	1.574
Stroke risk awareness	0.263	<0.001	3.779	2.255	6.333
Perception	0.253	0.272	1.321	0.804	2.169
Self-efficacy	0.261	<0.001	0.370	0.222	0.617
Living together	0.278	0.893	1.038	0.602	1.792

and information about stroke (41.4%). Plenty of information can affect or increase someone's knowledge and with knowledge creates awareness that eventually someone will behave in accordance with the knowledge they have.

Self-efficacy is an important precondition for successful self-management. Many studies have shown that self-efficacy is the strongest determinant of smoking cessation, increasing physical activity, and healthy diet [11], [12], [13]. Thus, increasing self-efficacy is an effective method to support healthy behavioral change in patients with stroke.

Respondents also indicated that the majority has a low stroke risk awareness. Many patients are not aware that hypertension is a risk of stroke so they ignore stroke prevention behavior. This is in line with the research of Hertz *et al.* which stated that 7.6% had the chance to become stroke, 46.4% had no chance to become a stroke, and 46.0 were unsure [14]. These findings indicate that self-awareness about stroke risk is poor. This disease attitude can inhibit efforts to raise awareness about stroke in society.

Most people think that stroke is more serious than heart attack. However, because stroke symptoms appear in variety of ways, they are not easily recognized. People with stroke report that they are confused by the symptoms such as migraine-like, headache, ear disease, or food poisoning. This kind of confusion can be one of the causes of delays to the hospital. Studies suggest that myocardial infarction patients with atypical symptoms lead to increased delay in many patients [15], [16]. Dracup *et al.* found the people who thought that they were simply having gas discomfort or indigestion significantly delayed seeing a doctor [17]. This study also revealed that people who had a low stroke self-awareness had a 3.78 times risk of having poor stroke prevention behavior. Therefore, increasing public awareness about the risk of stroke is an important factor in the stroke risk patients management.

Conclusion

Most respondents stroke prevention behaviors are bad. Respondent characteristics that significantly

influence behavior include gender, age, marital status, and income. Factors related to stroke prevention behavior include self-efficacy, perception, and awareness of stroke risk. The dominant factor influencing behavior is awareness of stroke risk. It is necessary to intervene to increase stroke prevention behavior in risk groups by increasing self-efficacy, perception, and awareness of stroke risk.

Ethical Aspect and Declaration of Interest Statement

The study has been declared ethical by the Dr. Moewardi Hospital's health research ethics commission number: 789/VI/HREC/2019. Informed consent was given to the participants before study.

Authors' Contributions

Concept and study design (Upoyo, Sari, Isworo), data collection, data analysis and interpretations (Taufik, Sumeru, Anam), processing the draft of the manuscript (Upoyo, Isworo), critical revision of the manuscript (Sari, Upoyo, Isworo), and article finalization (Upoyo, Isworo).

References

1. Venketasubramanian N, Yoon BW, Pandian J, Navarrod JC. Stroke epidemiology in South, East, and South-East Asia: A review. *J Stroke*. 2017;19(3):286-94. <https://doi.org/10.5853/jos.2017.00234> PMID:29037005
2. Ministry of Health of the Republic of Indonesia. The Reports of the Results of Basic Health Research; 2019. Available from: <https://www.litbang.kemkes.go.id/laporan-riset-kesehatan-dasar-risikesdas>. [Last accessed on 2020 Mar 17].
3. Slark J, Sharma P. Risk awareness in secondary stroke prevention: A review of the literature. *JRSM Cardiovasc Dis*. 2014;3:1-6. DOI: 10.1177/2048004013514737. PMID:24570838
4. Boehme AK, Esenwa C, Elkind MS. Stroke risk factors, genetics, and prevention. *Circ Res*. 2017;120(3):472-95. <https://doi.org/10.1161/circresaha.116.308398> PMID:28154098
5. Upoyo AS, Setyopranoto I, Pangastuti HS. The modifiable risk factors of uncontrolled hypertension in stroke: A systematic review and meta-analysis. *Stroke Res Treat*. 2021;2021:6683256. <https://doi.org/10.1155/2021/6683256> PMID:33680423
6. Hanchaiphiboolkul S, Pongvarin N, Nidhinandana S,

- Suwanwela NC, Puthkhao P, Towanabut S, *et al.* Prevalence of stroke and stroke risk factors in thailand: Thai epidemiologic stroke (TES) study. *J Med Assoc Thai.* 2011;94(4):427-36. <https://doi.org/10.1016/j.jstrokecerebrovasdis.2012.05.013> PMID:21591527
7. Ghani L, Mihardja LK, Delima D. Dominant risk factors of stroke in Indonesia. *Buletin Penelitian Kesehatan.* 2016;44(1):49-58. <https://doi.org/10.22435/bpk.v44i1.4949.49-58>
 8. Goldstein LB, Adams R, Albert MJ, Appel LJ, Brass LM, Bushnell CD, *et al.* Primary prevention of ischemic stroke. *Stroke.* 2006;37:1583-633.
 9. Erkoc SB, Isikli B, Metintas S, Kalyoncu C. Hypertension knowledge-level scale (HK-LS): A study on development, validity and reliability. *Int J Environ Res Public Health.* 2012;9(3):1018-29. <https://doi.org/10.3390/ijerph9031018> PMID:22690180
 10. Michie S, Atkins L, West R. *The Behaviour Change Wheel: A Guide to Designing Interventions.* London: Silverback Publishing; 2014.
 11. Garcia K, Mann T. From “i wish” to “i will”: Social-cognitive predictors of behavioral intentions. *J Health Psychol.* 2003;8(3):347-60. PMID:14670213
 12. Sniehotta FF, Scholz U, Schwarzer R. Bridging the intention-behaviour gap: Planning, self-efficacy, and action control in the adoption and maintenance of physical exercise. *Psychol Health.* 2005;20(2):143-60. <https://doi.org/10.1080/08870440512331317670>
 13. Sol BG, van der Bijl JJ, Banga JD, Visseren FL. Vascular risk management through nurse-led self-management programs. *J Vasc Nurs.* 2005;23(1):20-4. <https://doi.org/10.1016/j.jvn.2004.12.003> PMID:15741961
 14. Hertz JT, Madut DB, William G, Maro VP, Crump JA, Rubach MP. Perceptions of stroke and associated health-care-seeking behavior in Northern Tanzania: A community-based study. *Neuroepidemiology.* 2019;53(1-2):41-7. <https://doi.org/10.1159/000499069> PMID:30986785
 15. Moser DK, Dracup K. Gender differences in treatment-seeking delay in acute myocardial infarction. *Prog Cardiovasc Nurs.* 1993;8(1):6-12. PMID:8372090
 16. Schwarz B, Schoberberger R, Rieder A, Kunze M. Factors delaying treatment of acute myocardial infarction. *Eur Heart J.* 1994;15(12):1595-8. <https://doi.org/10.1093/oxfordjournals.eurheartj.a060439> PMID:7698126
 17. Dracup K, McKinley SM, Moser DK. Australian patients' delay in response to heart attack symptoms. *Med J Aust.* 1997;166(5):233-6. <https://doi.org/10.5694/j.1326-5377.1997.tb140101.x> PMID:9076265

Open Access Macedonian Journal of Medical Sciences

About ▾ Announcements Metrics ▾ Current ▾ Archives EU ▾ Scopus ▾ NCBI ▾ CABI ▾ Indexation ▾

🔍 Search

Membership ▾

Reduced Publication Fee for Ukraine

📅 2022-04-22



New Category "G - Nursing" was Opened

📅 2021-05-11

Open Access Macedonian Journal of Medical Sciences today migrated on the new URL and new OJS version!

📅 2021-03-08

Open Access Macedonian Journal of Medical Sciences, volume 8, 2020 was completed!

📅 2021-02-27

T1 - Thematic Issue "Coronavirus Disease (COVID-19)" Was Completed and Published in Open Access Macedonian Journal of Medical Sciences!

📅 2021-02-26

Information

For Readers

For Authors

For Librarians

Open Journal Systems

Make a Submission

Browse

Categories

Activate Windows

Go to Settings to activate Windows.

[Home](#) / [Editorial Team](#)

Editorial Team

Journal Manager

Eli Djulejic, MD. Open Access Macedonian Journal of Medical Sciences, Belgrade, Serbia
elispiroska@yahoo.com
 ORCID iD: <https://orcid.org/0000-0002-1736-9029>
 Web of Science ResearcherID: [AAC-3225-2020](#)
 Scopus Author ID: [33767574400](#)

Editor-in-Chief

Mirko Zhivko Spiroski, MD, PhD. Scientific Foundation SPIROSKI, Rajko Zhinzifov No 48,
 1000 Skopje, Republic of Macedonia
mspiroski@yahoo.com
 ORCID iD: <https://orcid.org/0000-0001-5559-7981>
 Web of Science ResearcherID: [F-7606-2012](#)
 Scopus Author ID: [6602348069](#)

Section Editors (Deputy Editors-in-Chief)

Stoleski Sasho, MD, PhD. Institute for Occupational Health of Republic of Macedonia -
 Skopje, WHO Collaborating Center, GA2LEN Collaborating Center, II Makedonska brigada
 43, 1000 Skopje, Republic of Macedonia
ssstoleski@yahoo.com
 ORCID iD: <https://orcid.org/0000-0003-1278-903X>
 Web of Science ResearcherID: [B-2304-2015](#)
 Scopus Author ID: [9943962300](#)

Slavica Hristomanova-Mitkovska, MD, MSc. Institute of Human Genetics, University
 Medical Center Göttingen, Heinrich-Düker-Weg 12, 37079 Göttingen, Germany
cacka_h@yahoo.com
 ORCID iD: <https://orcid.org/0000-0002-7770-5055>
 Web of Science ResearcherID: [G-8801-2012](#)
 Scopus Author ID: [33767823700](#)

Igor Spiroski, MD. University Clinic of Cardiology, Faculty of Medicine, Ss Cyril and
 Methodius University of Skopje, Skopje, Republic of Macedonia
ispiroski@gmail.com
 ORCID iD: <https://orcid.org/0000-0002-3448-0685>
 Web of Science ResearcherID: [AAA-5652-2020](#)
 Scopus Author ID: [23971606200](#)

Ksenija Bogoeva-Kostovska, MD, PhD. PHO Prof Bogoev, Skopje, Republic of Macedonia
ksenijabogoeva@gmail.com

Sinisha Stojanovski, MD, PhD. Medical Faculty, Saints Cyril and Methodius University in
 Skopje,
 Skopje, Republic of Macedonia
sinisa.stojanovski@hotmail.com
 ORCID iD: <https://orcid.org/0000-0002-4967-2102>
 Web of Science ResearcherID: [AFL-9514-2022](#)
 Scopus Author ID: [6504203064](#)

Filip Koneski, Doctor of Dental Medicine, MSc. University Clinic for Maxillofacial Surgery in
 Skopje, Ss. Cyril and Methodius University in Skopje, Faculty of Dental Medicine, Skopje,
 Republic of Macedonia
dr.koneski@gmail.com
 ORCID iD: <https://orcid.org/0000-0003-2412-7594>
 Web of Science ResearcherID: [J-9194-2019](#)
 Scopus Author ID: [57073856400](#)

Aleksandar Iliev, Doctor of Dental Medicine, PhD. Department of Maxillofacial Surgery,
 Faculty of Dentistry, The Saints Cyril and Methodius University of Skopje, Skopje, Republic
 of Macedonia
aleksandar.iliev@gmail.com
 ORCID iD: <https://orcid.org/0000-0001-5348-9143>
 Web of Science ResearcherID: [ACV-7553-2022](#)
 Scopus Author ID: [57021392000](#)

Information

[For Readers](#)
[For Authors](#)
[For Librarians](#)

Open Journal Systems

[Make a Submission](#)

Browse

Categories

A - Basic Sciences
 B - Clinical Sciences
 C - Case Reports
 D - Dental Sciences
 E - Public Health
 F - Review Articles
 G - Nursing
 T - Thematic Issues
 T1 - "Coronavirus Disease (COVID-19)"
 T2 - "Public Health and Nutrition Sciences in the Current Millennial Era"
 T3 - "Neuroscience, Neurology, Psychiatry and General Medicine"
 T4 - "Contribution of Nurses on Sustainable Development Goals (SDGs)"
 T5 - "Re-Advancing Nursing Practice, Education and Research in the Post Covid"
 T6 - "The Challenges and Opportunities for Nurses in The New Era Adaptation"
 T7 - "Neuroscience, Neurology, Psychiatry and Other Health Sciences - 2022"
 T8 - "APHNI: Health Improvement Strategies Post Pandemic Covid-19"

PKP Publishing Services

Part of the
 PKP Publishing Services Network

Layout Editor and Electronic Publishing

Ivo Spiroski, Eng, MSc. Scientific Foundation SPIROSKI, Rajko Zhinzifov No 48, 1000 Skopje, Republic of Macedonia
ivospiroski@gmail.com
[ORCID iD: https://orcid.org/0000-0003-4518-2138](https://orcid.org/0000-0003-4518-2138)
[Web of Science ResearcherID: F-2799-2014](#)

Prtoduction Editor

Copyeditor

Sinjore, India

Evidence Based Medicine

Prof. Dr. Katarina Stavric, Children Hospital Skopje, Macedonia, Vodnjanska 17, University "Ss Cyril and Methodius", Skopje, Republic of Macedonia

Editorial Board

DDS, MS, PhD, Associate Professor Nikola Angelov, Director of the Pre-Doctoral Periodontics Clinic, Loma Linda University School of Dentistry, Department of Periodontics. Loma Linda, CA, 92350, United States

Assist. Prof. Dr. Ramush Bejqi, University Clinical Centre of Kosovo, Paediatric Clinic, Albania

Prof. Semra Ćevaljuga, Department of Epidemiology and Biostatistics, Faculty of Medicine, Sarajevo, Bosnia and Herzegovina

MD Pei-Yi Chu, Diagnostic and research pathologist, Department of Surgical Pathology, Changhua Christian Hospital, Taiwan. Address: 135 Nan-Shiao Street, Changhua 500-06,, Taiwan, Province of China

MD, PhD Ivo Donkov, Staff Urologist, Lincoln County Hospital, United Kingdom

MD, PhD Andrew J. Dwork, Departments of Pathology and Cell Biology and Psychiatry, College of Physicians and Surgeons of Columbia University; Division of Molecular Imaging and Neuropathology, New York State Psychiatric Institute, Unit 62, 722 West 168th Street, New York, NY 10032, United States

Adriana Galan, Department of Health Programmes and Health Promotion, Institute of Public Health, Bucharest, Romania

Prof. Tania Santos Giani, Estacio de Sa University, in Health Sciences, Brazil

PhD Iva Ivanovska, Harvard Medical School, Department of Genetics, 77 Avenue Louis Pasteur, NRB room 239, Boston, MA 02115, United States

MD, PhD Jerzy Jablęcki, Associate Professor, Division of General Surgery St. Jadwiga of Silesia Hospital, Trzebnica; Head, Subdepartment of Hand Surgery an Replantation St Jadwiga of Silesia Hospital, Trzebnica; Professor, Department of Public Health, State Higher Professional Medical School, Opole, Poland. 55-100 Trzebnica, ul. Prusicka 53, Poland

MD Mehrdad Jalalian Hosseini, Khorasan-e Razavi Blood Center, Mashhad, Iran, Islamic Republic of

PhD Radka Kaneva, Department of Medical Chemistry and Biochemistry, Medical University - Sofia, Bulgaria

Prof. Dr. Kostandina Leonida Korneti-Pekevsk, Ss Cyril and Methodius University of Skopje, Faculty of Medicine, Skopje, Republic of Macedonia

MD, PhD Branko Malenica, Department of Immunology, Clinical Hospital Center Zagreb, Zagreb University School of Medicine, Zagreb, Croatia

Prof. Dr. Elida Mitevka, Institute of Histology and Embriology, Faculty of Medicine, Ss Cyril and Methodius University of Skopje, Skopje, Republic of Macedonia

MD, PhD Marija Mostarica-Stojković, Institute of Microbiology and Immunology, University of Belgrade School of Medicine, Belgrade, Serbia

PhD Vesna Nikolova-Krstevski, Harvard Institutes of Medicine, HIM-201, 4 Blackfan Circle, Boston, MA, 02134, United States

Prof. Dr. Nikola Panovski, Institute of Microbiology and Parasitology, Faculty of Medicine, Skopje, Republic of Macedonia

MD, BIDMC Iva Petkovska, Beth Israel Deaconess Medical CenterRadiology W CC - 3 330 Brookline Ave. Boston, MA 02215, United States

Prof. Dr. Gordana Petrusevska, Institute of Pathology, Medical Faculty, University of "Ss. Cyril and Methodius" Skopje, Republic of Macedonia

Prof. Enver Roshi, Dean of Faculty of Public Health, Medical University of Tirane, Chief of Epidemiological Observatory, National Institute of Public Health. Address: Rruga e Dibrës, Str. 371, Tirana, Albania

MD, PhD Gorazd B. Rosoklija, Professor at Columbia University and member of the Macedonian Academy of Sciences and Arts, United States

Prof. Dr. Aleksandar Sikole, University Clinic for Nephrology, Faculty of Medicine, Ss Cyril and Methodius University of Skopje, Skopje, Republic of Macedonia

MD, FESC Gianfranco Sinagra, Department of Cardiology, "Ospedali Riuniti" and University of Trieste, Ospedale Cattinara "Strada di Fiume, 447, 34149 Trieste, Italy

MD, PhD Rumen Stefanov, Information Centre for Rare Diseases and Orphan Drugs (ICRDOD), Bulgaria; Department of Social Medicine, Medical University of Plovdiv, Bulgaria

Prof. Dr. Vesna Velikj Stefanovska, Department of Epidemiology and Biostatistics with Medical Informatics, Medical Faculty, UKIM, Skopje, Republic of Macedonia

MD, MBA Milenko Tanasijevic, Director, Clinical Laboratories Division and Clinical Program Development, Pathology Department, Brigham and Women's Hospital, Dana Farber Cancer Institute, Associate Professor of Pathology, Harvard Medical School, United States

MD, FRCPC Kiril Trpkov, Associate Professor, University of Calgary, Department of Pathology and Laboratory Medicine, Calgary Laboratory Services. 7007 14 st, Calgary SW, Canada

MD, PhD Igor Tulevski, Department of Cardiology, Academic Medical Center, Amsterdam, 1100 DD, T 020 707 2930; F 020 707 2931, Netherlands

Past Members of Editorial Team

Prof. Dr. Doncho Donev, Institute of Social Medicine, Medical Faculty, Ss Cyril and Methodius University of Skopje, Skopje, Republic of Macedonia

Prof. Dr. Olivera Stojceva Taneva, University Clinic of Nephrology, Republic of Macedonia

Prof. Dr. Petar Miloshevski, from 2008-2014, Institute of Preclinical and Clinical Pharmacology with Toxicology, Faculty of Medicine, Skopje, Republic of Macedonia

Prof. Dr. Sonja Topuzovska, Institute of Medical and Experimental Biochemistry, Faculty of Medicine, Skopje, Republic of Macedonia

Prof. Dr. Aleksandar Dimovski, Institute of Pharmaceutical Chemistry, Faculty of Pharmacy, University "Ss Kiril and Metodij", Skopje, Republic of Macedonia

PhD Mirko Trajkovski, ETH Zurich, Wolfgang-Pauli-Str. 16/HPT D57, 8093 Zurich-CH, Switzerland

PhD Zoran Zdravkovski, Institute of Chemistry, Faculty of Natural Sciences and Mathematics, Ss Cyril and Methodius University of Skopje, Skopje, Republic of Macedonia

Published by



About us

[About the Journal](#)
Editorial Team
Author Fees
Online Payments
Bibliographic Information
Journal History

Information

[For readers](#)
[For authors](#)
[For librarians](#)
[For Reviewers](#)
Transparency

Contact

Publisher
Co publisher
Registration
Contact
Privacy statement

[Home](#) / [Archives](#) / Vol. 9 No. E (2021): E - Public Health

Vol. 9 No. E (2021): E - Public Health

**Published:** 2021-01-10

Epidemiology

Specific and Sensitive Nutrition Interventions with Nutritional Status of Toddlers as Prevention of Stunting in the Coronavirus Disease 2019 Pandemic in Sigi District, Indonesia

Putu Candriasih, Metrys Ndama, Anna Veronica Pont (Author)

415-418



PDF

Determinants of Coronary Heart Disease Incidence among Indonesian Hajj Pilgrims Hospitalized in Saudi Arabia in 2019

Juniarty Naim, Wahiduddin Wahiduddin, Masni Masni, Ridwan Amiruddin, Irwandy Irwandy, M. Nadjib Bustan (Author)

798-804



PDF

Behavior of the Use of Mosquito Net as a Prevention of Malaria in Ondorea Village, Nanga Panda Sub-district

Yustina PM Paschalia, Anatolia K. Doondori, Irfan Irfan, Norma Tiku Kambuno (Author)

620-623



PDF

Motivation Interview Effectiveness and Optimism Efficiency on the Quality of Life of Type 2 Diabetes Mellitus

Ridwan Amiruddin, Nurhaedar Jafar, Jumriani Ansar, Zhanaz Tasya (Author)
402-405



A One-year COVID-19 Pandemic Effect on the Orthopaedic Field in Indonesia: A Cross-sectional Multi-center Study

Pamudji Utomo, Abdurrahman Afa Haridhi, Mochammadisyah Beizar Yudistira (Author)
805-811



Benzene Exposure Analysis through S-Phenylmercapturic Acid in Urine at Platelet Levels in Footwear Workers in Sukajaya Village, Bogor Regency

Siti Norjannah, Ririn Arminsih Wulandari, Al Asyary (Author)
260-264



Determinant Factors Stroke Prevention Behavior among Hypertension Patient in Indonesia

Arif Setyo Upoyo, Atyanti Isworo, Yunita Sari, Agis Taufik, Annas Sumeru, Akhyarul Anam (Author)
336-339



Enrichment of Biscuits with Andaliman and Fermented Buffalo Milk Biscuits for the Intervention of Malnutrition in Children Under Five

Diana Sembiring, Evawany Y. Aritonang, Amir Purba, Rahayu Lubis (Author)
670-674



High-density Lipoprotein Cholesterol as a Risk Factor of Health-Related Quality of Life in 50-70-Year-Old Community-Dwelling Women

Raditya Wratsangka, Elly Herwana, Yenny Yenny, Endrico Xavieress, Aditya Krishnamurti (Author)
1092-1096



Effect of Vitamin A Supplementation, Cooking Oil Fortification, and Nutrition Education to Postpartum Mother on Breast Milk Retinol Levels

Abdul Salam, Dodik Briawan, Drajat Martianto, Abdul Razak Thaha, Devintha Virani (Author)

823-827



Household Food Insecurity, Level of Nutritional Adequacy, and Nutritional Status of Toddlers in the Coastal Area of Central Tapanuli Regency

Herta Masthalina, Heru Santosa, Etty Sudaryat, Fikarwin Zuska (Author)

1371-1375



Profile of Double of Undernutrition Problem, Coexistence with Anemia among Pregnant Women Indonesia 2018: A Cross Sectional Survey

Ina Kusriani, Donny Kristanto Mulyantoro, Dwi Hapsari Tjandrarini, Hadi Ashar (Author)

1250-1255



Percentage of Injuries, and Related Factors Among a Group of Medical Students in Cairo University: A Cross-Sectional Study

Hend Ali Sabry, Alaa Abou Zeid, Marwa Salem (Author)

675-680



Evaluation of the Effect of Sociocultural Factors on the Children Stature in Langkat Regency, Indonesia

Siti Saidah Nasution, Bina Melvia Girsang, Hariati Hariati (Author)

461-466



The Spatial Distribution of Pulmonary Tuberculosis in Kabanjahe District, Karo Regency, Indonesia

Risnawati Tanjung, Eka Lestari Mahyuni, Nelson Tanjung, Oster Suriani Simarmata, Jernita Sinaga, Helfi R. Nolia (Author)

817-822



Assessment of Antepartum Depression and its Effect on Pregnancy Outcome in Two Primary Health Care Units in Qaliobia Governorate, Egypt

Ola Mostafa, Mervat El-Rafie, Eman T. Al Sayed, Mohamed A. Khalil, Sherry M. Zaki (Author)

447-454



SJR

Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

[Home](#)

Journal Rankings

Country Rankings

Viz Tools

Help

About Us



Nursing Research Paper

Submit Your Nursing and Midwifery Research With Hinc

Hindawi

Open Access Macedonian Journal of Medical Sciences

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
Macedonia	Medicine Medicine (miscellaneous)		23
 Universities and research institutions in Macedonia			

PUBLICATION TYPE	ISSN	COVERAGE
Journals	18579655	2014-2021

SCOPE

Information not localized

 Join the conversation about this journal

Advances in Civil Eng. Papers

High Quality Open Access Original Research In Advances in Civil Engineering. Submit Now

Hindawi


FIND SIMILAR JOURNALS ?








1 Journal of Clinical and Diagnostic Research IND 64% similarity	2 Journal of International Medical Research GBR 61% similarity	3 Medicina (Lithuania) CHE 61% similarity	4 Advances in Clinical Experimental Medicine POL 58% similarity
--	--	---	---

Advances in Civil Eng. Papers

High Quality Open Access Original Research In Advances in Civil Engineering. Submit Now

Hindawi






← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç
```

SCImago Graphica

Explore, visually communicate and make sense of data with our [new data visualization tool](#).





Source details

Open Access Macedonian Journal of Medical Sciences

Formerly known as: Macedonian Journal of Medical Sciences

Scopus coverage years: from 2014 to Present

Publisher: Scientific Foundation SPIROSKI

E-ISSN: 1857-9655

Subject area: Medicine: General Medicine

Source type: Journal

CiteScore 2021

1.3



SJR 2021

0.257



SNIP 2021

0.603



[View all documents >](#)

[Set document alert](#)

[Save to source list](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

i Improved CiteScore methodology



CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore 2021

$$1.3 = \frac{4,420 \text{ Citations 2018 - 2021}}{3,303 \text{ Documents 2018 - 2021}}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022

$$1.2 = \frac{3,672 \text{ Citations to date}}{3,139 \text{ Documents to date}}$$

Last updated on 05 September, 2022 • Updated monthly

CiteScore rank 2021

Category	Rank	Percentile
Medicine		
General Medicine	#416/826	49th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)

About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

ELSEVIER

[Terms and conditions](#) ↗ [Privacy policy](#) ↗

Copyright © [Elsevier B.V](#) ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the [use of cookies](#) ↗.

