



Proceedings

Proceedings of the 1st Jenderal Soedirman International Medical Conference in conjunction with the 5th Annual Scientific Meeting (Temilnas) Consortium of Biomedical Science Indonesia

November 28-28, 2020, in Purwokerto, Indonesia



Editors: Joko Mulyanto¹; Teguh Haryo Sasongko²; Delvac Oceandy³; Dwi Utami Anjarwati¹; Wahyu Siswandari¹; Hutomo Suryo Wasisto⁴ and Motoyuki Sugai⁵

Affiliations: ¹ Faculty of Medicine Universitas Jenderal Soedirman, Indonesia ; ² Perdana University Malaysia, Malaysia ; ³ The University of Manchester, United Kingdom ; ⁴ Technische Universität Braunschweig, Germany ; ⁵ Hiroshima University, Japan

ISBN: 978-989-758-499-2

Conference Link: <http://jimc.fk.unsoed.ac.id>

Foreword: Nanoscience is a science that involves the study of materials on an ultra-small scale and the novel properties that these materials demonstrate. Nanoscience has the potential to reshape the world around us. It could lead to revolutionary breakthroughs in fields ranging from manufacturing to health care. At this time, we also have entered the era of genomic medicine (genomic medicine), which begins with the success in mapping the human genome. Even human genetic research in recent decades has led to the application of genomics for the benefit of patients (translational research). Genetic studies are now beginning to apply genetic information to achieve personalized medicine, which is an innovative approach to patient management by considering individual differences based on genes, environment, and lifestyle. This 1st Jenderal Sudirman International Medical Conference” in conjunction with 5th Annual Scientific Meeting (Temilnas) Consortium of Biomedical Science Indonesia (KIBI 2020) **(More)**

Volumes:

Vol. 1 - 978-989-758-499-2

The Role of Fibroblast Proliferation in Wound Healing by Different Plants: An Experimental Study

P. 5 - 9

Marisa Rilian , Indra Kusuma , Abdul Halim , Aliya Muhammad ,
Akbar Fitrianto and Ida Bagus Eka Narendra

DOI:10.5220/0010486300050009

Formative Research to Develop a Prototype of Sugar-sweetened Beverages Monitoring Application for Obesity Management among Indonesian Adolescents

P. 10 - 14

Eka Febriyanti , Ratih Yulistika Utami and Febrina Dewi Pratiwi Lingga

DOI:10.5220/0010486900100014

Chlorogenic Acid Ameliorates Vascular Remodeling and Perivascular Fibrosis in Kidney Fibrosis Model in Mice

P. 15 - 20

Gabriella Bamba Ratih Lintin , Nur Arfian , Dwi Cahyani Ratna Sari , Gina
Andyka Hutasoit , Mohammad Salman and Muhammad Mansyur Romi

DOI:10.5220/0010487000150020

Characteristics, Effective Reproduction Number (Rt), and Spatial Modelling of Covid-19 Spread at Magetan District, East Java, Indonesia

P. 21 - 27

Yudhi Wibowo , Nendyah Roestijawati , Atik C. Hidayah , Agoes Y. Purnomo
and Tutiek Herlina

DOI:10.5220/0010487100210027

Suppression of MnSOD by Andrographolide and its Relation to Oxidative Stress and Viability of Breast Cancer Stem Cells Treated with Repeated Doxorubicin Administration

P. 28 - 34

Angie Tara Rachman , Ayu Suraduhita , Resda Akhra Syahrani , Melva Louisa
and Septelia Inawati Wanandi

DOI:10.5220/0010487200280034

Toxicity of Self-nanoemulsifying Drug Delivery System Formulation of Nigella Sativa L. Seed Oil against Adult Danio rerio

P. 35 - 42

Yanti Tri Utami and Isnatin Miladiyah

DOI:10.5220/0010487300350042

Optimization of Immunohistochemical Staining with Anti Protein Gene Product 9,5 (PGP 9,5) Antibodies to Detecting Intraepidermal Nerve Fiber

P. 43 - 46

David Pakaya , Yustina Andwi Ari Sumiwi , Sri Herwiyanti and Rina Susilowati

DOI:10.5220/0010487400430046

Immunohistochemistry of KRAS Protein in Colorectal Cancer

P. 47 - 51

Dody Novrial , Kamal Agung Wijayana and Hanif Kun Cahyani

DOI:10.5220/0010487500470051

The Effect of Ethanol Extract of Lingzhi Mushroom (Ganoderma lucidum) on Caspase-3 Expression in Oral Cancer Cells

P. 52 - 57

Irfan Dwiandhono , Fadli Ashar , Arsa Hadiyatama Waskito Aji and Meta
Anjay Firmansyah

DOI:10.5220/0010487600520057

The Role of Chlorogenic Acid on Mice with Unilateral Ureteral Obstruction Model: A Study toward Tubular Injury

P. 58 - 62

Low-dose of Acetylsalicylic Acid Upregulates Expression of eNOS mRNA and Downregulates Interleukin-6 (IL-6) and Transforming Growth Gactor- β 1 (TGF- β 1) mRNA in Rat Kidney of Preeclampsia Model

P. 63 - 68

Yuyun Nailufar , Rul Afiah Syarif , Andi Fitriani Kusuma , Farmita Chairani ,
Nia Marlina , Charolina Vivi Vienetta , M. Hadri Ar-Ridho , Totok Utoro and
Nur Arfian

DOI:10.5220/0010487800630068

Dose-dependent Decaffeinated Green Tea Extract Administration Improved Hyperglycemia through Modulation of IRS-1 and GLUT-4 Genes Expression in Metabolic Syndrome Rat Model

P. 69 - 74

Dwi Adi Nugroho , Mifetika Lukitasari , Marlita Marlita , Mohammad
Saifur Rohman , Nashi Widodo , Inggita Kusumastuty and Nur Ida Panca
Nugrahini

DOI:10.5220/0010487900690074

Expression of HSA-MIR-155-5P and mRNA Suppressor of Cytokine Signalling 1 (SOCS1) on Plasma at Early-stage and Late-stage of Nasopharyngeal Carcinoma

P. 75 - 80

Recita Indraswary , Sofia Mubarika Haryana and Agus Surono

DOI:10.5220/0010488000750080

Cost Analysis Monotherapy of Type 2 Diabetes Mellitus Patients in RSUD. Prof. Dr. Margono Soekarjo

P. 81 - 83

Fajar Wahyu Pribadi , Afifah and Catharina Widiartini

DOI:10.5220/0010488100810083

Antibiofilm Activity of Aloe barbadensis Miller Extract Against Staphylococcus aureus

P. 84 - 90

Kanesti Ismirajna and Irena Agustiningtyas

DOI:10.5220/0010488200840090

The Renal Protective Potential Effect of Infusion of Anti-urolithiasis Formula in Urolithiasis Patients: A Randomized Clinical Study

P. 91 - 97

Ulfatun Nisa , Peristiwa Ridha Widhi Astana , Saryanto , Tyas Friska Dewi and
Enggar Wijayanti

DOI:10.5220/0010488300910097

In Vitro Study of Reduction of Oral Enterococcus faecalis Biofilm on Application of Combination of Chrysomya megacephala Maggot Extract and Sodium Hypochlorite

P. 98 - 103

Rizka Hidayati , Ari Asnani , Muhamad Salman Fareza and Dwi Utami Anjarwati

DOI:10.5220/0010488400980103

Effects of 8-Week Vitamin E (α Tocopherol) Supplementation on Reduced Insulin Resistance in Non-diabetic Obese Subjects

P. 104 - 108

Erlina Marfianti and Isnatin Miladiyah

DOI:10.5220/0010488501040108

The Activities of Streptomyces W-5A as Antibacterial and Antibiofilm towards Methicillin-resistant Staphylococcus aureus 2983

P. 109 - 115

Annisa Permata Dinda , Ari Asnani and Dwi Utami Anjarwati

DOI:10.5220/0010488601090115

Association of Epstein Barr Virus Infection to Prognosis Factors of Invasive Breast Cancer in Semarang Kariadi General Hospital Population P. 116 - 120

Wasisto Dwi Yudisaputro , Djoko Handojo , Yan Wisnu Prajoko and

Dik Puspasari

DOI:10.5220/0010488701160120

Analysis of the Effectiveness of Chrysomya sp. Maggot Extract in Inhibiting the icaA and icaD Genes Regulator Staphylococcus epidermidis Biofilm P. 121 - 125

Dylan Tamalsir , Anton Budhi Darmawan and Dwi Utami Anjarwati

DOI:10.5220/0010488801210125

Correlation between Hepcidin and Ferritin with Insulin and Hba1C as Biochemical Markers of Pancreas Damage in β -Thalassemia Patients P. 126 - 130

Irbath Hamdani , Qodri Santosa , Joko Setyono , Hernayanti and Lantip Rujito

DOI:10.5220/0010488901260130

Chitosan Nanoparticle as a Delivery System of miRNA 217 for Suppressing Hepatocellular Carcinoma Progressivity by Targeting AEG-1/P53 P. 131 - 138

Ulfatun Nisa , Indwiani Astuti , Ronny Martien , Dhani Rinaldi Maulana and

Ysrafil

DOI:10.5220/0010489001310138

DRD2 Gene Polymorphisms in Schizophrenia Patients P. 139 - 142

Zieske Maritska , Miranti Dwi Hartanti , Benediktus Wicaksono Widodo , Fitri ,

Bintang Arroyantri Prananjaya , Irsan Saleh and Nita Parisa

DOI:10.5220/0010489101390142

The Use of Autogenous Tooth Bone Graft Block (ABTB) in Post-Extraction Socket of Dental Implant Treatment Procedure: A Literature Review P. 143 - 147

Bambang Tri Hartomo , Ambar Delfi Mardiunti , Dewi Sartieka Putri ,

Irma Khoerunisa and Pratitis Widi Seno

DOI:10.5220/0010489201430147

Comparison of Purple Passion Juice (*Passiflora edulis* var. *edulis*) and Simvastatin on Lipid-lowering Effect of Hyperlipidemic Rats Model P. 148 - 154

Alfi Muntafiah , Tisna Sendy Pratama , Anisa Rachmawati , Dewi

Wahyu Wulandari and Qodri Santosa

DOI:10.5220/0010489301480154

Comparison of Tear Interleukin-6 level, Conjunctival Inflammation, and Tear Film Function in Daily Conventional Hydrogel and Extended Silicone Hydrogel Contact Lens Wear P. 155 - 163

Tri Rahayu , Adisti , Weni Puspitasari and Umar Mardianto

DOI:10.5220/0010489401550163

The Effect of Jackfruit (*Artocarpus heterophyllus*) Leaf Ethanolic Extract Gel on Superoxide Dismutase and Interleukin-1 β Levels in Wound Healing after Tooth Extraction in Diabetic Rats P. 164 - 169

Almasyifa Herlingga Rahmasari Amin , Yusuf Asto Pamasja , Christiana

Cahyani Prihastuti , Haris Budi Widodo and Hernayanti

DOI:10.5220/0010489501640169

Sources of Information and Knowledge of Gonorrhea in Indonesia P. 170 - 174

Annisa Tristifany , Fatihah Arifah Rahmawati , Bella Maharany Syahfira ,

Adhitama Noor Idninda , Muhammad Maulana Malik , Nina Tiwi Handayani

and Sani Rachman Soleman

DOI:10.5220/0010489601700174

Different Contribution of Estrogen Receptors, ET-1/ETBR and Superoxide Dismutase and in eNOS Availability based on Sexual Dimorphism in Early Stage of Kidney Diabetic Rats

P. 175 - 183

Anisa Fatwa , Dwi Cahyani Ratna Sari , Wiwit Ananda Wahyu Setyaningsih , Andrew Nobiantoro and Nur Arfian
DOI:10.5220/0010489701750183

Description of Methicillin-resistant Staphylococcus aureus Infection at Arifin Achmad General Hospital, Riau Province from 2015-2019

P. 184 - 189

Dewi Anggraini , Dewi Santosaningsih , Kuntaman , Muhammad Ihsan and Syifa Farhani Ridha
DOI:10.5220/0010489801840189

Dietary Intake of Obese Adult Men in Jakarta

P. 190 - 193

Karima Yudhistina , Erfi Prafiantini and Novi S. Hardiany
DOI:10.5220/0010489901900193

Iron Status of Newborns in Maternal Inflammation Status Differences

P. 194 - 201

Qodri Santosa , Alfi Muntafiah and Lantip Rujito
DOI:10.5220/0010490001940201

The Effects of Aqueous Extract of Jamaican Cherry (Muntingia calabura) on D-galactose-induced Liver Damage in BALB/c Mice (Mus musculus)

P. 202 - 207

Cempaka Jaga Pramudita , Dwi Nur Ahsani , Ika Fidianingsih and Evy Sulistyoningrum
DOI:10.5220/0010490102020207

The Effect of Caffeine towards Zebrafish (Danio rerio) Juvenile Working Memory Exposed by Unpredictable Chronic Stress (UCS)

P. 208 - 213

Alya Ayu Tazkia and Zainuri Sabta Nugraha
DOI:10.5220/0010490202080213

Iron for Human Brain Development: A Fulfill Strategy in the First 1,000 Days of Life

P. 214 - 222

Qodri Santosa
DOI:10.5220/0010490302140222

Expression of MicroRNA-155 and Suppressor of Cytokines Signaling 1 (SOCS1) mRNA in Plasma Breast Cancer Patients

P. 223 - 230

Dwi Nur Indah Sari , Dina Rahmina , Sumadi Lukman , Artanto Wahyono , Indwiani Astuti , Sofia Mubarika and Teguh Aryandono
DOI:10.5220/0010490402230230

The Effects of Administering Jicama Concentrate (Pachyrhizus erosus) and Kefir Grains as a Synbiotic Drink on Malondialdehyde and Superoxide Dismutase Levels in the Testicles of Hyperlipidemic Rats

P. 231 - 236

Mohammad Alvian Subhakti , Miranti Dewi Pramaningtyas , Rafik Prabowo and Rokhima Lusiantari
DOI:10.5220/0010490502310236

The Effect of Addition of Blood Cockles (Anadara granosa) Shell Nano-Hydroxyapatite on Hardness of Heat Cured Acrylic Resin

P. 237 - 240

Dian Noviyanti Agus Imam , Ryana Budi Purnama and Aris Aji Kurniawan
DOI:10.5220/0010490602370240

Drug-induced Hypersensitivity Syndrome in a Breast Cancer Patient: A Case Report

P. 241 - 244

Antimicrobial Potential Activity of Extract *Selaginella plana* (Desv. Ex Poir.) Hieron against the Growth of *Staphylococcus aureus* ATCC 25922 and Methicillin-Resistance *Staphylococcus aureus* (MRSA)

P. 245 - 253

Juen Carla Warella , Agung Dwi Wahyu Widodo , Rebekah Juniati Setiabudi , Retno Indrawati Roestamadji , Maftuchah Rochmanti and Pudji Lestari

DOI:10.5220/0010490802450253

The Lingzhi Mushroom (*Ganoderma lucidum*) Extract Reduce the Number of Lymphocyte in Diabetics Rats with Periodontitis: In Vivo Experimental on Sprague dawley Rats

P. 254 - 259

Pratitis Widi Seno , Amilia Ramadhani , Saryono , Haris Budi Widodo and Christiana C. Prihastuti

DOI:10.5220/0010490902540259

Reversible Imiquimod Effects on Skin Tissue of Psoriasis Mice Model: An Experimental Study

P. 260 - 266

Thianti Sylviningrum , Afifah , Dody Novrial , Brian Wasita , Bambang Purwanto and Harijono K. Sentono

DOI:10.5220/0010491002600266

Efficacy of Topical Binahong Leaf Ethanolic Extract Administration on Serum TGF- β 1 in Wistar Rats with *Staphylococcus aureus*-infected Wounds

P. 267 - 272

Rizki Andini Nawawi , Muhammad Totong Kamaluddin and Theodorus

DOI:10.5220/0010491202670272

The Oral Health Status, Salivary Flow Rate and pH in Hypertensive Patients Who Consume Antihypertensive Drugs in Puskesmas Kasihan I Yogyakarta

P. 273 - 277

Nurvita Risdiana and Eka Nuraeni

DOI:10.5220/0010491302730277

Hard Tissue Surgical Treatment with Embedded Dental Condition of Tuberosity Maxillary

P. 278 - 283

Bambang Tri Hartomo and Rizka Rachmatika Dewi

DOI:10.5220/0010491402780283

Blood Transfusion, Serum Total Iron Binding Capacity and Iron in Hemodialysis Patients Margono Soekarjo Hospital

P. 284 - 288

Vitasari Indriani , Aditiawarman , Yunanto Dwi Nugroho and Fania Salsabila

DOI:10.5220/0010491502840288

Secretion of IFN- γ and IL-17 after Stimulation of ESAT-6-CFP10 (EC610) Fusion Antigen from PBMC in Groups Active TB and Latent TB

P. 289 - 297

Nika Andriani , Nova Kurniati , Muhammad Irsan Saleh , Eddy Mart Salim , Zen Hafy , Jusak Nugraha , Kemas Ya'kub Rahadiyanto and Francisca Srioetami Tanoerahardjo

DOI:10.5220/0010491602890297

Antibacterial Activity of Synthesized Silver Nanoparticle using Langsung Leaf Extract (*Lansium domesticum* var. *pubescens* Kooders et Valeton) as Bioreductor against *Escherichia coli* and *Staphylococcus aureus*

P. 298 - 304

Khairunnida Rahma , Agung Dwi Wahyu Widodo , Rebekah Juniati Setiabudi , Retno Indrawati Roestamadji , Maftuchah Rochmanti and Pudji Lestari

DOI:10.5220/0010491702980304

The Effectiveness of Mesenchymal Stem Cell and Colostrum Bovine Combination in Post Hepatectomy Liver Failure with Liver Fibrosis Animal Model

P. 305 - 310

Dimas Aryo Kusumo , Agung Putra , Albertus Ari Adrianto , Erik Prabowo and Ignatius Riwanto

DOI:10.5220/0010491803050310

Prev 1 2 Next



SCITEPRESS - SCIENCE AND TECHNOLOGY PUBLICATIONS, LDA.

RESOURCES

[Proceedings](#)

[Papers](#)

[Authors](#)

[Ontology](#)

CONTACTS

Science and Technology
Publications, Lda
Avenida de S. Francisco Xavier,
Lote 7 Cv. C,
2900-616 Setúbal, Portugal.

Phone: +351 265 520 185
Fax: +351 265 520 186
Email: info@scitepress.org

EXTERNAL LINKS

[PRIMORIS](#)

[INSTICC](#)

[SCITEVENTS](#)

[CROSSREF](#)

PROCEEDINGS SUBMITTED FOR INDEXATION BY:

[dblp](#)

[Ei Compendex](#)

[SCOPUS](#)

[Semantic Scholar](#)

[Google Scholar](#)

[Microsoft Academic](#)

In Vitro Study of Reduction of Oral *Enterococcus faecalis* Biofilm on Application of Combination of *Chrysomya megacephala* Maggot Extract and Sodium Hypochlorite

Rizka Hidayati¹ , Ari Asnani² , Muhamad Salman Fareza³  and Dwi Utami Anjarwati⁴ 

¹Magister of Biomedicine, Faculty of Medicine, Universitas Jenderal Soedirman, Purwokerto, Indonesia

²Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia

³Department of Biochemistry, Department of Pharmacy, Faculty of Health Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia

⁴Department of Microbiology, Faculty of Medicine, Universitas Jenderal Soedirman, Purwokerto, Indonesia

Keywords: Antibiofilm, Root Canal Irrigant, Blowfly Larvae, Endodontic Treatment

Abstract: The tooth's infected root canal relates to the bacteria invasion, such as *Enterococcus faecalis*. The eradication of the bacteria using single root canal irrigants becomes difficult because of the formed biofilm. We aimed to investigate the effectiveness of a combination of *C. megacephala* maggot extract with the common irrigant, sodium hypochlorite 3%, on the biofilm reduction of *E. faecalis*. The *C. megacephala* maggot extract was tested at concentrations 25%, 50%, and 100%, and the combination of each extract concentration with sodium hypochlorite in a volume ratio of 1:1; 1:2; 1:3; 2:1, and 3:1. All treatments were performed three times of replication with incubation time for 1 hour and 3 hours. Antibiofilm effect was measured with crystal violet staining and the optical density reading. Data was analyzed with the Statistical Package for the Social Sciences Statistic Version 22. The least biofilm formation was observed in combination of maggot extract 25% with sodium hypochlorite (2:1) for 1 hour incubation ($p=0,05$) and combination of maggot extract 25% (1:1) for 3 hours incubation ($p=0,000$). This combination effectively inhibits the biofilm of *E. faecalis*. This study identified the protease enzymes in *C. megacephala* maggot extract and investigated *C. megacephala* maggot extract's antibiofilm effect combine with the other root canal irrigant.

1 INTRODUCTION

One of the bacteria found in the root canal of the infected tooth is *Enterococcus faecalis*. These bacteria are persistent and can form biofilms. Therefore they are difficult to be removed. The bacteria in the root canal treatment are eradicated by applying root canal irrigant. The agent commonly used as root canal irrigant is sodium hypochlorite (NaOCl), with 1,25-5% (Mulyawati, 2011). NaOCl effectively removes planktonic bacteria, but it is less effective in reducing biofilms produced by bacteria (Dunavant et al., 2006). NaOCl is usually combined with other irrigation solution such as chlorhexidine

(CHX) to increase the antibacterial and antibiofilm effects.

Several researchers have also carried out the combination of root canal irrigant with natural agents. Geethapriya et al. (2016) combined chitosan with EDTA against *E. faecalis* biofilms with a ratio of 1: 1 in their study, which showed effective results in inhibiting *E. faecalis* biofilm formation (Geethapriya et al., 2016). Other studies combined chitosan with chlorhexidine (CHX), whose results showed the same effect with 5% NaOCl in inhibiting *E. faecalis* biofilm (Jaiswal et al., 2017). Other studies regarding the combination of NaOCl with natural agents have never been done.

^a  <https://orcid.org/0000-0001-5351-752X>

^b  <https://orcid.org/0000-0002-8569-2565>

^c  <https://orcid.org/0000-0003-2414-1249>

^d  <https://orcid.org/0000-0001-8394-2543>

Maggot extract is material from natural sources that have not been studied as root canal irrigant. Some maggots, such as maggot *Lucilia sericata* contain proteolytic enzyme components that can degrade bacterial biofilms' extracellular matrix (Chen et al., 2012). Chymotrypsin in *L. sericata* secretion can affect bacterial biofilms' adhesion (Harris et al., 2013). Other types of maggots, such as *Chloroprocta* sp. contain protease enzymes that can reduce the extracellular biofilm matrix in *Staphylococcus epidermidis* (Anjarwati et al., 2017). Another maggot extract, such as *Chrysomya megacephala* maggot extract, has an excretory and secretory. This product contains serine (trypsin and chymotrypsin), an antibacterial effect on *Escherichia coli*, *Staphylococcus aureus*, and *Bacillus subtilis* (El-Ebiarie et al., 2012; Mohamed 2015a, 2015b).

In this study, the maggot was taken from the maggot *Chrysomya megacephala*, a local maggot that is quite common in Indonesia (Putri, 2018). Combination of maggot extract *C. megacephala* with 3% NaOCl as root canal irrigant is expected to increase the reduction of *E. faecalis* biofilm.

2 METHODS

2.1 Material

This experimental laboratory study conducted the *E. faecalis* biofilm test. The materials used were *C. megacephala* maggot extract with a concentration of 25%, 50%, 100% and a combination of each concentration of maggot extract with 3% NaOCl with a volume ratio of 1: 1, 1: 2, 1: 3, 2: 1 and 3: 1, 3% NaOCl as a positive control, tryptic soy broth (TSB) and sterile phosphate-buffered saline (PBS) as negative controls. Each treatment was replicated three times. Biofilm reduction was measured by added 1% crystal violet with optical density measured at a wavelength of 595 nm (OD595) using a microtiter plate reader at the Cancer and Stem Cell Research Center Laboratory, Muhammadiyah University, Purwokerto.

2.2 Bacterial Strains

The sample used in this study is that the colony is *E. faecalis* ATCC 29212, a moderate biofilm-producing bacteria. The colony has been isolated and cultured with Mueller Hinton Agar (MHA) media in the Microbiology Laboratory of the Faculty of Medicine, Universitas Jenderal Soedirman, Purwokerto. The powder of MHA with 12 grams was solved with 240

mL of aquadest and poured into the culture tray (20 mL/tray). The media was sterilized in an autoclave for 15 minutes (2 atm, 121°C), then put in an incubator for 24 hours to check if there is contamination or not. The *E. faecalis* were cultured in sterile media with an anaerobic environment. The bacteria's growth colony was taken and diluted with NaCl 0,9% until the concentration equal to 106 CFU/mL (CFU: Colony Forming Unit) or 0,5 Mc Farland Standard (Howarto et al., 2015).

2.3 Rearing Maggot and Collecting Maggot Extract

The process of collecting *C. megacephala* maggot extract was carried out at the Faculty of Medicine, Universitas Jenderal Soedirman. The rearing of the maggot *C. megacephala* was done by installing traps of flies that have been given raw fish (fish waste). After one day, the fly eggs that have been collected in fish waste were transferred to fresh raw fish and left to grow into a maggot. After growing into a maggot, the maggot at the end of the second instar and the beginning of the third instar was collected in a vessel and washed using ethanol and distilled water three times (Arora et al., 2010).



Figure 1: Soaking *C. megacephala* maggot in Phosphate Buffered Saline (PBS).

Every 1 gram of maggot was immersed in 1 ml of sterile PBS solution for 1 hour at room temperature in a dark room (Figure 1). Continued Soaking was for 24 hours (moved to room light). Then maggot in PBS was incubated at 37°C for 48 hours. Then the maggot and liquid were separated, then the liquid was centrifuged at 25°C, with 10,000 rpm for 15 minutes.

Table 1: The result of optical density reading in 1 hour and 3 hours incubation.

No.	Treatment	Mean of absorbance	
		1 hour	3 hours
1.	TSB with <i>E. faecalis</i> (control of bacteria)	0,851	3,064
2.	<i>C. megacephala</i> maggot extract 25%	0,253	2,966
3.	<i>C. megacephala</i> maggot extract 50%	0,319	2,902
4.	<i>C. megacephala</i> maggot extract 100%	0,424	2,927
5.	Combination of <i>C. megacephala</i> maggot extract 25%, and NaOCl 3% (1:1)	0,214	0,176
6.	Combination of <i>C. megacephala</i> maggot extract 50%, and NaOCl 3% (1:1)	0,756	2,809
7.	Combination of <i>C. megacephala</i> maggot extract 100%, and NaOCl 3% (1:1)	0,619	2,915
8.	Combination of <i>C. megacephala</i> maggot extract 25%, and NaOCl 3% (1:2)	0,268	0,294
9.	Combination of <i>C. megacephala</i> maggot extract 50%, and NaOCl 3% (1:2)	0,501	0,211
10.	Combination of <i>C. megacephala</i> maggot extract 100%, and NaOCl 3% (1:2)	0,190	2,998
11.	Combination of <i>C. megacephala</i> maggot extract 25%, and NaOCl 3% (1:3)	0,209	0,212
12.	Combination of <i>C. megacephala</i> maggot extract 50%, and NaOCl 3% (1:3)	0,232	0,216
13.	Combination of <i>C. megacephala</i> maggot extract 100%, and NaOCl 3% (1:3)	0,176	2,830
14.	Combination of <i>C. megacephala</i> maggot extract 25%, and NaOCl 3% (2:1)	0,155	0,294
15.	Combination of <i>C. megacephala</i> maggot extract 50%, and NaOCl 3% (2:1)	0,498	2,925
16.	Combination of <i>C. megacephala</i> maggot extract 100%, and NaOCl 3% (2:1)	0,664	2,273
17.	Combination of <i>C. megacephala</i> maggot extract 25%, and NaOCl 3% (3:1)	0,343	2,918
18.	Combination of <i>C. megacephala</i> maggot extract 50%, and NaOCl 3% (3:1)	0,273	2,962
19.	Combination of <i>C. megacephala</i> maggot extract 100%, and NaOCl 3% (3:1)	1,395	2,920
20.	PBS with <i>E. faecalis</i>	0,317	2,982
21.	NaOCl 3% with <i>E. faecalis</i>	0,157	0,194

Table 2: The calculating of the value of MBRC₅₀ and MBRC₈₀ in biofilm test with 1 hour and 3 hours incubation.

Time of incubation	OD of Bacterial control	OD of Blank	MBRC ₅₀ (OD of bacterial control – OD of blank) x 50%	MBRC ₈₀ (OD of bacterial control – OD of blank) x 20%
1 hour	0,851	0,097	0,394	0,151
3 hours	3,064	0,109	1,477	0,591

The supernatant obtained from the centrifuge process was collected and sterilized with a 0.2 µm membrane filter. Maggot extract obtained from this process was stored at -20 °C (Honda et al., 2011).

2.4 Biofilm Reduction Measurement by Administering Chrysomya megacephala Maggot Extract and Its Combination with Sodium Hypochlorite

The biofilm of *E. faecalis* (ATCC 29212) was measured using 96-well microtiter plates. The bacteria were transferred from Mueller Hinton Agar (MHA) media into Tryptic Soy Broth (TSB) and incubated for 24 hours under anaerobic conditions, which remained at 37 °C. The culture was diluted with 1: 100 on the medium. Then 20 µl of bacterial culture was inoculated with 200 µl TSB in each well, dispensed into 96-well microtiter plates with a flat bottom. After anaerobic incubation at 37 °C for 24

hours, the planktonic bacteria from each well were disposed of carefully by using a micropipette slowly until the bottom of the well looked clear. Each well was washed with 300 µl phosphate-buffered saline (PBS) 2 times slowly.

As many as 100 µL of maggot extract filled into each well of the 96-well microtiter plate was filled with at different concentrations (25%, 50%, 100%) and a combination of maggot extract and 3% NaOCl with a ratio of 1: 1, 1: 2, 2: 1, 1: 3, and 3: 1. The 96-well microtiter plate that has been inserted maggot extract and a combination of maggot extract and NaOCl are then incubated for 1 hour and 3 hours. After incubation, the biofilm is examined by giving a dye of 200 µL of 1% crystal violet solution in water for 30 minutes and washed with distilled water. Wells are reversed and dried on paper towels and dry air. Then, 200 µL of 5% acid isopropanol was added to each well to remove biofilm colour. The optical density was measured at 595 nm (OD₅₉₅) using a microtiter plate reader (Pierce et al., 2010).




Figure 2. The effect of *Chrysomya megacephala* maggot extract on biofilm reduction of *Enterococcus faecalis* ATCC 29212 for 1-hour incubation (a) and 3 hours incubation (b)

2.5 Statistical Analysis

The effects of various maggot extract concentrations and their combinations with sodium hypochlorite in reducing biofilms were analyzed using one-way ANOVA and Post hoc LSD (Least Significant Difference) tests by using The Statistical Package for the Social Sciences Statistic Version 22.

3 RESULTS

From all treatment groups with 1-hour incubation, the most extensive treatment group in reducing *E. faecalis* biofilm was a combination of 25% maggot extract and sodium hypochlorite with a ratio of 2: 1 (reducing 81.75% of biofilm production produced from the bacterial control, $p < 0,05$). The combination of maggot extract (25%) and sodium hypochlorite (2:1) reduced biofilm better than 3% sodium hypochlorite only. The Post hoc LSD test showed that the combination of 25% maggot extract and sodium hypochlorite with ratio 2:1 does not have a different effect than 3% sodium hypochlorite in reducing the biofilm of *E. faecalis* ($p > 0,05$).

In 3 hours of incubation, the most extensive treatment group in reducing *E. faecalis* biofilm was a combination of 25% maggot extract and sodium

hypochlorite with a ratio of 1: 1 (reducing 94.25% of biofilm production produced from bacterial control, $p < 0.001$) (Table 1). The 25% maggot extract and sodium hypochlorite application (ratio 1:1) showed more effect than 3% sodium hypochlorite only. The Post hoc LSD test also showed that the combination of 25% maggot extract and sodium hypochlorite with ratio 1:1 does not have a different effect than 3% sodium hypochlorite in reducing the biofilm of *E. faecalis* ($p > 0,05$).

In 1 hour incubation, the value of MBRC50 in this research is 0,394 and MBRC80 is 0,151 (Table 2). In 1 hour incubation, MBRC50 could be found in maggot extract 25%, 50%, combination of maggot extract 25% (1:1), 25% (1:2), 100% (1:2), 25% (1:3), 50% (1:3), 25% (2:1), 25% (3:1), and 50% (3:1). Meanwhile, the value of MBRC80 could not be found in all the treatment in 1 hour incubation (Figure 1). In 3 hours incubation, the value of MBRC50 is 1,477 and the value of MBRC80 is 0,591 (Table 2). In 3 hours incubation, MBRC50 and MBRC80 could be found in the same treatment which are the combination of maggot extract 25% (1:1), 25% (1:2), 50% (1:2), 25% (1:3), 50% (1:3), and 25% (2:1) (Figure 2).

Figure 2 showed that in 1-hour incubation, MBRC50 could be reached of maggot extract application (concentration 25%, 50%, the

combination of 25% maggot extract (1:1), 25% maggot extract (1:2), 100% maggot extract (1:2), 25% maggot extract (1:3), 50% maggot extract (1:3), 25% maggot extract (2:1), 25% maggot extract (3:1), and 50% maggot extract (3:1)). Meanwhile, MBRC80 could not be found in 1-hour of incubation.

4 DISCUSSION

Compared to the results of biofilm reduction at 1-hour incubation (maggot extract 25% (2:1)), there was a decrease in the concentration of maggot extract needed to reduce biofilms for a longer time (3 hours), namely with extract concentration of 25% (1:1). Due to a decrease in maggot extract protease activity, the possibility is as time increases, therefore at 3 hours incubation. The possibility of more significant antibiofilm activity is due to the combination of 3% sodium hypochlorite and maggot extract (Laba et al., 2010).

Compared with the effect of giving 3% sodium hypochlorite, the combination of 25% (1:1) maggot extract (1 hour) and a combination of 25% (2: 1) (3 hours) maggot extract had no different effects on *Enterococcus faecalis* biofilm reduction ($p>0.05$). This result shows that an evaluation in the research process is needed to increase the antibiofilm effect to be more maximal. In PBS solvent with *E. faecalis*, compared with a combination of 25% 2: 1 maggot extract (1-hour incubation) and 25% (1:1) (3 hours incubation) combination of maggot extract, there were no significant differences in antibiofilm effects. This result showed that the presence of PBS as a solvent in making maggot extract is not a factor that influences the reduction of bacterial biofilms. However, the antibiofilm effect produced comes from the extract of the maggot.

This study also showed that the *C. megacephala* maggot extract's antibiofilm effect was more effective at 3 hours incubation ($p<0.001$). In other studies, incubation time also significantly affected the reduction of extracellular biofilm matrix after maggot extract. This is thought to be due to proteases requiring time to break down proteins into dissolved proteins in exopolysaccharide (EPS) on bacterial walls. Protease is one type of enzyme contained in maggot extract. The effect of enzyme damage is directly proportional to the length of time the interaction of environmental exposure to the enzyme. The longer the exposure to the environment will damage the enzyme's structure, so that a decrease in enzyme activity (Laba et al., 2010)

5 CONCLUSIONS

C. megacephala maggot extract and its combination with sodium hypochlorite affect the reduction of biofilms produced by *E. faecalis*. Further research is needed to develop *C. megacephala* maggot extract, such as identifying other protease enzymes in *C. megacephala* maggot extract to reduce other bacterial biofilms and determine the antibacterial and antibiofilm effects of *C. megacephala* maggot extract with a combination of other root canal irrigation materials.

ACKNOWLEDGEMENTS

This paper can be arranged by support from many people. Therefore, we are grateful to Dr. Warsinah, M.Si., Apt, Dr. Oedjijono, M.Sc, and Dr. dr. Lantip Rujito, M.Si. Med for the advice to make this paper better.

REFERENCES

- Anjarwati DU, Nuryastuti T, Riwanto I, Wahyono H. Effects of *Chloroprocta* sp. maggot filtrates on extracellular matrix reduction and embedded *Staphylococcus epidermidis* viability. *Malaysian Journal of Microbiology*. 2017; 3(3) : 235-243.
- Arora S, Lim CS, Baptista C. Antibacterial activity of *Lucilia cuprina* maggot extracts and its extraction techniques. *International Journals of Integrative Biology*. 2010; 9(1): 43-48.
- Chen Y, Norde W, Mei HC, Busscher, HJ. Bacterial Cell Surface Deformation under External Loading. *American Society For Microbiology*. 2012; 3(6): 1-7.
- Dunavant TR, Regan JD, Glickman GN, Solomon ES, Honeyman AI. Comparative evaluation of endodontic irrigants against *Enterococcus faecalis* biofilms. *J. Endod*. 2006; 32 (6): 527-531.
- El-Ebiarie AS, Taha N. Molecular characterization of serine proteases from both first and third larval instars of *Chrysomya megacephala*. *Life Science Journal*. 2012; 9(3): 2086-2093.
- Geethapriya N, Subbiya A, Padmavathy K, Mahalakshmi K, Vivekanandan P, Sukumaran VG. Effect of chitosan-ethylenediamine tetraacetic acid on *Enterococcus faecalis* dentinal biofilm and smear layer removal. *J. C. D*. 2016; 19: 472-477.
- Harris LG, Nigam Y, Sawyer J, Mack D, Pritchard DI. *Lucilia sericata* Chymotrypsin Disrupts Protein Adhesin-Mediated *Staphylococcal* Biofilm Formation. *Appl. Environ. Microbiol*. 2013; 79(4): 1393-1395.
- Honda, K, Okamoto, K, Mochida Y, Ishioka K, Oka M, Maesato K., et al. A novel mechanism in maggot

- debridement therapy: protease in excretion/secretion promotes hepatocyte growth factor production. *Am. J. Physiol.* 2011; 301: C1423-C1430.
- Howarto SM, Wowor PM, Mintjelaskan CN. Uji efektivitas antibakteri minyak atsiri sereh dapur sebagai bahan medikamen saluran akar terhadap bakteri *Enterococcus faecalis*. *Jurnal e-Gigi.* 2015; Vol.3 (2), 432-438.
- Jaiswal N, Sinha DJ, Singh UP, Singh K, Jandial UA, Goel S, Evaluation of antibacterial efficacy of chitosan, chlorhexidine, propolis, and sodium hypochlorite on *Enterococcus faecalis* biofilm: an in vitro study. *J. Clin. Exp. Dent.* 2017; 9 (9): e1066-1074.
- Łaba W, Rodziewicz A. Keratinolytic Potential of Feather-Degrading *Bacillus polymyxa* and *Bacillus cereus*. *Polish Journal of Environmental Studies.* 2010; 19(2): 371-378.
- Mohamed NT. Activity of two serine proteases and antibacterial studies on both excretory/secretory products and whole body homogenates of third instar larvae of *Chrysomya megacephala*. *International Journal of Advanced Research.* 2015b; Vol. 3 (4), 451-459.
- Mohamed NT. Fractionation and purification of bioactive peptides in excretory/ secretory products of third instar larvae of *Chrysomya megacephala* (Calliphoridae: Diptera). *International Journal of Applied and Pure Science and Agriculture.* 2015a; Vol.01, 129-136.
- Mulyawati E. Peran bahan desinfeksi pada perawatan saluran akar. *Maj. Ked. Gi.* 2011; 18 (2): 205-209.
- Pierce, CG, Uppuluri P, Tummala S, Lopez-Ribot JL. A 96 Well Microtiter Plate-based Method for Monitoring Formation and Antifungal Susceptibility Testing of *Candida albicans* Biofilms. *Journal of Visualized Experiments.* 2010; 44: 1-4.
- Putri YP. Taksonomi lalat di pasar induk Jakabaring kota Palembang. Sainmatika: *Jurnal Ilmiah Matematika dan Ilmu Pengetahuan Alam.* 2018; 15(2): 105-111.



Dwi Utami Anjarwati <dwi.anjarwati@unsoed.ac.id>

Decision Letter For Acceptance

JIMC UNSOED <jimc@unsoed.ac.id>

2 Februari 2021 05.54

Kepada: dwi.anjarwati@unsoed.ac.id

Dear Ms. Dwi Utami Anjarwati

The scientific committee of JIMC 2020 thanks to the authors for submitting the full-paper manuscript for publication in JIMC 2020 conference proceeding. The paper has been distributed to two different peer-reviewers to be blindly reviewed. Following the comments from the reviewers, we are happy to inform you that the submitted paper is **accepted**. To be published in the JIMC 2020 conference proceeding please refer to the attached letter for further detailed information.

Yours sincerely,

Tirta Wardana, S.Si, M. Biotech

Program Manager the 1st Jenderal Soedirman Medical Conference (JIMC)

Faculty of Medicine, Universitas Jenderal Soedirman



Dwi Utami Anjarwati.pdf

298K