

PREFACE • **OPEN ACCESS**

10th Joint Conference on Chemistry

To cite this article: 2016 *IOP Conf. Ser.: Mater. Sci. Eng.* **107** 011001

View the [article online](#) for updates and enhancements.

Related content

- [11th Joint Conference on Chemistry in Conjunction with the 4th Regional Biomaterials Scientific Meeting](#)
- [Organizer of International Conference on Science and Applied Science \(Engineering and Educational Science\) 2016](#)
- [Students' conceptions analysis on several electricity concepts](#)
D E Saputro, S Sarwanto, S Sukarmin et al.



IOP | ebooks™

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.

Editorial Materials (preface and photograph)

PREFACE

The 10th Joint Conference on Chemistry is an international conference organized by 4 chemistry departments of 4 universities in central java, Indonesia. The universities are Sebelas Maret University, Diponegoro University, Semarang State University and Soedirman University. The venue was at Solo, Indonesia, at September 8-9, 2015. The total conference participants are 133 including the invited speakers. The conference emphasized the multidisciplinary chemical issue and impact of today's sustainable chemistry which covering the following topics:

- Material innovation for sustainable goals
- Development of renewable and sustainable energy based on chemistry
- New drug design, experimental and theoretical methods
- Green synthesis and characterization of material (from molecule to functionalized materials)
- Catalysis as core technology in industry
- Natural product isolation and optimization

The conference speakers are:



Prof. Takuji
Ogawa

*Single molecular electronic
devices and their integration,
Physical Organic Chemistry,
Osaka University*
Osaka University, Japan.
[Research](#)



Prof. Dr.
Evamarie. Hey-
Hawkins

*Catalysis- A powerful means for
sustainable developments,
Institute of Inorganic Chemistry
Leipzig University, Germany.*
[Research](#)



Prof. Masaaki
Nagatsu

*Plasma science and technology,
Shizuoka University*
Shizuoka University, Japan.
[Research](#)





Prof. Santiago
Gómez Ruiz

In search of novel sustainable systems based on nanostructured materials for different catalytic applications.,
Rey Juan Carlos University,
Móstoles (Madrid), Spain
[Research](#)



Dr. Henning
Störz

Current State and Future Prospects Including Monomer Preparation from Biomass
Institute of Agriculture Technology,
Johann Heinrich Von Thuenen Institute.
Braunschweig Germany.



Dr. Leny Yulianti

Photocatalytic hydrogen production,
Ibnu Sina Institute
University Teknologi Malaysia.
[Research](#)



Prof. Effendy,
PhD

Molecular Structure Analysis by Single X-ray Diffraction Method,



Dr.rer.nat
Hery.Haerudin

Catalytic and sustainable Processes in PERTAMINA,



Dr. Hasanudin
Abdurakhman

Advanced Material Research in Toray,
Toray Foundation



Dr. Fitria
Rahmawati

Materials for Solid Electrolytes,
Sebelas Maret University
. [Research1](#) [Research2](#)

OPEN ACCESS

Peer review statement

To cite this article: 2016 *IOP Conf. Ser.: Mater. Sci. Eng.* **107** 011002

View the [article online](#) for updates and enhancements.

Related content

- [Peer review statement](#)
- [Peer review statement](#)
- [Peer review statement](#)



IOP | ebooks™

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the **collection** - download the first chapter of every title for free.

Peer review statement

All papers published in this volume of *IOP Conference Series: Materials Science and Engineering* have been peer reviewed through processes administered by the proceedings Editors. Reviews were conducted by expert referees to the professional and scientific standards expected of a proceedings journal published by IOP Publishing.



NOTICE: Ukraine: Read IOP Publishing's statement.

Table of contents

Volume 107

2016

◀ Previous issue Next issue ▶

10th Joint Conference on Chemistry 8–9 September 2015, Solo, Indonesia

Accepted papers received: 21 December 2015

Published online: 05 February 2016

Open all abstracts

Preface

OPEN ACCESS 011001

10th Joint Conference on Chemistry

+ Open abstract  View article  PDF

OPEN ACCESS 011002

Peer review statement

+ Open abstract  View article  PDF

Papers

Catalysis

OPEN ACCESS 012001

Photocatalytic removal of phenol over titanium dioxide- reduced graphene oxide photocatalyst

Nor Shuhada Alim, Hendrik O. Lintang and Leny Yuliaty

+ Open abstract  View article  PDF

OPEN ACCESS 012002

The shape conversion of silver nanoparticles through heating and its application as homogeneous catalyst in reduction of 4- nitrophenol

H A Ariyanta and Y Yulizar

+ Open abstract  View article  PDF

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



OPEN ACCESS

012003

Preparation of TiO₂-ZnO and its activity test in sonophotocatalytic degradation of phenol

Is Fatimah and Novitasari

[+ Open abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012004

Chitosan and N-Alkyl chitosan as a heterogeneous based catalyst in the transesterification reaction of used cooking oil

T S Julianto and R A Mumpuni

[+ Open abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012005

High photocatalytic activity of mixed anatase-rutile phases on commercial TiO₂ nanoparticles

Wai Ruu Siah, Hendrik O Lintang, Mustaffa Shamsuddin and Leny Yuliaty

[+ Open abstract](#)[View article](#)[PDF](#)**Enviromental Chemistry****OPEN ACCESS**

012006

Synthesis of α -Bismuth oxide using solution combustion method and its photocatalytic properties

Y. Astuti, A. Fauziyah, S. Nurhayati, A.D. Wulansari, R. Andianingrum, A.R. Hakim and G. Bhaduri

[+ Open abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012007

Aliphatics hydrocarbon content in surface sediment from Jakarta Bay, Indonesia

M YAzis, L Asia, A Piram, BBuchari, P Doumenq and A D Syakti

[+ Open abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012008

Occurrence of aliphatic and polyaromatic hydrocarbons (PAHs) in Mytillus galloprovincialis from the traditional market in Marseille, France, by Gas Chromatography triplequadropole tandem Mass Spectrometry (GC-QQQ/MS)

M Y Azis, Yelmiza, L Asia, A Piram, B. Bucharil, P Doumenq and A D Syakti

[+ Open abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012009

Synthesis of thiazole silica hybrid from waste glass for adsorption of cadmium(II)

C. Azmiyawati and TaslimahVirkyanov

[+ Open abstract](#)[View article](#)[PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



OPEN ACCESS

012010

Alginate cryogel based glucose biosensor

Amin Fatoni, Dian Windy Dwiasi and Dadan Hermawan

[+ Open abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012011

Adsorption of remazol yellow FG from aqueous solution on chitosan-linked P-T-Butylcalix[4]Arene

Desi Suci Handayani, Candra Purnawan, Pranoto, Sri Hastuti and Diniyah Hilmiyana

[+ Open abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012012

Photocatalytic removal of 2,4-dichlorophenoxyacetic acid herbicide on copper oxide/titanium dioxide prepared by co-precipitation method

Shu Chin Lee, Norhasnita Hasan, Hendrik O. Lintang, Mustaffa Shamsuddin and Leny Yuliati

[+ Open abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012013

Analysis of diazinon pesticide using potentiometric biosensor based on enzyme immobilized cellulose acetate membrane in gold electrode

Mashuni, L O A N Ramadhan, M Jahiding and Herniati

[+ Open abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012014

Cross-linking of succinate-grafted chitosan and its effect on the capability to adsorb Pb(II) ion

Abu Masykur, Sri Juara Santosa and Dwi Siswanta dan Jumina

[+ Open abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012015

Development of electrokinetic remediation for caesium: A feasibility study of 2D electrode configuration system

Rudy Syah Putra

[+ Open abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012016

Improving quality of textile wastewater with organic materials as multi soil layering

Supriyadi, H Widiyanto, Pranoto and AK Dewi

[+ Open abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012017

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



Adsorption and desorption of Zn(II) and Cu(II) on Ca- alginate immobilized activated rice bran

A Suratman, N Z Kamalia and W A Kusumawati

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012018

Optimization of biosurfactant production in soybean oil by *rhodococcus rhodochrous* and its utilization in remediation of cadmium-contaminated solution

Venty Suryanti, Sri Hastuti and Dewi Andriani

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012019

Development of inorganic composite material based TiO₂ for environmental application

Sayekti Wahyuningsih, Ari Handono Ramelan, Edi Pramono, Candra Purnawan, Velina Anjani, Puji Estianingsih, Ludfiaastu Rinawati and Khusnan Fadli

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012020

Modification of chitosan by swelling and crosslinking using epichlorohydrin as heavy metal Cr (VI) adsorbent in batik industry wastes

B Hastuti, A Masykur and S Hadi

[+ Open abstract](#)
[View article](#)
[PDF](#)

Material Science and Engineering

OPEN ACCESS

012021

Solubility dynamic of methyl yellow and carbon black in microemulsions and lamellar liquid crystal of water, non ionic surfactants and cyclohexane system

A Amran, R Harfianto, W Y Dewi, D Beri and A Putra

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012022

Synthesis and characterization of geopolymer from bottom ash and rice husk ash

Ufafa Anggarini and Ndaru C Sukmana

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012023

Electrical conductivity of zirconia and yttrium-doped zirconia from Indonesian local zircon as prospective material for fuel cells

Karima Apriany, Ita Permadani, Dani G. Syarif, Syoni Soepriyanto and Fitria Rahmawati

[+ Open abstract](#)
[View article](#)
[PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.

012024



Optical simulation of surface textured TCO using FDTD method

I L Elviyanti, H Purwanto and Kusumandari

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012025

Effect of reaction time and (Ca+Mg)/Al molar ratios on crystallinity of Ca-Mg-Al layered double Hydroxide

E Herald, K D Nugrahaningtyas, F B Sanjaya, A A Darajat, D S Handayani and Y Hidayat

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012026

Intercalation of diclofenac in modified Zn/Al hydrotalcite-like preparation

E Herald, R W Suprihatin and Pranoto

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012027

Detection of nitrite and nitrate ions in water by graphene oxide as a potential fluorescence sensor

Ing Hua Tang, Rita Sundari, Hendrik O Lintang and Leny Yuliat

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012028

The study of FTO surface texturing fabrication using Argon plasma etching technique for DSSC applications

Lindha Jayanti, Kusumandari, Tjipto Sujitno and Risa Suryana

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012029

Degradation of self-compacting concrete (SCC) due to sulfuric acid attack: Experiment investigation on the effect of high volume fly ash content

S A Kristiawan, Sunarmasto and G P Tyas

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012030

Solvothermal and electrochemical synthetic method of HKUST-1 and its methane storage capacity

Witri Wahyu Lestari, Marisa Adreane, Candra Purnawan, Hamzah Fansuri, Nurul Widiastuti and

Sentot Budi Rahardjo

[+ Open abstract](#)[View article](#)[PDF](#)

OPEN ACCESS

012031

The site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



Changes in grammage, tearing resistance, and water vapor transmission rate of active paper incorporated with Cinnamaldehyde during storage at various temperatures

G J Manuhara, L U Khasanah and R Utami

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012032

Design of reflective optical fiber sensor for determining refractive index and sugar concentration of aqueous solutions

Ahmad Marzuki, Nila Wulan Sari and Riatun

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012033

Synthesis of $\text{LiNi}_x\text{Mn}_{2-x}\text{O}_4$ by low-temperature solid-state reaction and its microstructure

D Purwaningsih, R Roto and H Sutrisno

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012034

Synthesis and characterization of *bis*-(2-cyano-1-methyl-3- $\{2-\{(5\text{-methylimidazol-4-yl)methyl}\}$ thio}ethyl)guanidine copper(II) sulfate tetrahydrate

Sentot B Rahardjo, Teguh Endah Saraswati, Edy Pramono and Nur Fitriana

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012035

The single cell of low temperature solid oxide fuel cell with sodium carbonate-SDC (samarium-doped ceria) as electrolyte and biodiesel as fuel

F Rahmawati, A Nuryanto and K D Nugrahaningtyas

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012036

Physical and optical properties of lead doped tellurite glasses

Riyatun, Lita Rahmasari and Ahmad Marzuki

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012037

The influence of the composition of $\text{P3HT}:\text{TiO}_2$ on the characteristics of hybrid polymer solar cell

Erlyta Septa Rosa, Shobih, Hanik Munawaroh and Pranoto

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012038

Effect of cyclical thermal to mechanical properties of Hybrid Laminate Composites (HLG) with skin recycle polypropylene/natural fiber/halloysite and core PP/KF composites

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



Neng Sri Suharty, Hanafi Ismail, Desi Suci Handayani, Kuncoro Diharjo, Fajar Rachman Wibowo and
Margaretha Arnita Wuri

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012039

Physical characteristics of chitosan-silica composite of rice husk ash

Woro Sumarni, Retno Sri Iswari, Putut Marwoto and Endah F Rahayu

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012040

Study of the acidic properties of ZrO₂-pillared bentonite

Ahmad Suseno, Priyono, Karna Wijaya and Wega Trisunaryanti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012041

Effect of glycerol on mechanical and physical properties of silver-chitosan nanocomposite films

E Susilowati, I Kartini, S J Santosa and Triyono

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012042

Functionalization of C-4-methoxyphenylcalix[4]resorcinarene with several ammonium compounds

Suryadi B Utomo, Agung N C Saputro and Yudi Rinanto

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012043

The Effects of plasticizers and palmitic acid toward the properties of the carrageenan Film

Atmanto Heru Wibowo, Oktaviana Listiyawati and Candra Purnawan

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012044

Synthesis H-Zeolite catalyst by impregnation KI/KIO₃ and performance test catalyst for biodiesel production

W. Widayat, Adit Rizky Wicaksono, Lukman Hakim Firdaus and Ndaru Okvitarini

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012045

Nanocellulose prepared by acid hydrolysis of isolated cellulose from sugarcane bagasse

W P Wulandari, A Rochmad and I M Arcana
This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



[+ Open abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012046

Laboratory investigation on the role of tubular shaped micro resonators phononic crystal insertion on the absorption coefficient of profiled sound absorber

I Yahya, J I Kusuma, Harjana, R Kristiani and R Hanina

[+ Open abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012047

Preparation and properties of arenga starch-chitosan based edible film

Putri Rahmawati, Dwi Ni'maturrohman, Putri Luthfiana Sari, Putri Puspita Sari, Tetty Nur Indah Sari and Venti Suryanti

[+ Open abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012048

Synthesis and characterization of free-isocyanate polyurethane as renewable coating materials

O A Saputra, K Apriany, F R Putri and M Firdaus

[+ Open abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012049

Yeast fuel cell: Application for desalination

Ummy Mardiana, Christophe Innocent, Marc Cretin, Buchari Buchari and Suryo Gandasasmita

[+ Open abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012050

Fabrication of organic solar cells with design blend P3HT: PCBM variation of mass ratio

Agus Supriyanto, Amrina Mustaqim, Maya Agustin, Ari H Ramelan, Suyitno, Erlyta Septa Rosa, Yofentina and Fahru Nurosyid

[+ Open abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012051

X-ray Absorption Improvement of Single Wall Carbon Nanotube through Gadolinium Encapsulation

Alimin, I Narsito, Kartini and S J Santosa

[+ Open abstract](#)[View article](#)[PDF](#)**Medicinal Chemistry****OPEN ACCESS**

012052

HLA-DP, HLA-DQ, and HLA-DR-restricted epitopes in GRA5 of *Toxoplasma gondii* strains

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



S Haryati, Y Sari, A A Prasetyo and R Sariyatun

[+ Open abstract](#)

[View article](#)

[PDF](#)

OPEN ACCESS

012053

Virtual screening of Indonesian flavonoid as neuraminidase inhibitor of influenza a subtype H5N1

A A Parikesit, B Ardiansah, D M Handayani, U S F Tambunan and D Kerami

[+ Open abstract](#)

[View article](#)

[PDF](#)

OPEN ACCESS

012054

In silico modification of Zn²⁺ binding group of suberoylanilide hydroxamic acid (SAHA) by organoselenium compounds as *Homo sapiens* class II HDAC inhibitor of cervical cancer

Usman Sumo Friend Tambunan, Ridla Bakri, Arli Aditya Parikesit, Titin Ariyani, Ratih Dyah Puspitasari and Djati Kerami

[+ Open abstract](#)

[View article](#)

[PDF](#)

OPEN ACCESS

012055

Strain-specific variations in *Toxoplasma gondii* GRA1, GRA5, GRA6, GRA8, and GRA14

S Haryati, Y Sari, A A Prasetyo and R Sariyatun

[+ Open abstract](#)

[View article](#)

[PDF](#)

Natural Product

OPEN ACCESS

012056

Screening of Marine Actinomycetes from Segara Anakan for Natural Pigment and Hydrolytic Activities

A Asnani, D Ryandini and Suwandri

[+ Open abstract](#)

[View article](#)

[PDF](#)

OPEN ACCESS

012057

Characterization of active paper packaging incorporated with ginger pulp oleoresin

T Wiastuti, L U Khasanah, W Atmaka Kawiji, G J Manuhara and R Utami

[+ Open abstract](#)

[View article](#)

[PDF](#)

OPEN ACCESS

012058

Biological evaluation of certain substituted hydantoins and benzalhydantoins against microbes

Ika-Wiani Hidayat, Yee Yee Thu, David St. C Black and Roger W Read

[+ Open abstract](#)

[View article](#)

[PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.

012059

Ethyl-2-(3,5-Dihydroxyphenol): Phloroglucinol derivatives as potential anticancer material

Triana Kusumaningsih, Maulidan Firdaus, Muhammad Widyo Wartono, Anif Nur Artanti, Desi Suci Handayani and Angga Eryanto Putro

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012060

Acidity-controlled selective oxidation of alpha-pinene, isolated from Indonesian pine's turpentine oils (*pinus merkusii*)

Masruri, Mohamad Farid Rahman and Bagus Nurkam Ramadhan

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012061

The co-pigmentation of anthocyanin isolated from mangosteen pericarp (*Garcinia Mangostana L.*) as Natural Dye for Dye- Sensitized Solar Cells (DSSC)

H Munawaroh, G F adillah, L N M Z Saputri, Q A Hanif, R Hidayat and S Wahyuningsih

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012062

Addition reaction of methyl cinnamate with 2-amino-4- nitrophenol

Venty Suryanti, Fajar Rakhman Wibowo, Pranoto, Siti Robingatus Isnaeni, Meiyanti Ratna Kumala Sari and Sekar Handayani

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012063

Isolation and identification of curcumin and bisacurone from rhizome extract of temu glenyeh (*Curcuma soloensis*. Val)

Rista A Vitasari, Fajar R Wibowo, Soerya D Marliyana and M Widyo Wartono

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012064

Physical characteristics of cinnamon oil microcapsule

R F Hermanto, L U Khasanah, Kawiji, W Atmaka, G J Manuhara and R Utami

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012065

O-methylation of natural phenolic compounds based on green chemistry using dimethyl carbonate

N I Prakoso, P H Pangestu and T D Wahyuningsih

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.

The effect of polyethylene glycol on shellac stability

012066

Khairuddin, Edi Pramono, Suryadi Budi Utomo, Viki Wulandari, W A'an Zahrotul and Francis Clegg

[+ Open abstract](#)

[View article](#)

[PDF](#)

Theory, Characterization, Modelling

OPEN ACCESS

012067

The langmuir isotherm adsorption equation: The monolayer approach

Patiha, E Heraldly, Y Hidayat and M Firdaus

[+ Open abstract](#)

[View article](#)

[PDF](#)

OPEN ACCESS

012068

Theoretical study of adsorption and dissociation of NH₃ on pentanuclear Fe(111) surface

G Purwiandono, Triyono and K Wijaya

[+ Open abstract](#)

[View article](#)

[PDF](#)

OPEN ACCESS

012069

The effects of sintering temperature on dielectric constant of Barium Titanate (BaTiO₃)

DianisaKhoirum Sandi, Agus Supriyanto, Anif, Jamaluddin and Yofentina Iriani

[+ Open abstract](#)

[View article](#)

[PDF](#)

JOURNAL LINKS

[Journal home](#)

[Journal scope](#)

[Information for organizers](#)

[Information for authors](#)

[Contact us](#)

[Reprint services from Curran Associates](#)



This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



PAPER • OPEN ACCESS

Screening of Marine Actinomycetes from Segara Anakan for Natural Pigment and Hydrolytic Activities

To cite this article: A Asnani *et al* 2016 *IOP Conf. Ser.: Mater. Sci. Eng.* **107** 012056

View the [article online](#) for updates and enhancements.

Related content

- [Selective Isolation and Phenotypic Characterization of Bacteria and Actinomycetes from Oil-contaminated Soils](#)
C A Rodríguez, T C Gaviláñez, J P Chamorro *et al.*
- [Actinomycetes of *Orthosipon stamineus* rhizosphere as producer of antibacterial compound against multidrug resistant bacteria](#)
H Rante, R Yulianty, Usmar *et al.*
- [Effect of Rhizosphere on Sediment Microbial Numbers in Phytoremediation Process of Decabromodiphenyl Ether Contaminated Sediment](#)
Liangyuan Zhao, Weijie Guo, Weihua Zhao *et al.*

Recent citations

- [The Application of Red Pigments from *Streptomyces* K-4B and Dayak Onions \(*Eleutherine palmifolia* \(L.\) Merr.\) In Colouring Glycerine Soap](#)
Herlina *et al*



IOP | ebooks™

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.

Screening of Marine Actinomycetes from Segara Anakan for Natural Pigment and Hydrolytic Activities

A Asnani^{1*}, D Ryandini², Suwandri¹

¹Deptment of Chemistry, Faculty of Mathematic and Sciences, Jendral Sudirman University, Karangwangkal, North Purwokerto, Banyumas Regency, Central Java 53122

²Department of Biology, Faculty of Biology, Jendral Sudirman University Unsoed, Puwokerto Jl. Dr. Soeparno No. 63, Central Java 53122

*E-mail: ari.asnani@unsoed.ac.id

Abstract. Marine actinomycetes have become sources of great interest to natural product chemistry due to their new chemical entities and bioactive metabolites. Since April 2010, we have screened actinobacteria from five sites that represent different ecosystems of Segara Anakan lagoon. In this present study we focus on specific isolates, K-2C which covers 1) actinomycetes identification based on morphology observation and 16S rRNA gene; 2) fermentation and isolation of pigment; 3) structure determination of pigment; and 4) hydrolytic enzymes characterization; Methodologies relevant to the studies were implemented accordingly. The results indicated that K-2C was likely *Streptomyces fradiae* strain RSU15, and the best fermentation medium should contain starch and casein with 21 days of incubation. The isolate has extracellular as well as intracellular pigments. Isolated pigments gave purple color with λ_{max} of 529.00 nm. The pigment was structurally characterized. Interestingly, *Streptomyces* K-2C was able to produce potential hydrolytic enzymes such as amylase, cellulase, protease, lipase, urease, and nitrate reductase.

1. Introduction

Actinomycetes are known as a group of microorganisms that are most responsible for the production of crucial bioactive compounds, particularly antibiotics. The potency of actinomycetes as antibiotic producing bacteria has made these microbial groups become very important. However, the bioactive compounds produced by actinomycetes are not limited against bacteria, but also effective as anti-fungal agents, anti-cancer, and some of them are as anti-virus [1]. Until now, the search of actinomycetes producing new types of bioactive compounds continues.

The research exploration of marine actinomycetes is aimed to explore the indigenous species which are able to produce potentially bioactive compounds, such as pigments, enzymes, anti-bacterial and bioremediation agents. Samples of sediment/soil and water are obtained from the mangrove region as an area of mangrove forest is a source of organic matter of weathering and decomposition of litter that is very beneficial for heterotrophic microorganism such as actinomycetes. Asnani & Ryandini have reported 26 isolates of actinomycetes halotolerant of mangrove areas Segara Anakan that have potency as sources of anti-bacterial compounds. Based on the observation of the colony and cell morphology, actinomycetes isolates generally obtained have morphological properties of actinomycetes genera *Streptomyces* [2].



Certain actinomycetes show the various color of pigments in the hyphae, and various pigment diffused into the fermentation medium such as beige, white, red, green, and brown pigment. Gunasekaran & Poorniammal have reported that *Penicillium* sp capable of producing extracellular red pigment [3]. The red pigment with the molecular formula of anthraquinone produced by *Penicillium oxalicum* has now mass produced as a natural pigment. More recently, Mohanasrinivasan et al., [4] reported that *Streptomyces coelicolor* MSIS1 able to produce the extracellular blue pigment that turns red in acidic conditions. Following these findings, we continue exploring one specific isolate, namely K-2 because of its potency to produce color pigments as well as hydrolytic enzymes. Hence, this study was aimed to identify actinomycetes producing pigment, the isolation, and structure identification of pigment, and to evaluate the hydrolytic capability of isolates.

2. Materials and Methods

2.1. Sites of Sampling

The research was exploratory. Five different sites that represent diverse ecosystems of Segara Anakan Cilacap were selected for the present study. Sediment and water samples were taken in the area of mangrove vegetation with specifications Rhizophora area sp., grassy land, and open land areas. Each location was taken three sampling points. Soil and sludge samples were taken at a depth of soil about 5-8 cm and then put into sterile bottles.

2.2. Actinomycetes Isolation

1 g of sediment or 1 mL of water was mixed with 9 mL of sterile distilled water, shaken until homogeneous and become suspension with a dilution rate of 10^{-1} . From 10^{-1} dilution, the dilution was carried out with sterile seawater to obtain a suspension with a dilution rate of 10^{-2} , 10^{-3} , 10^{-4} and 10^{-5} . The suspension with 10^{-5} dilution was pipetted into 1 mL and cultured in casein nitrate medium in a sterile petri dish. Media for cultivation was Starch Casein Nitrate (SCN) agar (10 g soluble starch; 1 g Casein; 1 g KNO_3 ; 0.5 g K_2HPO_4 ; 0.5 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$; 0.5 g NaCl ; 0.01 g $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$; 20 g agar; pH 6.8-7.0) supplemented with nystatin [2]. Incubation was carried out at 28 °C for 7-9 days, or until the actinomycetes colonies grew. The characteristics observed in actinomycetes colonies grow rounded colonies, slow growth, colony surface (could) had hyphae, hard as dusty or velvety. Usually, there was a central point in the middle of the colony and typical colored of the colony.

2.3. Colony Identification

2.3.1. Morphology Identification. The colony morphology and cell identification were done by preparing Henrici's Slide Culture (HSC). The observations of the colony were intended to the form, surface, consistency, edges, color, growth radial and concentric growth. The observation of the presence or absence of cells was directed against the substrate mycelium and aerial mycelium, the appearance of the substrate mycelium, aerial mycelium appearance, presence of spores and structure, the color of spores, as well as the pigment that diffuses into the medium. The observations of the hyphae characteristics were then compared with Bergey's Manual of Determinative Bacteriology.

2.3.2. Species Identification. Identification of actinomycetes species was based on *16S rRNA*. DNA isolation used Presto™ Mini gDNA Bacteria Kit Geneaid Instruction Manual. DNA obtained was amplified with 27F and 1492R primers using KAPA Taq™ Extra Host start ReadyMix with dye. DNA electrophoresis used 1% of agarose gel submerged in 1x TAE buffer. Species determination used BLAST (Basic Local Alignment Search Tool).

2.4. Pigment Production.

The pigment production consisted of four stages, namely a) inoculum preparation, b) fermentation medium preparation, c) the pigment production, d) pigment isolation, and e) pigment extraction.

Actinomycetes isolated was a streak on SCN medium and incubated for 8 days at a temperature of 30 °C; then it served as the parent culture. Pigment production was done by inoculating five plugs sized of 0.5 mm from parent culture to 100 ml of liquid medium. The fermentation process was done for 14 to 21 days at room temperature. Media for fermentation was SCN broth (10 g soluble starch; 1 g Casein; 1 g KNO₃; 0.5 g K₂HPO₄; 0.5 g MgSO₄·7H₂O; 0.5 g NaCl; 0.01 g FeSO₄·7H₂O; pH 6.8-7.0) supplemented with nystatin. After incubation, the culture producing pigment was centrifuged at 3000 rpm for 30 minutes. The filtrate obtained was an extracellular pigment; while the mycelium obtained was extracted with ethanol to obtain the intracellular pigment [5].

2.5. Structure Determination of Pigment

Pigment extract obtained was purified with silica gel 60 GF254 followed by column chromatography eluted with chloroform: methanol (9: 1). The pure pigment was characterized by UV-Vis spectrophotometry (Shimadzu UV-1800), FTIR spectroscopy (Perkin Elmer Spectrum 100), and NMR Spectrometer (NMR Jeol 500 MHz).

2.6. Hydrolytic Enzymes Characterization

The physiology of the isolates was observed by a series of biochemical tests following the known procedure [6]. Biochemical tests included hydrolytic, enzymatic tests such as starch hydrolysis, CMC hydrolysis, casein hydrolysis, lipid hydrolysis, urea hydrolysis, and nitrate reduction.

3. Results and Discussion

Sediment and water samples were collected from Klaces (K) and Ujung Alang (U) areas in Segara Anakan. Sampling time was during the day with the local temperature of 28 °C and pH waters around 7-8 which corresponded to the growth of actinomycetes such as the temperature of 25-30 °C and a pH of about neutral pH. Actinomycetes cultivation was carried out on an agar medium Starch-casein nitrate (SCN) supplemented with antifungal nystatin to inhibit mold growth. SCN is a highly sensitive and specific medium for the growth of actinomycetes. Usha et al. managed to isolate 15 isolates of actinomycetes origin by using medium SCN soil [7]. Khan et al. also succeeded in isolating two actinomycetes isolate originating from the sponge using SCN medium supplemented with antifungal and antibacterial [8].

The population of actinomycetes which had colored mycelium was generally obtained from the litter under the mangrove vegetation [2]. This indicates that the litter of mangrove vegetation litter is rich in organic matter because the result of weathering leaf or root exudates. As estuarine areas, Segara Anakan region experiences water movement that causes a typical condition. Only microorganisms with high tolerance to a constantly changing of oxygen, temperature, salinity and humidity are able to survive.




Figure 1. Colony observation of actinomycetes K-2C: (a) top (b) bottom, (c) mycelium

3.1. Colony Identification

Actinomycetes are a group of bacteria that forms unicellular mycelium that has the appearance of morphology such as mold. Unicellular mycelium is formed consisting of hyphae branched hyphae with a diameter of 0.5-0.8 μm . There are two types of mycelium, the substrate mycelium (vegetative) growth in the medium and aerial mycelium (aerial) growing on the surface of the medium. Aerial hyphae can be white, gray, violet, red, yellow, brown, green or any other color due to the pigmentation. In observation with a light microscope on a glass slide, aerial mycelium was refractive and bright, while the substrate mycelium is transparent and dark [1].

Based on morphology observation, the colony of K-2C has a size of 6 mm, red pigmentation to medium, irregular shape, powdery surface, convex elevation, margin undulate, gray aerial mycelium, and brown substrate mycelium; Morphological properties of K-2C were suggested to be similar with actinomycetes genera *Streptomyces*. Vegetative hyphae form a mycelium was branched. Aerial hyphae forming three or more mature spore chains, and a spiral-shaped sectional existed (figure 1).

Determination of the species of the isolate K-2C was done using 16S rRNA. The isolated DNA was amplified using PCR and sequenced in two-way forward and reverse with 27F and 1492R primers to get the base sequence. Determination of the base sequence of K-2C was carried out by pairwise alignment method in Bioedit program. The results of Bioedit program is as follows:

```
CATGCAGTCGACGATGAACCACCTTCGGGTGGGGATTAGTGGCGAACGGGTGAGTAACA
CGTGGGCAATCTGCCCTGCACTCTGGGACAAGCCCTGGAAACGGGGTCTAATACCGGATA
CTGACCTGCCAAGGCATCTTGGCGGGTCGAAAGCTCCGGCGGTGCAGGATGAGCCCGCG
GCCTATCAGCTTGTTGGTGAGGTAATGGCTCACCAAGGCGACGACGGGTAGCCGGCCTGA
GAGGGCGACCGGCCACACTGGGACTGAGACACGGCCACACTCCTACGGGAGGCAGCAG
TGGGGAATATTGCACAATGGGCGCAAGCCTGATGCAGCGACGCCGCGTGAGGGATGACG
GCCTTCGGGTTGTAAACCTCTTTTCAGCAGGGAAGAAGCGAAAGTGACGGTACCTGCAGA
AGAAGCGCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGCGCAAGCGTTGT
CCGGAATTATTGGGCGTAAAGAGCTCGTAGGCGGCTTGTTCGCGTCGGTTGTGAAAGCCCC
GGGCTTAACCCCGGGTCTGCAGTCGATACGGGCAGGCTAGAGTTCGGTAGGGGAGATCG
GAATTCTTGGTGTAGCGGTGAAATGCGCAGATATCANGAGGAACACCGGTGGCGAAGGC
GGATCTCTGGGCCGATACTGACGCTGAGGAGCGAAAGCGTGTGGAGCGAACAGGATTAG
ATACCCTGGTAGTCCACGCCGTAAACGGTGGGCACTAGGTGTGGGCAACATTCCACGTTG
TCCGTGCCGCAGCTAACGCATTAAGTGCCCCGCTGGGGAGTACGGCCGCAAGGCTAAAC
TCAA
```

Total nucleotides	= 834 base pairs
Molecular weight	= 253152.00 Daltons, single stranded
Molecular	= 508419.00 Daltons, double stranded
Composition G+C	= 59.83%
Composition A+T	= 40.17%

Based on the BLAST analysis, isolate K-2C was identified to have similar sequences of 99% with a *Streptomyces fradiae* strain RSU15. Hangstrom et al., stated that isolates which have similarities sequences of the 16S rRNA gene could represent the same species while the sequence similarities between 93% -97% could represent the same identity at the level of genus, but different at the species level [9]. In other words, the difference between the 16S rRNA gene sequences of isolates stored in the data center if it is less than 3% showed the same species, while if more than 3% indicates a

different species from the data center. From BLAST searches, K-2C taxonomy can be seen, as follows:

Domain: Bacteria
Phylum: Actinobacteria
Class: Actinomycetes
Ordo: Actinomycetales
Familia: Streptomycetaceae
Genus: Streptomyces
Species: *Streptomyces fradiae* strain RSU15

3.2. Pigment of *Actinomycetes* K-2C

Aside from being able to produce antibiotics, actinomycetes are also able to produce the pigment that includes all rainbow colors. The bioactive pigments produced by microbes halotolerant has been reviewed. Those pigments have specific color and have an activity, such as anti-cancer, antibacterial, anti-inflammatory, antioxidant, as well as antibiotics. Asnani & Ryandini have also been reported marine actinomycetes expressing the color pigment. Observation of the pigment was observed on colony color and pigmentation to the substrate [5].

The microbes pigment can be isolated from three general stages, namely the cultivation of microbes, microbial fermentation to produce pigment, and pigment isolation. Measurement of the pigment concentration can be carried out using a UV spectrophotometer at the optimum wavelength (λ) for a particular color. For example, the yellow pigment can be observed on λ 390 nm, while the red pigment on λ 500 nm [10]. In this study, *Streptomyces* K-2 C produced purple pigment within 21 days of fermentation time, with optimum λ at 529.00 nm (figure 2). The wavelength in the range of 500 – 560 nm is the wavelength that can be absorbed by red and purple colors [11].




Figure 2. Pigment production from actinomycetes K-2C: (a) purple pigment, (b) optimum λ at 529.00 nm

The highest production of purple pigment was obtained with fermentation with modified casein nitrate medium, and the lowest production was obtained from that of yeast extract-malt extract medium. Microbial fermentation medium is used for microbial growth as well as for the pigment production. Optimization of the fermentation medium can be done by varying sources of carbon and nitrogen sources. Palanichamy et.al., reported that antibiotics actinorhodin red-bluish can be isolated from *Streptomyces* sp in the fermentation medium using glycerol and mannose as a source of C; and Na-casein peptone as a source of N [12]. Optimization of fermentation medium in order for *Penicillium* sp to excrete extracellular red pigment was using soluble starch as a source of C and peptone as a source of N, at a temperature of 30 °C and pH 9.0. The optimum condition of fermentation medium was able to increase the production of red pigment up to sevenfold [3].

FTIR analysis was conducted to determine the functional groups of isolated pigment. Infrared spectra of purple pigment were shown in Figure 3 while the data of wave number and related functional groups was presented in table 1. Interpretation of IR spectra indicated that *Streptomyces* K-2C pigment contained a cyclic core, carbonyl, hydroxyl, alkyl (CH₃ and CH₂), non-conjugated one and secondary NH.




Figure 3. IR spectrum of pigment from *Streptomyces* K-2C

Table 1. FTIR data of purple pigment

Wavenumber (cm ⁻¹)	Frequency range (cm ⁻¹)	Correlated functional groups
2853.22	2850-3000	C-H alkane
1742.26	1650-1780	C=O ester
1634.77	1620-1680	C=C non conjugated
1577.31	1550-1640	N-H secondary
1463.76	1450-1470	-CH ₂ -
1377.20	1350-1370	-CH ₃
1169.80	1000-1300	C-O ester
1040.27	990-1060	C-OH cyclic
721.22	650-1000	=C-H

¹H-NMR spectra analysis was done to describe different types of hydrogen atoms in the pigment molecule. The information that can be obtained from the ¹H-NMR spectrum is the chemical environment of hydrogen atoms, the number of hydrogen atoms in each environment and the cluster structures adjacent to each hydrogen atom [13]. ¹H-NMR spectrum of the pigment *Streptomyces* K-2C is shown in Figure 4.

¹H-NMR spectrum of the pigment *Streptomyces* K-2C showed a number of signals that appear in the chemical shift (δ) at 0.5 to 5.5 ppm. These data indicated aliphatic compounds. A chemical shift at 0.5 to 2.0 ppm was the signal for the H atoms bonded to C in a sp³ manner. Chemical shift at 0.87

ppm (3H) indicated the H atoms as $-\text{CH}_3$; $\delta = 1.29$ ppm (2H) indicated the H atoms as $-\text{CH}_2$; and $\delta = 1.62$ ppm (2H) indicated the H atoms as $-\text{CH}_2$ as well. Chemical shift at 2.34 ppm (2H) was a signal of H atom $-\text{CH}_2$ bonded to C heteroatoms. Chemical shift at 3.72 ppm (1H) indicated the H atoms as a form of $-\text{CH}$ bonded to C heteroatoms. Chemical shift at 5.35 ppm (1H) is a signal to H bound to C double bond ($-\text{C}=\text{H}$). Based on the interpretation of the ^1H -NMR spectrum, the purple pigment has H atoms in the form of methyl and methylene bonded to C Sp^3 ; H form methylene attached to the C-heteroatom; Metin H form attached to the C-heteroatom; and H attached to C double bond.



Figure 4. ^1H -NMR spectrum of pigment from *Streptomyces* K-2C


3.3. Hydrolytic Enzyme Characteristics

The other actinomycetes potency is its ability to produce hydrolytic enzymes. The results from biochemical tests revealed the ability of *Streptomyces* K-2C in hydrolyzing starch, casein, lipids, urea, cellulose and reduced nitrate. These positive results indicated that the isolate were able to produce hydrolytic enzymes such as amylase, cellulase, protease, lipase, urease and nitrate reductase (table 2).

The further quantitative study indicated that the cellulolytic ability of K-2C was higher (cellulase index of 20.06 mm) compared to the amylolytic ability (amylase index of 6.26 mm) (figure 5). Indeed, the areas of mangrove forest are sources of organic matter of weathering and decomposition of litter that is very beneficial for heterotrophic microorganism such as actinomycetes. Thus, marine actinomycetes obtained from mangrove areas in Segara Anakan, such as *Streptomyces* K-2C, allegedly able to act as decomposers organic materials. In this regard, the hydrolytic ability of *Streptomyces* K-2C will have a promising prospect for industrial application.

Table 2. Biochemical Tests of *Streptomyces* K-2C

	Results
Enzyme Production:	
- Amylase	+
- Protease	+
- Lipase	+
- Cellulase	+
- Nitrate reductase	+
Carbohydrate Utilization	
- Galactose	-
- Sucrose	+
- Raffinose	+
- Mannitol	+
- Arabinose	+
- Xylose	-



Starch hydrolysis (6,16mm) CMC hydrolysis (20,06mm)
Figure 5. Amylolytic and cellulolytic ability of *Streptomyces* K-2C

4. Conclusion

Marine actinomycetes K-2C was identified as *Streptomyces fradiae* strain RSU15. The isolate was able to produce purple pigment with λ_{max} of 529.00 nm within 21 days of fermentation. Partial structure of the pigment was identified using FTIR and NMR. However further characterization is needed to have a complete pigment structure. Interestingly, the isolate was also able to produce potential hydrolytic enzymes namely amylase, cellulase, protease, lipase, urease, and nitrate reductase.

Acknowledgement

The authors are grateful to Indonesian Higher Education for providing research grant under *Strategies National* scheme in 2015. The authors also want to appreciate Nurhasanah for her work on pigment purification.

References

- [1] Miyadoh S 1997 *Atlas of Actinomycetes*. Asakura Publishing Co. Ltd. Japan.
- [2] Asnani A, and Ryandini D 2011 Screening of marine actinomycetes from Segara Anakan Indonesia for antimicrobial activity Proceeding ICONS 2011 Shaker Verlag Germany ISBN 978-3-8440-1403-7 ISSN 1434-5536.
- [3] Gunasekaran S, and Poorniammal R 2008 Optimization of fermentation conditions for red pigment production from *Penicillium* sp. under submerged cultivation *Af. J. of Biotech.* **7** (12) pp 1894-1898.

- [4] Mohanasrinivasan V, SriramKalyan P, Nandi I, Subathradevi C, Selvarajan E, Suganthi V, and Jemimah N S 2013 Fermentative production of extracellular pigment from *Streptomyces coelicolor* MSIS1 Res. J. Biotec. **8** (4) pp. 31-41.
- [5] Asnani A, Ryandini D, and Suwandri 2014 Analysis of microbial pigment productivity in *Streptomyces* sp isolated from Segara Anakan. Joint Indonesia-UK Conference of Organic and Natural Product Chemistry, UGM, Yogyakarta.
- [6] Asnani A, Ryandini D, and Suwandri 2014 Analisis potensi amilolitik dan selulolitik dari isolat aktinomisetes laut Seminar Nasional Percepatan Desa Berdikari Melalui Pemberdayaan Masyarakat Dan Inovasi Teknologi LPPM Unsoed Prosiding ISBN: 978-602-1643-13.
- [7] Usha R, Ananthaselvi P, Venil C.K, and Palaniswamy M, 2010 Antimicrobial and antiangiogenesis activity of *Streptomyces parvulus* KUAP106 from mangrove soil *European J. of Biol. Sci.* **2**(4) 77-83.
- [8] Khan S.T, Motohashi K, Kozonel, Takagi M, and Shin-ya K, 2010 *Streptomyces* associated with a marine sponge *Haliclona* sp; biosynthetic genes for secondary metabolites and product. *Environ. Microbiol.* **13**(2) 391-403.
- [9] Hagstrom A, Pinhassi J, and Zweifel U.L, 2000 Biogeographical diversity among marine bacterioplankton. *Aquatic Microbial Ecology* **21**(3) 231-244.
- [10] Kasim E, Astuti S, Nurhidayat N, 2005 Karakterisasi pigmen dan kadar lovastatin beberapa isolat *Monascus purpureus* *Biodiversitas* **6** (4) hal. 245-247.
- [11] Sahin N, 2011, Significance of absorption spectra for the chemotaxonomic characterization of pigmented bacteria *Turk. J. Biol.* **35** pp 167-175.
- [12] Palanichamy V, Hundet A, Mitra B and Reddy N, 2011 Optimization of cultivation parameters for growth and pigment production by *Streptomyces* spp. isolated from marine sediment and rhizosphere soil *Int. J. of Plant and Animal Environm. Sci* **1** (3) pp 158-170.
- [13] Juliana V.A, Aisyah S, Mustapha I, 2010 Isolasi dan Karakterisasi Senyawa Turunan Terpenoid dari Fraksi n-Heksan *Momordica charantia* L. *J. Sains dan Tek. Kim.* **1**(1) hal. 88-93.

Full paper submission: Ari Asnani

2 messages

Ari Asnani <asnani@gmail.com>
To: Chemistry Uns <jccuns@gmail.com>

Mon, Sep 14, 2015 at 1:20 PM

Dear Organizing Committee

We have sent our full paper on Sept 1st, both via email and online.
However, I sent them back again as the attachment bellow.

Thank you very much for your kind consideration,

Best Wishes,
Ari Asnani,
Kimia-Unsoed

----- Forwarded message -----

From: Ari Asnani <asnani@gmail.com>
Date: Tue, Sep 1, 2015 at 9:12 AM
Subject: Full paper submission: Ari Asnani
To: jccuns@gmail.com

Dear. The Committee
10th JCC 2015

Attached is our full paper.
We really apologize for the very late submission.
Thank you very much for your kind consideration.

with kind regards
Ari Asnani

2 attachments

 **Ari Asnani.pdf**
301K

 **Ari Asnani.doc**
5019K

10th JCC 2015 <jccuns@gmail.com>
To: Ari Asnani <asnani@gmail.com>

Wed, Sep 16, 2015 at 1:30 PM

Dear Ari Asnani,

Many thanks for submitting your full paper. We have received them and will proceed accordingly.

Best wishes,
JCC-10 Organizer
[Quoted text hidden]

Words Version RE: 10JCC notification for paper 47

1 message

Ari Asnani <ari.asnani@unsoed.ac.id>

Sun, Nov 22, 2015 at 10:21 PM

To: 10JCC <10jcc@easychair.org>, "fitria@mipa.uns.ac.id" <fitria@mipa.uns.ac.id>

Cc: "asnani@gmail.com" <asnani@gmail.com>

Dear Editor,

Attached is our Word version from our article (paper 47).
We really apologize for not sending it before the date-line.

Thank you very much for your kind consideration.

Best Wishes,
Ari Asnani,
Chemistry Department
UNSOED Purwokerto

-----Original message-----

From: 10JCC <10jcc@easychair.org>**Sent:** Tuesday 10th November 2015 12:20**To:** Ari Asnani <ari.asnani@unsoed.ac.id>**Subject:** 10JCC notification for paper 47

Dear Prof./Dr. A Asnani,

Based on the reports of the referees, the Editors have determined that your manuscript:

Number: 47

Title: Screening of Marine Actinomycetes from Segara Anakan Cilacap for Natural Pigment and Hydrolytic Activities

is accepted and will be published in IOP series: Material Science and Engineering.

For layouting purpose, please send the WORD version of your accepted article. Please note that the deadline of submission will be Nov 16, 2015.

If you have any questions or wish to comment on your revisions, please E-mail the Editor.

Best regards,
Editors

Fitria Rahmawati (Email: fitria@mipa.uns.ac.id)Teguh Endah Saraswati (Email: teguh@mipa.uns.ac.id)10th JCC 2015 website: <http://jcc.mipa.uns.ac.id/>**Ari Asnani.doc**

5202K