

PAPER • OPEN ACCESS

Scanning SNPs of Diabetes Mellitus related genes; HNF4A, PTPN, KCNJ11, PPAR gamma; among Thalassemia Patients: a Preliminary Study

To cite this article: L Rujito *et al* 2019 *IOP Conf. Ser.: Earth Environ. Sci.* **255** 012008

View the [article online](#) for updates and enhancements.



IOP | ebooks™

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.

Scanning SNPs of Diabetes Mellitus related genes; HNF4A, PTPN, KCNJ11, PPAR gamma; among Thalassemia Patients: a Preliminary Study

L Rujito¹, F Fauziyah², E F Azizah³, QSantosa⁴, A T Hapsari⁵, D U Anjarwati⁶, F Arjadi⁷

- ^{1.} Department of Molecular Biology, Faculty of Medicine, Universitas Jenderal Soedirman, Purwokerto, email : l.rujito@unsoed.ac.id
- ^{2.} Department of Molecular Biology, Faculty of Medicine, Universitas Jenderal Soedirman, Purwokerto, email : fauziyahfrida@gmail.com
- ^{3.} Department of Molecular Biology, Faculty of Medicine, Universitas Jenderal Soedirman, Purwokerto, email: azizahesaa@gmail.com
- ^{4.} Department of Pediatrics, Faculty of Medicine, Universitas Jenderal Soedirman, Purwokerto, email : qodri.santosa@unsoed.ac.id
- ^{5.} Department of Pediatrics, Faculty of Medicine, Universitas Jenderal Soedirman, Purwokerto, email : ariadne.hapsari@unsoed.ac.id
- ^{6.} Department of Microbiology, Faculty of Medicine, Universitas Jenderal Soedirman, Purwokerto, email : dwi.utami.anjarwati@unsoed.ac.id
- ^{7.} Department of Anatomy, Faculty of Medicine, Universitas Jenderal Soedirman, Purwokerto, email : fitranto.arjadi@unsoed.ac.id

Email : l.rujito@unsoed.ac.id. Telp : 62281622022, Fax : 62281624990

Abstract. Thalassemia is one of health problem which is a burden on the Indonesian government. Thalassemia problems include wide-spreading, and clinical conditions, therapy management, and complications. One of the significant difficulties that cause mortality is due to Diabetes Mellitus (DM). This preliminary study aimed to explore the genotyping of SNPs on the genes related with DM in thalassemia patients in Indonesia. The study used a cross-sectional design involving 103 samples from stored DNA patients in Banyumas. There were four genes about DM; HNF4A, PTPN, KCNJ11, and PPAR gamma; which characterized using the PCR-RFLP based technique with appropriate restriction enzymes. The results showed that there was existence of SNP related to diabetes mellitus in thalassemia patients. The T130I (rs1800961) of HNF4A gene has a proportion of genotypes 79% CC, 16% CT, and 5% TT with minor allele frequency (MAF) 13%. The genotype distribution of PTPN gene 1023 C>A were 68% for CC, 30% CA, 2% AA and MAF 17%. PTPN gene of 467 T>C has percentage of CC 90%, CT 10%, and MAF 5%. The KCNJ11 gene at position rs 5219 had a portion of glutamic acid 46% and lysine 54%. Whereas, Pro12Ala (rs18012282) of PPAR gamma; gene has a genotype proportion of CC 94%, CG 5% , GG 1% and MAF 4%. The genotyping of DNA patients revealed the presence of DM-related genes in thalassemia. The coexistence of mutants may relate with the clinical condition. Further study, especially in genotype-phenotype relation is essential to explore the influence of the mutants in worsening the affected people.

Keywords: thalassemia, diabetes mellitus, HNF4A, PTPN, KCNJ11, PPAR gamma



1. Introduction

Thalassemia is a monogenic hematology disease that still the most endemic on a global, regional and national scale. The Southeast Asian population is reported to have a carrier frequency of Hemoglobinopathy and Thalassemia of 45.5% with 1.34 children of 1000 births born with moderate to severe clinical conditions [1]. In Indonesia, the number of the carrier has an average frequency of 3-10% of overall properties [2]. Thalassemia problems are very widely dispersed, ranging from the number of mutations involved in the pathogenesis of the disease, the large number of budgets in management, to the complexity of prevention problems. At present, there are at least more than 800 variations in the globin genes responsible to the pathogenesis of hemoglobinopathy and thalassemia [3], with 300 of them being β thalassemia mutations [4]. Diabetes Mellitus (DM) is the highest hallmark of death due to endocrine abnormalities in thalassemia patients [5]. Efforts to detect prediabetic conditions are an essential part of the management system of thalassemia patients so that the patient's morbidity and mortality can be adequately controlled. The diagnosis of pre-condition is closely related to individual genetic susceptibility. Previous thalassemia studies showed that with knowledge of gene modifier *Xmn1* and *BCL11A* could classify patients who require routine and non-regular transfusions [6]. which have an impact on the better management of thalassemia. The purpose of this study is to identify and characterize genetic markers that underlie the complications of thalassemia patients in endocrine (DM) cases. Currently, the genetic marker and bio panel markers for DM complications in thalassemia are still very limited, so research and data need to be done to reveal more evidence of the genetic potential of these markers. Genetic markers in question are *HNF4A*; the genes found in the linkage study, Peroxisome proliferator-activated gamma receptor (*PPAR γ*), SNPs in the *PTPN1* gene and *KCNJ11* [7-10].

2. Materials and Methods

The study used a cross-sectional design with a whole subject of 100 thalassemia major patients. Patient DNA was tested using PCR-RFLP method to determine the genotype of *Hnf4a* (T130I rs1800961), *Ptpn* 1023 C> A and *PTPN* 467 T> C, *KCNJ11* E23K (rs5219), *PPAR* Pro12Ala (rs18012282). Table 1 provides the primers and enzymes used to determine genotyping of these genes.

Table 1. Primers and Enzyms used in PCR-RFLP Experiment for Genotyping of Genes Related with Diabetes melitus.

Gene	Primers		enzim restriksi
	Forward 5'-3'	Reverse 5'-3'	
<i>Hnf4a</i>	CCACCCCCTACTCCATCCCTGT	CCCTCCCGTCAGCTGCTCCA	BsmI
<i>Ptpn</i> 1023	TAGCAGAAACCGAGTTTCACC	CCTGGGTAACAGAATCAGACC	BclI
<i>PTPN</i> 467	TTCATTCTGCAGCACCC AAG	GTTGAGTCACAGAGTGAGTGG	AvaI
<i>KCNJ11</i>	GACTCTGCAGTGAGGCCCTA	ACGTTGCAGTTGCCTTTCTT	BanII
<i>PPAR</i>	CCAATTCAAGCCCAGTCCTTTC	CAGTGAAGGAATCGCTTTCCG	BstU-I

PCR method used serial temperature in the following steps: Initial denaturation 95⁰ C for 1 minute, and followed by 30 cycles for denaturation 95⁰ C '30 seconds, annealing temperature 53-57⁰ C 30 seconds, elongation temperature 72⁰ C for 30 seconds, closed with final elongation temperature 72⁰ C for 30 2 minutes. Genotypic data was then tabulated to obtain minor allele frequencies and genotype distribution.

3. Results and Discussion

A total of 100 DNA samples from major thalassemia patients underwent a genotyping process which showed in Table 2.

Table 2. Genotyping of The Genes Related Diabetes Melitus ; HNF4A, PTPN, PPAR- γ , KCNJ11, among Indonesia patients.

Gene name	SNP locus	Genotype			Allele			HWE (χ^2)*
		base	frequency	Proportion	base	frequency	Proportion	
HNF4A	rs1800961	C/C	81	0.79	C	178	0.87	0.0093
		C/T	18	0.16	T	26	0.13	
		T/T	5	0.05				
PTPN 1023	-1023 C>A SNP	C/C	70	0.69	C	170	0.83	0.73
		A/C	30	0.29	A	34	0.17	
		A/A	2	0.002				
PTPN 467	-467 T>C SNP	C/C	92	0.9	C	194	0.95	1
		C/T	10	0.1	T	10	0.05	
		T/T						
PPAR- γ	rs1801282	C/C	96	0.94	C	197	0.97	0.1
		C/G	5	0.05	G	7	0.03	
		G/G	1	0.01				
KCNJ11	rs5219	K/K	29	0.28	E	93	0.46	0.69
		E/K	53	0.52	K	111	0.54	
		E/E	20	0.2				

Table 1 shows the distribution of minor HNF4A alleles by 13% with Hardy Weinberg Equilibrium (HWE) 0.009. Minor T alleles in HNF4A appear to deviate from allele distribution in other Asian populations. Data on the SNPS database shows that Asian people are in the range of 2-4%, but this population is reaching 13%. Data also confirmed by the HWE value in the non-equilibrium state (<0.05). The fact that some patients were coming from one descendant and Banyumas were in peripheral areas with local marriages, allowed the effect of an isolated gene. It caused the spreading of the alleles proportion is not in Equilibrium state. Percentage of T130I HNF4A polymorphism in thalassemia populations in Banyumas in the form of 16% heterozygotes and 5% homozygous mutants. T130I HNF4A polymorphism study by Zhu and colleagues in Japan obtained as much as 3.5% mutations in subjects with type 2 diabetes mellitus and 0.8% in non-diabetic subjects [11]. There were also 3 people suspected of MODY diabetes mellitus patients in Spain obtained results that the three patients were heterozygous T130I HNF4A polymorphisms and only one patient correlated with MODY [12]. The families of the three patients were measured for parameters of glucose metabolism and clinical symptoms leading to the T130I HNF4A polymorphism. The results showed that 74% of close family patients were heterozygous from the T130I polymorphism. Subjects with heterozygous T130I polymorphisms were diagnosed as diabetes mellitus by 50%, prediabetes by 21% and 29% as normoglycemia.

Based on the results of the study in table 1, the proportion of PTPN1 -1023C> A polymorphisms in the thalassemia population in Banyumas is heterozygous at 29% and homozygous mutants at 2%.

Allele C was obtained at 81% and A allele as much as 19%. The results of this study are in line with the previous study of Meshkani in Iran showed a proportion of heterozygotes in the polymorphism -1023C> A in the group of diabetic patients as much as 13.2% and the proportion of C alleles as much as 93.4% and A alleles as much as 6.6% [13]. Another study reported that the two groups of subjects who got the C allele results were more prevalent in the group of diabetic patients (93%) than the control group (82%). In addition, the study showed that there were significant differences in the parameters of insulin sensitivity, signs of obesity, hypertension and several metabolic parameters such as lipid profiles between groups of diabetic patients and the control group [14].

Population diversity data of PTPN1 -1023C> A polymorphism published by the National Center of Biotechnology Information (2018) states that the global MAF (minor allele frequency) allele A is 0.1288 / 645. The proportion of A allele in 5 large populations is the EAS (East Asian) population of 9%, the SAS (South Asian) population is 22%, the EUR (European) population is 8.5%, the AMR (Ad Mixed American) population is 6.7% and AFR (African) by 16%. The similarity between the proportions of the SAS population (population of Pakistan, India, Bangladesh) and thalassemia patients in Banyumas is possible because of racial similarity in the two populations due to the process of trade and migration from the Mongoloid race to the Indonesian mainland. The polymorphism in the PTPN gene at both locus 1023 and 467 shows the same value as the Asian population in general. This polymorphism at the locus 1023 and 467 in the PTPN gene is an important note because this PTPN functions at the formation of the protein tyrosine phosphatase-1B which has negative feedback on insulin release and also has an inverse effect on leptin [15]. The presence of this polymorphism can cause changes in insulin regulation in thalassemia patients, which may result in a decrease in the function of insulin in addition to the result of excessive iron deposits.

PPAR- γ and KCNJ11 gene polymorphism detection showed a positive value in 6 people for PPAR gamma and more than 50 people had a constitution of Proline or Alanine proteins at position 23 or rs5219. The presence of mutants in these genes indicates that thalassemia patients can experience a state of diabetes mellitus because the mutants of these genes can also exacerbate an iron deposit disorder in the pancreas which interferes with insulin action. PPAR- γ and KCNJ11 polymorphisms and previous genes can disrupt and accelerate the incidence of diabetes mellitus in thalassemia patients [16, 17]. Scanning and following up on these genes are, therefore, something necessary for clinical consideration.

4. Conclusion

There is some mutant locus in the Diabetes Mellitus related genes of HNF4A, PTPN, KCNJ11, and PPAR gamma among Thalassemia Patients in Banyumas. Further study must be addressed to elucidate the mechanism of these genes on Insulin metabolism.

References.

- [1] De Sanctis V, Kattamis C, Canatan D, Soliman A T, Elsedfy H, Karimi M, Daar S, Wali Y, Yassin M, Soliman N, Sobti P, Al Jaouni S, El Kholy M, Fiscina B and Angastiniotis M 2017 β -Thalassemia Distribution in the Old World: an Ancient Disease Seen from a Historical Standpoint *Mediterranean journal of hematology and infectious diseases* **9** e2017018-e
- [2] Rujito L, Basalamah M, Mulatsih S and Sofro A S M 2015 Molecular Scanning of β -Thalassemia in the Southern Region of Central Java, Indonesia; a Step Towards a Local Prevention Program *Hemoglobin* **39** 330-3
- [3] Giardine B, Borg J, Viennas E, Pavlidis C, Moradkhani K, Joly P, Bartsakoulia M, Riemer C, Miller W, Tzimas G, Wajcman H, Hardison R C and Patrinos G P 2014 Updates of the HbVar database of human hemoglobin variants and thalassemia mutations *Nucleic acids research* **42** D1063-D9
- [4] Sumera A, Radhakrishnan A, Baba A A and George E 2015 Review: Beta-thalassemia and molecular chaperones *Blood Cells, Molecules, and Diseases* **54** 348-52

- [5] Li M-J, Peng S S-F, Lu M-Y, Chang H-H, Yang Y-L, Jou S-T, Lin D-T and Lin K-H 2014 Diabetes mellitus in patients with thalassemia major *Pediatric blood & cancer* **61** 20-4
- [6] Rujito L, Basalamah M, Siswandari W, Setyono J, Wulandari G, Mulatsih S, Sofro A S M, Sadewa A H and Sutaryo S 2016 Modifying effect of XmnI, BCL11A, and HBS1L-MYB on clinical appearances: A study on β -thalassemia and hemoglobin E/ β -thalassemia patients in Indonesia *Hematology/Oncology and Stem Cell Therapy* **9** 55-63
- [7] Zyriax B-C, Salazar R, Hoepfner W, Vettorazzi E, Herder C and Windler E 2013 The Association of Genetic Markers for Type 2 Diabetes with Prediabetic Status - Cross-Sectional Data of a Diabetes Prevention Trial *PloS one* **8** e75807
- [8] Garca-Chapa E, Guadalupe, Leal-Ugarte E, Peralta-Leal V, Durn-Gonzalez J and Meza-Espinoza J P 2017 Genetic Epidemiology of Type 2 Diabetes in Mexican Mestizos *BioMed Research International* **2017** 10
- [9] Lall K, Magi R, Morris A, Metspalu A and Fischer K 2017 Personalized risk prediction for type 2 diabetes: the potential of genetic risk scores *Genet Med* **19** 322-9
- [10] Karachanak-Yankova S, Dimova R, Nikolova D, Nesheva D, Koprinarova M, Maslyankov S, Tafradjiska R, Gateva P, Velizarova M, Hammoudeh Z, Stoynev N, Toncheva D, Tankova T and Dimova I 2015 Epigenetic alterations in patients with type 2 diabetes mellitus *Balkan Journal of Medical Genetics : BJMG* **18** 15-24
- [11] Zhu Q, Yamagata K, Miura A, Shihara N, Horikawa Y, Takeda J, Miyagawa J and Matsuzawa Y 2003 T130I mutation in HNF-4 α gene is a loss-of-function mutation in hepatocytes and is associated with late-onset Type 2 diabetes mellitus in Japanese subjects *Diabetologia* **46** 567-73
- [12] Anık A, Çatlı G, Abacı A and Böber E 2015 Maturity-onset diabetes of the young (MODY): an update. In: *Journal of Pediatric Endocrinology and Metabolism*, p 251
- [13] Meshkani R, Taghikhani M, Mosapour A, Larijani B, Khatami S, Khoshbin E, Ahmadvand D, Saeidi P, Maleki A, Yavari K, Nasoohi N and Adeli K 2007 1484insG Polymorphism of the PTPN1 Gene Is Associated with Insulin Resistance in an Iranian Population *Archives of Medical Research* **38** 556-62
- [14] Mohamed Abdulaziz Al D, Asirvatham Alwin R, Rim B, Ayman Abdallah Al H, Abdulghani Al S, Rania Ahmed A and Fahad Sulaiman Al S 2016 Diabetes Mellitus in Saudi Arabia: A Review of the Recent Literature *Current Diabetes Reviews* **12** 359-68
- [15] Gurzov E N, Stanley W J, Brodnicki T C and Thomas H E 2015 Protein tyrosine phosphatases: molecular switches in metabolism and diabetes *Trends in Endocrinology & Metabolism* **26** 30-9
- [16] Haghvirdizadeh P, Mohamed Z, Abdullah N A, Haghvirdizadeh P, Haerian M S and Haerian B S 2015 KCNJ11: Genetic Polymorphisms and Risk of Diabetes Mellitus *Journal of diabetes research* **2015** 908152-
- [17] Kim H-S, Hwang Y-C, Koo S-H, Park K S, Lee M-S, Kim K-W and Lee M-K 2013 PPAR- γ activation increases insulin secretion through the up-regulation of the free fatty acid receptor GPR40 in pancreatic β -cells *PloS one* **8** e50128-e