


Jurnal Ilmiah Kimia

MOLEKUL

Volume 17, Issue 1, March 2022



Indexed by Scopus

Department of Chemistry
Faculty of Mathematics and Natural Sciences
Universitas Jenderal Soedirman
Purwokerto

Molekul	Vol. 17	No. 1	pp. 1-144	March 2022	P-ISSN 1907-9761	E-ISSN 2503-0310
---------	---------	-------	-----------	---------------	---------------------	---------------------

[Search](#)[HOME](#) / [Editorial Team](#)

Editor-in-Chief

- [Dr. Amin Fatoni](#), Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia, Indonesia

Editorial Board

- [Prof. Shin-ichiro Suye](#), Department of Applied Chemistry and Biotechnology, University of Fukui, Fukui, Japan
- [Prof. Dr. Wan Aini Wan Ibrahim](#), Universiti Teknologi Malaysia, Malaysia
- [Prof. Pierre Doumeng](#), Aix Marseille University, France.
- [Prof. Dr. Kristian H Sugijarto](#), Department of Chemistry, Faculty of Mathematics and Natural Sciences, Yogyakarta State University, Indonesia
- [Prof. Dr. drh. Maria Bintang](#), Department of Biochemistry, Faculty of Mathematics and Natural Sciences, Institut Pertanian Bogor,, Indonesia
- [Prof. Dr. Hassan Y Aboul-Enein](#), National Research Centre, Cairo, Egypt
- [Prof. Dr. Hadi Nur](#), Centre for Sustainable Nanomaterials, Ibnu Sina Institute for Scientific and Industrial, Universiti Teknologi Malaysia, Malaysia
- [Dr. Arief Cahyo Wibowo](#), King Fahd University of Petroleum & Minerals, Saudi Arabia
- [Dr. Adi Darmawan](#), Chemistry Department, Faculty of Sciences and Mathematics, Diponegoro University Jl. Prof. Soedarto, SH., Tembalang, Semarang, Indonesia
- [Assist. Prof. Dr. Apon Numnuam](#), Department of Chemistry, Faculty of Science, Prince of Songkla University, Hat Yai, Thailand
- [Dr. Dadan Hermawan](#), Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia
- [Dr. Uyi Sulaeman](#), Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia, Indonesia
- [Dr. Jas Raj Subba](#), Royal University of Bhutan, Bhutan
- [Dr. Hartiwi Diastuti](#), Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia

- [Dr. Saluma Samanman](#), Faculty of Science and Technology, Princess of Naradhiwas University, Narathiwas, Thailand
- [Mardiyah Kurniasih](#), Department of Chemistry, Faculty of Mathematic and Natural Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia
- [Zusfahair](#), Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia

Layout Editor

- [Ely Setiawan](#), Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia
- [Irmanto Irmanto](#), Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia

MAKE A SUBMISSION

FOCUS AND SCOPE

GUIDE FOR AUTHORS



VIEW ARTICLES




EDITORIAL BOARD

PUBLICATION ETHICS

PEER REVIEW PROCESS

PLAGIARISM POLICY

CONTACT



Indexing and Abstracting:

Scopus®

INDEX COPERNICUS
INTERNATIONAL

sinta
Science and Technology Index

DOAJ
DIRECTORY OF
OPEN ACCESS
JOURNALS

CAS®
A DIVISION OF THE
AMERICAN CHEMICAL SOCIETY

Google
scholar

Crossref



Statistic Summary

Molekul



This work is licensed under



Jurnal Ilmiah Kimia
Department of Chemistry, Faculty of Mathematics and
Natural Sciences,
Universitas Jenderal Soedirman, Purwokerto, Indonesia

a [Creative Commons](#)
[Attribution 4.0](#)
[International License.](#)

Platform & workflow by

OJS

creating a better future

PKP | PUBLIC KNOWLEDGE PROJECT

[Search](#)[HOME](#) / [ARCHIVES](#) / Vol 17 No 1 (2022)**PUBLISHED:** 2022-03-16

ARTICLES

Molecular Docking of Xanthone Derivatives as Therapeutic Agent for Covid-19

Emmy Yuanita, Sudirman Sudirman, Ni Komang Tri Dharmayani, Maria Ulfa, Saprizal Hadisaputra, Jufrizal Syahri
1-9

DOI: <https://doi.org/10.20884/1.jm.2022.17.1.5600>

[PDF](#)

The enrichment of silver ions in Ag₃PO₄ through the morphology changes and their photocatalytic activities

Uyi Sulaeman, Khusnul Afifah, Hartiwi Diastuti, Shu Yin
10-18

DOI: <https://doi.org/10.20884/1.jm.2022.17.1.5606>

[PDF](#)

Biodegradation Kinetic Study of Cassava & Tannia Starch-Based Bioplastics as Green Material in Various Media

Marcelinus Christwardana, Ismojo Ismojo, Sidik Marsudi
19-29

DOI: <https://doi.org/10.20884/1.jm.2022.17.1.5591>

[PDF](#)

Voltammetric Determination of Paracetamol with Carbon Paste Electrode Modified with Molecularly Imprinted Electropolymer

I Wayan Tanjung Aryasa, Indra Noviandri
30-38

DOI: <https://doi.org/10.20884/1.jm.2022.17.1.5595>

[PDF](#)

Computational Study Potency of Eugenol and Safrole Derivatives as Active Sunscreen Material

Yusthinus Thobias Male, I Wayan Sutapa, Mirella Fonda Maahury, Muhamad Jamal, Dominggus Male

39-48

DOI: <https://doi.org/10.20884/1.jm.2022.17.1.5574>

**PDF****Tannic acid modified iron oxide nanoparticles and its application in protein adsorption: isotherm, kinetic, and thermodynamic study**

Hans Kristianto, Joshua Alexander, Susiana Prasetyo, Asaf K Sugih

49-59

DOI: <https://doi.org/10.20884/1.jm.2022.17.1.5571>

**PDF****Azadirone-Type Limonoids from the Fruit of Chisocheton lasiocarpus and Their Cytotoxic Activity Against MCF-7 Breast Cancer Lines**

Erina Hilmayanti, Nurlelarsari Nurlelarsari, Dewa Gede Katja, Ace Tatang Hidayat, Sofa Fajriah, Akhmad Darmawan, Unang Supratman, Mohamad Nurul Azmi, Yoshihito Shiono

60-67

DOI: <https://doi.org/10.20884/1.jm.2022.17.1.5593>

**PDF****Preparation and characterization of nanopowder of Acalypha hispida Leaves Extract Using Planetary Ball Milling**

Hamzah Alfarisi, Siti Sa'diah, Berry Juliandi, Tutik Wresdiyati

68-75

DOI: <https://doi.org/10.20884/1.jm.2022.17.1.5601>

**PDF****Terpenoids from the Stem bark of Aglaia elaeagnoidea and their cytotoxic activity against HeLa and DU145 Cancer Cell lines**

Dini Oktaviani, Winda Sukmawati, Kindi Farabi, Desi Harneti, Nurlelarsari Nurlelarsari, Darwati Darwati, Rani Maharani, Tri Mayanti, Agus Safari, Unang Supratman

76-84

DOI: <https://doi.org/10.20884/1.jm.2022.17.1.5594>

**PDF****Lipase Activity, Hematological and Blood Biochemistry of Osphronemus gouramy Fed with Supplementation of Spirulina platensis and Chlorella vulgaris**

Sorta Basar Ida Simanjuntak, Hana Hana, Riska Yunida, Maditaningtyas Hawwa Zuwanda

85-97

DOI: <https://doi.org/10.20884/1.jm.2022.17.1.5602>

**PDF****Discrimination of Metabolite Profiles of Gayo Roasted Arabica and Robusta Coffees**

Nizar Happyana, Yana Maolana Syah, Euis Holisotan Hakim

98-106

DOI: <https://doi.org/10.20884/1.jm.2022.17.1.5603>



PDF

Selective Colorimetric Detection of Mercury(II) using Silver Nanoparticles-Chitosan

Monica Avissa, Mohammad Alauhdin

107-115

DOI: <https://doi.org/10.20884/1.jm.2022.17.1.5597>



PDF

Antimicrobial and Antibiofilm Activities of Bacterial Strain P-6B from Segara Anakan Against MRSA 2983

Ari Asnani, Annisa Permata Dinda, Abdi Rahman Nursyamsi, Dwi Utami Anjarwati

116-124

DOI: <https://doi.org/10.20884/1.jm.2022.17.1.5575>



PDF

Leucaena-Derived Biochar for Biodiesel Production

Jutaporn Chanathaworn, Chokchai Yatongchai, Saluma Samanman

125-133

DOI: <https://doi.org/10.20884/1.jm.2022.17.1.5604>



PDF

Rubber (Hevea brasiliensis) Seed shell Activated Carbon Preparation, Characterization and Antibacterial Activity Studies

Saluma Samanman, Nuttaporn Tepprom, Inun Yipong, Yasmee Saleah, Kholeeyah Navani, Nifanusi Samae, Amri Hj Mohammed, Azharin Shah Abd Aziz

134-144

DOI: <https://doi.org/10.20884/1.jm.2022.17.1.5605>



PDF

[MAKE A SUBMISSION](#)[FOCUS AND SCOPE](#)[GUIDE FOR AUTHORS](#)[VIEW ARTICLES](#)[EDITORIAL BOARD](#)[PUBLICATION ETHICS](#)

Antimicrobial and Antibiofilm Activities of Bacterial Strain P-6B from Segara Anakan Against MRSA 2983

Ari Asnani^{1,*}, Annisa Permata Dinda¹, Abdi Rahman Nursyamsi², Dwi Utami Anjarwati³

¹Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia

²Department of Biology, Faculty of Biology, Universitas Jenderal Soedirman, Purwokerto, Indonesia

³Department of Microbiology, Faculty of Medicine, Universitas Jenderal Soedirman, Purwokerto, Indonesia

*Corresponding author: ari.asnani@unsoed.ac.id

Received September 04, 2021; Accepted November 05, 2021; Available online March 20, 2022

ABSTRACT. Methicillin-resistant *Staphylococcus aureus* (MRSA) infection has become a worldwide concern due to the increasing MRSA resistance to antibiotics. This condition encourages the continuous exploration of new anti-MRSA compounds. Thus, this study explored the effect of carbon (starch, glucose, sucrose, lactose) and nitrogen (casein, yeast extract, urea, peptone) sources for the fermentation medium to produce antibacterial and antibiofilm compounds from bacterial strain P-6B against MRSA 2983. The results showed that six-day incubation produced bioactive extracts with the highest antimicrobial activity (11.361 ± 0.860 mm) and the highest activity to inhibit biofilm formation ($89.159 \pm 0.293\%$). In comparison, three-day incubation produced a bioactive extract with the highest activity to degrade biofilm ($86.450 \pm 3.260\%$). Glucose was the best carbon source to produce an extract with antibacterial and biofilm inhibition activities. However, starch was the best carbon source to produce an extract with biofilm degradation activity. Urea was the best nitrogen source to produce extracts with antimicrobial and antibiofilm activities. Species identification based on the 16S rRNA gene revealed a 99.78% similarity with *Ochrobactrum intermedium* strain NBRC 15820 (NR_113812.1). These findings suggested the potency of *Ochrobactrum* P-6B from Segara Anakan Cilacap as indigenous sources of bioactive compounds with anti-MRSA activities.

Keywords: antibacterial, antibiofilm, MRSA, *Ochrobactrum*

INTRODUCTION

Methicillin-resistant *Staphylococcus aureus* (MRSA) is a specific *S. aureus* bacteria resistant to methicillin antibiotics (Kemung et al., 2018). Nosocomial infection by MRSA occurs by forming a biofilm layer as a pathogenetic mechanism to infect the human body (Rewatkar & Wadher, 2013). The MRSA biofilms on the surface of tissues are aggregated by an extracellular matrix consisting of protein, DNA, and polysaccharides in the form of polysaccharide intercellular adhesin (PIA) (Boles, Thoendel, Roth, & Horswill, 2010). The colonization of MRSA in the formation of biofilms in the blood circulation can cause bacteremia. Prolonged local bacteremia by MRSA infection eventually causes sepsis (Tissot et al., 2014).

Bacteria that form biofilms are expected to be more resistant to antimicrobial exposure. The resistance occurs because antimicrobial substances' ability to penetrate is reduced due to the formation of aggregates cells, a slower metabolic state of bacteria, and horizontal exchange of resistance genes that are easier to occur between biofilm-forming cells (Khameneh, Diab, Ghazvini, & Fazly Bazaz, 2016). Indeed, WHO (2017) has ranked MRSA as a high

priority in the pathogen list for research and development of new antibiotics. The grouping of WHO pathogens list is according to the species and the type of resistance. Thus, exploration for new compounds with antibiofilm activities, particularly against MRSA, becomes necessary.

One strategy that can be used to overcome biofilm formation is to screen bacteria that can produce metabolites as anti-MRSA agents. A thorough review by Kemung et al. (2018) highlights *Streptomyces*' potency to produce various anti-MRSA compounds. Recently, indigenous microbes isolated from mangrove sediment Segara Anakan have been reported to have antibacterial activity against MRSA (Asnani, Amaliyah, & Yuniaty, 2020; Dinda, Asnani, & Anjarwati, 2021). We observed that strain P-6B suggested a potency to produce bioactive extract against MRSA based on qualitative screening. Species identification of P-6B indicated the genus *Ochrobactrum*. It was predicted to have one gene cluster in region 1.4 involved in the biosynthesis of ambactin from the known non-ribosomal peptide (NRP) cluster (Asnani et al., 2020). Genus *Ochrobactrum* is found under a wide variety of environmental conditions and is primarily studied for

its potential to degrade heavy metal pollutants (Sharma & Shukla, 2021). *Ochrobactrum* strain MZV101, a strong oil removal bacteria, exhibits antimicrobial activity against various bacterial pathogens (Zarinviarsagh, Ebrahimipour, & Sadeghi, 2017).

Furthermore, Fan et al. (2020) reported that *Ochrobactrum intermedium* D-2 exhibited potency as quorum quenching bacteria against the pathogen. Quorum quenching has been proposed as a promising strategy to prevent and control quorum sensing-mediated bacterial infections. Accessory gene regulator (*agr*) quorum sensing systems are critical components of MRSA cellular processes. It is a target for inhibiting virulence factor production to treat MRSA infections (Lade & Kim, 2021). Following that report, our research aimed to further analyze the potency of P-6B isolate as antibacterial and antibiofilm towards MRSA 2983.

Production of specific bacterial metabolites occurs under particular fermentation conditions. In this case, determining the optimal incubation time is crucial because microbes will synthesize different metabolites in different growth phases. Chodkowski and Shade (2020) have reviewed that some secondary metabolites, such as antibiotics, increase production at the stationary phases. Additionally, carbon and nitrogen in the medium are essential nutrients for microbial growth and the production of primary and secondary metabolites (Singh et al., 2017). Therefore, optimization of medium is necessary to maximize secondary metabolite production. Our research focused on determining the optimum incubation time for the production of anti-MRSA extracts from the bacterial strain P-6B and exploring the fermentation medium with various carbon and nitrogen sources.

EXPERIMENTAL SECTION

Materials

Bacterial strain P-6B was isolated from mangrove sediments in Segara Anakan Cilacap. The isolate was maintained in starch casein nitrate (SCN) medium consisting of starch (Merck), casein (Merck), KNO₃ (Merck), KH₂PO₄ (Merck), MgSO₄·7H₂O (Merck), NaCl (Merck), FeSO₄·7H₂O (Merck), bacto agar (Oxoid) supplemented with 1% nystatin® (kandistatin). The clinical methicillin-resistance *Staphylococcus aureus* 2983 (MRSA 2983) was the collection of Faculty of Medicine, UNSOED, Purwokerto. The pathogen, MRSA 2983, was cultured in mannitol salt agar (Himedia). The antibiofilm tests used Nutrient Broth (Merck), Mueller-Hinton Agar (Himedia), and Brain Heart Infusion broth (Merck). Other chemicals used were glucose (Merck), sucrose (Merck), lactose (Merck), yeast extract (Merck), urea (Merck), and peptone (Oxoid). DNA isolation used Quick-DNA Fungal/Bacterial Micropep Kit (Zymo Research). PCR was performed using MyTaq™ HS Red Mix (Bioline)

with primer 27F (AGAGTTTGATCMTGGCTCAG) and 1492R (TACGGYTACCTTGTTCAGCTT).

The Cultivation of Microbes

Cultivation of P-6B isolate followed the procedure described by (Asnani et al., 2020). The isolate was inoculated with a continuous streak on Starch casein nitrate agar supplemented with 1% nystatin®. The culture was incubated for seven days at room temperature before being used. The preparation of MRSA 2983 followed the procedure described by Dinda et al. (2021). MRSA 2983 was inoculated in Nutrient Broth and incubated for 18 to 24 hours. The culture suspension was adjusted to 0.5 units of standard McFarland with a 0.9% NaCl solution.

Determination of the Optimum Incubation Time

Bacterial strain P-6B was grown in a liquid SCN medium and incubated at 90 rpm for seven days at room temperature for inoculum. Next, 10% of the inoculum was inoculated into the new SCN medium and set at 90 rpm for 15 days, with interval sampling of three days (0, 3, 6, 9, 12, and 15 days). After each incubation time, the cultures were centrifuged at 4000 rpm for 10 minutes at 4°C. The mycelium obtained was dried and weighed as the dry weight of the mycelium (g). The supernatant obtained was analyzed for antibacterial activity, inhibition of biofilm formation, and biofilm degradation. The incubation times with the optimum activities for all analyzed parameters were then used for subsequent procedures.

Variation of Carbon and Nitrogen Sources

The procedure to determine the optimum carbon source followed Kannan et al. (2014), with modifications. A 10% inoculum was added liquid SCN medium containing 1% of four different carbon sources, which were starch, glucose, sucrose, and lactose. Cultures were incubated at 90 rpm, room temperature. The fermentation was carried out at the optimum incubation time that produces extract with optimum anti-MRSA activity. A similar procedure was applied to determine the optimum nitrogen sources. A 10% inoculum was added liquid SCN medium containing 1% of four different nitrogen sources, which were casein, yeast extract, urea, and peptone. After the optimal incubation time, the supernatant obtained was tested for antibacterial activity, inhibition of biofilm formation, and biofilm degradation.

Antibacterial Activity Test

The extract's ability as an antibacterial compound was tested using the disc paper diffusion method on Mueller-Hinton Agar following the procedure described in CLSI (2019). A total of 30 µl of the extract was added to sterile (6 mm) disc paper, then placed in MHA media that had been inoculated with MRSA. Cultures were incubated at 37°C for 24 hours. The antibacterial activity of each treatment was evaluated by measuring the zone of inhibition around the well with a vernier caliper.

Inhibition of Biofilm Formation

The extract's ability to inhibit biofilm formation was evaluated using a microplate semi-quantitative test following Suzuki et al. (2015) with modification. MRSA was inoculated in Brain Heart Infusion broth supplemented with 1% glucose (BHI-Glu) and then incubated at 37°C for 18 hours. Subsequently, the culture turbidity levels were adjusted to the McFarland standard of 0.5. The biofilm formation inhibitor test was carried out by adding 10 µL of MRSA culture with 100 µL of extract to the microplate and then incubated at 37°C for 24 hours. Biofilm staining used 50 µL of 0.1% crystal violet solution. Measurement of optical density (OD) value was carried out at 595 nm. The OD value obtained was used to calculate the percentage of biofilm formation inhibition using the following formula.

$$\% \text{ Inhibition} = \frac{\text{OD control} - \text{OD test}}{\text{OD control}} \times 100\%$$

Biofilm Degradation

The ability of the extract to degrade biofilm was also evaluated with a microplate method based on Suzuki et al. (2015) with modifications. A total of 10 µL of MRSA cultures on BHI-Glu medium were cultivated on a microplate for 24 hours at 37°C to form the biofilm. Next, 100 µL of the extract was added, and the incubation was continued for 24 hours at 37°C to expose the extract to the biofilm formed on the walls of microplate. Biofilm staining used 50 µL of 0.1% crystal violet solution. The measurement of the OD value was carried out at 595 nm. The OD value obtained was then used to calculate the percentage of inhibition of biofilm formation using the following formula.

$$\% \text{ Degradation} = \frac{\text{OD control} - \text{OD test}}{\text{OD control}} \times 100\%$$

Species observation and Identification

Macromorphological observations of bacterial strain P-6B focused on colony characters. The observation used a stereo microscope with a magnification of 30x. Micromorphological observations used the Heinrich Slide Culture method using a light microscope with a magnification of 400x. Species identification of P-6B was based on the 16S rRNA gene following the procedure described elsewhere (Asnani et al., 2020). DNA isolation was carried out using the Quick-DNA Fungal/Bacterial Micropep Kit. The DNA obtained was amplified using MyTaq™ HS Red Mix with primers 27F and 1492R. Amplicon DNA was sequenced in two directions with the Sanger sequencing method. The sequences obtained were analyzed with BioEdit 7.2, then aligned to other sequences in GenBank through BLAST nucleotide. The phylogenetic tree construction used MEGA X with construct Maximum Likelihood method and Kimura-2 parameter model with bootstrap 100 (Kumar, Stecher, Li, Knyaz, & Tamura, 2018).

Data Analysis

The research data was in the form of the inhibition index for the antibacterial activity test, the percent inhibition of biofilm formation, and the percent biofilm degradation. Data were analyzed using analysis of variance (ANOVA) at 95% significance levels. The Duncan test followed the results of ANOVA analysis that were significantly different.

RESULTS AND DISCUSSION

MRSA is an infection-causing bacteria that can defend itself by forming a protective layer called a biofilm. MRSA biofilms develop rapidly and form colonies on moist and nutrient-rich surfaces. Biofilm formation is one of the virulence factors of MRSA that contributes to its resistance to antibiotics.

Determination of Optimum Incubation Time

The fermentation process to produce anti-MRSA extract was carried out to determine the optimum incubation time. The growth curve of P-6B (**Figure 1**) shows that the exponential phase occurred from day 6 and reached maximum on day 12. The antibacterial activity of the P-6B extract increased from day 0 and reached the highest activity on day six with an inhibition zone diameter of 11.361 ± 0.860 mm. After that, the antibacterial activity decreased until the 15th day. The results of the variance analysis (Sig. <0.05) indicated that the incubation time significantly affected the diameter of the inhibition zone. The Duncan test further confirmed that antibacterial activity with an incubation time of six days was the highest value and significantly different. Based on this analysis, the 6-day incubation time was suggested for the fermentation process to produce extract with the highest antibacterial activity.

The percentage inhibition of biofilm formation by P-6B extract increased from day 0 to optimal incubation on day six, with a percent inhibition of $89.159 \pm 0.293\%$ (**Figure 2**). After reaching the optimal incubation time, the percentage of inhibition decreased. The analysis of variance (Sig. <0.05) showed that the incubation time was significantly affected the inhibition of biofilm formation. Further, the Duncan test concluded that some interactions showed no significant difference. However, the best treatment was the six-day incubation time, which gave the highest inhibition value of biofilm formation. Based on this analysis, the 6-day incubation time was proposed for the fermentation process to produce extract with the highest percentage of inhibition of biofilm formation. The percentage of biofilm degradation by the P-6B extract reached its optimum on the third day of incubation with a percentage of degradation of $86.450 \pm 3.260\%$ (**Figure 2**). After getting the optimum incubation time, the degradation decreased. The variance analysis (Sig. <0.05) indicated that the incubation time significantly affected biofilm degradation. Further Duncan test suggested that some interactions were not significantly different.




Figure 1. Production of antibacterial extract with incubation times. The dried mycelium weight (g) corresponds with antibacterial activity measured as the diameter of inhibition zone (mm);




Figure 2. Production of antibiofilm extract with incubation times. The inhibition of biofilm formation and biofilm degradation activities of the texttracts was measured as a percentage (%).

However, the best treatment was the incubation time of three days that yielded the highest biofilm degradation. Based on this analysis, the 3-day incubation time was recommended for the fermentation process to produce extract with the highest percentage of biofilm degradation.

The mangrove ecosystem in Segara Anakan Cilacap is rich in microbial diversity. The present study results suggested the potency of P-6B to produce antimicrobial and antibiofilm extract against MRSA 2983. The ability of the P-6B extract as antibacterial also suggested its ability to inhibit biofilm formation. Memariani et al. (2019) stated that compounds with antibacterial activity against microbes that form biofilms could also be used to inhibit the formation of those microbial biofilms. Indeed, the mechanisms of inhibition of biofilm formation are mostly the same as an antibacterial mechanism. On the contrary, biofilm

degradation has a different mechanism. It leads to an enzymatic activity that can break down the biofilm matrix, weakening microbial aggregation on biofilms (Lee et al., 2016).

The difference in incubation time is in accordance with the bioactive compounds produced. In this study, the extracts for antibacterial and biofilm inhibition were produced by incubation for six days, while the extract for biofilm degradation was produced only for three days incubation. These results were consistent with those of Xie et al. (2019). They stated that the production of antibacterial compounds by actinomycetes takes longer than the production of compounds that can degrade the extracellular layer of MRSA. Oja et al. (2015) reported that *Streptomyces violaceoruber* (DSM-40701) produced antibiofilm compounds against *S. aureus* with an incubation time of four days. Suzuki et al. (2015) reported that the

Streptomyces sp. strain MC11024 produced antibiofilm compounds against *S. aureus* and MRSA N315 with three incubation days. In other reports, *Streptomyces* sp. SBT343 has antibiofilm compounds against staphylococcal bacteria with an incubation time of ten days (Balasubramanian et al., 2017).

Determination of Optimum Carbon and Nitrogen sources

The best carbon and nitrogen sources for the fermentation medium were determined based on the highest antibacterial activity, inhibition of biofilm formation, and biofilm degradation. The production of P-6B extract with antibacterial activity was carried out

with an incubation time of six days. The results of the antibacterial activity test showed that glucose was the best carbon source (10.617 ± 0.646 mm) and urea was the best nitrogen source (12.426 ± 2.152 mm) in producing antibacterial extracts (Figure 3). The variance analysis results (Sig. <0.05) showed that carbon and nitrogen sources significantly affected the inhibition zone's diameter. Further, the Duncan test concluded that antibacterial activities were significant differences between treatments. The highest inhibition zone was produced by fermentation medium using glucose as a carbon source and urea as a nitrogen source.




Figure 3. Antibacterial activity of extract P-6B with different sources of carbon and nitrogen.




Figure 4. Inhibition of biofilm formation and biofilm degradation from the extract P-6B with different carbon and nitrogen sources.

Production of P-6B extracts with MRSA 2983 potential to inhibit the biofilm formation of MRSA 2983 was carried out with an incubation time of six days. The results showed that the highest inhibition of biofilm formation was produced from extracts using glucose ($97.968 \pm 0.151\%$) as a carbon source and urea ($97.144 \pm 0.516\%$) as a nitrogen source (**Figure 4**). The results from the variance analysis (Sig. <0.05) showed that carbon and nitrogen sources significantly affected biofilm formation inhibition. Further Duncan's test indicated that some interactions were not significantly different. However, the best treatment with the highest inhibition of biofilm formation was glucose as a carbon source and urea as a nitrogen source.

The production of P-6B extract, which could degrade the biofilm of MRSA 2983, was carried out with an incubation time of three days. The results showed that the highest percentage of biofilm degradation was obtained from fermentation medium using starch ($86.158 \pm 2.395\%$) as a carbon source and urea ($72.349 \pm 7.939\%$) as a nitrogen source (**Figure 4**). The results of the variance analysis (Sig. <0.05) indicated that the sources of carbon and nitrogen significantly affected biofilm degradation. Further, the Duncan test showed significant differences between treatments, with the highest biofilm degradation coming from starch as a carbon source and urea as a nitrogen source.

The composition of the fermentation medium must provide all the required components of the compound. The main components that must be present are carbon and nitrogen for microorganism cell metabolism, cell growth, and to produce specific metabolites. However, the difference in carbon and nitrogen sources in the fermentation medium consequently provides different activities. In this study, glucose was the best carbon source for producing antibacterial extracts and inhibition of biofilm formation, while starch showed the highest biofilm degradation activity. Similarly, Kannan et al. (2014) reported that glucose is the best carbon source to produce antibacterial compounds from the actinomycetes group. Likewise, Nandhagopal et al. (2017) also use starch to produce bioactive compounds for MRSA biofilm degradation.

Nitrogen sources from complex organic compounds function for microbial growth and

increase the production of bioactive compounds (González, 2018). Nitrogen is a macronutrient that plays a role in protein biosynthesis to prepare cell structure and cell function. Nitrogen in the fermentation medium can also induce specific enzymes according to the nitrogen source used. Thus, the nitrogen source in the fermentation medium will differ depending on the type of bacteria and bioactive compounds desired. In this study, the treatment of variations in nitrogen sources suggested that urea is the best nitrogen source for the fermentation medium. Urea is known to have the ability to lyse cells and tissues and damage biofilms originating from clinical isolates (Rasmussen et al., 2016). Sanawar et al. (2018) also reported that urea reduced protein and carbohydrate concentration in the EPS biofilm matrix by $> 50\%$. Urea destroys biofilms by breaking hydrogen bonds between amides and carbonyl groups in proteins. Differently, Suzuki et al. (2015) used casein in the fermentation medium *Streptomyces* sp. strain MC11024 to inhibit the *S. aureus* biofilm Streptorubin B as an MRSA biofilm inhibitor. Balasubramanian et al. (2017) used yeast extract to produce bioactive compounds that inhibit the Staphylococcal biofilm by *Streptomyces* sp. SBT343. Sharma and Manhas (2019) used casein as a nitrogen source in fermentation medium to produce antibacterial compounds from *Streptomyces* strain M7 against MRSA and VRE.

Species Observation and Identification

Bacterial strain P-6B was isolated from sediment mangrove in Kalipanas area, Segara Anakan, Cilacap. The bacterium was obtained from sediment samples which were pretreated with 1.5% phenol (Asnani & Oedjijono, 2019). The colonies of the P-6B bacteria strain grew slowly and were tightly attached to the agar medium (**Figure 5**). The colonies were circular with powdery surfaces. The edges of the colonies tended to be flat with raised elevation. The colony surface was gray, the reverse colony was dark brown, and colored pigmentation to the medium was observed. Based on the 16S ribosomal RNA gene, the P-6B isolate has 99.78% similarity to the *Ochrobactrum intermedium* strain NBRC 15820 (NR_113812.1).




Figure 5. Observation of *Ochrobactrum* P-6B, (a) colony surface, (b) single colony with stereo microscope 30x.




Figure 6. The phylogenetic tree of *Ochrobactrum* P-6B and related species were obtained from the Maximum Likelihood method and the Kimura-2 parameter model. The numbers on the branch node were bootstrap values expressed as a percentage of 100 replications.

The construction of the phylogenetic tree validated the relationship of *Ochrobactrum* P-6B with other species. Evolutionary analyzes were conducted in MEGA X (Kumar et al., 2018). The evolutionary relationship was inferred using the neighbor-joining method. The optimal tree with the sum of branch length = 1.13588203 is shown in Figure 6. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (100 replicates) are shown next to the branches. The tree is drawn to a 0.050 scale, with branch lengths in the same units as the evolutionary distances used to infer the phylogenetic tree. The evolutionary distances were computed using the Kimura 2-parameter method and are in the units of the number of base substitutions per site. This analysis involved 21 nucleotide sequences. There were a total of 1594 positions in the final dataset. All ambiguous positions were removed for each sequence pair (pairwise deletion option).

CONCLUSIONS

Our research highlights the potency of indigenous marine *Ochrobactrum* P-6B from the sediment mangrove in Segara Anakan Cilacap as antibacterial and antibiofilm sources against MRSA 2983. Indeed, the mangrove forest offers a unique environment with diverse antibiotic-producing microbes. Further research is required to purify and characterize the bioactive compounds with antibiofilm activities from

Ochrobactrum P-6B as an alternative therapy for MRSA infection.

ACKNOWLEDGEMENTS

The authors gratefully acknowledged the Ministry of Research and Technology/National Agency for Research and Innovation of the Republic of Indonesia, who funded the Fundamental Research Grant, contract No. T/1641/UN23.18/PT.01.03/2020 through the Directorate of Research and Community Service.

REFERENCES

- Asnani, A., Amaliyah, R., & Yuniaty, A. (2020). Screening anti-MRSA activities of indigenous microbes and prediction of the Biosynthetic Gene Clusters. *Journal of Physics: Conference Series*, 1665. <https://doi.org/10.1088/1742-6596/1665/1/012001>
- Asnani, A., & Oedjijono. (2019). *Exporation of actinomycetes in mangrove area Segara Anakan (Eksplorasi aktinomisetes di kawasan mangrove Segara Anakan)* (D.U. Anjarwati (ed.)). UNSOED Press Purwokerto.
- Balasubramanian, S., Othman, E. M., Kampik, D., Stopper, H., Hentschel, U., Ziebuhr, W., Oelschlaeger, T. A., & Abdelmohsen, U. R. (2017). Marine sponge-derived *Streptomyces* sp. SBT343 extract inhibits *Staphylococcal*

- biofilm formation. *Frontiers in Microbiology*, 8, 236. <https://doi.org/https://doi.org/10.3389/fmicb.2017.00236>
- Barrios-González, J. (2018). Secondary metabolites production: Physiological advantages in solid-state fermentation. In *Current Developments in Biotechnology and Bioengineering* (pp. 257–283). Elsevier. <https://doi.org/10.1016/B978-0-444-63990-5.00013-X>
- Boles, B. R., Thoendel, M., Roth, A. J., & Horswill, A. R. (2010). Identification of genes involved in polysaccharide-independent *Staphylococcus aureus* biofilm formation. *PLoS ONE*, 5(4), e10146. <https://doi.org/10.1371/journal.pone.0010146>
- Chodkowski, J. L., & Shade, A. (2020). Exometabolite dynamics over stationary phase reveal strain-specific responses. *MSystems*, 5(6). <https://doi.org/10.1128/msystems.00493-20>
- CLSI. (2019). Performance Standards for Antimicrobial Susceptibility Testing. In 29th Ed. CLSI supplement M100. Wayne, PA. https://doi.org/10.1007/978-3-662-48986-4_300418
- Dinda, A. P., Asnani, A., & Anjarwati, D. U. (2021). The Activities of *Streptomyces* W-5A as antibacterial and antibiofilm towards methicillin-resistant *Staphylococcus aureus* 2983. *Proceedings of the 1st Jenderal Soedirman International Medical Conference in Conjunction with the 5th Annual Scientific Meeting (Temilnas) Consortium of Biomedical Science Indonesia*, 109–115. <https://doi.org/10.5220/0010488601090115>
- Fan, X., Ye, T., Li, Q., Bhatt, P., Zhang, L., & Chen, S. (2020). Potential of a quorum quenching bacteria isolate *Ochrobactrum intermedium* D-2 against soft rot pathogen *Pectobacterium carotovorum* subsp. *carotovorum*. *Frontiers in Microbiology*, 11(May), 1–14. <https://doi.org/10.3389/fmicb.2020.00898>
- Kannan, R. R., Iniyar, A. M., & Vincent, S. G. P. (2014). Production of a compound against methicillin resistant *Staphylococcus aureus* (MRSA) from *Streptomyces rubrolavendulae* ICN3 & its evaluation in zebrafish embryos. *The Indian Journal of Medical Research*, 139(6), 913.
- Kemung, H. M., Tan, L. T.-H., Khan, T. M., Chan, K.-G., Pusparajah, P., Goh, B.-H., & Lee, L.-H. (2018). *Streptomyces* as a prominent resource of future Anti-MRSA drugs. *Frontiers in Microbiology*, 9, 2221. <https://doi.org/10.3389/fmicb.2018.02221>
- Khameneh, B., Diab, R., Ghazvini, K., & Fazly Bazzaz, B. S. (2016). Breakthroughs in bacterial resistance mechanisms and the potential ways to combat them. *Microbial Pathogenesis*, 95, 32–42. <https://doi.org/10.1016/j.micpath.2016.02.00>
- Kumar, S., Stecher, G., Li, M., Knyaz, C., & Tamura, K. (2018). MEGA X: Molecular Evolutionary Genetics Analysis across computing platforms. *Molecular Biology and Evolution*, 35(6), 1547–1549. <https://doi.org/10.1093/molbev/msy096>
- Lade, H., & Kim, J. S. (2021). Bacterial targets of antibiotics in methicillin-resistant *Staphylococcus aureus*. *Antibiotics*, 10(4). <https://doi.org/10.3390/antibiotics10040398>
- Lee, J.-H., Kim, Y.-G., Lee, K., Kim, C.-J., Park, D.-J., Ju, Y., Lee, J.-C., Wood, T. K., & Lee, J. (2016). *Streptomyces* -derived actinomycin D inhibits biofilm formation by *Staphylococcus aureus* and its hemolytic activity. *Biofouling*, 32(1), 45–56. <https://doi.org/10.1080/08927014.2015.1125888>
- Memariani, H., Memariani, M., & Ghasemian, A. (2019). An overview on anti-biofilm properties of quercetin against bacterial pathogens. *World Journal of Microbiology and Biotechnology*, 35(9), 143. <https://doi.org/10.1007/s11274-019-2719-5>
- Nandhagopal, S., Iniyar, A. M., Kannan, R. R., & Vincent, S. G. P. (2017). In vivo evaluation of anti-MRSA compound from *Streptomyces collinus* ICN1 in zebrafish embryos. *Indian Journal of Geo-Marine Sciences*, 46(6), 1155–1161.
- Oja, T., San Martin Galindo, P., Taguchi, T., Manner, S., Vuorela, P. M., Ichinose, K., Metsä-Ketelä, M., & Fallarero, A. (2015). Effective antibiofilm polyketides against *Staphylococcus aureus* from the Pyranonaphthoquinone Biosynthetic Pathways of *Streptomyces* species. *Antimicrobial Agents and Chemotherapy*, 59(10), 6046–6052. <https://doi.org/10.1128/AAC.00991-15>
- Rasmussen, K., Reilly, C., Li, Y., & Jones, R. S. (2016). Real-time imaging of anti-biofilm effects using CP-OCT. *Biotechnology and Bioengineering*, 113(1), 198–205. <https://doi.org/10.1002/bit.25701>
- Rewatkar, A. R., & Wadher, B. J. (2013). *Staphylococcus aureus* and *Pseudomonas aeruginosa*- Biofilm formation methods. *IOSR Journal of Pharmacy and Biological Sciences*, 8(5), 34–60. <https://doi.org/10.9790/3008-0853640>
- Sanawar, H., Pinel, I., Farhat, N. M., Bucs, S. S., Zlopasa, J., Kruithof, J. C., Witkamp, G. J., van Loosdrecht, M. C. M., & Vrouwenvelder, J. S. (2018). Enhanced biofilm solubilization by urea in reverse osmosis membrane systems. *Water Research X*, 1, 100004. <https://doi.org/10.1016/j.wroa.2018.10.001>
- Sharma, B., & Shukla, P. (2021). A comparative analysis of heavy metal bioaccumulation and functional gene annotation towards multiple metal resistant potential by *Ochrobactrum*

- intermedium* BPS-20 and *Ochrobactrum ciceri* BPS-26. *Bioresource Technology*, 320, 124330. <https://doi.org/10.1016/j.biortech.2020.124330>
- Sharma, M., & Manhas, R. K. (2019). Purification and characterization of actinomycins from *Streptomyces* strain M7 active against methicillin resistant *Staphylococcus aureus* and vancomycin resistant *Enterococcus*. *BMC Microbiology*, 19(1), 44. <https://doi.org/10.1186/s12866-019-1405-y>
- Singh, V., Haque, S., Niwas, R., Srivastava, A., Pasupuleti, M., & Tripathi, C. K. M. (2017). Strategies for fermentation medium optimization: An in-depth review. *Frontiers in Microbiology*, 7, 2087. <https://doi.org/10.3389/fmicb.2016.02087>
- Suzuki, N., Ohtaguro, N., Yoshida, Y., Hirai, M., Matsuo, H., Yamada, Y., Imamura, N., & Tsuchiya, T. (2015). A Compound inhibits biofilm formation of *Staphylococcus aureus* from *Streptomyces*. *Biological and Pharmaceutical Bulletin*, 38(6), 889–892. <https://doi.org/10.1248/bpb.b15-00053>
- Tissot, F., Calandra, T., Prod'homme, G., Taffe, P., Zanetti, G., Greub, G., & Senn, L. (2014). Mandatory infectious diseases consultation for MRSA bacteremia is associated with reduced mortality. *Journal of Infection*, 69(3), 226–234. <https://doi.org/10.1016/j.jinf.2014.05.004>
- WHO. (2017). World Health Organization releases global priority list of antibiotic-resistant bacteria to guide research, discovery, and development of new antibiotics. https://www.who.int/medicines/publications/WHO-PPL-Short_Summary_25Feb-ET_NM_WHO.pdf
- Xie, T. T., Zeng, H., Ren, X. P., Wang, N., Chen, Z. J., Zhang, Y., & Chen, W. (2019). Antibiofilm activity of three Actinomycete strains against *Staphylococcus epidermidis*. *Letters in Applied Microbiology*, 68(1), 73–80. <https://doi.org/10.1111/lam.13087>
- Zarinviarsagh, M., Ebrahimipour, G., & Sadeghi, H. (2017). Lipase and biosurfactant from *Ochrobactrum intermedium* strain MZV101 isolated by washing powder for detergent application. *Lipids in Health and Disease*, 16(1), 1–13. <https://doi.org/10.1186/s12944-017-0565-8>

[JM] Submission Acknowledgement

1 message

Amin Fatoni <j.molekul@gmail.com>
To: Ari Asnani <ari.asnani@unsoed.ac.id>

Sat, Sep 4, 2021 at 9:10 AM

Ari Asnani:

Thank you for submitting the manuscript, "Antimicrobial and Antibiofilm Activities of Bacterial Strain P-6B from Segara Anakan against MRSA 2983" to Molekul. With the online journal management system that we are using, you will be able to track its progress through the editorial process by logging in to the journal web site:

Manuscript URL:
<https://ojs.jmolekul.com/ojs/index.php/jm/author/submission/837>
Username: asnani

If you have any questions, please contact me. Thank you for considering this journal as a venue for your work.

Amin Fatoni
Molekul
Editor
MOLEKUL - Scientific Journal in Chemistry

Revision Required for manuscript 837

2 messages

Jurnal Molekul <j.molekul@gmail.com>
To: Ari Asnani <asnani@gmail.com>

Mon, Oct 18, 2021 at 10:55 AM

Dear Dr. Ari Asnani,

We have received the comment from the reviewer below, please revise according to the suggestion.

Best Regards,
Amin Fatoni,
Editor

Reviewer 1 comment:


- The manuscript is well written. The idea was good, however, there is an ambiguous statement in the introduction concerning the genus *Ochrobactrum*. Author should make it clearly, the previous research only investigated from the morphology or molecular.
- Please check the comment in the attached file

Reviewer 2 comment:

- Please briefly explain the bacterial isolation from mangrove sediment, or refer to previous reports.
- What does the pathogen mean in the materials?
- The grade/brand of material used, e.g. starch (merck), pepton (oxoid), should be mentioned in the materials, instead of procedures.
- Why do researchers use the disk paper method for antibacterial activity?

Editor comment:

- please use dot (.) instead of comma (,) for decimal
- figure 1,3,4 text is not clear, please enlarge the text
- please add the x axis of figure 3 and 4

 **837-2557-1-RV-3.docx**
579K

Ari Asnani <asnani@gmail.com>
To: Jurnal Molekul <j.molekul@gmail.com>

Thu, Oct 21, 2021 at 9:04 PM


Dear Editor

Thank you very much for the review.
Hereby we would like to submit our revised article, our response to the reviewers, and also the Excel file for the graphs. We have also submitted these three files via Molekul ojs system.

Thank you very much,
Your attention and kind consideration is highly appreciated.

Best wishes,
Ari Asnani

[Quoted text hidden]

3 attachments **Author Response.docx**
21K **Data Article Molecule.xlsx**
153K **Author Rev 837-2557-1-RV-3.docx**
364K

Proof Read - Jurnal MOLEKUL Vol 17 No 1

2 messages

Jurnal Molekul <j.molekul@gmail.com>

Mon, Mar 7, 2022 at 12:06 PM

To: Ari Asnani <asnani@gmail.com>, ari.asnani@unsoed.ac.id

Yth. Dr. Ari Asnani, dkk

Sehubungan dengan artikel yang telah diterima untuk diterbitkan pada Jurnal Molekul Volume 17 No 1, Maret 2022, dengan ini maka:

1. Terlampir file artikel yang telah di layout, mohon untuk dicek kembali (*proofread*). Untuk memastikan artikel anda terbit pada waktunya, mohon jika ada koreksi harap diperbaiki dan dikembalikan dalam waktu **maksimal 2 hari**. Adanya kesalahan setelah artikel diterbitkan atau dicetak **tidak memungkinkan untuk direvisi** lagi. Jika dalam batas waktu yang ditentukan **tidak ada tanggapan**, maka kami anggap penulis **telah setuju dengan artikel tersebut** baik isi, susunan penulis, dan komponen artikel lainnya.
Perubahan / revisi dapat disertakan dalam lembar tersendiri atau komentar di file pdf yang dikirimkan.
2. Mohon melakukan pembayaran biaya publikasi sebagai berikut:
 - o Jumlah Rp. 1.800.000 (Versi online + 2 eksemplar versi cetak termasuk ongkos kirim). Penambahan versi cetak, Rp 250.000 per eksemplar.
 - o Cara Pembayaran :
 - No Rekening BNI : **1135395011**, a.n. : **Jurnal Ilmiah Kimia Molekul**
 - o Setelah melakukan pembayaran, harap konfirmasi melalui **Whatsapp** ke Ibu Zusfahair: **081323760903**
3. Artikel akan diterbitkan di website kami yang baru, mohon register dan submit di file terakhir ini di website kami yang baru: <http://jos.unsoed.ac.id/index.php/jm/>
4. Mohon mengisi form pernyataan terlampir

Demikian, jika ada informasi yang belum jelas, dapat dikonfirmasi melalui email ini.

Salam,

Editor Jurnal MOLEKUL

English version

The proof of your article, to be published in 'MOLEKUL', is now available.

Please use this proof only for checking the typesetting, editing, completeness and correctness of the text, tables and figures. To assist you with this, copy-editing changes have been highlighted in red with the original text in a pop-up box.

For article processing fee payment (\$150, please contact the editor to select the payment method).


To ensure fast publication of your paper please return your corrections within 48 hours.

Fail to reply to this email within the time, we assumed that you have agreed with the final version of your article attached.

Note that once your corrections have been received, your article is finalized and any further correction is no longer possible.

Yours sincerely,

MOLEKUL Editorial Team.

2 attachments **837-Ari Asnani-(116-124)-OK.pdf**
427K **Form_pernyataan-Maret2022-EN.docx**
85K

Ari Asnani <ari.asnani@unsoed.ac.id>

Tue, Mar 8, 2022 at 6:48 AM

To: Jurnal Molekul <j.molekul@gmail.com>

Dear Editor Jurnal Molekul

Terima kasih banyak sebelumnya atas kesempatan publikasi artikel kami di Jurnal Molekul.

- 1) Kami ada satu koreksi afiliasi author Ari Asnani, menjadi **Faculty of Mathematics and Natural Sciences** (koreksi terlampir).
- 2) Pembayaran telah kami lakukan dengan penambahan pesanan 2 eksemplar versi cetak. Bukti pembayaran terlampir dan sudah diinfokan juga ke bu Zusfahair.
- 3) registrasi ke situs baru segera dikerjakan.
- 4) Form pernyataan bersama ini kami lampirkan.

Terima kasih,

Wassalam,
Ari Asnani

[Quoted text hidden]

--
Ari Asnani, Ph.D.
Jurusan Kimia - Fakultas MIPA
Universitas Jenderal Soedirman
Purwokerto, Jawa Tengah, Indonesia

3 attachments



Pembayaran biaya publikasi.jpeg
44K



837-Ari Asnani-(116-124)-OK.pdf
436K



Ari Asnani Pernyataan-Maret2022-EN.pdf
89K