


Home (<https://smujo.id/biodiv/index>) / Archives (<https://smujo.id/biodiv/issue/archive>)
/ Vol. 18 No. 3 (2017)



(<https://smujo.id/biodiv/issue/view/157>)

Vol. 18 No. 3 (2017)

Articles

Diversity of endophytic actinobacteria isolated from medicinal plants and their potency as pancreatic lipase inhibitor 857-863
(<https://smujo.id/biodiv/article/view/1935>)

LENNI FITRI

PDF (<https://smujo.id/biodiv/article/view/1935/1814>)

Diversity and pattern of nest preference of bat species at bat-dwelling caves in Gombong Karst, Central Java, Indonesia 864-874
(<https://smujo.id/biodiv/article/view/2129>)

FAHMA WIJAYANTI, IBNU MARYANTO

pdf (<https://smujo.id/biodiv/article/view/2129/2002>)

Behavioral ecology of reintroduced Orangutans in the Bukit Batikap, Central Kalimantan, Indonesia 875-886
(<https://smujo.id/biodiv/article/view/1999>)

IKE NURJUITA NAYASILANA, SUWARNO HADISUSANTO, HERY WIJAYANTO, SRI SUCI UTAMI ATMOKO, DIDIK PRASETYO, JAMARTIN SIHITE, CAREL P. VAN SCHAIK

pdf (<https://smujo.id/biodiv/article/view/1999/1878>)

Short Communication: Abundance of Arbuscular Mycorrhiza associated with corn planted with traditional and more modern farming systems in Kupang, East Nusa Tenggara, Indonesia 887-892
(<https://smujo.id/biodiv/article/view/2000>)

Short Communication: Rapid and accurate genetic authentication of *Penthorum chinense* by improved RAPD-derived species-specific SCAR markers (<https://smujo.id/biodiv/article/view/2049>) 1243-1249

ZHIQIANG MEI, MD. ASADUZZAMAN KHAN, XIANQIN ZHANG, JUNJIANG FU

pdf (<https://smujo.id/biodiv/article/view/2049/1925>)

Iron (II) and other heavy-metal tolerance in bacteria isolated from rock varnish in the arid region of Al-Jafer Basin, Jordan (<https://smujo.id/biodiv/article/view/2050>) 1250-1257

SULAIMAN ALNAIMAT, SAQER ABU SHATTAL, OSAMA ALTHUNIBAT, EID ALSBOU, REDA AMASHA

pdf (<https://smujo.id/biodiv/article/view/2050/1926>)

Throughfall quantity and carbon input beneath canopy gaps of varying size in degraded tropical peatland forest of West Kalimantan, Indonesia (<https://smujo.id/biodiv/article/view/2051>) 1258-1264

DWI ASTIANI, LISA M. CURRAN, MUJIMAN MUJIMAN, RUSPITA SALIM

pdf (<https://smujo.id/biodiv/article/view/2051/1927>)

Characterization of *Bifidobacteria* from infant feces with different mode of birth at Purwokerto, Indonesia (<https://smujo.id/biodiv/article/view/2052>) 1265-1269

PANCRASIA MARIA HENDRATI, DYAH FITRI KUSHARYATI, DINI RYANDINI, OEDJIJONO OEDJIJONO

pdf (<https://smujo.id/biodiv/article/view/2052/1928>)

Short Communication: Weak delineation of *Styrax* species growing in North Sumatra, Indonesia by *matK* + *rbcl* gene (<https://smujo.id/biodiv/article/view/2053>) 1270-1274

ARIDA SUSILOWATI, HENTI HENDALASTUTI, CUT RIZLANI KHOLIBRINA, RINI RAMADHANI

pdf (<https://smujo.id/biodiv/article/view/2053/1929>)

Occurrence of chitinolytic bacteria in shrimp rusip and measurement of their chitin-degrading enzyme activities (<https://smujo.id/biodiv/article/view/2054>) 1275-1281

INDUN DEWI PUSPITA, ARI WARDANI, RR. OKI ARUM PUSPITASARI, PRIHATI SIH NUGRAHENI, MGS. MUHAMMAD PRIMA PUTRA, SRI PUDJIRAHARTI, USTADI USTADI

pdf (<https://smujo.id/biodiv/article/view/2054/1930>)

Isolation and screening of cellulolytic fungi from *Salacca zalacca* leaf litter 1282-1288 (<https://smujo.id/biodiv/article/view/2055>)

SITI LUSI ARUM SARI, RATNA SETYANINGSIH, NOVITA FITRIATUL AINI WIBOWO

pdf (<https://smujo.id/biodiv/article/view/2055/1931>)

Information

For Readers (<https://smujo.id/biodiv/information/readers>)

For Authors (<https://smujo.id/biodiv/information/authors>)

For Librarians (<https://smujo.id/biodiv/information/librarians>)

Usage Statistics Information

We log anonymous usage statistics. Please read the privacy information (<https://smujo.id/biodiv/usageStats/privacyInformation>) for details.

Journals List

Biodiversitas Journal of Biological Diversity (<https://smujo.id/biodiv>)

Nusantara Bioscience (<https://smujo.id/nb>)

Prosiding Seminar Nasional Masyarakat Biodiversitas Indonesia (<https://smujo.id/psnmbi>)

Asian Journal of Agriculture (<https://smujo.id/aja>)

Asian Journal of Ethnobiology (<https://smujo.id/aje>)

Asian Journal of Forestry (<https://smujo.id/ajf>)

Biofarmasi Journal of Natural Product Biochemistry (<https://smujo.id/jnpb>)

Bioteknologi Biotechnological Studies (<https://smujo.id/bbs>)

Bonorowo Wetlands (<https://smujo.id/bw>)

Cell Biology and Development (<https://smujo.id/cbd>)

Ocean Life (<https://smujo.id/ol>)

Tropical Drylands (<https://smujo.id/td>)

Reviewers List

Reviewers (<https://smujo.id/biodiv/reviewers/index>)

Visitor Statistics

Statistics (<https://smujo.id/info/stats>)

Visitors



(<https://info.flagcounter.com/JKAr>)



(<https://www.scopus.com/sourceid/21100332431>)



(<http://doaj.org/toc/3eda1e70aa014e1e8bd1fc367b4df956>)



(<http://scholar.google.co.id/citations?hl=id&user=rae-lrEAAAAJ>)



(<http://search.crossref.org/?q=biodiversitas&type=Journal>)



(<https://academic.microsoft.com/#/detail/2738269101>)

BIODIVERSITAS

Journal of Biological Diversity

[Contact Us](#)
[FAQ](#)
[Useful links](#)
[Editors-PDF](#)
[Home](#)
[Content and Archives](#)
[Aims and Scope](#)
[Editorial Board](#)
[Guidance for Authors](#)
[Ethical Guidelines](#)
[Charges](#)
[Membership](#)
[Previously Reviewers](#)
[Conference Events](#)

EDITOR-IN-CHIEF:

Sutarno

EDITORIAL MEMBERS:

English Editors: Graham Eagleton (grahameagleton@gmail.com), Suranto (surantouns@gmail.com); Technical Editor & Banking: Artini Pangastuti (pangastuti_tutut@yahoo.co.id), Solichatun (solichatun_s@yahoo.com); Distribution & Marketing: Rita Rakhmawati (oktia@yahoo.com); Webmaster: Ari Pitoyo (aripitoyo@yahoo.co.id)

MANAGING EDITORS:

Ahmad Dwi Setyawan (unsjournals@gmail.com)

EDITORIAL BOARD (COMMUNICATING EDITORS):

Abd Fattah N. Abd Rabou, Islamic University of Gaza, Palestine
Agnieszka B. Najda, University of Life Sciences in Lublin, Lublin, Poland
Ajay Kumar Gautam, Abhilashi University Mandi, Himachal Pradesh, India
Annisa, Padjadjaran University, Sumedang, Indonesia
Alan J. Lymbery, Murdoch University, Perth, Australia
Bambang Hero Saharjo, Institut Pertanian Bogor, Bogor, Indonesia
Daiane H. Nunes, State University of Londrina, Londrina, Brazil
Darlina Md. Naim, University Sains Malaysia, Penang, Malaysia
Ghulam Hassan Dar, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, Srinagar, India
Hassan Pourbabaei, University of Guilan, Somehsara, Guilan, Iran
Joko Ridho Witono, Center for Plant Conservation-Bogor Botanical Gardens, Indonesian Institute of Sciences, Bogor, Indonesia
Kartika Dewi, Research Center for Biology, Indonesian Institute of Sciences, Cibinong, Bogor, Indonesia
Katsuhiko Kondo, University of Missouri, Columbia, USA
Kusumadewi Sri Yulita, Research Center for Biology, Indonesian Institute of Sciences, Cibinong, Bogor, Indonesia
Livia Wanntorp, Naturhistoriska riksmuseet, Stockholm, Sweden
M. Jayakara Bhandary, Government Arts and Science College, Karwar, Karnataka, India
Mahdi Reyahi-Khoram, Islamic Azad University (Hamadan Branch), Hamadan, Iran
Mahendra Kumar Rai, SGB Amravati University, Maharashtra, India
Mahesh K. Adhikari, Adhikari Niwas, Kathmandu, Nepal
Maria Panitsa, University of Patras, Agrinio, Greece
Mochamad A. Soendjoto, Lambung Mangkurat University, Banjarbaru, Indonesia
Mohamed M.M. Najim, University of Kelaniya, Kelaniya, Sri Lanka
Mohib A. Shah, Nepean Telehealth Technology Centre, Sydney, Australia
Nurhasanah, Mulawarman University, Samarinda, Indonesia
Praptiwi, Research Center for Biology, Indonesian Institute of Sciences, Cibinong, Bogor, Indonesia
Rasool B. Tareen, University of Balochistan, Quetta, Pakistan
Seyed Aliakbar Hedayati, Gorgan University of Agricultural Sciences and Natural Resources, Iran
Seyed Mehdi Talebi, Arak University, Iran
Shahabuddin, Universitas Tadulako, Palu, Indonesia

Shahir Shamsir, Universiti Teknologi Malaysia, Skudai, Malaysia
Shri Kant Tripathi, Mizoram University, Aizawl, India
Sugeng Budiharta, Purwodadi Botanical Gardens, Indonesian Institute of Sciences, Pasuruan, Indonesia
Sugiyarto, Universitas Sebelas Maret, Surakarta, Central Java, Indonesia
Subash C. Santra, University of Kalyani, India
Taufiq Purna Nugraha, Research Center for Biology, Indonesian Institute of Sciences, Cibinong, Bogor, Indonesia
Yosep S. Mau, Nusa Cendana University, Kupang, Indonesia

ISSN 1412-033X | E-ISSN 2085-4722

PUBLISHER: Society for Indonesian Biodiversity

CO-PUBLISHER: Department of Biology, FMNS, Sebelas Maret University Surakarta

FIRST PUBLISHED: 2000

ADDRESS:

Sebelas Maret University
Jl. Ir. Sutami 36A Surakarta 57126. Tel. +62-271-7994097, Tel. & Fax.: +62-271-663375, email: editors@smujo.id

ONLINE: biodiversitas.mipa.uns.ac.id

[Download the Editorial Boards.](#)

[Home](#) | [Archives](#) | [Contact Us](#)

Copyright © Biodiversitas, 2012. All Rights Reserved

BIODIVERSITAS

Journal of Biological Diversity

[Contact Us](#)[FAQ](#)[Useful links](#)[Home](#)[Content and Archives](#)[Aims and Scope](#)[Editorial Board](#)[Guidance for Authors](#)[Ethical Guidelines](#)[Charges](#)[Membership](#)[Previously Reviewers](#)[Conference Events](#)

Aims and Scope "Biodiversitas, Journal of Biological Diversity" or Biodiversitas encourages submission of manuscripts dealing with all biodiversity aspects of plants, animals and microbes at the level of gene, species, and ecosystem.

Article types The journal seeks original full-length research papers, reviews, and scientific feedback (short communication) about material previously published.

[Home](#) | [Archives](#) | [Contact Us](#)

Copyright © Biodiversitas, 2012. All Rights Reserved

Characterization of Bifidobacteria from infant feces with different mode of birth at Purwokerto, Indonesia

PANCRASIA MARIA HENDRATI^{*}, DYAH FITRI KUSHARYATI, DINI RYANDINI, OEDJIJONO

Faculty of Biology, Universitas Jenderal Soedirman. Jl. Dr. Soeparno 63, Purwokerto, Banyumas 53122, Central Java, Indonesia.
Tel.: +62-281-638794, Fax.: +62-281-631700, ^{*}email: maria.hendrati@gmail.com

Manuscript received: 30 April 2017. Revision accepted: 30 July 2017.

Abstract. Hendrati PM, Kusharyati DF, Ryandini D, Oedjijono. 2017. Characterization of Bifidobacteria from infant feces with different mode of birth at Purwokerto, Indonesia. Biodiversitas 18: 1265-1269. Bifidobacteria belongs to the so-called beneficial intestinal flora. Before attempting to raise their intestinal levels to improve the health status of the host, it is importance to know about physiological variations in the Bifidobacterial colonization of the human intestine. Birth process influenced the diversity of Bifidobacteria in infant feces. This research was intended to isolate and characterize Bifidobacterium spp. as well to evaluate their presence in the feces of infants who were born by mode of normal, caesarean and premature. The research was conducted by survey method and data were analyzed with descriptive analysis. Bifidobacterium character was observed include colony and cell morphology. The biochemical test included catalase, oxidase, indole, Voges-Proskauer, different pH growth, and resistance to lysozyme. Bifidobacterium metabolites obtained tested its bacterial activity to Salmonella typhimurium and Escherichia coli. The result of this research showed that 35 isolates are suspected Bifidobacterium group and after API 20 A test showed 17 isolates are rally genera of Bifidobacterium spp. and all isolates come from infant feces with caesar and premature delivery. These isolates inhibited the growth of S. typhimurium and E. coli with different inhibitory capabilities. This finding is very important for science and medical point of view and could be developed with further research.

Keywords: Bifidobacteria, Characters, Mode of birth, Infant feces

INTRODUCTION

Probiotics are living microorganisms that provide health benefits to the host when ingested in adequate amounts (Bermudez-Brito et al. 2012). In recent years, many studies have shown that probiotics can cure a variety of diseases, such as diarrhea, colitis, urinary tract infection, hypertension, allergies, and cancer. Probiotics benefits are important from an early age. The previous few years, the safe test of probiotic bacteria are on the group of Bifidobacteria, i.e. *Bifidobacterium lactis*. The probiotics use for children was focused on the prevention of diarrhea and allergies as well as increasing the number of Bifidobacteria in the gut. Bifidobacteria is the dominance microorganism in the human digestive tract of adults and infants (Holzapfel et al. 2001). The dominance of Bifidobacteria in the human digestive tract is about 10% in adults and up to 90% in infants (Harmsen et al., 2000a) that were analyzed from stool samples. Bifidobacteria colonize the intestine of infants during birth and the first week after birth (Heidarpour et al. 2008). Narayanan and Subramonian (2015) showed that there are about 29 Bifidobacteria species that have been identified, including eleven species isolated from infant feces. The most frequently isolated Bifidobacteria species in infant feces are *B. bifidum*, *B. longum*, *B. infantis* and *B. breve*. These organisms are Gram-positive, non-motile, non-spore forming, anaerobic, and pleomorphic rods. Some researchers suspect that the presence of intestinal Bifidobacteria in infants is affected

by birth process (Mariat et al. 2009). During vaginal birth process (normal), the contact with vaginal and intestinal flora are important resources to begin the colonization in infants, while during a cesarean birth process, direct contact of infant's mouth with vaginal and intestinal microbiota are not happen, and the intestinal colonization is derived from the skin of his parents, workers, and medical equipment (Lif Holgerson et al. 2011).

The composition of the microbiota at the beginning of human life can affect the health for several months. Biasucci et al. (2008) stated that the composition of microbiota in infants plays an important role in the development of the post-partum immune system. The development of fecal microbiota in neonates is important because those bacteria are the first to colonize the sterile intestine of the neonates and, thus, have a significant effect on the host. Bifidobacteria are predominant in the fecal microbiota of infants, and, therefore, they are important to an understanding of how commensal Bifidobacteria is established in the intestine of infants (Mikami, Kimura, and Takahashi. 2012). The important of *Bifidobacterium* is the ability to produce antimicrobial compounds called bacteriocin (Aly et al. 2006).

The problem of this research is "how is the dominance of *Bifidobacterium* sp. from the feces of infants with a different birth process (normal, cesarean, and premature)?" Information of *Bifidobacterium* isolated from newborn feces is still lack in Indonesia. This reason encourages researcher to explore *Bifidobacterium* from infant feces

especially in Purwokerto. The research on *Bifidobacteria* as the agent of probiotics is currently being developed by isolating the potential isolate from infants feces with the different birth process. Vaginal birth process and breastfeeding will stimulate the presence of *Bifidobacteria* in the gastrointestinal tract of infants. The research was intended to isolate and characterize the *Bifidobacterium* spp. as well as to evaluate their presence in the feces of infants who were born by modes of normal, caesarean and premature.

MATERIALS AND METHODS

The research was conducted at Laboratory of Microbiology, Faculty of Biology, Jenderal Soedirman University using survey method. The feces were taken from 10 until 12 days old infant with the different birth process (normal, cesarean, and premature). Samples were taken randomly, i.e. feces from 2 normal, 2 cesar, and 1 premature birth process. The data was analyzed with descriptive analysis. Determination of *Bifidobacterium* spp. was based on the morphological, chemical, and sugar test. Manual characterization refers to Bergey's Determinative Bacteriology, Cowan and Steel's Manual for the Identification of Medical Bacteria, World Journal of Dairy & Food Sciences, International Journal of Food Microbiology, Applied and Environmental Microbiology, and followed by API 20A(Biomériux) test

Procedures

Isolation

Isolation of *Bifidobacterium* sp. from infants' feces used the Modification of Gronlund (Modification of Gronlund et al., 2009). Samples were taken aseptically from 5 feces of under one month-aged infant with the different birth process (normal, cesarean, and premature). A 5 g of feces was added with 45 mL of sterile distilled water, then homogenized and diluted for up to 10^{-4} . The last two dilutions were used to isolate the bacteria using de Man, Rogosa, Sharpe Agar (MRSA) Oxoid with a spread plate method in duplicate and were anaerobically incubated at 30°C for 24 hours.

Identification

Morphological characterization. Morphology of colony. Colony characterization was performed in MRSA, such as shape, size, surface, elevation and edge formation of the colony.

Cellular characterization. Gram staining. One loop of the single colony was taken, and then was smeared on a glass object. Crystal violet dye was dropped for 1 minute, and then was washed. Mordant solution (Lugol's iodine) was dropped as the second dye for one minute, and then was washed. Ethanol 96% was dropped gently, until it was lightly crisp, and then was washed. Safranin was dropped for one minute, and then was washed. The glass object was air-dried, then was examined under the microscope. The cells of Gram-positive bacteria were purple in color. Meanwhile, Gram-negative bacteria were red. Motility test. An ose of the single colony was stab inoculated on

Sulphide Indole Motility Agar (SIMA) medium, and then was incubated at room temperature for 2x24 hours. The spread of growth was observed. A positive result was interpreted with pellicle formation on the surface of the medium.

Biochemical test

Catalase test. One loop of the single colony was smeared on glass object, and then was dropped with the H_2O_2 reagent. A positive result was interpreted with gas bubbles formation.

Test indole. One loop of single colony was inoculated into indole medium (Tryptone Broth) for 2x24 hours at 35-36°C. Then, the medium was added with Kovac's Indole reagent. Positive the result was interpreted with pink rings formation on the surface of the medium.

Voges-Proskauer test. One loop of the single colony was inoculated into Methyl Red Voges Proskauer (MR-VP) medium, and then was incubated for 2x24 hours at 30°C. A total of 3 drops of 40% KOH and 2 drops of alpha-naphthol reagent were added to the medium. A positive result was interpreted as color change into the pink medium.

Oxidase test. One loop of the single colony was covered with a piece of filter paper, and then was added by 1-2 drops of reagent (tetramethyl-D-phenylenediamine dihydrochloride). Positive result was interpreted with a color change to maroon blue.

Measurement of Total Lactic Acid. A total of 2 mL of 24 hours bacterial culture was taken, and then was diluted with 20 ml of distilled water and was shaken. Then, 3 drops of phenolphthalein (pp 1%) were added and titrated with 0.1 N NaOH solution until a light pink color formation.

Resistance test to Lysozyme. MRSA medium was added with the white part of the egg containing lysozyme (200 mg.mL⁻¹). The medium was poured into a sterile dish and allowed to solidify. A 1 mL of bacterial culture was inoculated on *Bifidobacterium* medium, then was anaerobically incubated for 24 hours at 35-36°C. The bacterial resistance was shown by the growth of *Bifidobacterium* in the medium

The pH range Growth test. Strains isolated MRS broth was grown in media with pH 4,7,9 and was incubated for 24 hours at a temperature of 37°C.

API 20A (Biomériux) test

The colony isolated from infants feces was re-cultured on MRSA medium and anaerobically incubated for 24 hours, then 3 loops of the colony was taken and identified with kit of API 20A. Indole test was added with mineral and anaerobically incubated for 24 hours. Indole test interpretation needs the addition of XYL and EHR reagent. The sugar test, glucose-arabinose, and glycerol-trehalose need the addition of BCP reagent. Catalase test was added by H_2O_2 . The data was analyzed in the web of API 20

Inhibition test against pathogenic bacteria

Bacterial culture. *Salmonella typhimurium* and *Escherichia coli* were re-cultured in 100 mL Nutrient Broth (NB), and incubated in the shaker of incubator at 150 rotary per minute (rpm) for 24 hours.

Antibacterial activity. Suspension of *S. typhimurium* and *E. coli* were inoculated on 10 mL Nutrient Agar (NA) medium (spread plate method). 0.1 mL supernatant was dropped on Sterile paper disc with the diameter of 6 mm. Then, the paper disc was placed on the surface of NA medium (which had been inoculated with *S. typhimurium* and *E. coli*), then incubated for 24 hours at 30°C. Antimicrobial 1% of 24 hours culture of *Bifidobacterium* spp. was inoculated into MRSB medium, and centrifuged at 13,000 rpm for 15 minutes. The supernatant was adjusted with 1 N of NaOH to pH 6.5 (neutralized), and heated at 100°C for 15 minutes. Antimicrobial activity was determined by measuring the clear zone around the paper disc after incubation.

RESULTS AND DISCUSSION

Characterization test

Based on the results of the morphological and biochemical test, there are 35 isolates belonging to the genus *Bifidobacterium*. Morphology of colonies is similar in color, shape, surface and elevation, but quite different on the size and edge formation. The milky white colonies have a flat edge/ smooth (entire), some are uneven like wool, round shape, shiny surface, convex elevation with varying sizes. These isolates have morphological characters referring to genus *Bifidobacterium*, i.e. colony with milky white color or slightly creamy, rounded form with the diameter of 0.1-0.5 mm as well as Gram-positive (Wasilewska and Bielecka 2003; Lievin 2000; Hadadji et al. 2005).

Biochemical tests

The results of biochemical tests of 35 isolates by Gram staining test were confirmed as Gram-positive bacteria as the purple cell. Gram-positive bacteria do not undergo decolorization so it stays shiny purple with color of crystal violet staining and the end is not stained by safranin and Gram-positive bacteria contains a peptidoglycan thick. Motility test showed that all isolates showed no spreading pellicle formation on the surface of the medium and without turbidity; therefore it was interpreted as nonmotile bacteria. Garrity et al. (2005) showed that the motility test positive if growth isolates form a pellicle widespreading on the surface of the medium, while it didn't show the spread of pellicle on the medium surface, named nonmotile. The positive reaction of catalase test is indicated by the formation of O₂ gas bubbles as the breakdown of H₂O₂ by catalase enzyme produced by the bacteria. The negative reaction to catalase test is indicated by no formation of gas bubbles. According to Hadadji et al. (2005), *Bifidobacterium* is catalase negative-bacteria. Indole test showed a negative result, indicated by no formation of the pink ring on the surface of the medium. Zinedine and Faïd (2007) stated that *Bifidobacterium* is negative in indole test. Voges-Proskauer test showed the negative result, indicated by color change into pink or dark red after a few drops of 40% KOH and alpha-naphthol 5%. Venkatesh et al (2012) stated that *Bifidobacterium* shows a negative

result in VP tests, because the bacteria do not ferment mixed acid or butanediol fermentation. Oxidase test showed the positive result, indicated by blue-black color formation after a few drops of tetramethyl-D-phenylenediamine dihydrochloride. The color change is due to the cytochrome enzyme oxidase the reagent. According to Feliatra et al. (2004), *Bifidobacterium* shows a positive result in oxidase test. Our data of lactic acid concentration from 35 isolates were ranged from 0.05-0.224%. According to Silalahi and Ikhsan (2010), longer fermentation leads to the greater production of lactic acid. This is due to the increase length of fermentation time that increases the number of cell, therefore it increases the microbial activity to produce lactic acid.

Lysozyme test results showed that all isolates were able to grow on de Man, Rogosa and Sharpe Agar (MRSA) medium supplemented with the white part of the egg (containing lysozyme). According to Gagnon et al. (2004), the *Bifidobacterium* is more resistant to white-egg lysozyme or digestive tract than other bacteria, even at the concentration of 300 µg/ml which is higher than the concentration of intestinal lysozyme. The test results showed that the pH range for the better growth of *Bifidobacterium* spp. is at pH 7 than pH 4 and pH 9. According to Biavati et al. (2000), the *Bifidobacterium* is an acid-tolerant microorganism with an optimum pH range of 6.5-7.0

API 20A test

Indole test, after the addition of XYL reagent (2-3 min) and EHR (5 min), indicated a positive result with red color and the negative result with yellow color. Urease test, after the addition of XYL reagent (2-3 min) and EHR (5 min), indicated a positive result with red color and the negative result with yellow color. The test of Glucose, mannitol, lactose, saccharose, maltose, salicine, xylose, arabinose, glycerol, cellobiose, mannose, melezitose, raffinose, sorbitol, rhamnose, and trehalose, after the addition of BCP reagent, showed that the positive result was indicated by yellow or yellow-green, and the negative result was indicated by purple color. Gelatin positive test was indicated by the occurrence of black pigment with diffusion and negative result was without diffusion. Esulin positive test was shown in black brown and fluorescent, the negative result was shown in yellow without fluorescent. The result of API test showed that 17 of 35 isolates were confirmed as genus *Bifidobacterium* namely BC4, BC5, BC6, BC7, BC8, BC10, BC13, BC14, BC15, BC16, BC17, BC18, BC19, BP1, BP6, BP7, dan BP9 (Figure 1). In normal birth process, contact between infant and nonpathogenic bacteria in mother's vagina bacteria lead to colonization of probiotic bacteria, while in caesar and premature birth processes, infant doesn't have a chance of that contact, therefore colonization of probiotic bacteria doesn't occur. In this research, the feces of infants in normal birth process did not contain any *Bifidobacterium* in their fecal samples. This is presumably due to fecal samples of infants having mothers who live in rural areas that their modes of feeding do not fulfill the nutrition during pregnancy (Figure 1, 2 and 3).




Figure 1. Distribution of 35 isolates suspected of *Bifidobacterium* after API 20A test. Note: Nnb: normal birth, non *Bifidobacterium*, Nbi: normal birth, *Bifidobacterium*, Cnb: Caesarean birth, non *Bifidobacterium*, Cbi: Caesarean birth, *Bifidobacterium*, Pnb: Premature birth, non *Bifidobacterium*, Pbi: Premature birth, *Bifidobacterium*

Inhibitory activity test

Figure 2 and Figure 3 show that the inhibitory activity test of 17 isolates belonging into *Bifidobacterium* spp. against *E. coli* and *S. typhimurium*. The diameter of the inhibitory zone of supernatant with NaOH 1 N addition against *E. coli* ranges from 6.00-13.8 mm and diameter of the inhibitory zone of supernatant without the addition of NaOH 1 N against *E. coli* ranges from 6.8-12.5 mm. The diameter of the inhibitory zone of supernatant with NaOH 1 N addition against *S. typhimurium* ranges from 6.00-11.5 mm and diameter of the inhibitory zone of supernatant without the addition of NaOH 1 N against *S. typhimurium* ranges from 7.19-12 mm.




Figure 2. Diameter of inhibitory zone secondary metabolites of *Bifidobacterium* without the addition of NaOH




Figure 3. Diameter of inhibitory zone secondary metabolites of *Bifidobacterium* with the NaOH addition

Discussion

The results showed that the diameters of the inhibitory zone of supernatant without NaOH 1 N addition are varied. This result is due to the present of lactic acid as the antimicrobial agent. The present of Lactic acid from the fermentation of the *Bifidobacterium* spp. has several advantages, such as high solubility, low toxicity and as an effective antibacterial property (Lye et al.2009). The lactic acid can suppress the growth of pathogenic bacteria, and reduce the infection and carcinogenic effects (Venkatesan et al. 2012).

The clear zone formation indicates the ability of metabolite of *Bifidobacterium* spp. in inhibiting *E. coli* and *S. typhimurium*. The NaOH addition to pH 6.5 aims to eliminate the influence of the produced acid during metabolism, therefore the inhibition of *E. coli* and *S. typhimurium* is expected due to the activity of bacteriocins. Supernatant consisting of bacteriocins produces a clear zone with clear boundaries because bacteriocins are bactericidal which produce acid or other antimicrobial components which are bacteriostatic. According to Ray and Field (1992), bacteriocins produce a really clear zone. If there is no clear zone, it is estimated as the action of hydrogen peroxide, acid or diacetyl. Hydrogen peroxide is produced by lactic acid bacteria that can damage the structure of the microbial lipid membrane, thus will increase the membrane permeability, and then it will damage the structure of nucleic acids and proteins of the cell (De Vuyst and Leroy 2007).

Bacteriocin is secondary metabolites of lactic acid bacteria. Bacteriocins have the same way of working with antibiotic, which is able to inhibit the growth of certain bacteria. Bacteriocins in performing the antimicrobial activity will attack the cytoplasm (Naidu and Clemens, 2000). Mechanism of bactericidal activity of bacteriocins generally starts from a molecule bacteriocins that come into direct contact with the cell membrane, therefore the process of this contact causes the disturbance of membrane potential, i.e. destabilization depolarization of the cytoplasmic membrane, therefore the cells cannot survive. Membrane fluidity impacts the formation of holes or pores in the cell membrane through the disruption of Proton Motive Force (PMV) (Rodney et al. 2014).

In conclusion, there are 35 isolates expected to be *Bifidobacterium* group. After the API 20A test, 17 isolates from normal birth process, 14 isolates from cesarean birth process, and 3 isolates from premature were confirmed to be genus *Bifidobacterium* spp.. All these isolates inhibited the growth of *S. typhimurium* and *E. coli* and can also be used as potential inoculum in fermented healthy beverages.

REFERENCES

- Aly S, Ouattara Cheik AT, Bassole Imael HN, Traore SA. 2006. Bacteriocins and lactic acid bacteria-a minireview. *African J Biotechnol* 5 (9): 678-83.
- Bermudez-Brito M, Plaza-Díaz J, Muñoz-Quezada S, Gómez-Llrente C, Gil A. (2012). Probiotic mechanisms of action. *Ann Nutr Metab* 61: 160-174.

- Biasucci G, Benenati B, Morelli L, Bessi B, Boehm G. 2008. Cesarean delivery may affect the early biodiversity of intestinal bacteria. *J Nutr* 138 (9): 1796S-1800S. doi: 138/9S-II/1796S [pii].
- Gagnon M, Kheadr EE, Le Blay G, Fliss I. 2004. In vitro inhibition of *Escherichia coli* O157: H7 by bifidobacterial strains of human origin. *Intl J Food Microbiol* 92 (1): 69-78.
- Garrity GM, Brenner DJ, Kreig NR, Staley JT. 2005. *Bergey's Manual of Systematic Bacteriology*. Vol 2, The Proteobacteria. Springer, New York.
- Feliatra, Efendi I, Suryadi E. 2004. Isolation and identification probiotic bacteria kerapu macan fish in attempt fish feed efficiency in an effort. *Jurnal Natur Indonesia* 6 (2): 75-80. [Indonesian]
- Hadadji M, Benama R, Noureddine S, Saidi N, Henni DE, Kihal M. 2005. Identification of cultivable *Bifidobacterium* species isolated from breast-fed infants feces in West-Algeria. *African J Biotechnol* 4 (5): 422-340.
- Harmen HJ, Wildeboer-Veloo AC, Raangs GC, Wagendorp AA, Klijn N, Bindels JG, Welling GW. 2000. Analysis of intestinal flora development in breast-fed and formula-fed infants by using molecular identification and detection methods. *J Pediatr Gastroenterol Nutr* 30: 61-67.
- Holzappel W, Haberer P, Geisen R, Bjorkroth J, Schillinger U. 2001. Taxonomy and important features of probiotic microorganisms in food and nutrition. *Amer Soci Clin Nutr* 73: 365S-373S.
- Lievin V, Peiffer I, Hudault S, Rochat F, Brassart D, Neeser JR, Servin AL. 2000. *Bifidobacterium* strain from resident infant human gastrointestinal microflora exerts antimicrobial activity. *Gut* 47: 646-652.
- Lif Holgersson P, Harnevik L, Hernell O, Tanner ACR, Johansson I. 2011. Mode of Birth Delivery Affects Oral Microbiota in Infants. *J Dental Res* 90 (10): 1183-1188.
- Lye H, Kuan C, Ewe J, Fung W, Liong M. 2009. The improvement of hypertension by probiotics: effects on cholesterol, diabetes, renin, and phytoestrogens. *Intl J Mol Sci* 10 (9): 3755-3775.
- Mariat D, Firmesse O, Levenez F, Guimar VD, Sokol H, Doré J, Corthier G, Furet JP. 2009. The Firmicutes/Bacteroidetes ratio of the human microbiota changes with age. *BMC Microbiol*. 2009;9. DOI: 10.1186/1471-2180-9-123.
- Mikami K, Kimura M, Takahashi H. 2012. Influence of maternal bifidobacteria on the development of gut bifidobacteria in infants. *Pharmaceuticals* 5: 629-642.
- Naidu AS, Clemens. 2000. Probiotics. In: *Food Antimicrobial System AS* (ed). CRC Press, LLC
- Narayanan R, Subramonian BS. 2015. Effect of Prebiotics on Bifidobacterial Species Isolated from Infant Faeces. *Indian J. Trad Knowl* 14 (2): 285-89.
- Perez RH, Zendo T, Sonomoto K. 2014. Proceeding from 11 th International Symposium on Lactic Acid Bacteria. Egmond aan Zee, the Netherlands. 31 August-4 September 2014.
- Ray B, Field RA. 1992. Antibacterial effectiveness of a pediocin ach based biopreservatif against spoilage and pathogenic bacteria from vacuum package refrigerated meat. 38th International Congres of Meat Science and Technology. 4: 31-734
- Silalahi FY, Ikhsan M. 2010. Fruit-gurt Fermentation with variations rind. [Hon. Thesis]. Chemical Engineering Department, Faculty of Engineering, Diponegoro University, Semarang. [Indonesian].
- Venkatesan S, Kirithika M, Roselin I, Ganesan R, Muthuchelian K. 2012. Comparative invitro and invivo study of three probiotic organisms, *Bifidobacterium* sp., *Lactobacillus* sp., *S. cerevisiae* and analyzing its improvement with the supplementation of prebiotics. *Intl J Plant Anim Environ Sci* 2 (2): 94-106.
- Wasilewska E, Bielecka M. 2003. Isolation and identification of Bifidobacteria From Infant GUT Ewa. *J Food Nutr Sci* 12 (53): 90-94.
- Vuyst LD, Leroy F. 2007. Bacteriocins from lactic acid bacteria: production, purification, and food applications. *J Mol Microbiol Biotechnol* 13 (4): 194-99.
- Zinedine A, Faïd M. 2007. Isolation and characterization of strains of Bifidobacteria with probiotic proprieties in vitro. *World J Dairy Food Sci* 2 (1): 28-34.