


ISSN: 1412-033X
E-ISSN: 2085-4722

BIODIVERSITAS

Journal of Biological Diversity

Volume 22 - Number 3 - March 2021



GUIDANCE FOR AUTHORS

Aims and Scope *Biodiversitas*, *Journal of Biological Diversity* or abbreviated as *Biodiversitas* encourages submission of manuscripts dealing with all biodiversity aspects of plants, animals and microbes at the level of the gene, species, and ecosystem as well as ethnobiology.

Article types The journal seeks original full-length research papers, reviews, and short communication. Manuscript of original research should be written in no more than 8,000 words (including tables and picture), or proportional with articles in this publication number. Review articles will be accommodated, while, short communication should be written at least 2,000 words, except for pre-study.

Submission The journal only accepts online submission, through open journal system (<https://smujo.id/biodiv/about/submissions>) or email to the editors at unsjournals@gmail.com. Submitted manuscripts should be the original works of the author(s). The manuscript must be accompanied by a cover letter containing the article title, the first name and last name of all the authors, a paragraph describing the claimed novelty of the findings versus current knowledge. Submission of a manuscript implies that the submitted work has not been published before (except as part of a thesis or report, or abstract); and is not being considered for publication elsewhere. When a manuscript written by a group, all authors should read and approve the final version of the submitted manuscript and its revision; and agree the submission of manuscripts for this journal. All authors should have made substantial contributions to the concept and design of the research, acquisition of the data and its analysis; drafting of the manuscript and correcting of the revision. All authors must be responsible for the quality, accuracy, and ethics of the work.

Ethics Author(s) must obedient to the law and/or ethics in treating the object of research and pay attention to the legality of material sources and intellectual property rights.

Copyright If and when the manuscript is accepted for publication, the author(s) still hold the copyright and retain publishing rights without restrictions. Authors or others are allowed to multiply article as long as not for commercial purposes. For the new invention, authors are suggested to manage its patent before published.

Open access The journal is committed to free-open access that does not charge readers or their institutions for access. Readers are entitled to read, download, copy, distribute, print, search, or link to the full texts of articles, as long as not for commercial purposes. The license type is CC-BY-NC-SA.

Acceptance The only articles written in English (U.S. English) are accepted for publication. Manuscripts will be reviewed by editors and invited reviewers(double blind review) according to their disciplines. Authors will generally be notified of acceptance, rejection, or need for revision within 1 to 2 months of receipt. The manuscript is rejected if the content does not in line with the journal scope, does not meet the standard quality, inappropriate format, complicated grammar, dishonesty (i.e. plagiarism, duplicate publications, fabrication of data, citations manipulation, etc.), or ignoring correspondence in three months. The primary criteria for publication are scientific quality and biodiversity significance. **Uncorrected proofs** will be sent to the corresponding author by email as .doc or .docx files for checking and correcting of typographical errors. To avoid delay in publication, corrected proofs should be returned in 7 days. The accepted papers will be published online in a chronological order at any time, but printed in the early of each month (12 times).

A charge Starting on January 1, 2019, publishing costs waiver is granted to authors of graduate students from **Least Developed Countries**, who first publish the manuscript in this journal. However, other authors are charged USD 250 (IDR 3,500,000). Additional charges may be billed for language editing, USD 75-150 (IDR 1,000,000-2,000,000).

Reprints The sample journal reprint is only available by special request. Additional copies may be purchased when ordering by sending back the uncorrected proofs by email.

Manuscript preparation Manuscript is typed on A4 (210x297 mm²) paper size, in a single column, single space, 10-point (10 pt) Times New Roman font. The margin text is 3 cm from the top, 2 cm from the bottom, and 1.8 cm from the left and right. Smaller lettering size can be applied in presenting table and figure (9 pt). Word processing program or additional software can be used, however, it must be PC compatible and Microsoft Word based (.doc or .rtf, not .docx). **Scientific names** of species (incl. subspecies, variety, etc.) should be written in italic, except for italic sentence. Scientific name (genera, species, author), and cultivar or strain should be mentioned completely for the first time mentioning it in the body text, especially for taxonomic manuscripts. Name of genera can be shortened after first mentioning, except generating confusion. Name of the author can be eliminated after first mentioning. For example, *Rhizopus oryzae* L. UICC 524, hereinafter can be written as *R. oryzae* UICC 524. Using trivial name should be avoided, otherwise generating confusion. **Biochemical and chemical nomenclature** should follow the order of the IUPAC - IUB. For DNA sequence, it is better used Courier New font. Symbols of standard chemical and abbreviation of chemistry name can be applied for common and clear used, for example, completely written butilic hydroxyl toluene (BHT) to be BHT hereinafter. **Metric measurement** use IS denomination, usage other system should follow the value of equivalent with the denomination of IS first mentioning. Abbreviations set of, like g, mg, mL, etc. do not follow by dot. Minus index (m², L⁻¹, h⁻¹) suggested to be used, except in things like "per-plant" or "per-plot". **Equation of mathematics** does not always can be written

down in one column with text, in that case can be written separately. **Number** one to ten are expressed with words, except if it relates to measurement, while values above them written in number, except in early sentence. The fraction should be expressed in decimal. In the text, it should be used "%" rather than "percent". Avoid expressing ideas with complicated sentence and verbiage, and used efficient and effective sentence.

Title of the article should be written in compact, clear, and informative sentence, preferably not more than 20 words. Name of author(s) should be completely written. **Name and institution** address should also be completely written with street name and number (location), postal code, telephone number, facsimile number, and email address. Manuscript written by a group, author for correspondence along with address is required. First page of the manuscript is used for writing above information.

Abstract should not be more than 200 words. **Keywords** is about five words, covering scientific and local name (if any), research theme, and special methods which used; and sorted from A to Z. All important **abbreviations** must be defined at their first mention. **Running title** is about five words. **Introduction** is about 400-600 words, covering the background and aims of the research. **Materials and Methods** should emphasize on the procedures and data analysis. **Results and Discussion** should be written as a series of connecting sentences, however, for manuscript with long discussion should be divided into subtitles. Thorough discussion represents the causal effect mainly explains for why and how the results of the research were taken place, and do not only re-express the mentioned results in the form of sentences. **Concluding** sentence should be given at the end of the discussion. **Acknowledgments** are expressed in a brief; all sources of institutional, private and corporate financial support for the work must be fully acknowledged, and any potential conflicts of interest are noted.

Figures and Tables of maximum of three pages should be clearly presented. Title of a picture is written down below the picture, while title of a table is written above the table. Colored figures can only be accepted if the information in the manuscript can lose without those images; chart is preferred to use black and white images. Author could consign any picture or photo for the front cover, although it does not print in the manuscript. All images property of others should be mentioned source. **There is no appendix**, all data or data analysis are incorporated into Results and Discussions. For broad data, it can be displayed on the website as a supplement.

References Author-year citations are required. In the text give the authors name followed by the year of publication and arrange from oldest to newest and from A to Z. In citing an article written by two authors, both of them should be mentioned, however, for three and more authors only the first author is mentioned followed by et al., for example: Saharjo and Nurhayati (2006) or (Boonkerd 2003a, b, c; Sugiyarto 2004; El-Bana and Nijs 2005; Balagadde et al. 2008; Webb et al. 2008). Extent citation as shown with word "cit" should be avoided. Reference to unpublished data and personal communication should not appear in the list but should be cited in the text only (e.g., Rifai MA 2007, pers. com. (personal communication); Setyawan AD 2007, unpublished data). In the reference list, the references should be listed in an alphabetical order (better, if only 20 for research papers). Names of journals should be abbreviated. Always use the standard abbreviation of a journal's name according to the **ISSN List of Title Word Abbreviations** (www.issn.org/2-22661-LTWA-online.php). The following examples are for guidance.

Journal:

Saharjo BH, Nurhayati AD. 2006. Domination and composition structure change at hemic peat natural regeneration following burning: a case study in Pelalawan, Riau Province. *Biodiversitas* 7: 154-158.

Book:

Rai MK, Carpinella C. 2006. Naturally Occurring Bioactive Compounds. Elsevier, Amsterdam.

Chapter in book:

Webb CO, Cannon CH, Davies SJ. 2008. Ecological organization, biogeography, and the phylogenetic structure of rainforest tree communities. In: Carson W, Schnitzer S (eds) *Tropical Forest Community Ecology*. Wiley-Blackwell, New York.

Abstract:

Assaeed AM. 2007. Seed production and dispersal of *Rhazya stricta*. 50th annual symposium of the International Association for Vegetation Science, Swansea, UK, 23-27 July 2007.

Proceeding:

Alikodra HS. 2000. Biodiversity for development of local autonomous government. In: Setyawan AD, Sutarno (eds.) *Toward Mount Lawu National Park; Proceeding of National Seminary and Workshop on Biodiversity Conservation to Protect and Save Germplasm in Java Island*. Universitas Sebelas Maret, Surakarta, 17-20 July 2000. [Indonesian]

Thesis, Dissertation:

Sugiyarto. 2004. Soil Macro-invertebrates Diversity and Inter-Cropping Plants Productivity in Agroforestry System based on Sengon. [Dissertation]. Universitas Brawijaya, Malang. [Indonesian]

Information from internet:

Balagadde FK, Song H, Ozaki J, Collins CH, Barnet M, Arnold FH, Quake SR, You L. 2008. A synthetic *Escherichia coli* predator-prey ecosystem. *Mol Syst Biol* 4: 187. www.molecularsystemsbiology.com

BIODIVERSITAS

Journal of Biological Diversity
Volume 22 – Number 3 – March 2021

ISSN/E-ISSN:
1412-033X (printed edition), 2085-4722 (electronic)

EDITORIAL BOARD:

Abdel Fattah N.A. Rabou (Palestine), Agnieszka B. Najda (Poland), Ajay Kumar Gautam (India), Alan J. Lymbery (Australia), Annisa (Indonesia), Bambang H. Saharjo (Indonesia), Daiane H. Nunes (Brazil), Darlina Md. Naim (Malaysia), Ghulam Hassan Dar (India), Hassan Pourbabaee (Iran), Joko R. Witono (Indonesia), Kartika Dewi (Indonesia), Katsuhiko Kondo (Japan), Kusumadewi Sri Yulita (Indonesia), Livia Wanntorp (Sweden), M. Jayakara Bhandary (India), Mahdi Reyahi-Khoram (Iran), Mahendra K. Rai (India), Mahesh K. Adhikari (Nepal), Maria Panitsa (Greece), Mochamad A. Soendjoto (Indonesia), Mohib Shah (Pakistan), Mohamed M.M. Najim (Srilanka), Nurhasanah (Indonesia), Praptiwi (Indonesia), Rasool B. Tareen (Pakistan), Seyed Aliakbar Hedayati (Iran), Seyed Mehdi Talebi (Iran), Shahabuddin (Indonesia), Shahir Shamsir (Malaysia), Shri Kant Tripathi (India), Subhash C. Santra (India), Sugeng Budiharta (Indonesia), Sugiyarto (Indonesia), Taufiq Purna Nugraha (Indonesia), Yosep S. Mau (Indonesia)

EDITOR-IN-CHIEF: Sutarno

EDITORIAL MEMBERS:
English Editors: **Graham Eagleton** (grahameagleton@gmail.com), **Suranto** (surantouns@gmail.com); Technical Editor: **Solichatun** (solichatun_s@yahoo.com), **Artini Pangastuti** (pangastuti_tutut@yahoo.co.id); Distribution & Marketing: **Rita Rakhmawati** (oktia@yahoo.com); Webmaster: **Ari Pitoyo** (aripitoyo@yahoo.com)

MANAGING EDITORS:
Ahmad Dwi Setyawan (unsjournals@gmail.com)

PUBLISHER:
The Society for Indonesian Biodiversity

CO-PUBLISHER:
Department of Biology, Faculty of Mathematics and Natural Sciences, Sebelas Maret University, Surakarta

ADDRESS:
Jl. Ir. Sutami 36A Surakarta 57126. Tel. +62-271-7994097, Tel. & Fax.: +62-271-663375, email: editors@smujo.id

ONLINE:
biodiversitas.mipa.uns.ac.id; smujo.id/biodiv



Society for Indonesia
Biodiversity



Sebelas Maret University
Surakarta

Published by Smujo International for The Society for Indonesian Biodiversity and Sebelas Maret University Surakarta

Journal Article



Article Title

Morphological, molecular and resistance responses to soft-rot disease variability among plantlets of *Phalaenopsis amabilis* regenerated from irradiated protocorms

HALIDA ADISTYA PUTRI, AGUS PURWITO, SUDARSONO, DEWI SUKMA



Agronomic characters, genetic and phenotypic diversity coefficients, and heritability of 12 genotypes of rice

JAENUDIN KARTAHADIMAJA, SETYO DWI UTOMO, ERWIN YULIADI, ABDUL KADIR SALAM, WARSONO, ANUNG WAHYUDI



Short Communication: A comparison of stand structure, species diversity and aboveground biomass between natural and planted mangroves in Sikka, East Nusa Tenggara, Indonesia



Molecular identification of five species of family Chaetomiaceae (Sordariomycetes, Ascomycota) from Iraqi soil

MUSTAFA A. AL-DOSSARY, SHAYMAA S. RAHEEM, MAYSOON K. ALMIYAH



Short Communication: A small-scale coprological survey of the endoparasites in the Himalayan goral *Naemorhedus goral* (Hardwick, 1825) in Nepal

JAGAN NATH ADHIKARI, ROSHAN BABU ADHIKARI, BISHNU PRASAD BHATTARAI, TEJ BAHADUR THAPA, TIRTH RAJ GHIMIRE



✓ Extracellular protease from *Bacillus subtilis* 6315 with antagonistic activity against bacterial wilt pathogen (*Ralstonia solanacearum*) of chili

NUR PRIHATININGSIH, ARI ASNANI, HERU ADI DJATMIKO



A pilot study into ecological burning in forests as part of a species conservation plan

Prosiding Seminar Nasional Masyarakat Biodiversitas Indonesia

Asian Journal of Agriculture

Asian Journal of Ethnobiology

Asian Journal of Forestry

Biofarmasi Journal of Natural Product Biochemistry

Bioteknologi Biotechnological Studies

Bonorowo Wetlands

Cell Biology and Development

Ocean Life

Tropical Drylands

Reviewers List

Reviewers

Extracellular protease from *Bacillus subtilis* B315 with antagonistic activity against bacterial wilt pathogen (*Ralstonia solanacearum*) of chili

NUR PRIHATININGSIH^{1,✉}, ARI ASNANI², HERU ADI DJATMIKO¹

¹Faculty of Agriculture, Universitas Jenderal Soedirman. Jl. Dr. Suparno, Purwokerto Utara, Banyumas 53123, Central Java, Indonesia.

Tel./fax. +62-281-638791, ✉email: nur.prihatiningsih@unsoed.ac.id

²Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Jenderal Soedirman. Jl. Dr. Suparno, Purwokerto Utara, Banyumas 53123, Central Java, Indonesia

Manuscript received: 22 January 2020. Revision accepted: 14 February 2021.

Abstract. Prihatiningsih N, Asnani A, Djatmiko HA. 2021. Extracellular protease from *Bacillus subtilis* B315 with antagonistic activity against bacterial wilt pathogen (*Ralstonia solanacearum*) of chili. *Biodiversitas* 22: 1291-1295. Antagonistic bacteria isolated from a healthy potato are potential biocontrol agents against a number of bacterial pathogens. The capability of antagonistic bacteria to produce extracellular protease could be considered as an indicator of their antagonistic capacity. The objective of this research was to study the optimum activity of protease produced by *Bacillus subtilis* B315 to control *Ralstonia solanacearum* causing wilt of chili. In this research were optimum incubation time, pH and crude extract protease concentration of *B. subtilis* B315 examined for maximum production of extracellular protease. The results showed that *B. subtilis* B315 was capable of producing protease at optimum incubation time of 30 min, pH 9, and crude extract protease concentration 0.50% with an activity of 1.3185 U/mL. Protease-producing capability of *B. subtilis* B315 was shown by the formation of a clear zone around the colony of *B. subtilis* B315 grown on skim-milk agar medium. *B. subtilis* B315 was capable of inhibiting the growth of *R. solanacearum* in vitro as indicated by 32 mm clear zone. Furthermore, in chili application of *B. subtilis* showed that this bacteria was capable of controlling chili bacterial wilt caused by *R. solanacearum* with effectiveness of 60.89%. The prospective of *B. subtilis* B315 could be formulated as biopesticide.

Keywords: *Bacillus subtilis* B315, bacterial wilt, biocontrol, chili, protease activity

INTRODUCTION

Bacterial wilt caused by *Ralstonia solanacearum* is an important disease in plants that may include tomato, chili, eggplant, bananas, potatoes, ginger, and mulberries. In chili and tomato, bacterial wilt could reduce yields up to 80% (Vanitha et al. 2009). Currently, biological control using antagonistic bacteria has been widely studied, as it is more environmentally friendly and supports sustainable agriculture. Antagonistic bacteria could be explored from habitats of which the pathogen naturally lives.

Bacillus subtilis B315 isolated from a healthy potato rhizosphere has been reported to be capable of controlling bacterial wilt in potatoes with an effectiveness of 64.9% (Prihatiningsih 2013), and 74.6% in tomato, chili, and eggplant (Prihatiningsih and Djatmiko 2016). The potential of bacteria to be antagonist is shown by its ability to produce secondary metabolites (antibiotics and other compounds) and enzymes. Protease is a product of microbes that could function to control plant pathogens, either fungi or bacteria. *Bacillus* sp. and other bacteria such as: *B. stercorophilus*, *B. mojavensis*, *B. megaterium* and *B. subtilis* (Shumi and Anwar 2004) secrete extracellular enzymes including proteases. Saha et al. (2012) found two new strains of *B. subtilis* from eggplant rhizosphere that have the potential to be biocontrol agents which is shown by their capabilities of producing secondary metabolites, such as siderophore, and hydrolytic enzymes, such as chitinase, protease, lipase and amylase. *B. subtilis* was also

reported to be able to produce IAA (*Indole Acetic Acid*) ranging from 57.56-79.33 ppm (Prihatiningsih et al. 2020). *B. subtilis* has the ability to secrete amylase, protease, pullulanase, chitinase, xylanase, and lipase (Morikawa 2006; Lestari et al. 2017).

Protease is a hydrolytic enzyme that hydrolyses extracellular proteins and increases the ability of cells to absorb useful hydrolysis products. *B. subtilis* has important bacteria because of its capability to secrete several hydrolytic enzymes, including protease, into the culture medium (Almas et al. 2009). Antagonistic *Bacillus* sp. significantly colonize rice plants and secretes various biocontrol determinants like protease of 1.1±5.5 U/mg of soil or U/mL of hydroponic solution. *Bacillus* spp. strains KFP-5 and KFP-17 produce various quantities of siderophore, protease and glucanase in absence as well as in presence of rice blast pathogen *Pyricularia oryzae*. The protease production is ranging about 20.9-29.3 U/mL and 11.6-24 U/mL in KFP-5 and KFP-17, respectively (Rais et al. 2017).

Bacillus subtilis B315 has important potential to produce protease because this enzyme supports the crops for enhanced resistance to disease. Nevertheless, enzymatic treatment was secreted by *B. subtilis* B315 as antagonist bacteria resulted in degradation of the EPS (*extracellular polymeric substances*) components and significant eradication of the biofilm of bacteria pathogens. One of the hydrolytic enzymes can penetrate to biofilm, the use of protease can degrade pathogenic bacteria biofilms.

Therefore, the optimum protease production by *B. subtilis* B315 should be important to detection. Biofilm bacteria pathogens have a role in allowing cell-to-cell communication, and responsible for adhesion to surface, its advantage to survive in unfavorable environmental conditions and it's much less susceptible to the antimicrobial agents (Mitrofanova et al. 2017).

The objective of this research was to study the optimum activity of protease produced extracellularly by *B. subtilis* B315 to control *R. solanacearum* of chili. In this research optimum incubation time, pH, and concentration of crude extract protease *B. subtilis* B315 examined for maximum production of extracellular protease.

MATERIALS AND METHODS

Culture of *B. subtilis* B315 and *R. solanacearum* of chili for protease effect test

Culture stock of *B. subtilis* B315 in 20% glycerol was used in this study. This isolate was previously reported to be capable of controlling *R. solanacearum* of potato (Prihatiningsih et al. 2006; Prihatiningsih 2013). *R. solanacearum* was isolated from the diseased roots of chili suffered from bacterial wilt that was grown on CPG-TTC medium (Denny and Hayward 2001). *R. solanacearum* was characterized by creamy white fluidal colony with pink-red color in the middle, irregular or regular shape, flat surface with uneven edges (Singh et al. 2010).

Protease assay of *B. subtilis* B315 on skim milk agar medium

For qualitative test of protease produced by *B. subtilis* B315, the bacterial culture was grown on skim milk agar medium (casein 5g, yeast extract 2.5g, glucose 1g, agar 15g, distilled water 1000 mL, skim milk 7% was added as inducer (Majumdar and Chakraborty 2017). Protease assay of *B. subtilis* B315 was carried out through protease screening using a skim milk agar medium with scratched inoculated method of one loop full (Vijayaraghavan and Vincent 2013; Majumdar and Chakraborty 2017). Clear zone that appeared around scratch was protease secreted by *B. subtilis* B315.

Protease producing assay of *B. subtilis* B315

Protease activity was measured using a modified Kunitz method (Majumdar and Chakraborty 2017). For the assay, a total of 0.5 mL Casein substrate (0.6 w/v) was added in 0.1 M Tris-HCl buffer at pH 8.0. Enzymatic reaction was initiated by adding 0.1 mL of protease solution to the Tris-HCl buffer at 45 °C and incubation time of 30 minutes. The protease was prepared using ethyl acetate extraction method of Alnahdi (2012; Abu-Mejdad et al. 2013). The reaction was stopped by adding 0.5 mL cold TCA (*Trichloroacetic Acid*) followed by centrifugation at 4000 rpm at 4 °C for 15 minutes. The dissolved peptide in the supernatant was further measured using spectrophotometer at λ 275 nm. The control solution is made with the same treatment but the substrate and the cold TCA were first dissolved and then protease solution was added. The

standard solution used to measure proteolytic activity was tyrosine 1-120 μ g /mL. One unit of protease activity (U) is defined as the amount of enzyme needed to produce 1 μ g of tyrosine/minute mL of the enzyme solution from the Casein substrate at certain pH and temperature conditions, calculated by the formula

$$(a) - (b)$$

$$\text{Activity} = \frac{\text{-----}}{30 \text{ min.} \times \text{mL enzyme}} \times \text{dilution factor}$$

Note: a: μ g tyrosine/mL sample, b: μ g tyrosine/mL control, dilution factor = total volume / (enzyme volume + filtrate sample volume).

In this research 3 factors, namely: incubation periods, incubation pH, and crude extract protease concentration were studied to know the optimum condition for protease activity of *B. subtilis* B315. The incubation periods used were 15, 30, 45, and 60 min. The incubation pH: 7 of Phosphate buffer, pH 8 and 9 of Tris-HCl buffer, pH 10 of NaHCO₃, and pH 11 of NaOH buffer. While the crude extract protease concentrations were 0.1, 0.3, 0.5 and 0.7%. For determination of optimum temperature for enzyme activity, incubation condition was set following standard protocol (pH 8, 30 minutes incubation time, and 1 mL protease extract of *B. subtilis* B315). For determination of optimum pH, incubation condition was set following standard protocol (30 minutes incubation time, 45°C, and 1 mL protease extract of *B. subtilis*) except the pH was varied at pH 7, 8, 9, 10 and 11. For optimum enzyme concentration determination, incubation condition was set following standard protocol (pH 8, 30 minutes incubation time at 45 °C) except the protease extract concentration was varied at 0.1, 0.3, 0.5, and 0.7%.

Biocontrol potency test against *R. solanacearum* in vitro

Potency test of *B. subtilis* B315 as a biocontrol of *R. solanacearum* in vitro was done by the two-layer medium method as stated by Ghosh et al. (2007). *B. subtilis* B315 was inoculated on the YPGA medium (*yeast peptone glucose agar*), after 24 hours the Petri dish was turned over and dropped with 500 μ l chloroform, incubated for 3 hours until the chloroform evaporated. Pathogenic bacteria of *R. solanacearum* chilies isolates that had been grown on YPGA medium were harvested in 10 mL of sterile water, then 200 μ l are put into 4 mL of 0.6% water agar solution which slowly poured into the Petri dish. After 48 hours of incubation, an inhibition zone was formed.

Plant resistance of bacterial wilt on chili by *B. subtilis* B315

For testing control bacterial wilt of chili with *B. subtilis* B315, chili seeds were soaking in *B. subtilis* B315 suspension with a population density of 10⁸ cfu/mL before seedling was carried out, then watering with *B. subtilis* B315 suspension at the same concentration when reaching planting age 10, 20 and 30 DAP (days after planting) at a dose of 100 mL/plant. The observation of the effectiveness of wilt disease control was done by assessing the intensity of the disease using the formula $DI = a / b \times 100\%$, with a:

number of wilted plants; b: total number of plants observed (Aslam et al. 2015). Then, control effectiveness calculation was done to compare the intensity of control disease with the intensity of treatment disease according to Pawaskar et al. (2014): $E = (\text{control DI} - \text{treatment DI}) / \text{control DI} \times 100\%$.

RESULTS AND DISCUSSION

The qualitative test of protease

The qualitative protease produced by *B. subtilis* B315 showed clear zone around scratches which indicated the production of protease secreted by *B. subtilis* B315 (Figure 1). The role of this protease from antagonistic bacteria was controlling plant pathogens, promoting plant growth, and enhancing plant resistance by inhibiting biofilm plant pathogens (Jha and Bhattacharyya 2012; Mitrofanova et al. 2017).

The clear zone in the protease-producing test of *B. subtilis* B315 means that this microorganism is capable of producing proteases. It was indicated by a zone width ranging from 12-26 mm. In qualitative plate assay, isolate S1 exhibited the largest clear zone (30 ± 1.13 mm) in skim milk agar, and isolate S5 exhibited the lowest (18 ± 1.41 mm), which S1 and S5 isolates are Gram-positive bacteria (Bhowmik et al. 2015; Joshi and Ghike 2019).

The protease activity of *B. subtilis* B315

The protease activity of *B. subtilis* B315 was calculated based on the regression equation obtained from the absorbance value of the tyrosine standard at λ 275 nm, as follows: $Y = 0.0045x + 0.0016$ (Figure 2), the protease activity of *B. subtilis* B315 at various incubation period, pH and crude extract protease concentration are shown in Figures 3, 4 and 5. The activity of protease *B. subtilis* B315 showed at 30 minutes incubation period was the highest by 1.259 U/ml (Figure 3) and at pH incubation 9 was maximum protease activity by 1.422 U/ml (Figure 4).

The incubation time variations showed that 30 minutes duration was the best treatment with the protease activity 1.259 U/ml. Variations in pH incubation treatment showed that the highest protease activity was at pH 9 of 1.422 U/ml. Protease activity at various concentrations at pH 9 with 30 minute incubation time of 1.3185 U/ml was at a concentration of 0.5% solution. The effect of extract enzyme concentration on protease activity is shown in Figure 5 based on the best pH and incubation time (9 and 30 min.). *B. subtilis* is resistant to growth at a high temperature. The activity crude protease was at 40°C, pH 8 by 255 U/ml (Jadhav et al. 2014). The protease activity was 236.37 U/ml and 175,083 U/ml with carbon and nitrogen sources at pH 7.4 and incubation temperature 50°C respectively (Pant et al. 2015). The protease activity depends on the *Bacillus* strain, medium, physical and chemical factors (Queiroga et al. 2013; Tebyanian et al. 2018). *Bacillus* spp. strains KFP-5, KFP-7, and KFP-17 produced variable quantities of siderophore, protease, and glucanase in absence as well as presence of rice blast pathogen *Pyricularia oryzae*, protease (20.9 ± 29.3 U/mL) in

presence *P. oryzae*, and protease (11.6 ± 24 U/mL) in absence of *P. oryzae* (Rais et al. 2017).

Biocontrol potency test against *R. solanacearum* in vitro

The potency of *B. subtilis* B315 as a biocontrol of *R. solanacearum* in vitro, showed largest inhibition zone (32 mm) for *R. solanacearum* in vitro of chilli (Figure 6.). In vitro inhibition test results of *R. solanacearum* isolate chili was smaller than its suppress for potato isolates of *R. solanacearum* because *B. subtilis* B315 was isolated from healthy potato plants as their habitat, the bacterial have adaptive to the condition. *B. subtilis* B315 can suppress the *R. solanacearum* isolated from potato of 36 mm. However, *B. subtilis* B315 was found capable of controlling *R. solanacearum*.

Protease secreted by *B. subtilis* B315 affects the virulence of bacterial pathogens with combating biofilms formation, and induces stress tolerance in plants. *Bacillus* sp. which the genomics, proteomics, and metabolomics were required to elucidate the mechanism of *Bacillus*-plant interaction for biotic and abiotic stress management in crops Radhakrishnan et al. (2017). *Bacillus* is also known to produce an array of extracellular proteolytic enzymes. *Bacillus* proteases have high stability and low pathogenicity and can be easily purified and obtained in industrial quantities. These features make them promising matrix-degrading agents for combating bacterial biofilms (Mitrofanova et al. 2017). The effect of *B. subtilis* B315 treatment to inhibit *R. solanacearum* in vitro and in chili showed at Table 1 with plant resistance factor to bacterial wilt shown by the effectiveness.

Table 1. Inhibition effect of *Bacillus subtilis* B315 in vitro, in chili, and the effectiveness

Treatment	Inhibition (mm) in vitro	Disease intensity (%) in chili	Effectiveness (%)
Control	0	7.34	-
<i>B. subtilis</i> B315	36	2.87	60.89
Bactericide	24	3.06	58.31

Note: Control (without *B. subtilis* B315, without bactericide)



Figure 1. Clear zone around the scratches of *Bacillus subtilis* B315 on skim milk medium to show its potential to produce protease




Figure 2. Absorbance of tyrosine standard at λ 275 nm




Figure 3. Protease activity of *Bacillus subtilis* B315 at variation incubation periods




Figure 4. Protease activity of *Bacillus subtilis* B315 at variation pH




Figure 5. Protease activity of *Bacillus subtilis* B315 at variation crude extract protease concentration



Figure 6. The inhibition zone of *Bacillus subtilis* B315 against chili *Ralstonia solanacearum*

Protease from *Bacillus* sp. can promote plant resistance to pathogen seen by the decrease of disease intensity (Table 1). The plant resistance is indicated by phenol compound as an indicator of the increase of plant resistance to pathogens and the decrease of disease intensity with the effectiveness of more than 50%. The induced plant resistance by *B. subtilis* strain FZB24 could improve plant strength and furthermore avoid pathogens infection (Jha and Bhattacharyya 2012). *B. subtilis* B298 in microencapsulated formula that is applied to chili can reduce the bacterial wilt caused by *R. solanacearum* and induced systemic resistance by increased phenol compound up to 6.27% (Prihatiningsih et al. 2019).

In conclusion, the bacteria of potato rhizosphere *B. subtilis* B315 was capable of producing protease with an activity of 1.3185 U/mL. *B. subtilis* B315 effectively controlled bacterial wilt by 60.89% in chili. Protease from *B. subtilis* B315 could indicate the potency to control bacterial wilt. This enzyme increases *B. subtilis* B315 activity as biological control agents and induces plant varieties resistance. The future *B. subtilis* B315 formulated as biopesticide to support sustainable agriculture, and protease from *B. subtilis* B315 can be produced as well as induced to control bacterial wilt of Solanaceae.

ACKNOWLEDGEMENTS

The authors were grateful to the Indonesian Ministry of Research, Technology and Higher Education (DRPM Kemenristekdikti) for supporting this research in fundamental scheme.

REFERENCES

- Almas S, Hamed A, Shelly D, Mohan P. 2009. Purification and characterization of novel protease from *Bacillus* strain SAL 1. Afr J Biotechnol 8 (15): 3603-3609.
- Abu-Mejdad NMJ, Aaiz FL, Jassim OT. 2013. Antifungal activity of ethyl acetate extract of four species *Bacillus* isolated from soil. J Am Sci 9 (10): 172-176.

- Alnahdi HS. 2012. Isolation and screening of extracellular proteases produced by new isolated *Bacillus* sp. *J Appl Pharm Sci* 2 (9): 71-74. DOI: 10.7324/JAPS.2012.2915
- Aslam MN, Mukhtar T, Ashfaq M, Asad MJ, Hussain MA. 2015. Incidence and prevalence of bacterial wilt of chili in Punjab. *Pak Mycopath* 13 (1): 37-41
- Bhowmik S, Islam S, Ahmed MM, Hossain MB, Md. Hossain A. 2015. Protease producing bacteria and activity in gut of tiger shrimp (*Penaeus monodon*). *J Fish Aquatic Sci* 10 (6): 489-500. DOI: 10.3923/jfas.2015.489.500
- Denny TP, Hayward AC. 2001. Gram-negative bacteria, *Ralstonia*. In: Schaad NW, Jones JB, Chun W. (Eds.), *Laboratory Guide for Identification of Plant Pathogenic Bacteria*. APS Press, St Paul. Minnesota.
- Ghosh S, Sinha A, Sahu C. 2007. Isolation of putative probionts from the intestines of Indian major carps. *Isr J Aquacult.-Bamid* 59 (3): 127-132.
- Jadhav AA, Ismail KS, Harale MA, Gadre SV, Williamson MT. 2014. Study of protease enzyme from *Bacillus* species and its application as a contact lens cleaner. *British Biomed Bull* 2 (2): 293-302.
- Jha DK, Bhattacharyya PN. 2012. Plant growth-promoting rhizobacteria (PGPR): emergence in agriculture. *World J Microbiol Biotechnol* 28: 1327-1350. DOI 10.1007/s11274-011-0979-9
- Joshi PP, Ghike PB. 2019. Study of protease producing bacteria and their enzymatic activity at different parameters. *Int Res J Engineer Technol* 6 (6): 2239-2245.
- Lestari P, Prihatiningsih N, Djatmiko HA. 2017. Partial biochemical characterization of crude extracellular chitinase enzyme from *Bacillus subtilis* B298. *IOP Conf Ser Mater Sci Eng* 172: 012041. DOI: 10.1088/1757-899X/172/1/012041
- Majumdar S, Chakraborty U. 2017. Optimization of protease production from plant growth-promoting *Bacillus amyloliquefaciens* showing antagonistic activity against phytopathogens. *Int J Pharm Biol Sci* 8 (2): 635-642.
- Mitrofanova O, Mardanova A, Evtugyn V, Bogomolnaya L, Sharipova M. 2017. Effects of *bacillus* serine proteases on the bacterial biofilms. *Hindawi BioMed Res Int*. DOI: 10.1155/2017/8525912
- Morikawa M. 2006. Beneficial biofilm formation by industrial bacteria *Bacillus subtilis* and related species. *J Biosci Bioeng* 101 (1): 1-8.
- Pant G, Prakash A, Pavani JVP, Bera S, Deviram GVNS, Kumar A, Mitali Panchpuri, Prasuna RG. 2015. Production, optimization and partial purification of protease from *Bacillus subtilis*. *J Taibah Univ Sci* 9: 50-55.
- Pawaskar JR, Kadam JJ, Navathe S, Kadam JS. 2014. Response of chilli varieties and genotypes to bacterial wilt caused by *Ralstonia solanacearum* and its management. *Indian J Sci* 11 (29): 66-69.
- Prihatiningsih N, Soedarmono, Arwiyanto T, Hadisutrisno B. 2006. Pengendalian hayati penyakit layu bakteri kentang dengan *Bacillus* sp.: 1. Eksplorasi dan pengujian in vitro dan di rumah plastik. *Agrosains* 8 (1): 27-31. [Indonesian]
- Prihatiningsih N. 2013. Aktivitas antibiosis *Bacillus* sp. B315 sebagai agen pengendali hayati *Ralstonia solanacearum* pada kentang. [Disertasi]. Program Pascasarjana Fakultas Pertanian Universitas Gadjah Mada, Yogyakarta. [Indonesian]
- Prihatiningsih N, Djatmiko HA. 2016. Enzim amilase sebagai komponen antagonis *Bacillus subtilis* B315 terhadap *Ralstonia solanacearum* Kentang. *J Hama Penyakit Tumbuhan Tropika* 16 (1): 10-16. [Indonesian]
- Prihatiningsih N, Djatmiko HA, Lestari P. 2016. Karakter fisiologis *Bacillus subtilis* asal rizosfer kentang sebagai pemacu pertumbuhan tanaman Solanaceae. [Laporan Penelitian Fundamental]. Fakultas Pertanian Unsoed. Purwokerto. [Indonesian]
- Prihatiningsih N, Djatmiko HA, Erminawati. 2019. Bio-management of anthracnose disease in chilli with microencapsulates containing *Bacillus subtilis* B298. *IOP Conf Series: Earth Environ Sci* 250: 012041. DOI: 10.1088/1755-1315/250/1/012041
- Prihatiningsih N, Djatmiko HA, Lestari P. 2020. Screening of competent rice root endophytic bacteria to promote rice growth and bacterial leaf blight disease control. *JHPT Tro*. 20(1): 78-84 doi: 10.23960/j.hptt.12078-84
- Queiroga AC, Pintado ME, Malcata FX. 2013. Medium factors affecting extracellular protease activity by *Bacillus* sp. HTS 102-A novel wild strain isolated from Portuguese merino wool. *Nat Sci* 5 (6A): 44-53. DOI: 10.4236/ns.2013.56A007
- Radhakrishnan R, Hashem A, Abd_Allah EF. 2017. *Bacillus*: A biological tool for crop improvement through bio-molecular changes in adverse environments. *Front Physiol* 8: 667. DOI: 10.3389/fphys.2017.00667
- Rais A, Jabeen Z, Shair F, Hafeez FY, Hassan MN. 2017. *Bacillus* spp. a bio-control agent enhances the activity of antioxidant defense enzymes in rice against *Pyricularia oryzae*. *PLOS One* 12 (11): e0187412. DOI: 10.1371/journal.pone.0187412.
- Saha D, Purkayastha GD, Ghosh A, Isna M, Saha A. 2012. Isolation and characterization of two new *Bacillus subtilis* strain from the rhizosphere of eggplant as potential biocontrol agents. *J Plant Pathol* 94 (1): 109-118.
- Shumi WHT, Anwar MN. 2004. Proteolytic activity of a bacterial isolate *Bacillus fastidiosus*. *J Biol Sci* 4 (3): 370-374. DOI: 10.3923/jbs.2004.370.374
- Singh D, Sinha S, Yadav DK, Sharma JP, Srivastava DK, Lal HC, Mondal KK, Jaiswal RK. 2010. Characterization of biovar/ races of *Ralstonia solanacearum*, the incitant of bacterial wilt in solanaceous crops. *Indian Phytopath* 63 (3): 261-265.
- Tebyanian H, Mirhosseini SH, Bakhtiari A, Karami A, Dadseresht S, Otroshi B. 2018. Effect of physical and chemical factors in production of alkaline protease enzyme by *Bacillus* strains. *Int Lett Nat Sci* 71: 10-16. DOI: 10.18052/www.scipress.com/ILNS.71.10
- Vanitha SC, Niranjana SR, Mortensen CN, Umesha S. 2009. Bacterial wilt of tomato in Karnataka and its management by *Pseudomonas fluorescens*. *Biocontrol* 54 (5): 685-695.
- Vijayaraghavan P, Vincent SGP. 2013. A simple method for the detection of protease activity on agar plates using bromocresol green dye. *J Biochem Technol* 4 (3): 628-630.