




ISSN-0975-833X

INTERNATIONAL JOURNAL OF CURRENT RESEARCH

(PEER REVIEWED OPEN ACCESS JOURNAL: FAST PUBLICATION) ISSN: 0975-833X; IMPACT FACTOR: 7.992

Quality Scientific Publishing: Frequency: Monthly


[HOME](#) [ABOUT US](#) [EDITORIAL BOARD](#) [INSTRUCTION TO AUTHORS](#) [CURRENT ISSUE](#) [ARCHIVE](#) [SUBMIT ARTICLE](#) [CONTACT](#) [REMEMBER ME](#)

CATEGORIES

- Indexing & Abstracting
- Editorial Board
- Submit Manuscript
- Instruction to Author
- Current Issue
- Past Issues
- Call for papers February 2022
- Ethics and Malpractice
- Conflict of Interest Statement
- Statement of Informed Consent
- Final Proof Correction and Submission
- Publication Ethics
- Peer review process
- Cover image - December 2021
- Quick Response - Contact Us

CALL FOR PAPERS

Call For Paper

ARTICLE SUBMISSION

AUTHOR INFORMATION

- Copyright Form
- Join Editor & Peer Reviewers
- Paper Submission
- Impact Factor
- Policy of Screening

CERTIFICATE




IMPACT FACTOR 2019



SUBJECT AREA

- Life Sciences / Biology

WELCOME TO IJCR

SJIF Scientific Impact Factor: 7.992; ISI Impact Factor 2019-2020: 1.532; Doi: <https://doi.org/10.24941/ijcr.2017>

International Journal of Current Research, (IJCR) is an international double blinded refereed and peer-viewed monthly online academic research journal in all the streams. IJCR encourages new ideas and works in all the fields and it publishes high-quality original papers, theory-based empirical papers, review papers, case reports, conference reports/papers, technology reports, book reviews, commentaries, events, and news. We welcome papers from both academicians and practitioners on theories, business models, conceptual paradigms, academic research, and consultancy projects. All subject areas of **Biology and life sciences**: Agriculture, Anatomy, Astrobiology, Behavior, Bioacoustics and etc., **Computer and Information Sciences**: Artificial intelligence, Catalogs, Computer applications, Computer architecture, and etc., **Earth sciences**: Atmospheric science, Earth, Geochemistry, Geography, Geology and etc., **Ecology and environmental sciences**: Aquatic environments, Biogeochemistry, Biogeography, Bioindicators, Conservation science and etc., **Engineering and technology**: Aerospace engineering, Architectural engineering, Automotive engineering, Bioengineering, Chemical engineering and etc., **Medicine, health sciences and Dental**: Anatomy, Anesthesiology, Cardiology, Cardiovascular medicine and etc., **People and places**: Demography, Geographical locations, Population groupings and etc., **Physical sciences**: Astronomical Sciences, Chemistry, Materials science, Mathematics, Physics and etc., **Research and analysis methods**: Animal studies, Bioassays and physiological analysis, Biological cultures and etc., **Science policy**: Bioethics, Open science, Research funding and etc., **Social sciences**: Anthropology, Archaeology, Economics and etc. Authors are cordially invited to submit the full-length paper; Original and unpublished research articles, based on theoretical or experimental works, are solicited for publication in the journal. Submission of an article implies that the work described has not been published previously (except in the form of an abstract or academic thesis) and is not under consideration for publication elsewhere.

(ALL SUBJECT/MULTIDISCIPLINARY JOURNAL)

- Submission Date: Open
- Publication Date: Open

The features of our journal site www.journalcra.com

- Free full-text availability of articles
- Email Notification of new issue release
- Search Articles by author name, titles, issues and volumes
- Open Access Journal
- The facility of Online Submission of Articles on the website

What's News

- Volume 14, Issue 01 is now available online
- Online Publication any time for Every Month
- Submit your Research articles/papers/case study at journalcra@gmail.com

Important Notes:


- **SJIF Scientific Journal Impact Factor: 7.992**
- Fast Publication and Indexing Journal
- All changes in published articles may be made only 7 days after the online release
- Corresponding authors responsible for own paper contents and authors have to refer and cite those contents which come from other papers thus, in case of plagiarism journal doesn't have any responsibility
- Post-publication removal of articles from the website is impossible and all inquiries won't be answered



INTERNATIONAL JOURNAL OF CURRENT RESEARCH

(PEER REVIEWED OPEN ACCESS JOURNAL: FAST PUBLICATION) ISSN: 0975-833X; IMPACT FACTOR: 7.992

ISSN-0975-833X Quality Scientific Publishing: Frequency: Monthly



HOME ABOUT US EDITORIAL BOARD INSTRUCTION TO AUTHOR CURRENT ISSUE ARCHIVE SUBMIT ARTICLE CERTIFICATE REVIEWERS FORM CONTACT

Editorial Board Member's



Luai Farhan Zghair
Iraq



Hasan Ali Abed Al-Zu'bi
Jordanian



Fredrick OJIJA
Tanzanian



Firuza M. Tursunkhodjaeva
Uzbekistan



Faraz Ahmed Farooqi
Saudi Arabia




Elsadig Gasoom
Elbashir
Sudan



FadelAlla Eapen, Asha Sarah
United State



Dr. Arun Kumar A
India



Dr. Zafar Iqbal
Pakistan




Dr. SHAHERA S. PATEL
India



Dr. Recep TAS
Turkey



Dr. Rasha Ali Eldeeb
Egypt



Dr. Pralhad
Rahangdale
India




Kanhaiyalal DR. PATRICK D. CERNA
Philippines



Dr. Nicolas Padilla-Raygoza
Mexico



Dr. Muhammad shoaib Ahmedani
Saudi Arabia



DR. MUHAMMAD
MOHMAD
United State



ISMAIL DR. MAHESH SHIVAJI CHAVAN
India



DR. M. ARUNA
India



Dr. Lim Gee Nee
Malaysia



DR. IRAM BOKHARI
Pakistan



Dr. FARHAT NAZ RAHMAN
Pakistan



Dr. Devendra kumar Gupta
India



Dr. ASHWANI KUMAR DUBEY
India



Dr. Ali Seidi
Iran



Dr Ashok Kumar Verma
India



Thi Mong Diep NGUYEN
France



Dr. Muhammad Akram
Pakistan



Dr. Imran Azad
Oman



Dr. Meenakshi Malik
India



Anam Bhatti
Malaysia



Md. Amir Hossain
Bangladesh



Ahmet IPEKÇİ
Turkey



Mirzadi Gohari
Iran

Copyright © 2010 International Journal Of Current Research. All rights reserved






INTERNATIONAL JOURNAL OF CURRENT RESEARCH

(PEER REVIEWED OPEN ACCESS JOURNAL: FAST PUBLICATION) ISSN: 0975-833X; IMPACT FACTOR: 7.992

ISSN-0975-833X Quality Scientific Publishing: Frequency: Monthly



HOME ABOUT US EDITORIAL BOARD INSTRUCTION TO AUTHOR CURRENT ISSUE ARCHIVE SUBMIT ARTICLE CERTIFICATE REVIEWERS FORM CONTACT

CATEGORIES

Indexing & Abstracting

- Indexing & Abstracting
- Editorial Board
- Submit Manuscript
- Instruction to Author
- Current Issue
- Past Issues
- Call for papers February 2022
- Ethics and Malpractice
- Conflict of Interest Statement
- Statement of Informed Consent
- Final Proof Correction and Submission
- Publication Ethics
- Peer review process
- Cover image - December 2021
- Quick Response - Contact Us

CALL FOR PAPERS

Call For Paper

**ARTICLE
SUBMISSION**

AUTHOR INFORMATION

- Copyright Form
- Join Editor & Peer Reviewers
- Paper Submission
- Impact Factor
- Policy of Screening

CERTIFICATE



IMPACT FACTOR 2019



SUBJECT AREA

- Life Sciences / Biology

	Index Copernicus	ResearcherID	Google Scholar
Global Index Medicus	SJIF	ISI	ijome
IFSIJ	ESJI	J-Gate	Qaji
CiteFactor	DAIJ	IIFS	JOUR
Sherpa	SOA	Ulrich	Cosmos
POLSKA BIBLIOGRAFIA NAULOVA	Directory of Research Journal Indexing (DRJI)	ISI Factor	International Journal Impact Factor(IJIF)
Science Library Index	Scientific World Index	International Innovative Journal Impact Factor (IJIF)	Scholarsteer
General Impact factor	Global Impact Factor	Journal Quality Factor	ROAD
DocFoc	Open Access Library	Indian journals Index	Electronic
World Cat	BOAI	RIF	DMF
JIF	Research Junction	Root Indexing	Journal Factor
Geneva foundation for medical education and research			

Invited for
Call for p
Call for E
We an
Editorial
benefits



INTER
MEDI



CHIEF
Piosane C
ASSOCIA




Jean-Mai
Chief Sci



INTERNATIONAL JOURNAL OF CURRENT RESEARCH

(PEER REVIEWED OPEN ACCESS JOURNAL: FAST PUBLICATION) ISSN: 0975-833X; IMPACT FACTOR: 7.992

ISSN-0975-833X Quality Scientific Publishing: Frequency: Monthly



HOME ABOUT US EDITORIAL BOARD INSTRUCTION TO AUTHOR CURRENT ISSUE ARCHIVE SUBMIT ARTICLE CERTIFICATE REVIEWERS FORM CONTACT

CATEGORIES

- Indexing & Abstracting
- Editorial Board
- Submit Manuscript
- Instruction to Author
- Current Issue
- Past Issues
- Call for papers February 2022
- Ethics and Malpractice
- Conflict of Interest Statement
- Statement of Informed Consent
- Final Proof Correction and Submission
- Publication Ethics
- Peer review process
- Cover image - December 2021
- Quick Response - Contact Us

CALL FOR PAPERS

Call For Paper

ARTICLE SUBMISSION

AUTHOR INFORMATION

- Copyright Form
- Join Editor & Peer Reviewers
- Paper Submission
- Impact Factor
- Policy of Screening

CERTIFICATE



IMPACT FACTOR 2019



SUBJECT AREA

- Life Sciences / Biology

May 2019

- The cave opening types of the bakony region (transdanubian mountains, hungary)**
Márton Veress and Szilárd Vetési-Foith
DOI: <https://doi.org/10.24941/ijcr.34579.05.2019>
- Review on available feed resources and feeding practice of village chicken: Ethiopia**
Merhun Lamaro and Tagesse Sawo
DOI: <https://doi.org/10.24941/ijcr.35041.05.2019>
- Farmer's perception on climate change and its consequences in kapilvastu district, Nepal**
Amit Prasad Timilsina, Ghanashyam Malla, Bishnu Prasad Paudel, Aksh Sharma, Rameshwar Rimal, Achyut Gaire and Hemlal Bhandari
DOI: <https://doi.org/10.24941/ijcr.35138.05.2019>
- Ayurvedic management of axial spondyloarthritis - a case report**
Yogesh Kumar Pandey and Neha Kaushik
DOI: <https://doi.org/10.24941/ijcr.35143.05.2019>
- Analysis of the situation of pharmaceutical assistance management in a city in Bahia**
Nicio Roberto S. Santana, B. Pharm, Emily Bianca O. Farias, B. Pharm, Rodrigo S. Damascena, MSc, Nadielle S. Bidu, M. Pharm and Rafael Luiz de A. Rodrigues, SC
DOI: <https://doi.org/10.24941/ijcr.35181.05.2019>
- Development and validation of rp-hplc method for simultaneous estimation of zaltoprofen and paracetamol in bulk and tablet formulation**
Annasaheb S Gaikwad, Yogesh R Thorbare, Nisha S Mhaske and Mahendra B Dattar
DOI: <https://doi.org/10.24941/ijcr.35203.05.2019>
- Knowledge and determinants of postpartum family planning use among postpartum women in abakaliki south-east Nigeria**
Nwafor Johnbosco Ifunanya, Obi Chuka Nibert, Obi Vito Okwuchukwu, Ibo Chukwurenye Chukwu, Wgaji Darlington-Peter Chibuzor, Orwe Blessing and Egbuji Chima
DOI: <https://doi.org/10.24941/ijcr.35336.05.2019>
- Life style and health risk behavior of youth in Bangalore**
Mr. Shibu Puthenparamil and Dr. Baburaj
DOI: <https://doi.org/10.24941/ijcr.35361.05.2019>
- Carbohydrate consumption profile and insulin resistance in adults of both the sexes**
Maria Clara Pereira Teixeira, Arielly Rayanne Amaral Alves Santana, Anne Karolline Ferreira Santos, Beatriz Rocha Sousa, Larissa Alves Guimarães, Ramon Alves Pires, Mauro Fernandes Teles, Felipe Oliveira Bittencourt, Iago Raphael David and Stenio Ferman
DOI: <https://doi.org/10.24941/ijcr.35374.05.2019>
- Impact of using back belts and back physiotherapy on reducing low back pain among operative room nurses in north palestinian hospitals**
Mosab S. Amoudi and Rasmieh N. Anabtawi
DOI: <https://doi.org/10.24941/ijcr.35382.05.2019>
- Butterflies as an indicator species for the toxicity assessment of a commercial pesticide formulation containing d-tetramethrin and cyphenothrin**
Karunananda, H.T.A.R., Gunawardena, M.P. and Samarakoon, H.M.T.R.
DOI: <https://doi.org/10.24941/ijcr.35407.05.2019>
- Detection of risk of preeclampsia in pregnancy using the uterine artery doppler and cytokines**
Maja Pejkovska Ilieva and Vesna Antovska
DOI: <https://doi.org/10.24941/ijcr.35408.05.2019>
- Incidence of chronic disease among displaced people in sulaymaniyah province**
Maan Taqi Rafeeq
DOI: <https://doi.org/10.24941/ijcr.35413.05.2019>
- A study on internet addiction among adolescences**
Bindhu Joseph
DOI: <https://doi.org/10.24941/ijcr.35230.05.2019>
- Laboratory performance of cold-mix asphalt using epoxy resin modified-asphalt emulsion**
Lihong Guo, Shujian Zhang, Zhongda Chen, Yong Song, Xiaohu Niu, Mengzhuo Guo, Pengtao Xue and Xianjun Gu
DOI: <https://doi.org/10.24941/ijcr.35301.05.2019>

QUICK RE



INTER MED



CHIEF ED
Rosane C
ASSOCIA



Jean-Mia
Chief Sci

- DOI: <https://doi.org/10.24941/ijcr.35263.05.2019>
55. **Analysing fatty acids content in winged bean (*p. tetragonolobus* (L) dc) through half seed chromatography**
Nurtjahjo D Sasongko, Siti Samiansih and Nettiyanai Naipospos
DOI: <https://doi.org/10.24941/ijcr.35323.05.2019>
56. **Analysis of good governance system to improve government performance in the office of services municipais registo, notariado e cadastramento bobonaro**
Faustina da Costa, Lucas da Costa and Fernando Dias Gusmao
DOI: <https://doi.org/10.24941/ijcr.35335.05.2019>
57. **Human resource capability and work performance as improving effect for the public services quality in nacional de transportes e terrestres (dnrt) offices in Timor-Leste**
Romeu Marques Daci, Jose Cornelio Guterres and Leoneto Madeira Martins
DOI: <https://doi.org/10.24941/ijcr.35348.05.2019>
58. **Effect of different peroxide bleaching regimens and subsequent remineralization on the hardness of human enamel**
Dr. Kavita Sharma and Dr. Shahjad Pathak
DOI: <https://doi.org/10.24941/ijcr.35381.05.2019>
59. **Social relevance of snake charmer's songs in odisha – a historical analysis**
Prof. Bhagaban Sahu
DOI: <https://doi.org/10.24941/ijcr.35515.05.2019>
60. **To evaluate the importance of education and motivation in patients undergoing orthodontic treatment- a prospective study**
Dheerendra Kumar Singh, Komal Khatni, Anit Khatni, Pavan Kumar Singh, Sarajeet Kaur and Sourendra Bhaskar Maiti
DOI: <https://doi.org/10.24941/ijcr.35309.05.2019>
61. **Clinical evaluation of connective tissue graft in the treatment of Millers class II & III multiple adjacent gingival recessions in the esthetic zone**
Dr. Syed Saima, Dr. Dolly, Dr. Aqib Mushtaq and Dr. Mumtaz Ali
DOI: <https://doi.org/10.24941/ijcr.35331.05.2019>
62. **Root Coverage by Coronally Advanced Flap with and Without Platelet-Rich Fibrin: A Clinical Study**
Dr. Syed Saima, Dr. Mumtaz Ali, Dr. Aqib Mushtaq and Dr. Dolly
DOI: <https://doi.org/10.24941/ijcr.35360.05.2019>
63. **Relationship between interdental spacing and dental caries in primary dentition – a randomized control trial**
Dr. Savitha Sathiyaprasad, Dr. Krishnamoorthy, S.H., *Dr. Nikhil Das, K.R., Dr. Ramesh, R., Dr. Vinisha Vinod and Dr. Monika G Reddy
DOI: <https://doi.org/10.24941/ijcr.35493.05.2019>
64. **Analysis of the trends of climatic variables in southeast Nigeria**
Osugiri, I.I., Ben-Chendo, G.N., Okwara, M.O., Anyanwu, G.U. and Utazi, C.O
DOI: <https://doi.org/10.24941/ijcr.31970.05.2019>
65. **Effect of integrated nutrients management on kernel yield of chickpea cultivars**
R.A. Singh, R.K. Singh, P.V. Singh, Jitendra Singh, Asha Yadav and Jitendra Singh
DOI: <https://doi.org/10.24941/ijcr.34779.05.2019>
66. **Effects of different types of commercial feed on the zootechnical performance of larvae of the "Brazil" strain of tilapia of the Nile *Oreochromis niloticus* (Linnaeus, 1758).**
Ué Claver Zéa Bi, Nahoua Issa Ouattara and Siaka Berté
DOI: <https://doi.org/10.24941/ijcr.35368.05.2019>
67. **Stomatal abnormalities in the foliar epidermis of some scrophulariaceae**
Lahari P. S. and Hanumantha Rao B.
DOI: <https://doi.org/10.24941/ijcr.35414.05.2019>
68. **Biofilm detection in staphylococcal strains by two phenotypic methods**
Luis Núñez Oreza, Betty Sarabia Alcocer, Estephania Aganegui Robles, Paulino Tamay Segovia, Selene Blum Domínguez, Ruth López Alcántara, Baldemar Aké Canché, Javier Maldonado Rivas and Patricia M. Gamma Quen
DOI: <https://doi.org/10.24941/ijcr.35415.05.2019>
69. **Sensory assessment of cookie type biscuits produced with eggplant flour (*Solanum melongena*, L.)**
Elecia Rosa dos Santos Barros, Débora Maria Santos Azevedo, Janielle Oliveira Caldeira, Beatriz Rocha Sousa, Felipe de Oliveira Bittencourt, Viviane Figueiredo Vieira, Stenio Fernando Pimentel Duarte and Larissa Costa Fogaça
DOI: <https://doi.org/10.24941/ijcr.35426.05.2019>
70. **Huhs1015 reduces pkm2 protein due to autophagic degradation in gastric cancer cells**
Tomoyuki Nishizaki
DOI: <https://doi.org/10.24941/ijcr.35467.05.2019>
71. **Mining green gold: an optimization strategy on the implementation of coffee farmer policy in the fields of forestry department in Ijen District of Bondowoso**
Amrotus Soviah and Hary Yuswadi
DOI: <https://doi.org/10.24941/ijcr.35503.05.2019>
72. **Assessment of hematological status of selected women**

129. **Antibiotic susceptibility patterns of community and hospital acquired urinary isolates of *Escherichia coli***
Fouzia Haider, Bashayer Matar Quhimeia Al-huthali, Reem Ahmed Khelaf Alghuraybi, Ghadeer Jassim Mohammad Zahiri and Wejdan Mustafa Bakur Barnawi
DOI: <https://doi.org/10.24941/ijcr.35634.05.2019>
130. **Ultrasound Evaluation in Ophthalmology, comparative study**
Dr. Enas Sami Al- Modaris and Dr. Saba Hashim Mahmood
131. **Prevention of prolonged labor by active management of first stage using spasmolytic drugs a double blind randomized controlled clinical study**
Ahmed A. M. Nasr and Mahmoud Fouad
DOI: <https://doi.org/10.24941/ijcr.37611.05.2019>
132. **The edentulous mandible: an organised approach to two implant overdentures**
Dr. Nayar Islam, Dr. Nilishashukla, Dr. Arvyn Chamadia and Dr. Saniket Deskar
DOI: <https://doi.org/10.24941/ijcr.38094.05.2019>
133. **Antibacterial activities of papaya carica papaya l.**
Premalatha, M. and Lakshmi, S.
DOI: <https://doi.org/10.24941/ijcr.37749.05.2019>
134. **Attitudes of jazan university students towards khat abuse, KSA 2019**
Muath A.hazmi, Ali Suan Mugarbi, Abutalib Hassan Zarban, Nawaf Omer Hamzi, Miahdi Abdu Ayoub Hakami, Nasser Ibrahim Hakami, Wafaa Y. Hagawi, Muhessin Ibrahim Hakami, Yehia A. Maghlat and Ali. H. Muslimi
DOI: <https://doi.org/10.24941/ijcr.39440.05.2019>

EDITORIAL BOARD

Copyright © 2010 International Journal Of Current Research. All rights reserved





RESEARCH ARTICLE

ANALYSING FATTY ACIDS CONTENT IN WINGED BEAN (*P. tetragonolobus* (L) DC) THROUGH HALF SEED CHROMATOGRAPHY

*¹Nurtjahjo D Sasongko, ²Siti Samiarsih and ²Nettyani Naipospos

¹Laboratory genetics and molecular biology, Faculty of Biology-The University of n Jenderal Soedirman, Indonesia

²Laboratory of Plant Taxonomy, Faculty of Biology-The University of n Jenderal Soedirman, Indonesia

ARTICLE INFO

Article History:

Received 18th February, 2019

Received in revised form

09th March, 2019

Accepted 15th April, 2019

Published online 30th May, 2019

Key Words:

Mass selection, single cotyledon, gas Chromatography, putative genes.

*Corresponding author:

Nurtjahjo D Sasongko

ABSTRACT

Indonesia, the second largest mega diversity in the world. It has a wide array of plants species including Winged bean (*Psophocarpus tetragonolobus* (L) DC). Here, this crop is cultivated in many areas though they are not consumed daily leads to very limited effort to develop it. Developing this crop may attract people to cultivate and so consumption like as source for industrial oil. Selections for high fatty acids content seeds, so far are done by selecting good quality of seedlings through selection of the whole seeds. Current study initiated the use of half seed analysis on gas chromatography (GC) to collect data of high fatty acids content of winged bean. Six genotypes were collected from Sumatera (SMT 1 and 2), Java (1 and 2), Bali and West Nusa Tenggara (WNT), and used as the research material. The SMT genotypes are long pod genotypes (27 and 20,5 cm), Java have either long or short genotypes (27 cm length and 10.3 cm). Bali and WNT genotypes were not clear as they were collected as seeds instead of pods. The long pod genotype has total 8-22 seeds but only 6-8 seeds for short type. The SMT seeds germinated 100% but other genotypes showed lower rate of germination (65% and 55%). of those fatty acids, the Sumateran seeds contain 40-50% palmitic acids but only few numbers of seeds from other genotypes. Three individuals were high in palmitic and oleic acids content. The high palmitic acid content reflected the present of putative genes with with four or 3 dominant alleles (PPPP, PPPp) in few individuals. This data showed if half seed was able to be used as source for GC analysis.

Copyright © 2019, Nurtjahjo D Sasongko et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Nurtjahjo D Sasongko, Siti Samiarsih and Nettyani Naipospos, 2019. "Analysing fatty acids compound in winged bean (*P. tetragonolobus* (L) DC) through half seed chromatography", International Journal of Current Research, 11, (05), 3861-3865.

INTRODUCTION

Some annual and perennial crops are known to bear oil either in their seeds, fruits or mesocarp and nuts of up to 50% but contain only 15-30 % carbohydrates and 15-25% protein. FAO listed 21 primary oil crops (FAO, 2018). But not winged bean (*Psophocarpus tetragonolobus* (L) DC) which might due to limited data. In Indonesia, winged bean has long time been being cultivated in many areas from Western to Eastern parts and so have various local names/genotypes (Rukmana, 2000), however, they consume this plant as vegetable only which leads to no breeding efforts, for example by analysing its nutrition contents. Breeding needs a wide array of genetic variations, (Poespodarsono, 1988) suggested to collect various germ plasm which might obtained from (Fitriani et al., 2015) either local, national or overseas. Indonesian farmers produced 0.7 tons winged bean dry seeds which are much smaller than any Asean countries (Rismunandar, 1983); thus needs to change customers' preferences from utilizing it as vegetables to become as raw material in industrial oil (Rukmana, 2000); (Yeates et al., 2014) stated the matured winged bean seeds contain 17% fat consists of 35% MUFA (mono unsaturated fatty acids). The

MUFA might reduce cholesterol in blood thus reduce the possibility of coronary vascular blockage (Freese, 2001; Velasco et al., 1999). Winged bean seeds have also reported as raw material for "soy sauce", yoghurt, and "soy-milk" (Haryoto, 1995); these industries require huge number of winged bean seeds; this might be obtained through breeding. Plant composes fat/oil as food reservation and utilizes it during the initial phase of growth; started from producing fatty acids with different numbers of carbon (Cramer, 1990). Of this, the first product is palmitic acid C16:0 which might get 2 extra units carbon to form stearic acid (C18:0). The stearic acid might further undergo desaturation to form C18:1, C18:2 and C18:3 or to get another 2 carbon units to form C22:0 and C22:1 (Downey, 1987; Stumpf, 1980). Those steps run in four different phases and enzymes: β -ketoacyl-CoA-synthase (KCS) condenses Malonyl CoA to form β -ketoacyl-CoA; β -ketoacyl-CoA reductase to form β -hydroxyacyl-CoA. The β -hydroxyacyl-CoA reductase dehydrate it to form enoyl-CoA; last the trans-2,3-enoyl-CoA reductase changes enoyl-CoA and form acyl CoA (Millar and Kunst, 1997); and (Puyaubert et al., 2000). Current study was done to select parental plant candidates based on their fatty acids content as reflected by half seed gas chromatography (GC) results, and plant another

half as planting material (Sasongko, 2003). The GC, separates the material into 2 different phases (Day and dan, 1980); the first phase forms a stationer layer with wide surface, and the second forms a liquid phase which passes through the first phase (Montgomery *et al.*, 1983). Stated the compounds are split through processes like adsorbition, absorbition or both to pass the layer and might be used to analyse closed to complex compounds or very small amount of samples but more precise than those previous methods (Roth *et al.*, 1981).




Figure 1. Simplified biosynthesis of fatty acids [Slabas, 2001]

MATERIALS AND METHODS

Research Material: Current study applied 6 genotypes winged beans from Sumatera (1 and 2), Java 1 and 2, Bali and West Nusa Tenggara/WNT) to be analysed for their morphological characters like pod's length, total number of seeds per pod, seeds' colour, form and weight prior to GC analyses. The seeds of each genotypes was germinated and half of the seed/ one cotyledon was subjected for GC. The GC machine is Shimadzu type 2010 with a silica *capillary column* of DB-23, 0.25 x 30 m x 0.25 film thickness of (7%-W Scientific) Agilent Technology USA. The N2UHP (nitrogen ultra high pressure) gas was chosen as delivering gas material. Temperatures were set up as follows: *oven*, *detector*, and *injection* 90°C, 250°C, and 260°C consecutively and split to 1:50 at 100 kPa.

Research protocol

Seeds germination: 10 seeds of each genotype were selected based on their physiscal performances and placed in multi tray pot filled with smooth sand in the dark room. The seeds were positioned in a sequential number and sprayed with water daily. At the 10th days, the non-germinated seeds were categorized as rotting and replaced with the new ones. In contrast, prior to GC analyses, the cotyledon of germinated seeds will be dissected and placed in plastic clip then stored in minus 80°C compartment. Another half seed will be observed for their germination time, length of shoots and roots. Following to this, another half seed/one cotyledon was returned to the growing media for another 10 days before planted on the cultivation area.

Half seed's preparation for GC analyses: The dissected-cotyledone was put into 1 mL injection vial, added with 100 µl iso octan:iso propanol (9:1) and vortexed for 3 minutes and left over night.

The solvent was dried in warm air and added with 50 µl 0.5 N NaOH methanolic and vortexed for 3 minutes, left for 5 minutes. 100 µl BF3 20% methanol was added, vortexed for 3 minutes and left for 30 minutes to allow methylation. Following to this, 200 µl N hexane p.a. was added and vortexed for 3 minutes then added with 200 µl NaCl and re-vortexed. The top layer was taken by a hamilton syringe to be injected into the gas chromatography apparatus. Data of this step were fatty acids content showed in the GC machine.

Seed's planting: The left half sed/cotyledon was acclimatized in a polybag filled with cattle's manure for at least 2 weeks to reach approximately of 20 cm height prior to planting in the soil. Five plants were planted on each row which was set as follows: 3.5 m length, 50 cm within row and 70 cm between rows. At both edges, border crops of land-water spinach were planted. Each plant was given 3 m high pole to allow the plant twist to the pole and grow well. The plants which were not grow well then replaced with new seedlings. Data of this step were total number of plants grew well to harvesting time.

Plant's care: The growing plants were cared out by giving one garden-scoop organic fertilizer of cattle's manure weekly until approaching its generative then changed to one spoon urea per plant per week. Watering however, was done daily from planting to harvesting. Weeding was done weekly by pulling out the weeds.

Harvesting: Harvesting was done when the plants showed at least having one dry-cracked pod winged bean, this particular plant then pulled out. Data were obtained from observing the plants length, root legth, fresh and dry weight, total number of pods, pod's length, number of seeds per pod, seed's weight, total number of root nodules, and tubers.

RESULTS

Sampling siteS: From some environemtal factors were observed, all sampling sites were noticeably fit for cultivating winged bean. The temperature and rain falls of all sampling sites were quite similar but not Java which have more rain falls (Table 1). Those environemtal factors, however, do not parallel to winged bean production level from those particular areas due to small areas spent for cultivating it. Data of fatty acids compound (Table 3) obtained from individual half seed GC analysis then calculated and shown as group. The Java genotypes (1 and 2) contain the mediocre content of Palmitic acid (C16:0) of approximately of 35% and erucic acid (C22:1) approximately of 1.2%, but they were showing low level in Oleic (C18:1) of about 10.1% and Linoleic acid (C18:2) of 20% only. The Bali and WNT genotypes however, had the highest amount of palmitic which varied from 35 to 46% and oleic acids of 23 and 15.86%. The high palmitic acid content of 35 -60% in winged bean seeds genotypes Bali and WNT showed the presence of putative genes for palmitic acid as many as 4 or 3 dominant alleles dominant alleles (PPPP or PPPp) (Fig.2).

DISCUSSION

Environmental data of the sampling sites proofed that winged beans are cultivated in many areas of Indonesia from West to Eastern part thus indicating that Indonesia is quite favourable to cultivate them.

Table 1. Sampling sites parameters of rain falls, air temperature, and total cultivation area

Sites	Rain falls (mm)	Temperature (°C)	Total Area(Ha)
SMT 1	100-350	26-33	<1
SMT 2	100-350	26-33	<1
JAVA 1	>750	26-33	<1
JAVA 2	> 750	26-33	<1
Bali	600-900	25-33	<1
WNT	150-400	28-34	<1

Table 2. Average value of number of pod’s length, seeds per pod, 100 seed’s weight, number of seeds germinate, dormation period and shoot’s length

Genotypes	Pod’s length (cm)	Number of seeds pod ⁻¹	100 seed’s weight (gr)	Number of seeds germinate (unit)	Time to germinate (days)	Shoot’s length (cm)
SMT 1	27.00	14.39	47.7	10	15	13.1
SMT 2	20.50	12.77	40.7	10	14	12.5
JAVA 1	27.00	15.95	41.6	9	21	11.8
JAVA 2	10.63	6.95	43.2	4	21	12.0
BALI	NA	NA	39.2	5	27	17.3
WNT	NA	NA	37.4	6	28	9.25

Table 3. Genotype Winged bean plants containing high palmitic (C16:0) and other fatty acids

Source	C16:0	C18:1	C18:2	C22:1
Java 1	35.72	9.85	20.46	1.27
Java 2	35.51	10.54	20.2	1.28
Sumatra 1	40.90	7.77	26.88	2.32
Sumatra 2	40.30	5.28	18.24	2.32
Bali	35.26	23.12	6.20	2.33
WNT	45.85	15.86	13.23	2.03




Figure 2. Distribution of winged bean’s Palmitic acid (C16:0) with their putative gene for palmitic acid

The production rate, however, does not parallel with the environmental data but more related with the consumer’s preferences in utilizing them and reflected in small cultivation area spent for (Table 1). The small area of winged bean plantation were might due to the famers’ preferences to cultivate more higher economic value and so called as cash-crops e.g.: corn, peanuts, soy bean. Corn for example, in the mean time the Indonesian government put it as a new primadona and expected to export up to 1 million tons in the 2019. The Indonesian farmers also grow soy beans in a massive and extensive way to fullfill hgh deman on derivative products; currently it is cultivated in approximately 648,000 Hectare land with 2.75% incrase p.a. Increasing of the winged bean production, therefore, supposed to be started from changing the preferences in utilizing it, for example by utilizing mature seeds instead of young/immature pods (Rukmana, 2000). Stated that mature winged bean seeds contain all nutrition including fat/oil. Interestingly, the fat/oil contains in the seeds comprises of unsaturated fatty acids which is known as important compound in edible oil\

industries. The demands on edible oil is always increase parallel with people’s conciousness on healthy food (Murphy, 1994). (Fresee, 2001) stated the utilization of unsaturated fatty acid oil in daily intake will reduce the possibility of hearth coronary disease since this fatty acid might lowering cholesterol content in blood (Velasco *et al.*, 1999). Observations on morphological characters (Table 2) showed there were some variations among those genotypes collected. Three genotypes (Sumatera 1, 2, and Java 1) were classified as long pod’s type, however, they still varied in length (27 and 20.5 cm length). These genotypes also bearing different numbers of seeds (14.29 and 12.77 seeds for long and short pods consecutively). It was predicted that the farmers treated their plants in different ways. More interestingly the Javanese winged beans, the farmers also cultivated two different genotypes (27 and 10.63 cm length) which produced 15.95 and 6.95 seeds. From 100 seeds’ weight point of view, the Sumatera 1 genotype produced the largest seeds of 47.7 gr whereas the smallest came from WNT (37.4 gr). These differences might due to lesser rain falls in this area (Table 1) as

this crop require 600-1,500 mm p.a.as stated by (Davis, 1945), amount, distribution and intensity of rain falls affecting the peas pod formation. Current data were different from that reported by (Rukmana, 2000) stated the long pod genotype supposed to bear 8-22 seeds/pod and 6-8 seeds/pod for the short type. The difference in seeds bearing per pod might due to planting space applied in this study or long period to generative phase of 60-140 days after planting (DAP) and another 30 days for seed's filling. (23) stated that spacing took an important role in affecting the plants to stand in their generative phase. Spacing of 12.5 cm within row took longer time to form the first flower than 75 cm, whereas current study applied 50 cm apart. Space within row also important factor in green pods and seed's formation, however, winged beans are mostly cultivated in small areas in narrow-space (<15 cm). Alternatively, it might probably due to some environmental factors which were not conducive along the plant's growth as well as seeds filling in the pods (Sastrapradja *et al.*, 1981). Stated that winged beans have 2-10 flowers which will turn to pods per inflorescence. But in most of the cases, not all flowers turn to pod and not all pods filled by seed on each of its seta. Moreover, (Lambeth, 1950) reported that the low yield of lima beans was strongly correlated with high temperature, low relative humidity and soil moisture deficiencies. These three factors affecting total number of dropping flowers and young pods. The seed's filling in the pod might be affected by both external and internal factors. External factors such as pollinator's insects, as well as predators like acar, and grasshoppers are commonly noted in the plantation's areas. In line to this study, the low seeds formation in angiospermae was also quite low (Stephenson, 1981; Bawa and Buckley, 1989; Campbell and Halama, 1993).

There are three steps of fruit's formations namely: 1. evocation 2. Inisiation, and 3. Anthesis (Elisa, 2018). The longer time for seeds' filling the more possibility of pods abortion, because of one or combined-factors like: lack of soil nutrients, exceeding water during seed's filling, environment stress or pest and diseases attack (Asamoah *et al.*, 1981). Moderate to severe soil moisture stress affect pods formation and seeds filling. However, (Liu, 2004) suggested drought stress might also responsible to pod's filling. Apart from those factors, (Bennett *et al.*, 2011) suggested the following phytohormones like salicylic acids, ethylene, abscisic acids, gibberellin, and cytokinin take a role in pod's development (Habekotté, 1993). Stated that seed's formation much related with branch's number and the time of pod initiation during flowering. The seeds will germinate when their embryo release gibberellin actively, and push out aleuron of the endosperm, synthesis amylase, maltase and protease and changes reserve food to energy as well as cell's formation. The Sumatera genotypes needed shortest time (14 days) to germinate, indicated the pods collected from Sumatera genotypes were obtained from the newest harvesting time in this area. Whereas, the Balinese and WNT genotypes required double time to germinate, and Javanese in the middle (Table 2). The Sumateran genotypes had 100% germination rate while Bali and WNT showed only 55% germinated, indicated if they were obtained from different harvesting period. Alternatively, the differences in germination rate might due to level of seed's maturity as it affect dormation period. Others are total amount of water, temperature, light and soil. The seed's maturity level is important because indicating seed's physiological maturity as well as enough food reservation prior to germinate (Tianna, 2012) and so having higher viability than the younger seeds.

Apart from that, the author suggested also seed's size and dormant period as the factors affecting germination. Some seeds undergo a resting time and germinate when the environment are favour to them. Dormant period might also affected by thinnes or thickness of the seed's coat thus needs physical or chemical break to allow the seed start to germinate (Nawwaf, 2018). The low germination rate of the seeds collected from Java and Bali as well as WNT were probably because of those single or combined-factors. GC analyses of winged bean cotyledon (Table 3) showed some individuals of the Sumateran seeds containing high palmitic acids of (40-60%), followed by the Javanese whereas the lowest content was noted from both Bali and WNT seeds. These data were strongly against of those reported by (Rukmana, 2000) who stated that winged bean seeds contain <10% palmitic acid. This discrepancy might due to different methods of analysis, soxhleation against half seed gas chromatography. The previous methods of soxhleation, which is based on a serial steps of sieving, is the most common method on analyses of fats but this method takes much longer time and more labourious than that of gas chromatography. (Kramer *et al.*, 2010), however, reminded the use of a correct detector and standard solution, as moving phase in gas chromatography are two important factors of GC analysis.

This paper noted high palmitic acids content in winged bean seeds was closely to that of contain in palm oil plant. This suggested the importance effort on breeding activity to increase customers preferences of other oil sources crop. Selecting high in palmitic acid content seeds which might then used further as the plants candidate for either direct oil production or cross breeding. By doing so, the hybrids will produce high palmitic acid since this compound is important in either edible oils and oleochemical industries like: soap, cosmetics, and release agents following saponification of sodium palmitate (Fresee, 2001). Recently, paliperidone palmitate, a long-acting antipsychotic medication, has been used in the treatment of schizophrenia through injection intravenously (Bennett *et al.*, 2011). Apart from that, current data confirmed if single cotyledone contains fatty acids as much as in the whole seeds leads to a further seeds' analysis without discarding the germplasm which is mostly being applied in the previous time for seed's analysis. These data allow the breeders to select which seeds would be treated further planting as the parental candidates or discarding. This report, showed also the presence of putative dominant genes of P which caused high palmitic acid (C16:0) content of 40-60% with at least 3 dominant alleles but only 1 allele for recessive. In considering to this situation, winged bean oil might then be applied as raw material in oleochemical industries than that of edible oil industries such as: soaps, cosmetics, paintings, drugs, and chocolates. The utilization of palmitic oil as edible oil, need a serial steps like degumming.

Conclusion

Plant mass-selection which is based on their fatty acids content might be done by applying half seed (single cotyledon) analysis and another single seed is planted and used as candidate for particular interest. The Sumatera Winged bean genotypes contain high palmitic acid.

Acknowledgement

The authors would like to thank you the Dean of the Faculty, the Head of Research and Public Service the University of Jenderal Soedirman for their valuable help during the research

as well as preparing this manuscript. We also would like to thank you the Research and Public Service Directorate of Ministry for High Education, Research and Technology the Republic of Indonesia for providing fund under the contract No.: 1939/UN.23.14/PN/2018.

REFERENCES

- Asamoah, A.S. 2018. The growth and Yield of winged beans *Psophocarpus tetragonolobus* L. DC) as affected by soil moisture. *Agris.fao.org*.1981 (Jabatino, 2017 accessed on 12 October 2018).
- Bawa, K.S. and Buckley, D.P. 1989. Seeds: Ovule ratios, selective seed abortion and mating system in Leguminosae. In: Stirton C.H. Zarucchi ZL., (eds). *Advances in Legumes Biology. Monograph system Botany. Missouri Botanical Garden* 29: 243-262.
- Bennett, E.J., Roberts, J. A. and Wagstaff, C. 2011. The role of the pod in seed development: strategies for manipulating yields. *New Phytologist*. 190: 838-853.
- Chapman, H.R. and Halama, K.J. 1993. Resource and Pollen limitations to lifetime seed production in a natural plant population. *Ecology*, 74: 1043-1051.
- Cramer, N. 1990. *Raps: Anbau und Verwertung*. Verlag. Eugen Ulmer. Stuttgart.
- Davis, J.F. 1945. The effect of some environmental factors on the set of pods and yield of white pea beans. *Journal of Agricultural Research*, (70) 7:237-249.
- Day, R.A. dan A.L. and Underwood, 1980. Analisis Kimia Kuantitatif. Edisi 4. Diterjemahkan oleh R. Soendoro. Erlangga. Jakarta.
- Downey, R.K. 1987. Genetic manipulation of oilseed quality. In: P.K. Stumpf, J.B. Mudd. and W.D. Nes (Eds). *The Metabolism. structure. and functions of plant lipids*. Plenum Press. New York.
- Elisa, N. 2018. Pertumbuhan dan Perkembangan Tanaman <http://jai.staf.ipb.ac.id>. Diakses pada tanggal 30.
- FAO. 2018. *Oil bearing Crops and Derived Product*. *FAO.org/* accessed on 10 October.
- Fitriani, H., N. Rahman, N. Rahman and Sudarmonowati, E. 2015. Evaluasi Stabilitas Daya Hasil Ubi Kayu (*Manihot esculenta*) genotip lokal hasil kultur jaringan. *Proceeding Seminar Nasional, Masyarakat Biodiversitas Indonesia*, 12 September 2015. Jakarta.
- Freese, R. 2001. Low-erucic acid rapeseed oil and platelet function. *European Journal of Lipid Science and Technology*, 103: 483-489.
- Habekotté, B. 1993. Options for increasing seed yield of winter oilseed rape (*Brassica napus* L.): a simulation study. *Field Crops research*, (54) 2-3: 109-126. Germinate
- Haryoto, 1995. Tempe dan Kecap Kecap. Kanisius, Yogyakarta.
- Indonesia akan ekspor Jagung ke Filipina. *Republika.co.id*. 18/02/12 accessed on 28/09/18)
- Kramer, M., R. Litman, D., Hough, R., Lane, P., Lim, Y. and Liu, M. 2010. Eerdekens. Paliperidone palmitate, a potential long-acting treatment for patients with schizophrenia. Results of a randomized, double-blind, placebo-controlled efficacy and safety study.
- Lambeth, V. N. 1950. Some factors influencing pod set and yield of the Lima bean. *University of Missouri Columbia. Research Bulletin*.
- Lee, C.T. 1987. Effect of Plant Spacing on growth and yield of winged bean. *Agris.fao.org*.
- Liu, F. 2004. Physiological regulation of pod set in soy bean (*Glycine max* L. Merr) during drought at early reproductive stage. PhD dissertation. Københavns universiteit. Denmark.
- Millar, A.A. and Kunst, L. 1997. Very-long chain fatty acid biosynthesis is controlled through the expression and specification of the condensing enzyme. *The Plant Journal*, 1: 121-131.
- Montgomery, R., Dryer, R.L., Conway, T.W. and dan, A.A. 1983. *Spectro. Biokimia. Suatu Pelekat Berorientasi Kasus. Jilid 2. Edisi 4. Diterjemahkan oleh M Ismadidan S.D. Ismadi. GadjahMada University Press. Yogyakarta*.
- Murphy, D. J. 1994. *Designer Oil Crops, Breeding, Processing, and Biotechnology*. VCH Verlagsgesellschaft bH. Weinheim. Germany.
- Nawwaf, N.A. 2018. *Kromatografi Gas*. *Wordpress.com*. 2011, accessed.
- Poespodarsono, 1988. *Dasar-dasar Pemuliaan Tanaman*. PAU Institut Pertanian Bogor. Bogor.
- Puyaubert, J., B. Garbay. W., Dieryck. P., Costaglioli, T., Roscoe, C. and Casagne, R, 2000. *Lessire. Fatty acids biosynthesis: Elongation. Biochemical Society Transaction*, 28: 645-647.
- Rismunandar, 1983. *Kecapir Penghasil Protein dan Karbohidrat yang serbaguna*. Sinar Baru. Bandung.
- Roth, H.J. and dan, G. 1981. *Blaschke. Analisis Farmasi. Diterjemahkan oleh S. Kisman dan S. Ibrahim*. Gadjah Mada University Press. Yogyakarta.
- Rukmana, R. 2000. *Kecapir. Budidaya dan Pengolahan Pasca Panen*. Kanisius. Yogyakarta.
- Sastrapradja, S., Lubis, S.H.A. and Djajakusuma, E.S. 1981. *Sayur-sayuran*. Balai Pustaka-bekerjasama dengan LBN-LIPI. Jakarta.
- Slabas, A.R., Simon, J.W. and Brown, A.P. 2001. Biosynthesis and regulation of fatty acids and triglycerides in oil seed rape. Current status and future trends. *European Journal of Lipid Science and Technology*, 103:455-166. Sasongko. N.D. Increase of erucic acid content in oilseed rape (*Brassica napus* L.) through the combination with genes for high oleic acid. *Cuvillier Verlag. Goettingen. Germany*. 2003.
- Stephenson, A.G. 1981. Flower and fruit abortion: proximate causes and ultimate functions. *Annual Review Ecology Evolution system*. 12: 253-279.
- Stumpf, P.K. 1980. Biosynthesis of saturated and unsaturated fatty acids. In: P.K. Stumpf (Ed). *The Biochemistry of Plants: A comprehensive treatise*. Vol. 4. *Lipid Structure and Function*. Academic Press. New York. London. and San Francisco.
- Tianna, S. 2012. *Seed and seedling Biology-department of Agriculture and Biological Engineering. Pennsylvania University Cooperative extension*. 1-6.
- Velasco, L., Goffman, F.D. and Becker, H.C. 1999. Development of calibration equations to predict oil content and fatty acids composition in Brassicaceae germplasm by near infrared reflectance spectroscopy. *Journal of American Oil Chemists Society*, 76: 25-30.
- Yeates, CS., Ebrahimpour, A., Hamid, AA., Bakar, J., Muhammad, K. and Saari, N. 2014. Winged bean (*Psophocarpus tetragonolobus* (L.) DC) seeds as an underutilized source of proteolysate and biopeptide. *Journal of Food and Function*, DOI: 10.1039/C3FO60667H.