



Proceedings

Proceedings of the 1st Jenderal Soedirman International Medical Conference in conjunction with the 5th Annual Scientific Meeting (Temilnas) Consortium of Biomedical Science Indonesia

November 28-28, 2020, in Purwokerto, Indonesia



Editors: Joko Mulyanto¹; Teguh Haryo Sasongko²; Delvac Oceandy³; Dwi Utami Anjarwati¹; Wahyu Siswandari¹; Hutomo Suryo Wasisto⁴ and Motoyuki Sugai⁵

Affiliations: ¹ Faculty of Medicine Universitas Jenderal Soedirman, Indonesia ; ² Perdana University Malaysia, Malaysia ; ³ The University of Manchester, United Kingdom ; ⁴ Technische Universität Braunschweig, Germany ; ⁵ Hiroshima University, Japan

ISBN: 978-989-758-499-2

Conference Link: <http://jimc.fk.unsoed.ac.id>

Foreword: Nanoscience is a science that involves the study of materials on an ultra-small scale and the novel properties that these materials demonstrate. Nanoscience has the potential to reshape the world around us. It could lead to revolutionary breakthroughs in fields ranging from manufacturing to health care. At this time, we also have entered the era of genomic medicine (genomic medicine), which begins with the success in mapping the human genome. Even human genetic research in recent decades has led to the application of genomics for the benefit of patients (translational research). Genetic studies are now beginning to apply genetic information to achieve personalized medicine, which is an innovative approach to patient management by considering individual differences based on genes, environment, and lifestyle. This 1st Jenderal Sudirman International Medical Conference” in conjunction with 5th Annual Scientific Meeting (Temilnas) Consortium of Biomedical Science Indonesia (KIBI 2020) **(More)**

Volumes:

Vol. 1 - 978-989-758-499-2

The Role of Fibroblast Proliferation in Wound Healing by Different Plants: An Experimental Study

P. 5 - 9

Marisa Rilian , Indra Kusuma , Abdul Halim , Aliya Muhammad ,
Akbar Fitrianto and Ida Bagus Eka Narendra

DOI:10.5220/0010486300050009

Formative Research to Develop a Prototype of Sugar-sweetened Beverages Monitoring Application for Obesity Management among Indonesian Adolescents

P. 10 - 14

Eka Febriyanti , Ratih Yulistika Utami and Febrina Dewi Pratiwi Lingga

DOI:10.5220/0010486900100014

Chlorogenic Acid Ameliorates Vascular Remodeling and Perivascular Fibrosis in Kidney Fibrosis Model in Mice

P. 15 - 20

Gabriella Bamba Ratih Lintin , Nur Arfian , Dwi Cahyani Ratna Sari , Gina
Andyka Hutasoit , Mohammad Salman and Muhammad Mansyur Romi

DOI:10.5220/0010487000150020

Characteristics, Effective Reproduction Number (Rt), and Spatial Modelling of Covid-19 Spread at Magetan District, East Java, Indonesia

P. 21 - 27

Yudhi Wibowo , Nendyah Roestijawati , Atik C. Hidayah , Agoes Y. Purnomo
and Tutiek Herlina

DOI:10.5220/0010487100210027

Suppression of MnSOD by Andrographolide and its Relation to Oxidative Stress and Viability of Breast Cancer Stem Cells Treated with Repeated Doxorubicin Administration

P. 28 - 34

Angie Tara Rachman , Ayu Suraduhita , Resda Akhra Syahrani , Melva Louisa
and Septelia Inawati Wanandi

DOI:10.5220/0010487200280034

Toxicity of Self-nanoemulsifying Drug Delivery System Formulation of Nigella Sativa L. Seed Oil against Adult Danio rerio

P. 35 - 42

Yanti Tri Utami and Isnatin Miladiyah

DOI:10.5220/0010487300350042

Optimization of Immunohistochemical Staining with Anti Protein Gene Product 9,5 (PGP 9,5) Antibodies to Detecting Intraepidermal Nerve Fiber

P. 43 - 46

David Pakaya , Yustina Andwi Ari Sumiwi , Sri Herwiyanti and Rina Susilowati

DOI:10.5220/0010487400430046

Immunohistochemistry of KRAS Protein in Colorectal Cancer

P. 47 - 51

Dody Novrial , Kamal Agung Wijayana and Hanif Kun Cahyani

DOI:10.5220/0010487500470051

The Effect of Ethanol Extract of Lingzhi Mushroom (Ganoderma lucidum) on Caspase-3 Expression in Oral Cancer Cells

P. 52 - 57

Irfan Dwiandhono , Fadli Ashar , Arsa Hadiyatama Waskito Aji and Meta
Anjay Firmansyah

DOI:10.5220/0010487600520057

The Role of Chlorogenic Acid on Mice with Unilateral Ureteral Obstruction Model: A Study toward Tubular Injury

P. 58 - 62

Low-dose of Acetylsalicylic Acid Upregulates Expression of eNOS mRNA and Downregulates Interleukin-6 (IL-6) and Transforming Growth Gactor- β 1 (TGF- β 1) mRNA in Rat Kidney of Preeclampsia Model P. 63 - 68

Yuyun Nailufar , Rul Afiah Syarif , Andi Fitriani Kusuma , Farmita Chairani ,
Nia Marlina , Charolina Vivi Vienetta , M. Hadri Ar-Ridho , Totok Utoro and
Nur Arfian
DOI:10.5220/0010487800630068

Dose-dependent Decaffeinated Green Tea Extract Administration Improved Hyperglycemia through Modulation of IRS-1 and GLUT-4 Genes Expression in Metabolic Syndrome Rat Model P. 69 - 74

Dwi Adi Nugroho , Mifetika Lukitasari , Marlita Marlita , Mohammad
Saifur Rohman , Nashi Widodo , Inggita Kusumastuty and Nur Ida Panca
Nugrahini
DOI:10.5220/0010487900690074

Expression of HSA-MIR-155-5P and mRNA Suppressor of Cytokine Signalling 1 (SOCS1) on Plasma at Early-stage and Late-stage of Nasopharyngeal Carcinoma P. 75 - 80

Recita Indraswary , Sofia Mubarika Haryana and Agus Surono
DOI:10.5220/0010488000750080

Cost Analysis Monotherapy of Type 2 Diabetes Mellitus Patients in RSUD. Prof. Dr. Margono Soekarjo P. 81 - 83

Fajar Wahyu Pribadi , Afifah and Catharina Widiartini
DOI:10.5220/0010488100810083

Antibiofilm Activity of Aloe barbadensis Miller Extract Against Staphylococcus aureus P. 84 - 90

Kanesti Ismirajna and Irena Agustiningtyas
DOI:10.5220/0010488200840090

The Renal Protective Potential Effect of Infusion of Anti-urolithiasis Formula in Urolithiasis Patients: A Randomized Clinical Study P. 91 - 97

Ulfatun Nisa , Peristiwa Ridha Widhi Astana , Saryanto , Tyas Friska Dewi and
Enggar Wijayanti
DOI:10.5220/0010488300910097

In Vitro Study of Reduction of Oral Enterococcus faecalis Biofilm on Application of Combination of Chrysomya megacephala Maggot Extract and Sodium Hypochlorite P. 98 - 103

Rizka Hidayati , Ari Asnani , Muhamad Salman Fareza and Dwi Utami Anjarwati
DOI:10.5220/0010488400980103

Effects of 8-Week Vitamin E (α Tocopherol) Supplementation on Reduced Insulin Resistance in Non-diabetic Obese Subjects P. 104 - 108

Erlina Marfianti and Isnatin Miladiyah
DOI:10.5220/0010488501040108

The Activities of Streptomyces W-5A as Antibacterial and Antibiofilm towards Methicillin-resistant Staphylococcus aureus 2983 P. 109 - 115

Annisa Permata Dinda , Ari Asnani and Dwi Utami Anjarwati
DOI:10.5220/0010488601090115

Association of Epstein Barr Virus Infection to Prognosis Factors of Invasive Breast Cancer in Semarang Kariadi General Hospital Population P. 116 - 120

Wasisto Dwi Yudisaputro , Djoko Handojo , Yan Wisnu Prajoko and

Dik Puspasari

DOI:10.5220/0010488701160120

Analysis of the Effectiveness of Chrysomya sp. Maggot Extract in Inhibiting the icaA and icaD Genes Regulator Staphylococcus epidermidis Biofilm P. 121 - 125

Dylan Tamalsir , Anton Budhi Darmawan and Dwi Utami Anjarwati

DOI:10.5220/0010488801210125

Correlation between Hepcidin and Ferritin with Insulin and Hba1C as Biochemical Markers of Pancreas Damage in β -Thalassemia Patients P. 126 - 130

Irbath Hamdani , Qodri Santosa , Joko Setyono , Hernayanti and Lantip Rujito

DOI:10.5220/0010488901260130

Chitosan Nanoparticle as a Delivery System of miRNA 217 for Suppressing Hepatocellular Carcinoma Progressivity by Targeting AEG-1/P53 P. 131 - 138

Ulfatun Nisa , Indwiani Astuti , Ronny Martien , Dhani Rinaldi Maulana and

Ysrafil

DOI:10.5220/0010489001310138

DRD2 Gene Polymorphisms in Schizophrenia Patients P. 139 - 142

Zieske Maritska , Miranti Dwi Hartanti , Benediktus Wicaksono Widodo , Fitri ,

Bintang Arroyantri Prananjaya , Irsan Saleh and Nita Parisa

DOI:10.5220/0010489101390142

The Use of Autogenous Tooth Bone Graft Block (ABTB) in Post-Extraction Socket of Dental Implant Treatment Procedure: A Literature Review P. 143 - 147

Bambang Tri Hartomo , Ambar Delfi Mardiunti , Dewi Sartieka Putri ,

Irma Khoerunisa and Pratitis Widi Seno

DOI:10.5220/0010489201430147

Comparison of Purple Passion Juice (*Passiflora edulis* var. *edulis*) and Simvastatin on Lipid-lowering Effect of Hyperlipidemic Rats Model P. 148 - 154

Alfi Muntafiah , Tisna Sendy Pratama , Anisa Rachmawati , Dewi

Wahyu Wulandari and Qodri Santosa

DOI:10.5220/0010489301480154

Comparison of Tear Interleukin-6 level, Conjunctival Inflammation, and Tear Film Function in Daily Conventional Hydrogel and Extended Silicone Hydrogel Contact Lens Wear P. 155 - 163

Tri Rahayu , Adisti , Weni Puspitasari and Umar Mardianto

DOI:10.5220/0010489401550163

The Effect of Jackfruit (*Artocarpus heterophyllus*) Leaf Ethanolic Extract Gel on Superoxide Dismutase and Interleukin-1 β Levels in Wound Healing after Tooth Extraction in Diabetic Rats P. 164 - 169

Almasyifa Herlingga Rahmasari Amin , Yusuf Asto Pamasja , Christiana

Cahyani Prihastuti , Haris Budi Widodo and Hernayanti

DOI:10.5220/0010489501640169

Sources of Information and Knowledge of Gonorrhea in Indonesia P. 170 - 174

Annisa Tristifany , Fatihah Arifah Rahmawati , Bella Maharany Syahfira ,

Adhitama Noor Idninda , Muhammad Maulana Malik , Nina Tiwi Handayani

and Sani Rachman Soleman

DOI:10.5220/0010489601700174

Different Contribution of Estrogen Receptors, ET-1/ETBR and Superoxide Dismutase and in eNOS Availability based on Sexual Dimorphism in Early Stage of Kidney Diabetic Rats

P. 175 - 183

Anisa Fatwa , Dwi Cahyani Ratna Sari , Wiwit Ananda Wahyu Setyaningsih , Andrew Nobiantoro and Nur Arfian
DOI:10.5220/0010489701750183

Description of Methicillin-resistant Staphylococcus aureus Infection at Arifin Achmad General Hospital, Riau Province from 2015-2019

P. 184 - 189

Dewi Anggraini , Dewi Santosaningsih , Kuntaman , Muhammad Ihsan and Syifa Farhani Ridha
DOI:10.5220/0010489801840189

Dietary Intake of Obese Adult Men in Jakarta

P. 190 - 193

Karima Yudhistina , Erfi Prafiantini and Novi S. Hardiany
DOI:10.5220/0010489901900193

Iron Status of Newborns in Maternal Inflammation Status Differences

P. 194 - 201

Qodri Santosa , Alfi Muntafiah and Lantip Rujito
DOI:10.5220/0010490001940201

The Effects of Aqueous Extract of Jamaican Cherry (Muntingia calabura) on D-galactose-induced Liver Damage in BALB/c Mice (Mus musculus)

P. 202 - 207

Cempaka Jaga Pramudita , Dwi Nur Ahsani , Ika Fidianingsih and Evy Sulistyoningrum
DOI:10.5220/0010490102020207

The Effect of Caffeine towards Zebrafish (Danio rerio) Juvenile Working Memory Exposed by Unpredictable Chronic Stress (UCS)

P. 208 - 213

Alya Ayu Tazkia and Zainuri Sabta Nugraha
DOI:10.5220/0010490202080213

Iron for Human Brain Development: A Fulfill Strategy in the First 1,000 Days of Life

P. 214 - 222

Qodri Santosa
DOI:10.5220/0010490302140222

Expression of MicroRNA-155 and Suppressor of Cytokines Signaling 1 (SOCS1) mRNA in Plasma Breast Cancer Patients

P. 223 - 230

Dwi Nur Indah Sari , Dina Rahmina , Sumadi Lukman , Artanto Wahyono , Indwiani Astuti , Sofia Mubarika and Teguh Aryandono
DOI:10.5220/0010490402230230

The Effects of Administering Jicama Concentrate (Pachyrhizus erosus) and Kefir Grains as a Synbiotic Drink on Malondialdehyde and Superoxide Dismutase Levels in the Testicles of Hyperlipidemic Rats

P. 231 - 236

Mohammad Alvian Subhakti , Miranti Dewi Pramaningtyas , Rafik Prabowo and Rokhima Lusiantari
DOI:10.5220/0010490502310236

The Effect of Addition of Blood Cockles (Anadara granosa) Shell Nano-Hydroxyapatite on Hardness of Heat Cured Acrylic Resin

P. 237 - 240

Dian Noviyanti Agus Imam , Ryana Budi Purnama and Aris Aji Kurniawan
DOI:10.5220/0010490602370240

Drug-induced Hypersensitivity Syndrome in a Breast Cancer Patient: A Case Report

P. 241 - 244

Antimicrobial Potential Activity of Extract *Selaginella plana* (Desv. Ex Poir.) Hieron against the Growth of *Staphylococcus aureus* ATCC 25922 and Methicillin-Resistance *Staphylococcus aureus* (MRSA)

P. 245 - 253

Juen Carla Warella , Agung Dwi Wahyu Widodo , Rebekah Juniati Setiabudi , Retno Indrawati Roestamadji , Maftuchah Rochmanti and Pudji Lestari

DOI:10.5220/0010490802450253

The Lingzhi Mushroom (*Ganoderma lucidum*) Extract Reduce the Number of Lymphocyte in Diabetics Rats with Periodontitis: In Vivo Experimental on Sprague dawley Rats

P. 254 - 259

Pratitis Widi Seno , Amilia Ramadhani , Saryono , Haris Budi Widodo and Christiana C. Prihastuti

DOI:10.5220/0010490902540259

Reversible Imiquimod Effects on Skin Tissue of Psoriasis Mice Model: An Experimental Study

P. 260 - 266

Thianti Sylviningrum , Afifah , Dody Novrial , Brian Wasita , Bambang Purwanto and Harijono K. Sentono

DOI:10.5220/0010491002600266

Efficacy of Topical Binahong Leaf Ethanolic Extract Administration on Serum TGF- β 1 in Wistar Rats with *Staphylococcus aureus*-infected Wounds

P. 267 - 272

Rizki Andini Nawawi , Muhammad Totong Kamaluddin and Theodorus

DOI:10.5220/0010491202670272

The Oral Health Status, Salivary Flow Rate and pH in Hypertensive Patients Who Consume Antihypertensive Drugs in Puskesmas Kasihan I Yogyakarta

P. 273 - 277

Nurvita Risdiana and Eka Nuraeni

DOI:10.5220/0010491302730277

Hard Tissue Surgical Treatment with Embedded Dental Condition of Tuberosity Maxillary

P. 278 - 283

Bambang Tri Hartomo and Rizka Rachmatika Dewi

DOI:10.5220/0010491402780283

Blood Transfusion, Serum Total Iron Binding Capacity and Iron in Hemodialysis Patients Margono Soekarjo Hospital

P. 284 - 288

Vitasari Indriani , Aditiawarman , Yunanto Dwi Nugroho and Fania Salsabila

DOI:10.5220/0010491502840288

Secretion of IFN- γ and IL-17 after Stimulation of ESAT-6-CFP10 (EC610) Fusion Antigen from PBMC in Groups Active TB and Latent TB

P. 289 - 297

Nika Andriani , Nova Kurniati , Muhammad Irsan Saleh , Eddy Mart Salim , Zen Hafy , Jusak Nugraha , Kemas Ya'kub Rahadiyanto and Francisca Srioetami Tanoerahardjo

DOI:10.5220/0010491602890297

Antibacterial Activity of Synthesized Silver Nanoparticle using Langsung Leaf Extract (*Lansium domesticum* var. *pubescens* Kooders et Valeton) as Bioreductor against *Escherichia coli* and *Staphylococcus aureus*

P. 298 - 304

Khairunnida Rahma , Agung Dwi Wahyu Widodo , Rebekah Juniati Setiabudi , Retno Indrawati Roestamadji , Maftuchah Rochmanti and Pudji Lestari

DOI:10.5220/0010491702980304

The Effectiveness of Mesenchymal Stem Cell and Colostrum Bovine Combination in Post Hepatectomy Liver Failure with Liver Fibrosis Animal Model

P. 305 - 310

Dimas Aryo Kusumo , Agung Putra , Albertus Ari Adrianto , Erik Prabowo and Ignatius Riwanto

DOI:10.5220/0010491803050310

Prev 1 2 Next



SCITEPRESS - SCIENCE AND TECHNOLOGY PUBLICATIONS, LDA.

RESOURCES

[Proceedings](#)

[Papers](#)

[Authors](#)

[Ontology](#)

CONTACTS

Science and Technology
Publications, Lda
Avenida de S. Francisco Xavier,
Lote 7 Cv. C,
2900-616 Setúbal, Portugal.

Phone: +351 265 520 185
Fax: +351 265 520 186
Email: info@scitepress.org

EXTERNAL LINKS

[PRIMORIS](#)

[INSTICC](#)

[SCITEVENTS](#)

[CROSSREF](#)

PROCEEDINGS SUBMITTED FOR INDEXATION BY:

[dblp](#)

[Ei Compendex](#)

[SCOPUS](#)

[Semantic Scholar](#)

[Google Scholar](#)

[Microsoft Academic](#)

The Activities of Streptomyces W-5A as Antibacterial and Antibiofilm towards Methicillin-resistant *Staphylococcus aureus* 2983

Annisa Permata Dinda¹^a, Ari Asnani¹^b and Dwi Utami Anjarwati²^c

¹Department of Chemistry, Faculty of Mathematics and Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia

²Department of Microbiology, Faculty of Medicine, Universitas Jenderal Soedirman, Purwokerto, Indonesia

Keywords: MRSA, Streptomyces, antibiofilm, antibacterial

Abstract: Methicillin-resistant *Staphylococcus aureus* (MRSA) causes nosocomial infection worldwide. MRSA can defend itself by forming a biofilm layer, thereby increasing the virulence factor. WHO categorizes MRSA as high risk on the Priority Pathogen List for searching for new antibiotics. Recently, we have reported the potency of Streptomyces W-5A as anti-MRSA based on qualitative screening. Thus, this research aimed to analyze the potency of Streptomyces W-5A as antibacterial and antibiofilm towards MRSA 2983. Cultivation of Streptomyces W-5A used Starch Casein Nitrate (SCN) agar, and anti-MRSA extract production used SCN broth. Samples were taken at incubation times of 0, 3, 6, 9, 12, and 15 days. Each sample was tested for antibacterial activity with the Kirby-Bauer method, whereas inhibition of biofilm formation and biofilm degradation with microtiter method. The results showed that the optimum antibacterial activity was achieved after nine days of incubation with an inhibition zone of 9.589 ± 0.521 . The optimum % inhibition of biofilm formation was $77.806 \pm 13.595\%$ after 12 days of incubation. The optimum % degradation of biofilm was $80.465 \pm 7.586\%$ after nine days of incubation. These findings suggest that Streptomyces W-5A has the potency to produce antibacterial and antibiofilm compounds against MRSA 2983.

1 INTRODUCTION

Methicillin-resistant *Staphylococcus aureus* (MRSA) is an *S. aureus* bacterium resistant to methicillin and several other beta-lactam antibiotics (Boucher & Corey, 2008). MRSA is an infection-causing bacterium that can defend itself by forming a protective layer called a biofilm. *S. aureus* biofilms develop rapidly and form colonies on moist and nutrient-rich surfaces (Tarver, 2009). The ability to form biofilms is one of *S. aureus*' virulence factors, which causes antibiotic resistance to bacteria (Høiby et al., 2010; Lee et al., 2013).

Infection by microbes is estimated to be 80% related to the formation of biofilms, which contribute to the nature of antibiotic resistance (Archer et al., 2011). Biofilms consist of microbial cells and a matrix of extracellular polymeric substance (EPS). As much as 50-90% of the EPS matrix's main ingredients are organic carbon consisting of polysaccharides,

proteins, nucleic acids, lipids, phospholipids, and humic substances, and 15% are bacterial cells (Deshpande & Joshi, 2011). These biofilms cause antibiotics cannot reach the target, bacterial cells; hence the bacteria that cause infection cannot be inhibited or destroyed using antibiotics. Therefore, exploration of antibiofilm compounds becomes necessary. The antibiofilm compounds are expected to inhibit biofilm formation and degrade biofilms (Konai & Haldar, 2017). The inhibitory activity of biofilms occurs because these compounds have antibacterial activity, while biofilms' degradation activity occurs because they can depolymerize complex compounds in the biofilm matrix.

Actinobacteria, particularly the genus Streptomyces, has been reported to produce bioactive compounds with anti-MRSA activity. Streptomyces albus, granaticin B from *Streptomyces violaceoruber*, and streptorubin B from Streptomyces sp. MC11204 has been reported to inhibit biofilm formation and

^a <https://orcid.org/0000-0002-1992-6685>

^b <https://orcid.org/0000-0002-8569-2565>

^c <https://orcid.org/0000-0001-8394-2543>

damage the *S. aureus* biofilm (Oja et al., 2015; Suzuki et al., 2015). Bhakyashree and Krishnan (2018) reported that *Streptomyces* sp. Strain VITBKA3 showed anti-MRSA activity. Balasubramanian et al. (2017) also reported that ethyl acetate extract from *Streptomyces* sp. strain SBT343 could reduce biofilm formation of several *Staphylococcal* species, including MRSA USA300.

Recently, Asnani et al. (2020) have reported the potency of *Streptomyces* W-5A as anti-MRSA based on qualitative screening. *Streptomyces* W-5A was isolated from mangrove areas in Segara Anakan Cilacap, a source of indigenous marine actinobacteria (Asnani et al., 2016). Following that report, this research aimed to analyze the potency of W-5A isolates as antibacterial and antibiofilm towards MRSA 2983.

2 MATERIALS AND METHODS

The research was conducted from January to September 2020 in the Biochemistry Laboratory and Research Laboratory in UNSOED, Purwokerto. *Streptomyces* W-5A was isolated from mangrove sediment in Segara Anakan Cilacap. The MRSA 2983 is known as a bacterium capable of producing biofilm matrix. It was isolated from a clinical specimen from the pus of female patient in Prof. Dr Margono Soekarjo Hospital, Banyumas regency, Indonesia.

2.1 Cultivation of Isolate W-5A

Streptomyces W-5A was cultivated using continuous streak on Starch Casein Nitrate (SCN) following the procedure described by Asnani et al. (2016). Starch Casein Nitrate (SCN) agar [starch, casein, KNO_3 , KH_2PO_4 , $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, NaCl , $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, agar] was added 1 μL nystatin for every 10 mL of SCN medium. The cultures were incubated for seven days at room temperature.

2.2 Production of Anti-MRSA Extract

Production of anti-MRSA extract used SCN broth following the procedure described by Asnani et al. (2020). A total of 10 plugs (6 mm diameter) of *Streptomyces* W-5A were inoculated into 100 mL of SCN broth. The culture was incubated at 90 rpm and room temperature until it reached the exponential phase for inoculum. Next, 10% of the inoculum was inoculated into a new SCN broth to produce anti-MRSA compounds. The cultures were incubated using an orbital shaker at a speed of 90 rpm. Samples

were taken at incubation times of 0, 3, 6, 9, 12, and 15 days. Each sample was separated by centrifugation at 4.000 rpm for 10 minutes at 4°C, then filtered to obtain the crude extract. Each extract was tested for antibacterial activity, inhibition of biofilm formation, and biofilm degradation.

2.3 Antibacterial Test

The extract's antibacterial activity was evaluated using the disc paper diffusion method on the Mueller Hinton Agar (MHA) medium following CLSI (2019). A total of 30 μL of the extract was added to disc paper (6 mm), then placed on the MHA medium that had been inoculated by MRSA 2984 using the spread plate method. The test culture was incubated at 37°C for 24 hours. A clear zone around the disc paper indicated a positive result of antibacterial activity against MRSA. Hence, the parameter observed was the diameter of the inhibition zone.

2.4 Biofilm Formation Inhibition Test

The potency of the extract to inhibit biofilm formation was tested using the microtiter plate method described by Suzuki et al. (2015) with modification. The MRSA 2983 was inoculated in Brain Heart Infusion (BHI) medium with 1% glucose (BHI-Glu) and incubated at 37°C for 24 hours. Then, the culture was adjusted for its turbidity level using a 0.5 McFarland standard and diluted in BHI-Glu with a ratio of 1:100. The inhibition of biofilm formation was carried out by adding a total of 10 μL of diluted MRSA and 100 μL of extract to the microplate and incubated at 37°C for 24 hours. After incubation, planktonic cells were removed carefully, washed twice with PBS, stained with 0.1% crystal violet solution, and incubated for 30 minutes. Then, the microplate was washed with water to remove excess crystal violet and dried. HCl in isopropanol (1:20) was then added, and the optical density (OD) value was measured using a microplate reader at a wavelength of 595 nm. The percentage of biofilm formation inhibition was calculated using the following formula (Pratiwi et al., 2015).

$$\% \text{Inhibition} = \frac{OD_{\text{control}} - OD_{\text{test}}}{OD_{\text{control}}} \times 100\% \quad (1)$$

2.5 Biofilm Degradation Test

The potency of extract to degrade biofilms was tested using the microtiter plate method described by Suzuki

et al. (2015) with modification. The MRSA 2983 was prepared as the previous procedure. The biofilm degradation test was carried out by inoculating diluted MRSA in each microplate well and incubated at 37°C 24 hours. After incubation, planktonic MRSA cells were carefully removed, then the extract was added into each well. The mixtures were incubated at 37°C for 24 hours, exposing the extract to biofilm formation at the bottom of the well. After incubation, the microplate was treated similarly to the previous procedure, and the OD value was measured at 595 nm. The percentage of biofilm degradation was calculated using the following formula (Pratiwi et al., 2015).

$$\% \text{Degradation} = \frac{OD_{\text{control}} - OD_{\text{test}}}{OD_{\text{cpntrol}}} \times 100\% \quad (2)$$

2.6 Data Analysis

The research data were the diameter of the inhibition zone, %inhibition, and %degradation. All data were analyzed using one-factor analysis of variance (ANOVA) at a 95% confidence level. If the ANOVA analysis results have a significant effect, it continued with the Duncan test.

3 RESULTS

3.1 Antibacterial Activity

The extract was tested for antibacterial activity using the paper disc diffusion method. This method is expected to determine the sensitivity of a microbe to various antibiotics. The experimental parameters were the presence or absence of a clear zone formed around the disc paper, which showed MRSA's growth inhibition. The research result indicated that the extract has antibacterial activity (Figure 1). The antibacterial activity of Streptomyces W-5A's extract increased from day 0 and reached the highest activity on day 9th with the diameter of the inhibition zone was 9.589 ± 0.521 mm. The zone of inhibition showed potent inhibition of extract against MRSA 298.

The one-way ANOVA test analysis results showed a significant effect (Sig. <0.05) between the incubation time and the inhibition zone diameter (Table 1). Further tests using the Duncan test showed that the 9-day incubation time had an antibacterial activity significantly different from other incubation times. This finding indicated that the optimum

incubation time for producing antibacterial compounds was nine days.




Figure 1: Antibacterial activity of Streptomyces W-5A extract against MRSA 3983.

3.2 Inhibition of Biofilm Formation

The inhibition of biofilm formation aimed to determine the potency of extract to inhibit MRSA biofilms' formation. Compounds with antibacterial activity against microorganisms that form biofilms can be used to inhibit biofilm formation (Memariani et al., 2019). The tests used a microplate and BHI medium with glucose. BHI medium is often used to grow Gram-positive and Gram-negative bacteria, whereas glucose to the BHI medium aimed to induce MRSA biofilms formation (You et al., 2014).




Figure 2: The inhibition formation of biofilm MRSA 2983 by Streptomyces W-5A extract.

The research result is presented in Figure 2. The extract inhibited the formation of biofilm MRSA 2983. The percentage inhibition of biofilm formation by the extract increased from 0 days until it reached the optimum incubation time, 12 days, with a per cent inhibition of $77.806\% \pm 13.595\%$.

Table 1: Results of one-way ANOVA for antibacterial and antibiofilm activities

	Sum of Squares	df	Mean Square	F	*Sig.
(1) Antibacterial Activity					
Between Groups	17.209	5	3.442	6.307	0.004
Within Groups	6.549	12	0.546		
Total	23.758	17			
(2) Inhibition of biofilm formation					
Between Groups	3766.922	5	753.384	2.856	0.630
Within Groups	3165.493	12	263.791		
Total	6932.415	17			
(3) Biofilm degradation					
Between Groups	4655.138	5	931.028	3.084	0.051
Within Groups	3622.848	12	301.904		
Total	8277.986	17			
*Significant value = 0.05					

After it reached the optimum incubation time, the rate of inhibition of biofilm formation decreased.

The analysis results with the one-way ANOVA test showed no significant effect (Sig. <0.05) between the incubation time treatment on the percentage of inhibition of biofilm formation (Table 1). However, 50% inhibition of biofilm formation was achieved after six days of incubation and reached a maximum of 12 days of incubation.

3.3 Biofilm Degradation

The research result indicated that the extract could degrade the biofilm of MRSA 2983 (Figure 3). The percentage of biofilm degradation increased from day 0 until it reached the optimum incubation time, which was the 9th day, with the percentage of biofilm degradation was $80.465\% \pm 7.586\%$. After it reached the optimum incubation time, the rate of biofilm degradation decreased.




Figure 3: The degradation of biofilm MRSA 2983 by Streptomyces W-5A extract.

The analysis results using the one-way ANOVA test showed no significant effect (Sig. <0.05) between the incubation time treatment on the

percentage of biofilm degradation (Table 1). However, 50% of biofilm degradation was achieved after three days incubation and reached a maximum at nine days incubation.

The tests used the microplate method with BHI-Glu medium. The clinical MRSA 2983 was added to a well-filled medium, then incubated for biofilm formation. After incubation, planktonic cells formed on the surface, while biofilms formed and stuck tightly to the well-walls. This adhesion was accompanied by a build-up of organic materials covered by an extracellular polymer matrix. This matrix was a structure of threads crossed with each other and acted as an adhesive for the biofilm. The planktonic cells formed were removed by washing, then the Streptomyces W-5A extract was added to the well. After incubation, all wells were washed to remove planktonic bacteria and added crystal violet to colour the biofilm. Besides using crystal violet, biofilm quantification can also use resazurin, safranin, and tryptan blue dyes (Peeters et al., 2008; Sandasi et al., 2010). In this research, crystal violet dye was used because it was easy to obtain and economical in price.

4 DISCUSSION

Actinobacteria have been reported to have antibacterial activity against MRSA. Bister et al. (2004) wrote that Abyssomicin C obtained from actinobacteria has potential as anti-MRSA. León et al. (2011) reported that the dichloromethane extract from actinobacteria isolates (I-400A, B1-T61, M10-77) have high antibacterial activity against MRSA ATCC 43300 and VRE ATCC 51299. Rajan & Kannabiran (2014) also reported 2,4-dichloro-5-sulfamoyl benzoic acid (DSBA) extracted from marine

Streptomyces sp. VITBRK2 has anti-MRSA activity. Furthermore, Bhakyashree & Krishnan (2018) wrote that *Streptomyces* sp. VITBKA3 is potential as an anti-MRSA compound against MRSA strains ATCC43300 and ATCC700699.

Isolate at a particular incubation time that produces the highest inhibition zone is considered the optimum time for making antibacterial compounds. The duration of the production phase of each microbe varies depending on genetic factors and environmental conditions. Susilowati et al. (2007) reported that actinobacteria produced optimal antibacterial compounds at the optimum incubation time of 72 hours (isolate A3.5) and 96 hours (isolate F6.1) against Enteropathogen bacteria. Dhananjeyan et al. (2012) used an incubation time of five days to produce antibacterial compounds from actinobacteria against *Escherichia coli* MTCC 50, *Pseudomonas aeruginosa* MTCC 424, and *Bacillus subtilis* MTCC 441. Bhakyashree & Krishnan (2018) used an incubation time of seven days to produce antibacterial compounds from actinobacteria against MRSA strains ATCC 43300 and ATCC 700699.

The clinical isolate MRSA 2983 is known as a biofilm producer. This research used an initial MRSA concentration of 1.5×10^8 CFU/mL (0.5 McFarland standard). According to Skogman et al. (2016), biofilms can be formed with an initial bacterial concentration of 10^6 - 10^8 CFU/mL. MRSA was added to the well containing the medium, and then the extract was added so that the biofilm growth would coincide with the presence of extract. If the extract has inhibitory activity, there will be inhibition of MRSA biofilm formation.

After incubation, the well was washed to remove planktonic bacteria, then stained with crystal violet. Crystal violet can bind to proteins and polysaccharides in the bacterial extracellular matrix (Peeters et al., 2008). In this research, if, after staining with crystal violet, a purple colour was formed, the amount of binding dye was assumed to be the same as the number of biofilm matrices in the well (O'Toole, 2011).

Microbes capable of producing biofilms generally have the potential for resistance to antibiotics. Bacteria in the biofilm can withstand antibiotics because the antibiotics fail to penetrate to destroy the biofilm. Biofilm degradation test is known as an approach to dealing with biofilms by adding bioactive compounds that can trigger biofilm destruction (Belbase et al., 2017; Bjarnsholt et al., 2013; Schierle et al., 2009). Thus, the biofilm degradation test aims to determine the potency of extract to degrade biofilm MRSA.

Bioactive compounds to inhibit biofilm formation were highly produced at 12 days of incubation in this research, whereas bioactive compounds to degrade biofilm were highly made at nine incubation days. Indeed, the production of anti-MRSA compounds has various optimum incubation time. Oja et al. (2015) reported that *Streptomyces violaceoruber* (DSM-40701) produced antibiofilm compounds against *S. aureus* with an incubation time of four days. Suzuki et al. (2015) said that *Streptomyces* sp. strain MC11024 made antibiofilm compounds against *S. aureus* and MRSA N315 with an incubation time of three days. Balasubramanian et al. (2017) reported that *Streptomyces* sp. SBT343 produces antibiofilm compounds against Staphylococcal bacteria with an incubation time of ten days.

5 CONCLUSIONS

Streptomyces W-5A isolated from Segara Anakan Cilacap indicated the potency as anti-MRSA. To our knowledge, this is the first report of indigenous *Streptomyces* with three major anti-MRSA activities, including antibacterial, biofilm formation inhibition, and biofilm degradation. This finding highlights Segara Anakan Cilacap as an essential source of indigenous microbe for pharmaceutical purposes. Further research is necessary, particularly for the isolation and characterization of bioactive compounds.

ACKNOWLEDGEMENTS

The authors would like to express profound gratitude for the fundamental research grant contract No. T/1641/UN23.18/PT.01.03/2020 from the Directorate of Research and Community Service, Ministry of Research and Technology/National Agency for Research and Innovation of the Republic of Indonesia.

REFERENCES

- Archer, N. K., Mazaitis, M. J., Costerton, J. W., Leid, J. G., Powers, M. E., & Shirtliff, M. E. (2011). *Staphylococcus aureus* biofilms: properties, regulation, and roles in human disease. *Virulence*, 2(5), 445–459. DOI: 10.4161/viru.2.5.17724.
- Asnani, A., Ryandini, D., & Suwandri. (2016). Screening of marine actinomycetes from Segara Anakan for natural pigment and hydrolytic activities. IOP

- Conference Series: Materials Science and Engineering, 107, 12056. DOI: 10.1088/1757-899x/107/1/012056.
- Asnani, A., Luviriani, E., & Oedjijono. (2020). Activity of actinomycetes isolated from mangrove Segara Anakan Cilacap toward Methicillin-Resistant *Staphylococcus aureus* (MRSA). *Jurnal Kimia Sains Dan Aplikasi*, 23(1), 1–7. DOI: 10.14710/jksa.23.1.1-7.
- Balasubramanian, S., Othman, E. M., Kampik, D., Stopper, H., Hentschel, U., Ziebuhr, W., Oelschlaeger, T. A., & Abdelmohsen, U. R. (2017). Marine sponge-derived Streptomyces sp. SBT343 extract inhibits Staphylococcal biofilm formation. *Frontiers in Microbiology*, 8, 236. DOI: 10.3389/fmicb.2017.00236.
- Belbase, A., Pant, N. D., Nepal, K., Neupane, B., Baidhya, R., Baidya, R., & Lekhak, B. (2017). Antibiotic resistance and biofilm production among the strains of *Staphylococcus aureus* isolated from pus/wound swab samples in a tertiary care hospital in Nepal. *Annals of Clinical Microbiology and Antimicrobials*, 16(1), 15–19. DOI: 10.1186/s12941-017-0194-0.
- Bhakyashree, K., & Krishnan, K. (2018). Actinomycetes mediated targeting of drug-resistant MRSA pathogens. *Journal of King Saud University - Science*, 32(1), 260–264. DOI: 10.1016/j.jksus.2018.04.034.
- Bister, B., Bischoff, D., Ströbele, M., Riedlinger, J., Reicke, A., Wolter, F., Bull, A. T., Zähler, H., Fiedler, H.-P., & Süßmuth, R. D. (2004). Abyssomicin C-A polycyclic antibiotic from a marine Verrucosipora strain as an inhibitor of the p-aminobenzoic acid/tetrahydrofolate biosynthesis pathway. *Angewandte Chemie*, 43(19), 2574–2576. DOI: 10.1002/anie.200353160.
- Bjarnsholt, T., Alhede, M., Alhede, M., Eickhardt-Sørensen, S. R., Moser, C., Kühl, M., Jensen, P. Ø., & Høiby, N. (2013). The in vivo biofilm. *Trends in Microbiology*, 21(9), 466–474. DOI: 10.1016/j.tim.2013.06.002.
- Boucher, H. W., & Corey, G. R. (2008). Epidemiology of methicillin-resistant *Staphylococcus aureus*. *Clinical Infectious Diseases*, 46(5), 344–349. DOI: 10.1086/533590.
- CLSI. (2019). *Performance standards for antimicrobial susceptibility testing*, 29th Edition. CLSI supplement M100. CLSI: Wayne, PA.
- Deshpande, J., & Joshi, M. (2011). Antimicrobial resistance: The global public health challenge. *International Journal of Students Research*, 1(2), 41–44. DOI: 10.5549/IJSR.1.2.41-44.
- Dhananjeyan, V., Selvan, N., & Dhanapal, K. (2012). Isolation, characterization, screening and antibiotic sensitivity of actinomycetes from locally (near MCAS) collected soil samples. *Journal of Biological Sciences*, 10(6), 514–519.
- Høiby, N., Bjarnsholt, T., Givskov, M., Molin, S., & Ciofu, O. (2010). Antibiotic resistance of bacterial biofilms. *International Journal of Antimicrobial Agents*, 35(4), 322–332. DOI: 10.1016/j.ijantimicag.2009.12.011.
- Konai, M. M., & Halder, J. (2017). Fatty acid comprising lysine conjugates: anti-MRSA agents that display in vivo efficacy by disrupting biofilms with no resistance development. *Bioconjugate Chemistry*, 28(4), 1194–1204. DOI: 10.1021/acs.bioconjchem.7b00055.
- Lee, J.-H., Park, J.-H., Cho, H. S., Joo, S. W., Cho, M. H., & Lee, J. (2013). Anti-biofilm activities of quercetin and tannic acid against *Staphylococcus aureus*. *Biofouling*, 29(5), 491–499. DOI: 10.1080/08927014.2013.788692.
- León, Q. J., Aponte-Ubillus, J., Rojas, R., Cuadra, D., Ayala, N., Tomás, G., & Guerrero, M. (2011). Study of marine actinomycetes isolated from the central coast of Peru and their antibacterial activity against methicillin-resistant *Staphylococcus aureus* and vancomycin-resistant *Enterococcus faecalis*. *Revista Peruana de Medicina Experimental y Salud Publica*, 28(2), 237–283. DOI: 10.1590/S1726-46342011000200010.
- Memariani, H., Memariani, M., & Ghasemian, A. (2019). An overview on anti-biofilm properties of quercetin against bacterial pathogens. *World Journal of Microbiology and Biotechnology*, 35(143). DOI: 10.1007/s11274-019-2719-5.
- O'Toole, G. A. (2011). Microtiter dish biofilm formation assay. *Journal of Visualized Experiments*, (47). DOI: 10.3791/2437.
- Oja, T., Galindo, P. S. M., Taguchi, T., Manner, S., Vuorela, P. M., Ichinose, K., Metsä-Ketelä, M., & Fallarero, A. (2015). Effective antibiofilm polyketides against *Staphylococcus aureus* from the pyranonaphthoquinone biosynthetic pathways of Streptomyces species. *Antimicrobial Agents and Chemotherapy*, 59(10), 6046–6052. DOI: 10.1128/AAC.00991-15.
- Peeters, E., Nelis, H., & Coenye, T. (2008). Resistance of planktonic and biofilm-grown *Burkholderia cepacia* complex isolates to the transition metal gallium. *The Journal of Antimicrobial Chemotherapy*, 61(5), 1062–1065. DOI: 10.1093/jac/dkn072.
- Pratiwi, S. U. T., Lagendijk, E., Weert, S., Idroes, R., Hertiani, T., & Hondel, C. (2015). Effect of *Cinnamomum burmannii* Nees ex Bl. and *Mossioa aromatica* Becc. essential oils on planktonic growth and biofilm formation of *Pseudomonas aeruginosa* and *Staphylococcus aureus* in vitro. *International Journal of Applied Research in Natural Products*, 8(2), 1–13.
- Rajan, B. M., & Kannabiran, K. (2014). Extraction and identification of antibacterial secondary metabolites from marine Streptomyces sp. VITBRK2. *International Journal of Molecular and Cellular Medicine*, 3(3), 130–137.
- Sandasi, M., Leonard, C. M., & Viljoen, A. M. (2010). The in vitro antibiofilm activity of selected culinary herbs and medicinal plants against *Listeria monocytogenes*. *Letters in Applied Microbiology*, 50(1), 30–35. DOI: 10.1111/j.1472-765X.2009.02747.x.
- Schierle, C. F., De la Garza, M., Mustoe, T. A., & Galiano, R. D. (2009). Staphylococcal biofilms impair wound healing by delaying reepithelialization in a murine cutaneous wound model. *Wound Repair and Regeneration*, 17(3), 354–359. DOI: 10.1111/j.1524-475X.2009.00489.x.

- Skogman, M. E., Vuorela, P. M., & Fallarero, A. (2016). A platform of anti-biofilm assays suited to the exploration of natural compound libraries. *Journal of Visualized Experiments*, (118), 54829. DOI: 10.3791/54829.
- Susilowati, D. N., Hastuti, R. D., & Yuniarti, E. (2007). Isolasi dan karakterisasi aktinomisetes penghasil antibakteri Enteropatogen *Escherichia coli* K1.1, *Pseudomonas pseudomallei* 02 05, dan *Listeria monocytogenes* 5407. *Jurnal AgroBiogen*, 3(1), 15–23. DOI: 10.21082/jbio.v3n1.2007.p15-23.
- Suzuki, N., Ohtaguro, N., Yoshida, Y., Hirai, M., Matsuo, H., Yamada, Y., Imamura, N., & Tsuchiya, T. (2015). A compound inhibits biofilm formation of *Staphylococcus aureus* from Streptomyces. *Biological and Pharmaceutical Bulletin*, 38(6), 889-892. DOI: 10.1248/bpb.b15-00053.
- Tarver, T. (2009). Biofilms: A threat to food safety. In *Food Technology*, 5(63).
- You, Y., Xue, T., Cao, L., Zhao, L., Sun, H., & Sun, B. (2014). *Staphylococcus aureus* glucose-induced biofilm accessory proteins, GbaAB, influence biofilm formation in a PIA-dependent manner. *International Journal of Medical Microbiology* 304(5–6), 603–612. DOI: 10.1016/j.ijmm.2014.04.003.





Ari Asnani Unsoed <ari.asnani@unsoed.ac.id>

Decision Letter for Revision JIMCA58

1 message

JIMC UNSOED <jimc@unsoed.ac.id>
To: Ari Asnani <ari.asnani@unsoed.ac.id>

Wed, Dec 23, 2020 at 9:33 PM

Dear Mrs Ari Asnani

The scientific committee of JIMC 2020 thanks to the authors for submitting the full paper manuscript for publication in JIMC 2020 conference proceeding. The paper has been distributed to two different peer-reviewers to be blindly reviewed. Following the comments from the reviewers, we would like to inform you that the submitted paper **needs to be revised before accepted for publication**. Therefore, we invite you to revise and resubmit your manuscript to be published in the JIMC 2020 conference proceeding.

Please refer to the attachment letter for further detail information.

Yours sincerely,

Tirta Wardana, S.Si, M. Biotech

Program Manager the 1st Jenderal Soedirman Medical Conference (JIMC)

Faculty of Medicine, Universitas Jenderal Soedirman

4 attachments



Template Paper Scitepress.docx
48K



Paper guideline Scitepress.pdf
142K



DECISION LETTER FOR REVISION_JIMC A58.docx
15K



A58.docx
14K



Ari Asnani Unsoed <ari.asnani@unsoed.ac.id>

Decision Letter For Acceptance

1 message

JIMC UNSOED <jimc@unsoed.ac.id>
To: Ari Asnani <ari.asnani@unsoed.ac.id>

Tue, Jan 19, 2021 at 2:24 AM

Dear Ms. Ari Asnani

The scientific committee of JIMC 2020 thanks to the authors for submitting the full-paper manuscript for publication in JIMC 2020 conference proceeding. The paper has been distributed to two different peer-reviewers to be blindly reviewed. Following the comments from the reviewers, we are happy to inform you that the submitted paper is **accepted**. To be published in the JIMC 2020 conference proceeding please refer to the attached letter for further detailed information.

Yours sincerely,

Tirta Wardana, S.Si, M. Biotech

Program Manager the 1st Jenderal Soedirman Medical Conference (JIMC)

Faculty of Medicine, Universitas Jenderal Soedirman



Decision Letter for Acceptance Ari Asnani.pdf

335K

DECISION LETTER FOR FULL-PAPER SUBMISSION

Purwokerto, January 19th 2021

Dear Ms Ari Asnani

The corresponding author of The Activities of Streptomyces W-5A as Antibacterial and Antibiofilm towards Methicillin-resistant Staphylococcus aureus 2983

The scientific committee of JIMC 2020 thanks the authors for submitting the full-paper manuscript for publication in JIMC 2020 conference proceeding. The paper has been distributed to two different peer-reviewers to be blindly reviewed. Following the comments from the reviewers, we are happy to inform you that the submitted paper is **accepted**. To be published in the JIMC 2020 conference proceeding, the manuscript should fulfill the following conditions:

1. Follow the format of conference proceeding by SCITEPRESS (template and guideline are attached with this letter)
2. All the authors should include ORCID ID in the manuscript.
3. Publication fee has been paid.

Please submit your edited manuscript and payment receipt of publication fee via this link <https://bit.ly/FullPaperJIMC>, no later than two weeks from the date of this letter.

We congratulated you again for the acceptance of your paper, and looking forward to publish your manuscript in our conference proceeding.

On behalf of the scientific committee JIMC 2020,

Tirta Wardana, S.Si, M. Biotech

Program Manager the 1st Jenderal Soedirman Medical Conference (JIMC)

Faculty of Medicine, Universitas Jenderal Soedirman