

PAPER • OPEN ACCESS

## The Control Of Geological Structure At Songgo River Dyke Intrusion In Nanggulan Formation, Western Part Of Yogyakarta-Central Java

To cite this article: Asmoro Widagdo *et al* 2020 *IOP Conf. Ser.: Mater. Sci. Eng.* **982** 012046

View the [article online](#) for updates and enhancements.

# The Control Of Geological Structure At Songgo River Dyke Intrusion In Nanggulan Formation, Western Part Of Yogyakarta-Central Java

Asmoro Widagdo<sup>1,3</sup>, Rachmad Setijadi<sup>1</sup> and Eko Bayu Purwasatriya<sup>1</sup>

<sup>1</sup> Geological Engineering Department, Jenderal Soedirman University (UNSOED), Purwokerto, Indonesia

<sup>3</sup>asmoro.widagdo@unsoed.ac.id

**Abstract.** This study to explain the presence of intrusive rocks in Nanggulan Formation was carried out with a series of geological research methods. The method used in this research is through reference studies from previous research, observing the image of the study area and its surroundings, direct observation of Paleogenic rock bodies, and measurement of fault structures, joints, and folds in the field. Measurement of the elements of geological structure is used to determine the main structure and stress forming structure in Nanggulan Formation rock which is of Paleogene age and surrounding rocks which are Neogene age. The igneous rocks found in the study area is a dyke that cut the sedimentary rock layers of Nanggulan Formation. Dyke igneous rock distribution in the study area forms a northwest-southeast (NW-SE) direction. The distribution of igneous rocks is controlled by geological structures with the main stress direction of northwest-southeast which creates extensional vertical fractures in the direction of the main compressional stress source.

## 1. Introduction

The study area is in the Nanggulan area, the western part of the Yogyakarta Special Region-Central java (Figure 1). This area is very geologically interesting because it is one of the oldest sedimentary rocks found on the island of Java. This rock is the Nanggulan Formation which is an Eocene age rock according to [1] and [2]. In the western part of Yogyakarta, this rock forms a base for Oligocene - Miocene volcanic and sedimentary rocks at the top.

The presence of Nanggulan Formation rocks in this area has attracted many researchers including Van [3], [4], [5], [6], [7], [8] and [9]. Research by [10] described the presence of fold structure, strike-slip faults, and normal faults in the Songgo River area and surroundings. Research by [11], map the distribution of the Nanggulan Formation, and describe north-south trending anticlines and basalt intrusions in the Songgo River area.






**Figure 1.** Research area location in Java Island.

## 2. Regional Stratigraphy

The Nanggulan Formation has a type location on the Songgo River, Nanggulan area. [3], explains that this formation is the oldest rock in the western part of Yogyakarta. The constituent lithology consists of sandstones with lignite, sandy marl, claystone with limonite, marl and limestone concretions, sandstones, tuffs rich in foraminifera and mollusks, estimated thickness up to 350 m. Based on the study of planktonic foraminifera, the Nanggulan Formation has an age range between the Middle Eocene to the Oligocene (Figure 2).



**Figure 2.** Stratigraphic column of study area according to [13] with modifications.

The volcanic rocks referred to by [3] as the Old Andesite Formation (OAF) are the same as those referred to as the Kebo Butak Formation (Figure 2) by [12] and [13]. [13] mentions the constituent rocks of the Kebo Butak Formation are andesite breccias, tuffs, lapilli tuffs, agglomerates and inserts andesite lava flows. Research of [14] describes the rocks that make up the Old Andesite Formation including volcanic breccias, lava, and intrusion. Volcanic breccias consist of volcanic breccias associated with lava; associate with lapilituff and tuff; with volcanic sandstone lenses; and alternating with finely layered sandstones. Lava is found around the center of the eruption (feeder) of the existing Oligo-Miosen volcano. Intrusion rocks consist of andesite, dacite, and basal intrusion.

Publication of [15] states, this volcanic group are covered unconformably by the shallow marine deposits of the Jonggrangan Formation and Sentolo Formation. Based on [16] states that the first carbonate sedimentation occurred at the top of the Early Miocene and occupied high areas such as Kulon Progo and Karang Bolong (Kebumen) resulting in the Sentolo and Jonggrangan Formations of Early Middle Miocene. Reefs are built on the heights of former volcanic bodies. Result of [17] states that carbonate sedimentation formed at the top of the Early Miocene and took place in high areas such as at the height of the early Middle Miocene Kulon Progo.

### 3. Research Method

The study was conducted through a series of reference studies from previous research, observation of the image of the study area and its surroundings, direct observation of the rock body, and measurement of fault structures, joints, and folds in the field. Measurement of the elements of geological structure is used to determine the main structure and the main stress of structure formation in the Nanggulan Formation rock which is of Paleogene age and the surrounding rocks which are of Neogene age.


### 4. Research Result

Nanggulan folds develop on the Nanggulan Formation rocks (Figure 3) in the Girimulyo and Kalibawang areas, Kulon Progo. The anticline fold has a Southwest-Northeast-directed axis. Anticline limb facing to the southeast can be seen in the coal seam on the Watupuru River with dipping  $25^\circ$ . The limb that leads to the northwest has a slope between  $7^\circ$ - $17^\circ$ . Limb in the southwest is found with dipping  $35^\circ$  and  $24^\circ$ .




**Figure 3.** Outcrops of Nanggulan fold limb. a. The northwestward view of the anticline limb on sandstone was inserted by claystone dipping to the NW. b. Westward view of the southeast anticline limb on claystone intercalated by siltstone.

The results of the stereographic analysis of the Nanggulan Folds data (Figure 4) resulted in the position of the northwest limb is  $N228^\circ E / 13^\circ$ . Southeast Limb domiciled  $N4^\circ E / 29^\circ$ . The Nanggulan fold is formed by the main stress ( $T_1$ )  $7^\circ / N105^\circ E$ . The formed axis has the position  $N196^\circ E / 83^\circ$  with the axis line  $07^\circ / N015^\circ E$ . Based on the results of the analysis, the Naggulan Fold is an Upright Horizontal Fold type according to Rickard's Classification, 1971. This fold is formed by the maximum main stress of  $T_1$ :  $7^\circ / 105^\circ E$  or stress from the southeast with an almost horizontal of plunge.




**Figure 4.** Stereographic analysis of Nanggulan folds.



**Figure 5.** Geological map of the study area according to [15]. Songgo River dyke is in the center of the map.

Field observations show the distribution of igneous rocks as shallow intrusions (Figures 5 and 6). These rocks are slightly light gray to dark gray, fine afanite-porphyrific textures, vesicular structures (the structure of holes in dyke), and scoria structures. The vesicular structure only develops in the middle part of the body of intrusion rock that forms a columnar joint.





**Figure 6.** The appearance of basalt igneous rock in the study area [7].


## 5. Discussion

Based on fault kinematics, dyke is possible to form due to maximum extension, which allows magma to flow into the low-pressure zone. The relationship between the dyke pathway and the anticline structure shows its relationship with normal faults or extensional fracture and magmas fill spaces in the strain zone produced by the fault [18]. Magmatism activities are closely related to local area tectonic activities. As long as the area's tectonic activities are still ongoing, it can cause reactivation of magmatism activities [19].

The direction tendency of dyke is likely to be concentrated in the direction of the maximum compressional stress (T1) with the opening towards the minimum compressional stress (T3). The T1 direction is usually horizontal in the contractional tectonic region which is characterized by the presence of a strike-slip fault or thrust fault that is not in line with the vertical dyke intrusion. The trajectory can be controlled by the arrangement of stress-field lithospheric and local stress fields and fractures caused by such as normal faults, thrust faults, and strike-slip faults [20]. According to [21], [22] and [23] mentioned the presence of anticline and thrust faults as controllers for the emergence of the Nanggulan Formation in western Yogyakarta.


Figure 7 shows the movement of magma in the crust along the plane of the thrust fault, illustrating that magma can spread to the surface in a tectonic regime characterized by contractional deformation. If the thrust fault is below the surface with a gentle angle in a flat view, vertical tension fractures can form in the shallow part of the hanging wall subparallel to the imposed shortening. These structures can be formed due to local loading, due to uplifted areas in regional stress fields [24].

Dyke which intrudes into an area can be simplified into a vertical plane, better known as the dyke plane, as the main magma escape route. Normal-tension is perpendicular to the dyke plane. Volumetric changes around the magma sac may be related to the influence of magma overpressure as a source of magma. Active fault zone can cause wider swelling in the magma reservoir which allows volcanic eruptions [25].



**Figure 7.** Intrusion models in contractional tectonic settings [24]. Shows the movement of magma under the surface along the plane of the thrust fault.

Magmatism in the study area is of the same age as northwest-southeast-directed local compression. The maximum main stress is horizontal T1 and the minimum main stress is vertical T3 creating a horizontal / almost horizontal thrust fault structure. Magma flows through the composite system of sub-horizontal tension fractures and thrust fault planes.



**Figure 8.** Schematic block diagram showing the uplift process of the Nanggulan Formation due to strike-slip and thrust fault.

The block diagram scheme (Figure 8) shows the thrust fault and anticline that gave rise to the Nanggulan Formation. In this picture, the rising of the Nanggulan (Eocene) Formation rock is drawn up to cut the young rock of the Sentolo Formation (Middle Miocene). To the north and south of the Nanggulan Formation, the Sentolo Formation is found. So it can be said that laterally the Naggulan Formation is flanked by the Sentolo Formation in the north and south. While in the west there is contact with the Old Andesite Formation [3] or the Kebo Butak Formation [13] or the Gajah Volcano rock according to [26] and [15].

## 6. Conclusion

The igneous rock found in the study area is a dyke that cuts anticline structures of the Nanggulan Formation sedimentary rocks. Dyke igneous rock distribution in the study area forms a northwest-southeast direction. The distribution of igneous rocks is controlled by geological structures with the main direction of stress is southeast-northwest which creates an extensional vertical fracture in the direction of forming stress.

## References

- [1] Premonowati, 1994. Informasi Paleobatimetri Asosiasi Fosil Moluska. Preceedings Geologi dan Geotektonik Pulau Jawa Sejak Akhir Mesozoik Hingga Kuarter, Yogyakarta.
- [2] Harley and Morley 1995. Ultrastructural studies of some fossil and extant palm pollen, and the *reconstruction* of the biogeographical history of subtribes Iguanurinae and Calaminae, Review of Palaeobotany and Palynology **85** 153-182.
- [3] Van Bemmelen R W 1970 The Geology of Indonesia. Vol. IA, General Geology of Indonesia and Adjacent Archipelago, Government Printing Office, The Hague.
- [4] Prasetyadi C 2008 Provenan Batupasir Eosen Jawa Bagian Timur. Proceeding of annual meeting IAGI **37**.
- [5] Rahmad B, Maha M, and Rodhi A 2008 Reflektan Vitrit dan Komposisi Maseral Seam Batubara Eosen Formasi Nanggulan Daerah Kalisonggo, Kecamatan Girimulyo, Kabupaten Kulon Progo, DIY. Prosiding Pertemuan Ilmiah Tahunan IAGI **37**
- [6] Winardi, S, Toha, B, Imron M, and Amijaya D H 2010 The Potency Of Eocene Shale Of Nanggulan Formation As Hydrocarbon Source Rock, Proceedings PIT IAGI Lombok.
- [7] Hartono H G and Pambudi S 2015 Gunung Api Purba Mujil, Kulon Progo, Yogyakarta: Suatu Bukti dan Pemikiran. National Seminar ReTII Ke-**10**, STTNAS, Yogyakarta.
- [8] Pambudi, S and Sujono 2016 Konfigurasi Cekungan Purba Formasi Nanggulan di Daerah Nanggulan, Kulon Progo, Daerah Istimewa Yogyakarta, Proceeding of **11**
- [9] Hartono H G dan Sudradjat A 2017 Nanggulan Formation and Its Problem As a Basement in Kulonprogo Basin, Yogyakarta, Indonesian Journal on Geoscience **4** 71-80.
- [10] Astuti B S, Humantoro R, Hidayat M and Kusuma H D 2016 Analisis Struktur Geologi Jalur Kali Watupuru Dan Kali Songgo Daerah Degan Kulonprogo, Dan Implikasinya Terhadap Penyebaran Batupasir Kuarsa Formasi Nanggulan Yang Berpotensi Sebagai Reservoir. National Seminar, UNPAD, Bandung.
- [11] Saputra R and Akmaluddin 2015 Biostratigrafi Nannofosil Gampingan Formasi Nanggulan Bagian Bawah Berdasarkan Batuan Inti dari Kec. Girimulyo dan Kec. Nanggulan, Kab. Kulon Progo, D.I. Yogyakarta. Prosiding Seminar Nasional Kebumihan Ke-**8**, FT-UGM.
- [12] Rahardjo W, Sukandarrumidi, and Rosidi H M D 1995 Peta Geologi Lembar Yogyakarta. Pusat Penelitian dan Pengembangan Geologi, Bandung.
- [13] Rahardjo W, Sukandarrumidi, and Rosidi H M D 2012 Peta Geologi Lembar Yogyakarta. Pusat Survey Geologi-Badan Geologi-Kementrian ESDM.
- [14] Suroso, Rodhi A, dan Sutanto, 1987 Usulan Penyesuaian Tata Nama Litostratigrafi Kulon Progo, Daerah Istimewa Yogyakarta. Kumpulan Makalah Pertemuan Ilmiah Tahunan XV Ikatan Ahli Geologi Indonesia, Volume **1**, IAGI-Yogyakarta.
- [15] Harjanto A 2011 Vulkanostratigrafi di Daerah Kulon Progo dan Sekitarnya, Daerah Istimewa Yogyakarta. Jurnal Ilmiah MTG, Vol. **4** Yogyakarta.
- [16] Satyana A H and Purwaningsih M E M 2003 Oligo-Miocene Carbonates of Java: Tectonic setting and effects of Vulcanism. Proceedings of Joint Convention Jakarta 2003, The **32nd** IAGI and **28th** HAGI Annual Convention and Exhibition.
- [17] Satyana, A H 2005 Oligo-Miocene Carbonates of Java, Indonesia : Tectonic-Volcanic Setting and Petroleum Implications. Proceedings Indonesian Petroleum Association.



- [18] Alvarez S A, Samaniego A F N, Zenteno D J M, and Aldave L A 2002 Rhyolitic volcanism in extension zone associated with strike-slip tectonics in the Taxco region, southern Mexico. *Journal of Volcanology and Geothermal Research* **118** 1-14.
- [19] Bronto S 2010 *Geologi Gunung Api Purba*. Special publication, Badan Geologi, Kementrian Energi dan Sumberdaya Mineral, Bandung.
- [20] Thouret 1999 Volcanic geomorphology-an overview. *Earth-Science Reviews* **47** 95–131.
- [21] Widagdo A, Pramumijoyo, S P, Harijoko A, Setiawan A 2016 *Kajian Pendahuluan Kontrol Struktur Geologi Terhadap Sebaran Batuan-Batuan Di Daerah Pegunungan Kulonprogo-Yogyakarta*, Proceeding Seminar Nasional Kebumian ke-9 TG FT-UGM, Yogyakarta.
- [22] Widagdo A, Pramumijoyo S P, and Harijoko A 2018 Morphotectono-Volcanic of MenorehGajah-Ijo Volcanic Rock In Western Side of Yogyakarta-Indonesia, *Journal of Geoscience, Engineering, Environment, and Technology (JGEET)* **03**. <http://journal.uir.ac.id/index.php/JGEET>; DOI: 10.24273/jgeet.2018.3.3.1715
- [23] Widagdo A, Pramumijoyo S P, and Harijoko A 2019 The Structural Geology Constellation of Western Boundary of Yogyakarta Basin. *Proceeding Seminar Nasional Kebumian* **12**, 5-6
- [24] Tibaldi A, Pasquare F, and Tormey D 2010 Volcanism in Reverse and Strike-Slip Fault Settings. In: S. Cloetingh, J. Negendank (eds.), *New Frontiers in Integrated Solid Earth Sciences*, International Year of Planet **315**, Earth, DOI 10.1007/978-90-481-2737-5\_9
- [25] Hosono M, Mitsui Y, Ishibashi H, and Kataoka J, 2016 Elastostatic Effects Around A magma reservoir and pathway Due To Historic Earthquake : A Case Study Of Mt. Fuji, Japan, *Progress In Earth and Planetary Science*, **3** 33, DOI 10.1186/s40645-016-0110-9 .
- [26] Barianto D H, Kuncoro P, and Watanabe K, 2010 The Use of Foraminifera Fossils for Reconstructing the Yogyakarta Graben, Yogyakarta, Indonesia. *Journal of South East Asian Applied Geology* **2** 138-143.

PAPER • OPEN ACCESS

## Preface

To cite this article: 2020 *IOP Conf. Ser.: Mater. Sci. Eng.* **982** 011001

View the [article online](#) for updates and enhancements.

## PREFACE

The International Conference on Engineering, Technology and Innovative Researches (ICETIR) 2020 is held in September 2<sup>nd</sup> 2020, organized by the Faculty of Engineering, Universitas Jenderal Soedirman. This conference is aimed to communicate the results of research in the field of engineering and to jointly contribute to society through innovative research. In 2020, we highlight the theme “**Sustainable Engineering and Technology for Reshaping our Community**”.

However, due to the covid19 and travel restriction, we held the conference virtually through zoom meeting, inviting the keynote speakers from the various fields of engineering, along with the accepted papers, which consist of some tracks, i.e., information system, software engineering, computer engineering, mechanical engineering, geological engineering, industrial engineering, electrical power, telecommunication, control systems, agriculture engineering, and civil engineering. On this occasion, we have received submissions from authors originated from 5 countries (Indonesia, Taiwan, Malaysia, Saudi Arabia, Japan), with the attendance number were around 100 participants. We had plenary and parallel session for each presenter with 10 minutes talk, followed by 4 minutes of discussion (question answer session).

We would like to acknowledge all supports for the success of this conference, especially, the board of reviewers, the board of editors, the organizing committee, the sponsors, and all conference participants. We hope that the 2<sup>nd</sup> ICETIR 2020 has been an excellent discussion forum in the field of engineering.

Sugeng Waluyo

The Chairman of ICETIR 2020



### Scientific Committee

1. Prof. Silvio Ranise (Fondazione Bruno Kessler, Italy)
2. Prof. Hitoshi Habe (Kindai University, Japan)
3. Prof. Toshihiko Noguchi (Shizuoka University, Japan)
4. Dr. Albert Causo (Nanyang Technological University, Singapore)
5. Prof. Imam Sadisun (Bandung Institute of Technology, Indonesia)
6. Prof. Budi Santosa (Sepuluh Nopember Institute of Technology, Indonesia)
7. Dr.-Ing. Supriyanto (Advantech Europe, Jerman)
8. Dr. Adie Wahyudie (United Arab Emirates University, United Arab Emirates)
9. Dr. Suroso (Universitas Jenderal Soedirman, Indonesia)
10. Prof. Retno Supriyanti (Universitas Jenderal Soedirman, Indonesia)
11. Dr. Gito Sugiyanto (Universitas Jenderal Soedirman, Indonesia)
12. Dr. Eng. Agus Maryoto (Universitas Jenderal Soedirman, Indonesia)

### Organizing Committee

1. Sugeng Waluyo (Universitas Jenderal Soedirman, Indonesia)
2. Asmoro Widagdo (Universitas Jenderal Soedirman, Indonesia)
3. Rani Aulia Imran (Universitas Jenderal Soedirman, Indonesia)
4. FX Anjar Trilaksono (Universitas Jenderal Soedirman, Indonesia)

### Editors

Editor in Chief : Maulana Rizki Aditama  
Aini Hanifa

Editor :

1. Yogie Indra Kurniawan
2. FX Anjar Trilaksono
3. Januar Aziz Zaenurrochman
4. Huzaely Latief Sunan
5. Ahmad Khalil Gibran
6. M. Syaiful A'alim
7. Ari Fadli
8. Rani Aulia I.

9. Tigar Dian P. A.
10. Amanda Sofiana
11. Katon Muhammad
12. Indro Prakoso
13. Ayu Anggraeni
14. Yudi Syahrullah
15. Dani Nugroho
16. Redityo Januardi



PAPER • OPEN ACCESS

## Peer review declaration

To cite this article: 2020 *IOP Conf. Ser.: Mater. Sci. Eng.* **982** 011002

View the [article online](#) for updates and enhancements.

## Peer review declaration

All papers published in this volume of *Material Science and Engineering: Conference Series* have been peer-reviewed through processes administered by the proceedings Editors. All papers have been processed using the double-blind system, considering the topics, contents and the plagiarism indexes through Turnitin. All papers which passed the editing process is then given to the two different reviewers, to meet the standard of quality for an international publication at this level. Expert referees conducted reviews to the professional and scientific standards expected of a proceedings journal published by IOP Publishing.

- **Type of peer review: Double blind**
- **Conference submission management system: Submission is done through the system**
- **Number of submissions received: 83**
- **Number of submissions sent for review: 83**
- **Number of submissions accepted: 67**
- **Acceptance Rate (Number of Submissions Accepted / Number of Submissions Received X 100): 80.7**
- **Average number of reviews per paper: 5**
- **Total number of reviewers involved: 17**
- **Any additional info on review process: No**
- **Contact person for queries: [maulanarizkia@unsoed.ac.id](mailto:maulanarizkia@unsoed.ac.id) and [sugeng.waluyo@unsoed.ac.id](mailto:sugeng.waluyo@unsoed.ac.id)**

