


Volume 6, Issue 1, April 2021



Published by:
Faculty of Biology
Universitas Gadjah Mada
In collaboration with:



Vol 6, No 1 (2021)

April

Full Issue

[View or download the full issue](#)

Table of Contents

Short Communications

Biomass Estimation of Eaglewood (*Aquilaria filaria* (Oken) Merr.) in the Karst Ecosystem of West Papua

Andes Hamuraby Rozak, Zaenal Mutaqien, Destri Destri

jtbb59221

10.22146/jtbb.59221 Abstract views : 810 | views : 937

Antibacterial Activity of Kecombrang Flower (*Etlingera elatior* (Jack) R.M. Sm) Extract against *Staphylococcus epidermidis* and *Propionibacterium acnes*

Vilya Syafriana, Renita Noviani Purba, Yayah Siti Djuhariah

jtbb58528

10.22146/jtbb.58528 Abstract views : 682 | views : 666

Seed and Germination Study of a New Guinea Endemic Plant Species *Grevillea papuana* Diels.

Farid Kuswantoro

jtbb61115

10.22146/jtbb.61115 Abstract views : 146 | views : 138

Identification of *Dendrobium* (Orchidaceae) in Liwa Botanical Garden Based on Leaf Morphological Characters

Mahfut Mahfut, Tundjung Tripeni Handayani, Sri Wahyuningsih, Sukimin Sukimin

jtbb59423

10.22146/jtbb.59423 Abstract views : 187 | views : 178

Research Articles

Screening and Identification of Mushrooms Growth Promoting Bacteria on Straw Mushrooms (*Volvariella volvacea*)

Indah Juwita Sari, I Nyoman Pugeg Aryantha

jtbb60618

10.22146/jtbb.60618 Abstract views : 657 | views : 585

Antifungal Activity of Bacterial Isolates from Straw Mushroom Cultivation Medium against Phytopathogenic Fungi

Masrukhin Masrukhin, Ade Lia Putri, Tri Ratna Sulistiyani, Muhammad Ilyas, Ismu Purnaningsih, Iwan Saskiawan, Muhammad Yusrun Niam

jtbb59235

10.22146/jtbb.59235 Abstract views : 870 | views : 819

Avifaunal Diversity and Community Structure in Universitas Brawijaya Forest, East Java, Indonesia

Agus Nurrofik, Muhammad Fathoni, Agung Sih Kurnianto, Luhur Septiadi, Nia Kurniawan

jtbb58335

10.22146/jtbb.58335 Abstract views : 627 | views : 629

Callus Induction and secondary metabolite profile from *Elephantopus scaber* L.

Junairiah Junairiah, Diah Ayu Wulandari, Edy Setiti Wida Utami, Nabillah Istighfari Zuraidassanaaz

jtbb59234

10.22146/jtbb.59234 Abstract views : 193 | views : 175

Article
template

Sciwheel



MENDELEY



...Bibliographies Made Easy



grammarly



Username

Password

☐ Remember me

JOURNAL CONTENT

Search

Search Scope

All

Browse

By Issue

By Author

By Title

Other Journals

The Use of DNA Barcoding and Phylogenetic Analysis to Improve Identification of *Usnea* spp. Based on ITS rDNA

jtbb58635

Miftahul Jannah, Muhammad Rifqi Hariri, Rina Sri Kasiandari, Niken Satuti Nur Handayani

10.22146/jtbb.58635 Abstract views : 263 | views : 212

Update on New Species and Record of Fishes in the Coral Triangle Region for the Last 10 Years (2008-2019)

jtbb59230

Fione Yukita Yalindua, Teguh Peristiwady, Putri Saphira Ibrahim

10.22146/jtbb.59230 Abstract views : 146 | views : 134

The Abundance of Fish Species in Branched (*branching*) and Table (*tabulate*) Coral Habitat on Tanjung Beach, Muna Island, Southeast Sulawesi

jtbb59481

Muhamad Tasyian, Muhammad Sirih

10.22146/jtbb.59481 Abstract views : 67 | views : 136

Potential Screening of Bacteriocinogenic-Lactic Acid Bacteria from Mangrove Sediment of Logending Beach for Fisheries Product Preservation

jtbb61927

Dyah Fitri Kusharyati, Taruna Dwi Satwika, Afifah Mariana, Anwar Rovik

10.22146/jtbb.61927 Abstract views : 42 | views : 65

Ethnobotanical and Phytochemical Study of Bayur (*Pterospermum javanicum* Jungh.) on Sasak Tribe around Mount Rinjani National Park, West Lombok as a Conservation Effort

jtbb61008

Putri Sri Andila, Tri Warseno, I Putu Agus Hendra Wibawa, I Gede Tirta

10.22146/jtbb.61008 Abstract views : 72 | views : 66

Predicting the Distribution of Sunda Pangolin (*Manis javanica* Desmarest, 1822) in Way Canguk Research Station, Bukit Barisan Selatan National Park, Lampung

jtbb58612

Silvi Dwi Anasari, Wulan Pusparini, Noviar Andayani

10.22146/jtbb.58612 Abstract views : 25 | views : 18

Review Articles

Nutrigenomic and Biomolecular Aspect of *Moringa oleifera* Leaf Powder as Supplementation for Stunting Children

jtbb60113

Agus Indra Yudhistira Diva Putra, Nyoman Budhi Wirananda Setiawan, Made Indira Dianti Sanjiwani, Ida Ayu Ika Wahyuniari, Agung Wiwiek Indrayani

10.22146/jtbb.60113 Abstract views : 1021 | views : 1191

Editorial address:

Faculty of Biology, UGM

Jl. Teknik Selatan, Sekip Utara, Yogyakarta, 55281, Indonesia

ISSN: 2540-9581 (online)

NOTIFICATIONS

View

Subscribe

CURRENT ISSUE



ISSN BARCODES



KEYWORDS

Alas Purwo National Park
Bali Diversity East Java
LAB Malang Moringa
oleifera RAPD abundance
antioxidant behaviour
biodiversity biomass
conservation extract
flowering fruiting
identification orchid
phenetic relationship
species diversity

59301

View My Stats



Dr. Phoon Lee Quen

Department of Allied Health Sciences, Faculty of Science, Universiti Tunku Abdul Rahman, Malaysia
 Link: [Google Scholar](#), [ORCID](#), [Academic Profile](#)



Sukirno, M.Sc., Ph.D.

Faculty of Biology, Universitas Gadjah Mada, Indonesia
 Link: [Google Scholar](#), [Scopus](#), [Academic Profile](#)



Dr. rer. nat. Andhika Puspito Nugroho

Faculty of Biology, Universitas Gadjah Mada, Indonesia
 Link: [Google Scholar](#), [Scopus](#), [Academic Profile](#)



Assoc. Prof. Dr. Ruqiah Ganda Putri Panjaitan

Biology Education Program, Universitas Tanjungpura, Pontianak, Indonesia
 Link: [Google Scholar](#), [Scopus](#), [Academic Profile](#)



Dr. Abdul Razaq Chasani

Faculty of Biology, Universitas Gadjah Mada, Indonesia
 Link: [Google Scholar](#), [Academic Profile](#)



Dr. Ratna Stia Dewi

Faculty of Biology, Universitas Jenderal Soedirman, Purwokerto, Indonesia
 Link: [Google Scholar](#), [Scopus](#), [Academic Profile](#)

Editorial address:

Faculty of Biology, UGM

Jl. Teknik Selatan, Sekip Utara, Yogyakarta, 55281, Indonesia

ISSN: 2540-9581 (online)

NOTIFICATIONS

► [View](#)

► [Subscribe](#)

ISSN BARCODES



9 772540 958004

KEYWORDS

Alas Purwo National Park
 Bali Diversity East Java
 LAB Malang Moringa
 oleifera RAPD abundance
 antioxidant behaviour
 biodiversity biomass
 conservation extract
 flowering fruiting
 identification orchid
 phenetic relationship
 species diversity

59301

[View My Stats](#)

Menu

Home About the Journal Editorial Policies Focus and Scope Section Policies Peer Review Process Publication Frequency Open Access Policy Audience Publication Ethics Screening for Plagiarism Digital Archiving Statistic Download Article

Home > About the Journal > Editorial Policies

Editorial Policies

- [Focus and Scope](#)
- [Section Policies](#)
- [Peer Review Process](#)
- [Publication Frequency](#)
- [Open Access Policy](#)
- [Audience](#)
- [Publication Ethics](#)
- [Screening for Plagiarism](#)
- [Digital Archiving](#)
- [Statistic Download Article](#)

Focus and Scope

Journal of Tropical Biodiversity and Biotechnology (JTBB) is an open access, peer-reviewed journal dedicated to the publication of novel research concerned with the advancement of tropical biology studies in the Southeast Asia Region.

We invite authors to submit articles in the fields It publishes original scientific work related to a diverse range of fields in tropical biodiversity, functional biology, and biotechnology.

The journal invites research articles, short communication, and reviews concerning biodiversity in the Southeast Asia Region, in these following subjects:

- Descriptive and analytical studies of genetic, species, and community diversity
- Ecological and conservation studies
- Utilisation of tropical organisms in biotechnology
- Novel methods in studying biodiversity
- Ethnobotany and etnozoology
- Bioinformatics

Manuscripts that are not in accordance with the subjects will not be considered to be published. We also discourage manuscripts written by a single author.

Section Policies

Short Communications

Short communications are concise documents for providing brief novel findings that have inadequate content to satisfy a full-length research article. They are intended to be used for the reporting of preliminary studies or short descriptive studies.

☒ Open Submissions ☒ Indexed ☒ Peer Reviewed

Research Articles

Research articles are comprehensive research reports containing detailed descriptions of experimental work, clearly interpreting and discussing the theoretical and experimental results and data.

☒ Open Submissions ☒ Indexed ☒ Peer Reviewed

Review Articles

A systematic review is a review of a clearly formulated question that uses systematic and reproducible methods to identify, select and critically appraise all relevant research, and to collect and analyse data from the studies that are included in the review.

A systematic review should answer a focused research question, employs a comprehensive, reproducible search strategy, identifies all relevant studies (both published and unpublished), assesses all results for inclusion/exclusion, and for quality, presents an unbiased, balanced summary of findings.

☒ Open Submissions ☒ Indexed ☒ Peer Reviewed

ISSN 2540-9581 (online)

Home About the Journal Editorial Policies

Focus and Scope Section Policies

Peer Review Process

Publication Frequency

Open Access Policy

Audience Publication Ethics

Screening for Plagiarism

Digital Archiving

Statistic Download Article

Article template



Article
template

ISI/SCOPUS INDEXED

Sciwheel

MENDELEY

EndNote
...Bibliographies Made Easy

grammarly

JTBB

Username

Password

☐ Remember me

Login

JOURNAL CONTENT

Search

Search Scope

All

Search

Browse

By Issue

By Author

By Title

Other Journals

Menu

[Home](#) | [About](#) | [Contact](#) | [Register](#) | [Search](#) | [Submit](#) | [Articles](#) | [Instructions](#) | [Editorial](#) | [Privacy](#) | [Contact Us](#)
[Home](#) > [Indexing & Accreditation](#)

Indexing & Accreditation

This journal is accredited to [Sinta 1](#) by the Indonesian Government.

This journal is indexed in:

- [Scopus](#)
- [Directory of Open Access Journals \(DOAJ\)](#)
- [Crossref](#)
- [Google Scholar](#)
- [Science and Technology Index \(SINTA\)](#)
- [Garba Rujukan Digital \(GARUDA\)](#)

Other affiliations:

- [Publons](#)

Editorial address:

Faculty of Biology, UGM

Jl. Teknik Selatan, Sekip Utara, Yogyakarta, 55281, Indonesia

ISSN: 2540-9581 (online)



Article
template



Sciwheel



MENDELEY



grammarly

ASST

Username

Password

☐ Remember me

Login

JOURNAL CONTENT

Search

Search Scope

All

Search

Browse

► By Issue

► By Author

► By Title

► Other Journals

Research Article

Potential Screening of Bacteriocinogenic-Lactic Acid Bacteria from Mangrove Sediment of Logending Beach for Fisheries Product Preservation

Dyah Fitri Kusharyati^{1*}, Taruna Dwi Satwika¹, Affah Mariana¹, Anwar Rovik²

1) Laboratory of Microbiology, Department of Microbiology, Faculty of Biology, Universitas Jenderal Soedirman, Jl. DR. Soeparno No. 63, Grendeng, Purwokerto Utara, Banyumas, Jawa Tengah, Indonesia 53122

2) Centre of Tropical Medicine, Faculty of Medicine, Public Health, and Nursing, Universitas Gadjah Mada, Jl. Teknik Utara, Kocoran, Depok, Sleman, Yogyakarta, Indonesia 55281

* Corresponding author, email: dfitri.k@gmail.com

Submitted: 06 December 2020; Accepted: 08 March 2021; Published online: 01 April 2021

ABSTRACT

The meat and fisheries products have high nutritional content which is highly ideal for bacterial growth. Lactic Acid Bacteria (LAB) have several potential advantages as a bio-preservative agent in the food industry because they produce antimicrobial substances against pathogenic bacteria e.g. bacteriocin. Our previous study has succeeded in isolating and characterizing LAB from the mangrove sediments of Logending Beach, Kebumen. This present study aimed to determine the activity of bacteriocinogenic-LAB against food-borne pathogens and their potential for fisheries product preservation. The study consisted of five serial stages, as follows: screening of LAB isolates, cell-free supernatant production and its inhibition activity, extraction of partially purified bacteriocin, bacteriocin confirmation against proteolytic enzymes, and in-vitro test of partially-purified bacteriocin against *Listeria monocytogenes*, *Shigella flexneri*, and *Salmonella typhi*. A total of 25 out of 99 isolates were able to grow on MRSA+1% CaCO₃ medium. Initial screening showed that the cell-free supernatant of 14 LAB isolates was able to inhibit the growth of *S. typhi*, *S. flexneri*, and *L. monocytogenes*. There was an increased inhibitory activity of partially purified bacteriocin when compared with the cell-free supernatant which was statistically different ($p < 0.01$). It indicated that the purification was successfully performed. Bacteriocin expressed a lower inhibition against *S. typhi* than *L. monocytogenes* and *S. flexneri*. The ANOVA test showed that each indicator pathogenic-bacterium expresses a very significant sensitivity to the partially purified bacteriocin.

Keywords: Bacteriocin, fisheries product, LAB, mangrove sediments, food-borne pathogens

INTRODUCTION

The human population in the world is increasing rapidly which influences the high demand for food availability. In this modern era, the safety, quality, and health profile of food products are being a big concern. The meat and fisheries products have high nutritional content. On the other hand, they are highly ideal for bacterial growth e.g. *Listeria monocytogenes*, *Salmonella* spp., and *Shigella* spp. which cause food-borne diseases (Jennison & Verma 2004; Novotny et al. 2010).

Bacterial contamination of seafood and its products affects the nutritional properties or undesirable organoleptic changes. The contamination may take place during harvesting, handling, preparation, processing, transportation, and storage. Many food treatments have been applied to control bacterial contamination, such as salt administration, smoking, canning, freezing, and vacuum application. However, these treatments are not sufficient to kill pathogenic microorganisms since some bacteria have been reported to be resistant to high salt concentration, drying, freezing, and heat processes (Dupard et al. 2006).

Bio-preservation is a promising approach to control bacterial contamination and extend the shelf life of foods, including the application of living microorganisms and their metabolites (Moradi et al. 2020). Bacteriocin, a secondary metabolite of Lactic Acid Bacteria (LAB), has several potential advantages as a bio-preservative agent in the food industry including (a) generally recognized as safe substances, (ii) inactive and nontoxic on eukaryotic cells, (iii) inactivated by digestive proteases, (iv) usually pH and heat-tolerant, and (v) have a relatively broad antimicrobial spectrum against many food-borne pathogenic and spoilage bacteria (Siedler et al. 2019).

LAB are reported to be able to produce antimicrobial substances, such as carbon dioxide, diacetyl, acetaldehyde, ethanol, hydrogen peroxide, and bacteriocin (de-Vuyst & Leroy 2007; Liao & Nyachoti 2017). Bacteriocin has bactericidal or bacteriostatic effects (Cheong et al. 2014; Gerez et al. 2013; Yang et al. 2012) against various foodborne pathogens, such as *Staphylococcus aureus*, *Bacillus cereus*, *L. monocytogenes*, *Clostridium botulinum*, and *E. coli* (Aunpad et al. 2007; Dobson et al. 2012).

LAB are commonly isolated from fermented food products, milk, and the intestinal tract of humans or animals (Dobson et al. 2012; Kusharyati et al. 2020). Meanwhile, only a few studies isolated LAB from soils, water, and plants. Mangroves are unique ecosystems that harbor unique and diverse microorganism groups, such as fungi, actinomycetes, and bacteria. Our previous study has succeeded in isolating and characterizing LAB from the mangrove sediments of Logending Beach, Kebumen. This present study aimed to determine the activity of bacteriocinogenic-LAB against food-borne pathogens and their potential for fisheries product preservation.

MATERIALS AND METHODS

The study consisted of five serial stages, as follows: screening of LAB isolates, cell-free supernatant production and its inhibition activity, production of partially-purified bacteriocin, bacteriocin confirmation against proteolytic enzymes, and in-vitro test of partially purified bacteriocin against *L. monocytogenes*, *Shigella flexneri*, and *Salmonella typhi*.

Screening of bacteriocinogenic-LAB

One loop of isolates originating from mangrove sediments of Logending Beach Kebumen was grown on the de-Man Rogose Sharpe Agar (MRSA) medium (Oxoid) which was supplemented with 1% CaCO_3 . The spot inoculation was performed triplicate in each dish. Bacterial incubation was carried out for 48 hours at 37°C. Isolates that expressed a clear zone around the growing colony were assumed as Lactic Acid Bacteria. The bacterial characterization was performed following Hendrati et al. (2017). The selected isolates were stocked on MRSA slant medium. The pathogenic bacteria, *S. typhi*, and *S. flexneri* were re-cultured using Salmonella Shigella Agar (SSA) medium (Merck), meanwhile, *L. monocytogenes* using Nutrient Agar (NA) medium (Merck).

Pre-screening of LAB's inhibition activity

One loop of LAB isolate was inoculated in MRS Broth (MRSB) medium and incubated for 18 hours at 37°C. The bacterial culture was centrifuged for 10 minutes at 10,000 rpm to obtain a bacterial cell-free supernatant (CFS). Its inhibition activity was evaluated using Kirby's Bauer method by following Hendrati et al. (2017). The pathogenic bacteria were inoculated on Nutrient Broth (NB) medium (1% v/v), then was incubated for 8 hours at 37°C. The pathogenic bacteria were spread on NA medium. A 6 mm diameter paper disc was dropped by 20 µL of cell-free supernatant, then was placed into the pathogenic bacterial lawn-medium. The assay was performed triplicate in each dish. Incubation was carried out for 24 hours at 37°C. The formed inhibition zone was observed and calculated.

Production of partially-purified bacteriocin

One loop of LAB isolates was inoculated into 10 mL MRSB medium, then was incubated for 18 hours at 37°C. This culture was used as an inoculum. One milliliter (~ 10⁸ CFU/mL) of inoculum was inoculated into 100 mL MRSB medium, then was incubated at 37°C for 24 hours. The 24-hour LAB culture was cold-centrifuged (Thermo Scientific) at 10,000 rpm, 4°C for 10 minutes. The supernatant was salted out by adding ammonium sulfate. The mixture was homogenized using a magnetic stirrer. The 50% ammonium sulfate was slowly added to gradually precipitate the bacteriocin until the end of saturation. The precipitated bacteriocin was separated from the mixture by performing a cold-centrifugation at 10,000 rpm for 15 minutes. Then, the partially-purified bacteriocin was dissolved into 2 mL of 0.1 M phosphate buffer saline (PBS) with pH 5.3.

Bacteriocin confirmation against proteolytic enzymes

A total of 200 µL partially-purified bacteriocin was mixed with 20 µL of proteolytic enzyme solution (one gram of proteolytic 'papain' enzymes was dissolved into 1 mL of 0.1 M PBS with pH 5.3), then was incubated at 37°C for 2 hours. Its inhibition activity was evaluated using Kirby's Bauer method by following Hendrati et al. (2017). The degraded bacteriocin did not show a clear zone around the paper disc (Nithya et al. 2012).

In-vitro test of partially-purified bacteriocin against pathogenic bacteria

The inhibition activity of crude bacteriocin was evaluated using Kirby's Bauer method following by following Hendrati et al. (2017). The diameter of the inhibition zone expressed by CFS and partially-purified bacteriocin were then compared to evaluate the success of bacteriocin extraction.

Data analysis

The data of inhibitory activity of partially-purified bacteriocin against pathogenic bacteria was analyzed by using an analysis of variance (ANOVA) at 95% and 99% levels of confidence.

RESULTS AND DISCUSSION

The consumption of seafood in various countries is increasing since the global population is increasing rapidly (Elbashir et al. 2018). Food spoilage results in undesirable odor, texture, and appearance which make it unfit for consumption (Odeyemi et al. 2020). LAB is effective to enhance the shelf life of food products and inhibit undesirable microorganism growth in food (Xi et al. 2018). Therefore, LAB has been broadly applied in various food industries. Food industries commonly use individual and/or consortium of different LAB, including genera *Lactococcus* sp., *Streptococcus* sp., *Pediococcus* sp.,

Enterococcus sp., *Lactobacillus* sp., *Leuconostoc* sp., and *Weissella* sp. (Hamad et al. 2020; Mokhtar et al. 2016; Nithya et al. 2012).

In this present study, we screened and determined the inhibitory activity of bacteriocinogenic-LAB against food-borne pathogens. A total of 99 bacterial isolates originating from mangrove sediment of Logending Beach (unpublished data) were grown on MRSA+1% CaCO₃ as a selective medium for Lactic Acid Bacteria groups. A total of 25 isolates were able to grow on MRSA+1% CaCO₃ medium and expressed a clear zone around the colonies. It indicated the ability of bacteria to grow and dissolve the CaCO₃ in the medium (Mahulette et al. 2016). Gram-positive, non-spore-forming, and Catalase negative isolates were selected for further analysis (data not shown). Pringsulaka et al. (2012) described a LAB as Gram-positive, non-spore-forming, cocci, cocci-bacilli, or rods-shape cells, have anaerobic respiration, and Catalase-negative.




Figure 1. The diameter of the inhibition zone of cell-free supernatant against pathogenic bacteria.

Garrido et al. (2012) reported three major foodborne pathogens affecting people worldwide are *Salmonella* spp., *Shigella* spp., and *Listeria monocytogenes*. *Salmonella* genus is the leading cause of food-borne outbreaks and remains a major public health concern. *Salmonella* spp. are Gram-negative, rod-shaped, facultatively anaerobic, usually motile, Catalase-positive, and Oxidase-negative. *Salmonella* produces enterotoxins and causes inflammatory reactions and diarrhea. Symptoms often start 12-72 hours after the ingestion of contaminated food. *Salmonella* infections from the consumption of seafood products are most commonly associated with raw, undercooked, and poorly cooked seafood (Iwamoto et al. 2010).

Shigella spp. are reported as the third most frequently isolated bacteria from foodstuffs (WHO 2005). *Shigella flexneri* is reported as responsible for shigellosis outbreaks in developing countries (Shahin et al. 2018). *Shigella* spp. are Gram-negative, rod-shaped, non-motile, Oxidase-negative, and non-lactose fermenting bacteria. *Shigella* produces enterotoxin 1 and 2 which causes watery loose stool, fever, abdominal pain, and bloody diarrhea (Scallan et al. 2011).

Listeria monocytogenes is a Gram-positive, rod-shaped, non-spore-forming, Catalase-positive, and glucose fermenter. The bacterium can adapt to various environmental conditions such as a broad range of temperatures, pH, and high salt content (Aspri et al. 2017). Food including seafood

contamination by *L. monocytogenes* is recognized as a public health and food safety concern since early 1981. The transmission to humans is mainly through the consumption of ready-to-eat foods (Miya et al. 2010).

The LAB cell-free supernatant was tested against both Gram-negative and positive pathogenic bacteria i.e. *S. typhi*, *S. flexneri*, and *L. monocytogenes*. A total of 14 LAB isolates could inhibit all tested pathogenic bacteria. This is indicated by the presence of a clear zone or zone of inhibition around the paper disc (previously dropped by cell-free supernatant) on the bacterial lawn. The largest inhibition against *S. typhi* was shown by LAB isolate LG109 with an average diameter of 8.17 mm. The largest inhibition against *S. flexneri* was shown by LAB isolate LG7 with an average diameter of 13.00 mm. While the largest inhibition against *L. monocytogenes* was shown by LAB isolate LG10 with an average diameter of 8.67 mm. All LAB isolates express a broad and strong inhibition (Figure 1). Pan et al. (2009) categorized the more than 6 mm clear zone expresses a strong inhibition activity, a 3-6 mm clear zone as moderate, while a 0-3 mm clear zone as weak inhibitory. Some research reported the antimicrobial activity of bacteriocin against both Gram-negative and positive bacteria (Cotter et al. 2005; Duranti et al. 2017; Martinez et al. 2013).




Figure 2. Bacteriocin confirmation against proteolytic enzymes, no clear zone was shown around the paper disc (A: isolate LG50, B: isolate LG135, C: isolate LG73, D: isolate LG17).

The presence of bacteriocin was confirmed by mixing the cell-free supernatant with proteolytic enzymes. Confirmatory tests of CFS extracted from all LAB isolates showed no clear zone presence around the paper disc which indicated the loss of their inhibition activity against pathogenic bacteria (Figure 2). Bacteriocin is proteinaceous in nature which could be inactivated by a proteolytic enzyme (Nithya et al. 2012; Yang et al. 2012). Therefore, it was agreed that bacteriocin is present in the cell-free supernatant.

Cell-free supernatant may contain bacteriocins, organic acids, enzymes, alcohols, and low-molecular-mass substances which are the main metabolites responsible for the antimicrobial action of LAB (Chen et al. 2003). The salted-out method was performed to partially purify the bacteriocin and removing other substances. To evaluate the success of bacteriocin partial purification, we monitor their inhibition activity against pathogenic bacteria. The partially-purified bacteriocin showed a bigger clear zone than the cell-free supernatant (Figure 3). However, the total inhibition zone diameter was varied among isolates. The largest inhibition zone was shown by LAB isolate LG7 against




Figure 3. The inhibition zone of partially-purified bacteriocin against *S. typhi* (A), *S. flexneri* (B), *L. monocytogenes* (C), (black arrow pointed the formed clear zone).

S. flexneri with an average diameter of 14.67 mm, followed by LAB isolate LG71 against *L. monocytogenes* with 15.17 mm in diameter.

Bacteriocin can inhibit the growth of pathogenic bacteria due to the electrostatic interactions between positively charged bacteriocin and negatively charged cytoplasmic membrane lipids. The hydrophobic part of the bacteriocin will enter the cytoplasmic membrane by forming pores. This pore formation will cause the failure of the proton motive force (PMF). PMF is a proton that forms energy to be used in various cell activities including bacterial cell metabolism (Chen et al. 2003; Perez et al. 2014). On the other hand, the bacteriocin-producing bacteria has its immunity to the produced bacteriocin with a specific immune system (Martinez et al. 2013).

There was an increased inhibitory activity of all the partially purified bacteriocin when compared with the cell-free supernatant (Figure 4). It indicated that the purification was successfully performed. Also, these results were statistically significant ($p < 0.01$). The lower activity of cell-free supernatant might be due to the presence of various metabolites. Moradi et al. (2020) suggested that the biological activity of bacteriocin is attributed to the purification process. Moreover, different purification methods of bacteriocin might result in different levels of purification and yields.

The inhibitory action of bacteriocin can vary between identical species within- and inter-genera (Castellano et al. 2004). The data showed that bacteriocin expressed a lower inhibition against *S. typhi* than *L. monocytogenes* and *S. flexneri*. The ANOVA test showed that each pathogenic bacterium expresses a very significant sensitivity to the partially-purified bacteriocin (Table 1). The high sensitivity was shown by *S. flexneri*, *L. monocytogenes*, and *S. typhi*, respectively. Interestingly, the inhibitory activity of bacteriocin against *S. flexneri* (Gram-negative) was higher than *L. monocytogenes* (Gram-positive bacteria). Gram-negative bacteria are generally resistant to the bacteriocin produced by Gram-positive bacteria due to their outer membrane acts as an effective and protective barrier (Cao-Hoang et al. 2010). However, Caridi (2002) studied *Lactobacillus paracasei* and its strong activity of bacteriocin against *Escherichia coli*.

Table 1. DMRT test of bacterial sensitivity to partially-purified bacteriocin.

Pathogens	Average of total inhibition zone (mm)
<i>S. typhi</i>	8.83 ± 1.09^a
<i>L. monocytogenes</i>	9.30 ± 1.57^b
<i>S. flexneri</i>	10.38 ± 1.58^c




Figure 4. The comparison of inhibition zone of cell-free supernatant (CFS) and partially purified bacteriocin against *S. typhi* (a), *S. flexneri* (b), *L. monocytogenes* (c).

In general, Gram-positive bacteria are more sensitive to bacteriocin. The cell wall structure of Gram-positive bacteria has lipopolysaccharide, lipoprotein, and phospholipid composition which is lower than Gram-negative bacteria. The simpler cell wall structure of Gram-positive bacteria is facilitating the easier activity of bacteriocin (Cao-Hoang et al. 2010; Usmiati & Marwati 2007). Although, some Gram-positive bacteria have been reported resistant to LAB bacteriocins (e.g. nisin, lactocin, enterocin), such as *L. monocytogenes*, *L. innocua*, *Staphylococcus aureus*, *Clostridium botulinum*, and *Bacillus cereus* (Garsa et al. 2014; Vignolo et al. 2000).

This present study showed a wide range of antimicrobial activity against food-spoilage and/or food-borne pathogens, both Gram-positive and Gram-negative bacteria. It indicating that bacteriocin produced by LAB from mangrove sediments are suitable as bio-preservatives for food including fisheries products. Previous independent studies have reported the application of bacteriocinogenic-LAB from nature origin as bio-preservative, such as *Lactobacillus curvatus* BCS35 marine origin in fish bio-preservation (Gómez-Sala et al. 2016), *Leuconostoc* sp. application in ground meat to reduce *E. coli* contamination (Koo et al. 2015). Our present paper did not study the species-specific effects which possibly influence the bacteriocin's efficacy. Therefore, in-vivo studies are urged to perform. As well as, the study to confirm the type and classes of produced bacteriocins, since LAB produces a variety of bacteriocins (Elayaraja et al. 2014).

CONCLUSION

Our study found 14 bacteriocinogenic-LAB isolates originating from mangrove sediments, Logending Beach Kebumen had a wide range of antimicrobial activity against pathogenic bacteria *S. typhi*, *S. flexneri*, and *L. monocytogenes*. The ANOVA test showed a statistical difference of inhibitory activity between cell-free supernatant and partially-purified bacteriocin and their relationship to the type of indicator pathogenic-bacteria.

AUTHORS CONTRIBUTION

Conceptualization: DFK; Methodology: TDS, AM, and AR; Investigation: TDS and AM; Writing—original draft preparation: DFK, TDS, and AR; Writing—review and editing: AR.

ACKNOWLEDGMENTS

The author thanks the Institute of Research and Community Services, Universitas Jenderal Soedirman for funding support through the scheme of DIPA UNSOED 2020.

CONFLICT OF INTEREST

The author declares that there is no conflict of interest in this study.

REFERENCES

- Aspri, M. et al., 2017. Application of bacteriocin-producing *Enterococcus faecium* isolated from Donkey milk, in the bio-control of *Listeria monocytogenes* in fresh whey cheese. *International Dairy Journal*, 73, pp.1-9.
- Aunpad, R. et al., 2007. Bacteriocins with anti-MRSA activity produced by water and soil isolated bacteria. *Annals of Microbiology*, 57(1), pp.9-14.
- Cao-Hoang, L. et al., 2010. Potential of nisin-incorporated sodium caseinate films to control *Listeria* in artificially contaminated cheese. *Food Microbiology*, 27, pp.940-944.
- Caridi, A., 2002. Selection of *Escherichia coli* inhibiting strains of *Lactobacillus paracasei* subsp. *paracasei*. *Journal of Industrial Microbiology and Biotechnology*, 29, pp.303-308.
- Castellano, P. et al., 2004. The control of *Listeria innocua* and *Lactobacillus sakei* in broth and meat slurry with the bacteriocinogenic strain *Lactobacillus casei* CRL705. *Food Microbiology*, 21, pp.291-298.
- Chen, H. & Hoover, D.G., 2003. Bacteriocins and their food application. *Comprehensive Reviews in Food Science and Food Safety*, 2, pp.82-100.

- Cheong, E.Y.L. et al., 2014. Isolation of lactic acid bacteria with antifungal activity against the common cheese spoilage mold *Penicillium commune* and their potential as bio-preservatives in cheese. *Food Control*, 46, pp.91-97.
- Cotter, P.D. et al., 2005. Bacteriocins: developing innate immunity for food. *Nature Review of Microbiology*, 3, pp.777-778.
- De-Vuyst, L. & Leroy, F., 2007. Bacteriocins from lactic acid bacteria: production, purification and food applications. *Journal of Molecular Microbiology and Biotechnology*, 13, pp.194-199.
- Dobson, A. et al., 2012. Bacteriocin production: a probiotic trait?. *Applied Environmental Microbiology*, 78(1), pp.1-6.
- Dupard, T. et al., 2006. Antimicrobial effect of cetylpyridinium chloride on *Listeria monocytogenes* V7 growth on the surface of raw and cooked retail shrimp. *Journal of Food Science*, 71(7), pp.241-244.
- Duranti, S. et al., 2017. Maternal inheritance of bifidobacterial communities and bifido-phages in infants through vertical transmission. *Microbiome*, 5(1), pp.66.
- Elayaraja, S. et al., 2014. Production, purification and characterization of bacteriocin from *Lactobacillus murinus* AU06 and its broad antibacterial spectrum. *Asian Pacific Journal of Tropical Biomedicine*, 4, pp.305-311.
- Elbashir, S. et al., 2018. Seafood pathogens and information on antimicrobial resistance: a review. *Food Microbiology*, 70, pp.85-93.
- Garrido, A. et al., 2012. Development of a multiplex real-time PCR method for simultaneous detection of *Salmonella enterica*, *Shigella flexneri*, and *Listeria monocytogenes* in processed food samples. *European Food Research and Technology*, 234, pp.571-580.
- Garsa, A.K. et al., 2014. Bacteriocin production and different strategies for their recovery and purification. *Probiotics and Antimicrobial Protein*, 6, pp.47-58.
- Gerez, C.L. et al., 2013. Control of spoilage fungi by lactic acid bacteria. *Biological Control*, 64, pp.231-237.
- Gómez-Sala, B. et al., 2016. Strategies to increase the hygienic and economic value of fresh fish: bio-preservation using lactic acid bacteria of marine origin. *International Journal of Food Microbiology*, 223, pp.41-49.
- Hamad, G.M. et al., 2020. Commercial probiotic cell-free supernatants for inhibition of *Clostridium perfringens* poultry meat infection in Egypt. *Anaerobe*, 62, pp.102-181.
- Hendrati, P.M. et al., 2017. Characterization of bifidobacteria from infant feces with different modes of birth at Purwokerto, Indonesia. *Biodiversitas*, 18(3), pp.1265-1269.
- Iwamoto, M. et al., 2010. Epidemiology of seafood-associated infections in the United States. *Clinical Microbiology Review*, 23, pp.399-411.
- Jennison, A.V. & Verma, N.K., 2004. *Shigella flexneri* infection: pathogenesis and vaccine development. *FEMS Microbiology Review*, 28, pp.43-58.
- Koo, O.K. et al., 2015. Antimicrobial potential of *Leuconostoc* sp. against *E. coli* O157:H7 in ground meat. *Journal of the Korean Society for Applied Biological Chemistry*, 58(6), pp.831-838.
- Kusharyati, D.F. et al., 2020. Bifidobacterium from infant stool: the diversity and potential screening. *Biodiversitas*, 21(6), pp.2506-2513.
- Liao, S.F. & Nyachoti, C.M., 2017. Using probiotics to improve swine gut health and nutrient utilization. *Animal Nutrition*, 3, pp.331-343.
- Mahulette, F. et al., 2016. Isolation and characterization of lactic acid bacteria from Inasua. *Journal of Tropical Biodiversity and Biotechnology*, 1, pp.71.
- Martinez, F.A.C. et al., 2013. Bacteriocin production by *Bifidobacterium* spp.: a review. *Biotechnology Advances*, 31, pp.482-488.

- Miya, S. et al., 2010. Risk of *Listeria monocytogenes* contamination of raw ready-to-eat seafood products available at retail outlets in Japan. *Applied and Environmental Microbiology*, 76(10), pp.3383-3386.
- Mokhtar, M. et al., 2016. Effect of bacteriocins (from *Bifidobacterium* spp.) on the prevalence of some *Aeromonas* and *Pseudomonas* species in minced meat during cold storage. *Journal of Food and Nutritional Disorders*, 5(1).
- Moradi, M. et al., 2020. Postbiotics produced by lactic acid bacteria: the next frontier in food safety. *Comprehensive Reviews on Food Science and Food Safety*, 19, pp.3390-3415.
- Nithya, K. et al., 2012. Characterization of bacteriocin producing lactic acid bacteria and its application as a food preservative. *African Journal of Microbiology Research*, 6(6), pp.1138-1146.
- Novotny, L. et al., 2010. Morphology and distribution of granulomatous inflammation in freshwater ornamental fish infected with mycobacteria. *Journal of Fish Disease*, 33, pp.947-955.
- Odeyemi, O.A. et al., 2020. Understanding spoilage microbial community and spoilage mechanisms in foods of animal origin. *Comprehensive Review in Food Science and Food Safety*, 19, pp.311-331.
- Pan, X. et al., 2009. The acid, bile tolerance, and antimicrobial property of *Lactobacillus acidophilus* NIT. *Journal of Food Control*, 20(6), pp.598-602.
- Perez, R.H. et al., 2014. Novel bacteriocins from lactic acid bacteria (LAB): various structures and applications. *Microbial Cell Factories*, 13(1), pp.1-13.
- Pringsulaka, O. et al., 2012. Partial characterization of bacteriocins produced by lactic acid bacteria isolated from Thai fermented meat and fish products. *Food Control*, 23(2), pp.547-551.
- Scallan, E. et al., 2011. Foodborne illness acquired in the United States - major pathogens. *Emerging Infectious Disease*, 17(1), pp.7-15.
- Shahin, K. et al., 2018. Isolation, characterization and genomic analysis of a novel lytic bacteriophage vB_SsoS-IISF002 infecting *Shigella sonnei* and *Shigella flexneri*. *Journal of Medical Microbiology*, 67, pp.376-386.
- Siedler, S. et al., 2019. Bio-protective mechanisms of lactic acid bacteria against fungal spoilage of food. *Current Opinion in Biotechnology*, 56, pp.138-146.
- Usmiati, S. & Marwati, T., 2007. Seleksi dan optimasi proses produksi bakteriosin dari *Lactobacillus* sp. *Jurnal Pascapanen*, 4(1), pp.27-37.
- Vignolo, G. et al., 2000. Combined effect of bacteriocins on the survival of various *Listeria* species in broth and meat system. *Current Microbiology*, 41, pp.410-416.
- World Health Organization, 2005. Guidelines for the control of shigellosis, including epidemics due to *Shigella dysenteriae* type 1. WHO Press: Geneva.
- Xi, Q. et al., 2018. Purification and characterization of a bacteriocin produced by a strain of *Enterococcus faecalis* TG2. *Applied Biochemistry and Biotechnology*, 184, pp.1106-1119.
- Yang, E. et al., 2012. Antimicrobial activity of bacteriocin-producing lactic acid bacteria isolated from cheeses and yogurts. *AMB Express*, 2, pp.48.