



REINWARDTIA

A JOURNAL ON TAXONOMIC BOTANY, PLANT SOCIOLOGY AND ECOLOGY

ISSN 0034 – 365 X | E-ISSN 2337 – 8824 | Accredited 158/E/KPT/2021



2021 20 (2)

Editorial Team

Chief Editor

- [KARTINI KRAMADIBRATA](#), Herbarium Bogoriense, Botany Division, Research Center for Biology, Indonesian Institute of Sciences (LIPI), Indonesia

International Editors

- [EIZI SUZUKI](#), Department of Earth and Environmental Sciences, Kagoshima University, Japan
- [JUN WEN](#), National Museum of Natural History, Smithsonian Institution, United States
- [DEDY DARNAEDI](#), Herbarium Bogoriense, Botany Division, Research Center for Biology, Indonesian Institute of Sciences (LIPI), Indonesia
- [BARRY J. CONN](#), School of Life and Environmental Sciences, The University of Sydney, Australia
- [TUKIRIN PARTOMIHARDJO](#), Herbarium Bogoriense, Botany Division, Research Center for Biology, Indonesian Institute of Sciences (LIPI), Indonesia
- [DAVID FRODIN](#), Royal Botanic Gardens Kew, England, United Kingdom
- [GRAHAM EAGLETON](#), Wagstaffe, NSW, Australia, Australia

Editors

- [JOENI SETIJO RAHAJOE](#), Herbarium Bogoriense, Botany Division, Research Center for Biology, Indonesian Institute of Sciences (LIPI), Indonesia
- [HIMMAH RUSTIAMI](#), Herbarium Bogoriense, Botany Division, Research Center for Biology, Indonesian Institute of Sciences (LIPI), Indonesia
- [TOPIK HIDAYAT](#), Universitas Pendidikan Indonesia (UPI)- Bandung, Indonesia
- [MARLINA ARDIYANI](#), Herbarium Bogoriense, Botany Division, Research Center for Biology, Indonesian Institute of Sciences (LIPI), Indonesia
- [LULUT DWI SULISTYANINGSIH](#), Herbarium Bogoriense, Botany Division, Research Center for Biology, Indonesian Institute of Sciences (LIPI), Indonesia
- [EKA FATMAWATI TIHURUA](#), Herbarium Bogoriense, Botany Division, Research Center for Biology, Indonesian Institute of Sciences (LIPI), Indonesia, Indonesia

Administrator

- [MUHAMMAD RUSLAN BUKHORI](#), Botany Division, Research Center for Biology, Indonesian Institute of Sciences (LIPI), Indonesia

Layout Editor

- [Liana Astuti](#), Botany Division, Research Center for Biology, Indonesian Institute of Sciences (LIPI), Indonesia

Illustrator

- [WAHYUDI SANTOSO](#), Herbarium Bogoriense, Botany Division, Research Center for Biology, Indonesian Institute of Sciences (LIPI), Indonesia
- [ANNE KUSUMAWATY](#), Herbarium Bogoriense, Botany Division, Research Center for Biology, Indonesian Institute of Sciences (LIPI), Indonesia

Web administrator

- [YULIA ARIS KARTIKA](#), Research Center for Biology, Indonesian Institute of Sciences (LIPI), Indonesia

Indexing/ Partners



[Home](#) / [Archives](#) / [Vol 20, No 2 \(2021\)](#)

Vol 20, No 2 (2021)

Reinwardtia

The articles in this issue were authored/co-authored by authors from Indonesia, Malaysia, the Netherlands, and the United Kingdom.

Full Issue

View or download the full issue

[Volume 20 No 2](#)

Table of Contents

Front Cover Reinwardtia Vol. 20, No 2 (2021)



| Abstract views: 1 |  PDF views: 114

[PDF](#)

Preface Reinwardtia Vol. 20, No 2 (2021)



| Abstract views: 1 |  PDF views: 115

[PDF](#)

Articles

[THE CONFUSING TAXONOMY AND NOMENCLATURE OF SYZYGIUM CONFUSUM COMPLEX \(MYRTACEAE\)](#)



Pudji Widodo, Jan Frits Veldkamp
[doi> 10.14203/reinwardtia.v20i2.4160](#)



| Abstract views: 264 |  PDF views: 193

[PDF](#)

43-
49

[VANILLA YERSINIANA \(ORCHIDACEAE\), A NEW RECORD FOR PENINSULAR MALAYSIA](#)

 Muhammad Ikhwanuddin Mat Esa, Farah Alia Nordin, Rusea Go, Akmal Raffi

 [10.14203/reinwardtia.v20i2.4186](https://doi.org/10.14203/reinwardtia.v20i2.4186)

 | Abstract views: **234** |  PDF views: **214**

[PDF](#)

51-
55

[A NEW SPECIES OF DEPARIA FROM NEW GUINEA](#)

 Wita Wardani, Jaenudin Jaenudin, Ismail Apandi, Anne Kusumawaty, Wahyudi Santoso

 [10.14203/reinwardtia.v20i2.4231](https://doi.org/10.14203/reinwardtia.v20i2.4231)

 | Abstract views: **240** |  PDF views: **149**

[PDF](#)

57-
61

[ETLINGERA COMOSA, A NEW SPECIES \(ZINGIBERACEAE: ALPINIOIDEAE\) FROM CENTRAL SULAWESI](#)

 Marlina Ardiyani, Wisnu Handoyo Ardi, Prima Wahyu Kusuma Hutabarat, Axel Dalberg Poulsen

 [10.14203/reinwardtia.v20i2.4243](https://doi.org/10.14203/reinwardtia.v20i2.4243)

 | Abstract views: **353** |  PDF views: **205**

[PDF](#)

63-
68

[EVALUATING THE ECOPHYSIOLOGY OF SURVIVAL FOR MAPANIA CUSPIDATA \(MIQ.\) UTTIEN \(CYPERACEAE\) TRANSPLANTATION](#)

 Zinnirah Shabdin, Hollena Nori, Kalu Meekiong, Mohammad Fajaruddin Mohd Faiz

 [10.14203/reinwardtia.v20i2.4172](https://doi.org/10.14203/reinwardtia.v20i2.4172)

 | Abstract views: **200** |  PDF views: **150**

[PDF](#)

69-
75

Obituaries

Dr. Willem J.J.O. de Wilde and Dr. Ed de Vogel



 [10.14203/reinwardtia.v20i2.4289](https://doi.org/10.14203/reinwardtia.v20i2.4289)



| Abstract views: **157** |  PDF views: **131**

THE CONFUSING TAXONOMY AND NOMENCLATURE OF *SYZYGIUM CONFUSUM* COMPLEX (MYRTACEAE)

Received June 02, 2021; accepted August 02, 2021

PUDJI WIDODO

*Fakultas Biologi, Universitas Jenderal Soedirman, Jln. Dr. Soeparno 63, Purwokerto 53122, Indonesia
Biology Department, Institute Pertanian Bogor (IPB University), Jln. Raya Dramaga, Kampus IPB, Dramaga, Bogor 16680, Indonesia. Email: pwidodo@unsoed.ac.id*

JEFF F. VELDKAMP (†)

*Naturalis Biodiversity Center, Research Group of Tropical Botany, P.O. Box 9517, 2300 RA Leiden, the Netherlands.
Institute of Biology Leiden, Leiden University, P.O. Box 9505, 2300 RA Leiden, the Netherlands.*

ABSTRACT

WIDODO, P. & VELDKAMP, J. F. 2021. The confusing taxonomy and nomenclature of *Syzygium confusum* complex (Myrtaceae). *Reinwardtia* 20(2): 43 – 48. — The taxonomic and nomenclatural confusions surrounding the *Syzygium confusum* complex are elucidated. For that purpose, type specimens are designated and circumscriptions are presented for each species. Typifications, newly characterized descriptions and illustrations are presented for *Syzygium korthalsii* Widodo, *S. confusum* (Blume) Bakh f., *S. blumei* (Steudel) Merr. & L.M. Perry, *S. insigne* (Blume) Merr. & L.M. Perry. The new species *Syzygium sipirokense* Widodo & Veldkamp is described.

Key words: *Jambosa*, Malesia, Myrtaceae, Southeast Asia, *Syzygium*.

ABSTRAK

WIDODO, P. & VELDKAMP, J. F. 2020. Kekusutan taksonomi dan tata nama kompleks *Syzygium confusum* (Myrtaceae). *Reinwardtia* 20(2): 43 – 48. — Kekusutan taksonomi dan tata nama jenis seputar kompleks *Syzygium confusum* dicoba diuraikan dengan pemantapan penunjukan spesimen-spesimen tipenya. Selanjutnya perapihan batasan takson yang termasuk jenis kompleks juga telah dilakukan. Oleh karena itu pemantapan nama dan pertelaan serta ilustrasi *Syzygium korthalsii* Widodo, *S. confusum* (Blume) Bakh f., *S. blumei* (Steudel) Merr. & L.M. Perry, *S. insigne* (Blume) Merr. & L.M. Perry dan jenis baru *Syzygium sipirokense* Widodo & Veldkamp disajikan.

Kata kunci: Asia Tenggara, *Jambosa*, Malesia, Myrtaceae, *Syzygium*.

INTRODUCTION

In studying the Sumatran free petalled species of *Syzygium* one may have difficulty in identifying the species with narrow leaves, especially because some of their representatives are rare and hence poorly known (Backer & Bakhuizen van den Brink Jr., 1963:343). In April 1972, for the Flora Malesiana project, Bakhuizen van den Brink Jr. and van Steenis tentatively identified and annotated two specimens preserved in L (namely HLB no. 898.203-342 part of Herb *Blume* s.n. collected in Java without definite locality and HLB no. 898.203-344 collected around Bogor, Java by an unknown collector) as *Syzygium confusum* (Blume) Bakh.f. Another specimen (HLB no 898.203-345 collected in Mount Malintang, West Sumatra by Korthals) was tentatively identified by them as *Syzygium* cf. *confusum* (Blume) Bakh.f.

In 1846 Korthals had already identified his collection (HLB 898.203-345) as *Jambosa lanceolata* Korthals. Confusion arose when Blume (1849) proposed the name *Jambosa confusa* Blume for other material from Java and Sumatra, inferring that Korthals's name was superfluous, as *Jambosa lanceolaria* was an earlier name for Korthals's material, based on *Eugenia lanceolaria* Roxb.

(1832). To rectify this, Blume (1850) proposed *Jambosa korthalsii* Blume as a new name for *Jambosa lanceolata* Korthals. Although these specific epithets are similar, they do not mean exactly the same thing because *lanceolarius* (= small tip of a spear) and *lanceolatus* (= lancet-shaped) and they are not confusable under the ICNafp (Turland *et al.*, 2018). In this case, what is the nomenclatural status of Blume's proposed new name *Jambosa korthalsii* Blume? Is it a superfluous name? Korthals' specific epithet *lanceolatus* cannot be transferred to *Syzygium* because it is pre-empted by the combination *S. lanceolatus* (Lam.) Wight & Arn. (1834). It is clear, therefore, that there is a need to clarify this nomenclaturally confused situation.

In revising the taxonomy of the narrow leaved *Syzygium* in Sumatra (Widodo, 2011) we found that twigs, leaf shape, leaf base and apex more often than not offer valuable characters for delimiting species. Consequently, morphological variation in these characters in the *Syzygium confusum* complex will be given special attention.

During the course of this study, in BO we found specimens from Sumatra with characteristics very much like *Syzygium insigne* (Blume) Merr. & L.M. Perry and *S. blumei* (Steudel) Merr. & L.M.

Perry (species also related to the *Syzygium confusum* complex) but with consistently varying characteristics. We take this opportunity to describe these specimens and propose a species new to science.

MATERIALS AND METHODS

Materials used in this research are herbarium specimens from Sumatra, Java, and Borneo preserved in the herbaria of BO, L and K. Procedures and methods of observations used in this study mostly followed those elaborated by Rifai (1976), de Vogel (1987), Widodo (2011) and Widodo (2012).

RESULTS AND DISCUSSION

Results of our renewed observations of morphological characters of *Syzygium confusum* complex are presented in Table 1. We found that combinations of these characters are of assistance for delimiting closely related species as can be observed in Table 1.

1. *SYZYGIIUM KORTHALSII* Widodo. — Fig. 1. *Jambosa lanceolata* Korth. Ned. Kruidk. Arch. 1: 199. 1846. [non *Syzygium lanceolatum* (Lam.) Wight & Arn., 1834]. — *Jambosa korthalsii* Blume, Mus. Bot. Lugd.-Bat. 1: 101. 1849 [1850], nom. superfl. — *Syzygium korthalsii* Widodo, Reinw. 13(3): 235–240 (2012). — TYPE: INDONESIA, West Sumatra, Gunung Malintang, *Korthals s.n.* (Holotype: L!, HLB no. 898.203-345), designated by Widodo (2012).

Tree diameter unknown. Twigs usually 4-angled to 4-winged, with smooth and whitish pale brown bark. Leaves relatively long compared to width, the leaf form very narrowly ovate, 30–45 cm by 2.5–5 cm, brown above and milky brown below when dry; leaf base cordate, leaf apex long narrowly acuminate; petiole *ca.* 3 mm long, swollen and corky, drying pale brown; midrib channelled on the upper surface and raised on the lower surface, pale brown when dry; major lateral veins consists of *ca.* 25 pairs, 1–1.5 cm apart, at an angle of 60°–70°, sometimes curved near the midrib and straight near the intramarginal veins; minor lateral veins absent or present, oil dots between 2 major lateral veins less than 20 per cm²; intramarginal vein 1 or 2, faint, 1–3 mm from margin. Inflorescence a terminal cyme, but the flower with a pseudostipe 5–7 mm long, hypanthial cup funnel-shaped; sepals triangular, 5–6 mm long, 5 mm wide; petals unknown; style 35 mm long. Fruits unknown.

Distribution. *Syzygium korthalsii* is known from a limited area in West Sumatra, namely in Priaman and in Gunung Malintang.

Notes. *Syzygium korthalsii* can be readily distinguished from other Sumatran species by its leaf form which is very narrowly ovate and almost linear, reaching approximately 45 cm long and only around 3.5 cm wide on average.

2. *SYZYGIIUM CONFUSUM* (Blume) Bakh. f. — Fig. 2.

Jambosa confusa Blume, Mus. Bot. Lugd.-Bat. 1: 101. 1849 (non *J. confusa* Blume ex Miq., Anal. Bot. Ind. 1: 27. 1850, nom. inval., in syn. sub *E. microbotrya* Miq., non pert.). *Syzygium confusum* (Blume) Bakh. f. in Bakhuizen v/d Brink & Koster, Blumea 12: 61. 1963. — *Eugenia dolichophylla* Koord. & Valetton, Meded. Lands Plantentuin 40: 78. 1900, “*doligophylla*” non *Eugenia dolichophylla* Kiaersk., En. Myrt. Bras. 157. 1893, nec *Syzygium dolichophyllum* (Laut. & K. Schum.) Merr. & L.M. Perry, J. Arn. Arb. 23: 249. 1942. — *Eugenia malayana* Gagnep. in Lecomte, Fl. Indo-China 2: 838. 1921. *Syzygium malayanum* (Gagnep.) I.M. Turner, Gard. Bull. Singapore 47: 378. Jul 1997 (“Dec 1995”); Singapore Natl. Acad. Sci. 22–24: 21. Aug 1997 (“1996”), nom. superfl. — *Syzygium amshoffianum* Merr., Philipp. J. Sci. 79: 366. 1951 (“1950”), nom. superfl. — TYPE: INDONESIA, Java without definite locality. Herb. Blume *s.n.* (Holotype: L!, HLB no. 898.203-342), tentatively identified/annotated as *Syzygium confusum* (Blume) Bakh.f. in April 1972 by Bakhuizen van den Brink Jr. & van Steenis.

Tree to 8 m tall. Twigs terete and slightly compressed near the nodes. Leaves narrowly lanceolate, 20–30 cm by 3–5 cm tapered gradually from the middle to apex; upper surface blackish brown, lower surface reddish brown when dry; leaf base narrowly cuneate, apex long acute to acuminate; petiole 10–13 mm long, slender or swollen, scaly, peeling off; midrib rounded below, pale brown when dry; lateral veins very faint on both the upper and lower surfaces *ca.* 30 pairs, 1–2 cm apart, at an angle of 60°–70°, oil dots a few per cm²; intramarginal vein 1, very faint 1–2 mm from margin. Inflorescence simple or paniculate to 5 cm long, terminal, up to 21 flowers per inflorescence. Rachis terete and 4-angled, drying dark brown. Flowers with short ultimate inflorescence axis, pseudostipe and hypanthial cup 8–15 mm long, trumpet-shaped to turbinate. Sepals 4 free, semiorbicular, 3.5 mm long *ca.* 4 mm wide. Petals semiorbicular *ca.* 5.5 mm long and wide, a few gland dots. Stamens *ca.* 10 mm long. Style *ca.* 20 mm long. Ovary 2-locular. Fruits campanulate (immature).

Table 1. Morphological differences between species of *Syzygium confusum* complex

No	Characters	<i>S. korthalsii</i>	<i>S. confusum</i>	<i>S. blumei</i>	<i>S. insigne</i>	<i>S. sipirokense</i>
1	Twigs	4-angled to 4-winged	Terete and slightly compressed near nodes	terete	Terete and 4-angled near nodes	4-winged
2	Leaf form	Very narrowly ovate	Narrowly lanceolate	Quite narrowly ovate	Narrowly ovate	Quite narrowly ovate to almost oblong ovate
3	Leaf apex	Long narrowly acuminate	Acute to acuminate	Acute to acuminate	Acute	Acuminate to apiculate
4	Leaf size	30–45 cm by 2.5–5 cm	20–44 cm by 3–5 cm	15–20 cm by 2–3 cm	4–10 cm by 1–2.75 cm	10–15 cm by 3–5.5 cm
5	Leaf base	cordate	Almost narrowly cuneate	Rounded or subcordate	Subcordate or almost rounded	Rounded or subcordate
6	Inflorescens	Peduncle unknown	Peduncle unknown	Peduncle very short 2–5 mm or sessile	Peduncle unknown	Peduncle 4-angled, drying black
7	Locality	Sumatra, Mount Malintang	Java	Java	Borneo, Martapura	Aceh, North Sumatra

Distribution. Java. In Sumatra, *Syzygium confusum* is known only from Batam Island.

Notes. Koorders & Valetton (1900) realised that Blume's specific epithet *confusa* could not be combined with *Eugenia* because it was pre-empted by *E. confusa* DC. (1828), so that he proposed the new combination *E. doligophylla*. This, however, is an orthographic variant of the earlier *E. dolichophylla* Kiaersk. (1893) as can be seen when Koorders himself corrected it (1912). It is not a misprint as was suggested by Henderson (1949: 50), as the spelling is consequently used throughout in the 1900 paper. This combination is therefore also a later homonym and illegitimate.

Gagnepain (1921) proposed the new name *E. malayana* for this species (Govaerts, 2008), which Turner (1997a, b) used in *Syzygium*, overlooking the fact that *S. confusum* was required, and that this combination had already been made by Bakhuizen van den Brink Jr. & Koster (1963). Gagnepain's specimens (*Dussaud s.n.*, *Harmand 1314* and *Thorel s.n.*) and his description based on them actually refer to *Syzygium megacarpum* (Craib) Rathakr. & N.C. Nair (Wuu Kuang Soh, TCD, *in litt.*).

Unaware of Gagnepain's action Merrill (1951) proposed yet another name: *Syzygium amshoffianum*, which is superfluous.

3. *SYZYGIUM BLUMEI* (Steud.) Merr. & L.M. Perry. — Fig. 3.

Eugenia angustifolia Blume, *Flora* 7(1): 291 (1824), [nom. illeg., non *Eugenia angustifolia* Lam., *Encycl.* 3: 203 (1789)]. — *Myrtus hypericifolia* Blume, *Bijdr. Fl. Ned. Ind.*: 1082 (1826) [nom. illeg., non *Myrtus hypericifolia* Salisb., *Prodr. Stirp. Chap. Allerton*: 354 (1796)]. *Jambosa hypericifolia* (Blume) DC., *Prodr.* [A. P. de Candolle] 3: 287 (1828), nom. illeg. *Eugenia hypericifolia* (Blume) Koord. & Valetton, *Meded. Lands Plantentuin* 40, *Bijdr.* 6: 69 (1900) [nom. Illeg.] — *Eugenia blumei* Steudel, *Nomencl. Bot.* ed. 2. 1: 601 (1840). *Syzygium blumei* (Steudel) Merr. & L.M. Perry, *Mem. Amer. Acad. Arts* 18: 164 (1939). — TYPE: INDONESIA, Jawa, Bogor. (Holotype: L!. HLB no. 898.203-344). labelled as *Jambosa confusa*, and preidentified/annotated as *Syzygium cf. confusum* (Blume) Bakh. f. by Bakhuizen van den Brink f. & van Steenis in April 1972.

Habit unknown. Twigs terete, glabrous, drying yellowish. Leaves sessile, glabrous, narrowly ovate, 6–19 cm long by 1.5–3 cm wide; leaf base rounded to subcordate, leaf apex acute to acuminate; major lateral veins 6–9 pairs, very faint on both surfaces; leaves drying greyish above and yellowish below. Inflorescence terminal and in leaf axil. Pedicel terminally solitary, 1 flower

with shorter petals. Calyx 4 lobed, base attenuate. Fruit unknown.

Distribution. West Java, Mt. Salak.

Habitat & Ecology. Tropical rain forest.

4. *SYZYGIUM INSIGNE* (Blume) Merr. & L.M. Perry — Fig. 4.

Jambosa insignis Blume, Mus. Bot. Lugd.-Bat. 1: 100. 1849. *Syzygium insigne* (Blume) Merr. & L.M. Perry, Mem. Acad. Arts & Sci. 18: 163. 1939. Mem. Gray Herb. Harvard Univ. 4: 163. 1939; Masam., Enum. Phan. Born.: 530. 1942. — *Jambosa lancifolia* Miq., Anal. Ind. 1: 17. 1850; Fl. Ned. Ind. 1, 1: 427. 1855, nom. superfl. — TYPE: INDONESIA, Borneo, Martapoera. *Korthals s.n.* (Holotype: L!, HLB no. 898.203-347), accepted by Merrill (1921), Merrill & Perry (1939), and Masamune (1942).

Tree, height and diameter unknown. Twigs 4-angled. Leaves opposite, very shortly petiolate nearly two-ranked, narrowly ovate, leaf apex acute, base subcordate or almost rounded, 4–10 cm by 1–2.75 cm, transverse veins confluent in the inframarginal nerve, coriaceous, shiny above, and often impressed with inconspicuous dots, paler below. Racemes 1-3, terminal and solitary in the axils, short, few-flowered. Flowers showy, rather longly pedicellated, pink. Calyx about 2.5 cm, tube of the calyx turbinate, widened above, the limb with sort rounded flaps of 6-8-fid, lobes unequal, the outermost shorter, deciduous.

Distribution: Borneo.

Notes. Not to be confused with *Eugenia insignis* Thwaites from Sri Lanka, a “true” *Eugenia*. Miquel (1850: 17) did not mention *Eugenia insignis* Blume, but described *E. lancifolia* as new based on the same material, so that the name is superfluous. Though he realised it in 1855 (p. 427), he still retained his own epithet. Merrill (1921: 429) regarded *Jambosa insignis* and *J. lancifolia* as distinct species, repeating Blume’s suggestion that more than one collection would be involved. Merrill & Perry (1939: 163) joined the two, but did not mention *Jambosa lanceolata* at all, probably because they were aware that this was a Sumatran species, whereas they were dealing with Bornean material.

5. *Syzygium sipirokense* Widodo & Veldkamp spec. nov. — Fig. 5.

— TYPE: INDONESIA, Sumatra, Tapanuli Selatan, Cagar Alam Sipirok, Nagurguran. *EA Widjaja* 2012, 19 March 1983 (Holotype: BO!). Other specimens are Ketambe Research Station,

Aceh. *Kramadibrata* K 329, 333, 11 March 1982.

Small tree or shrub. Twigs winged near the nodes. Leaves sessile, opposite, lanceolate, 10–15 cm by 3–5.5 cm, leaves upper surface dark brown, lower surface brown when dry. Major lateral veins 10–14; leaf base rounded or cordate; leaf apex acuminate to apiculate; intramarginal veins one, 1–2 mm from margin, channelled above, raised below. Inflorescence arises from the leaf axil, peduncle four-angled, slightly winged, slender, black when dry. Flower unknown. Fruit ovoid to oval, 8–12 mm long, 5–7 mm diameter, green-red.

Distribution. Aceh Province, Aceh Tenggara Regency. North Sumatra, Tapanuli Selatan, Cagar Alam Sipirok, Nagurguran.

Habitat & Ecology. Primary forest 700 m alt.

Etymology. The epithet *sipirokense* came from one of the areas where this specimen was collected.

Conservation Status. This species is known from two locations, namely Sipirok Nature Reserve in North Sumatra and Ketambe Research Station in Aceh. The IUCN Assessment (IUCN, 2020) is categorized as Critically Endangered (CR).

Notes. *Syzygium sipirokense* resembles *S. blumei*. However, the leaves of *Syzygium sipirokense* dry dark brown above and pale brown below, instead of drying greyish above and yellowish below as in *S. blumei*. Twigs of *Syzygium sipirokense* are 4-winged, while the twigs of *S. blumei* are terete.

ACKNOWLEDGEMENTS

The first author has been supported by a grant from the Directorate General of Higher Education, the Republic of Indonesia. Acknowledgement is due to Prof. Dr. Alex Hartana and Prof. Dr. Tatik Chikmawati who have supervised the first author during his study and to Prof. Dr. Mien A. Rifai for improving the rewritten final manuscript. We would like to appreciate the Keepers of L, K, and BO for permission to observe the collections. Prof. Dr. Peter van Welzen is thanked for his assistance and service. Dr. Lyn Craven (†), CSIRO, Australia, is acknowledged for his advice. Prof. Dr. John Parnell, Trinity College Dublin is thanked for providing references. Dr. Rafael Govaerts, Royal Botanic Gardens Kew is appreciated for giving relevant information. Dr. Wu Kuang Soh, Trinity College Dublin kindly commented on Gagnepain’s new name. Grateful acknowledgement is due to Dr. Eve Lucas, Royal Botanic Garden Kew for her critical review of the manuscript.

Fig. 1. *Syzygium korthalsii*. Leafy twig.Fig. 2. *Syzygium confusum*. Leafy twig.

REFERENCES

- BAKHUIZEN VAN DEN BRINK f., R. C. & KOSTER, J. T. 1963. Notes on the Flora of Java VIII. Myrtaceae. *Blumea* 12: 61.
- BLUME, C. L. 1849. Ord. Myrtaceae. Subord. Myrteae. *Museum botanicum Lugduno-Batavum*. 1, 7: 100–101.
- DE CANDOLLE, A. P. 1828. *Prodromus Systematis Naturalis Regni Vegetabilis* 3: 279. Treutel & Würz, Paris.
- GAGNEPAIN, F. 1921. Myrtaceae. In: LECOMTE, P. H. *Flora of Indo-Chine* 2: 838. Masson & Cie, Paris.
- GOVAERTS, R., SOBRAL, M., ASHTON, P., BARRIE, F., HOLST, B. K., LANDRUM, L. L., MATSUMOTO, K., MAZINE, F. F., LUGHADHA, E. N., PROENÇA, C., SOARES-SILVA, L. H., WILSON, P. G., LUCAS, E. 2008. *World Checklist of Myrtaceae*. Kew Publishing. Royal Botanic Gardens, Kew.
- HENDERSON, M. R. 1949. The genus *Eugenia* (Myrtaceae) in Malaya. *Gardens' Bulletin Singapore* 12: 50–51.
- KOORDERS, S. H. 1912. *Exkursionsflora von Java* 2: 675. Fischer, Jena.
- KOORDERS, S. H. & VALETON, T. 1900. Bijdrage no. 6 tot de kennis der boomsoorten op Java. *Mededeelingen uit 's Lands Plantentuin* 40: 78–79.
- KORTHALS, P. W. 1846. *Bijdrage tot de kennis der Myrtaceae*. Nederlandsch kruidkundig archief. 1: 199.
- MASAMUNE, G. 1942. *Enumeratio phanerogamarum bornearum*: 530. Government of Taiwan, Taipei.
- MERRILL, E. D. 1921. A bibliographic enumeration of Bornean plants. *Journal of the Straits branch of the Royal Asiatic Society* Special Number: 428–429.
- MERRILL, E. D. 1951 (“1950”). Readjustments in the nomenclature of Philippine *Eugenia* species. *Philippine Journal of Science* 79: 366.



Figs. 3. *S. blumei* (A), *S. insigne* (B), *S. sipirokense* Widodo & Veldkamp spec. nov (C).

- MERRILL, E. D. & PERRY, L. M. 1939. The myrtaceous genus *Syzygium* Gaertner in Borneo. *Memorial Academic Arts & Science* 18: 163. ≡ *Memorial Gray Herbarium of Harvard University* 4: 163.
- MIQUEL, F. A. W. 1850. *Stirpes quaedam borneenses. Analecta Botanica Indica*. 1: 17, 27.
- MIQUEL, F. A. W. 1855. *Flora van Nederlandsch Indië* 1, 1: 427. C.G. van der Post, Amsterdam, C. van der Post Jr., Utrecht.
- RIFAI, M. A. 1976. *Sendi-Sendi Taksonomi Tumbuhan*. Herbarium Bogoriense, Bogor.
- TURLAND, N. J., WIERSEMA, J. H., BARRIE, F. R., GREUTER, W., HAWKSWORTH, D. L., HERENDEEN, P. S., KNAPP, S., KUSBER, W.-H., LI, D.-Z., MARHOLD, K., MAY, T. W., MCNEILL, J., MONRO, A. M., PRADO, J., PRICE, M. J. & SMITH, G. F. (Eds.). 2018. *International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017*. Regnum Vegetabile 159. Glashütten: Koeltz Botanical Books. DOI <https://doi.org/10.12705/Code.2018>
- TURNER, I. M. 1997a. (16 Jul 1997; "1995"). A catalogue of vascular plants of Malaya. *Gardens' Bulletin Singapore* 47: 378, 384.
- TURNER, I. M. 1997b. (Aug 1997; "1996"). What should the kelat trees of Malaya be called? *Journal of the Singapore National Academy of Science* 22–24: 21.
- VOGEL, E. DE (Editor). 1987. *Manual of Herbarium Taxonomy: Theory and Practice*. UNESCO Southeast Asia Regional Office, Jakarta.
- WIDODO, P. 2011. *Syzygium of Sumatra: The Free Petalled Species*. (Bessons Beatrice, Ed.) (1st ed., p. 119). Saarbrücken: LAP LAMBERT Academic Publishing GmbH & Co. KG Heinrich-Böcking-Str. 6-8 66121, Saarbrücken, Germany. Retrieved from www.lapublishing.com
- WIDODO, P. 2012. New nomenclature in *Syzygium* (Myrtaceae) from Indonesia and its vicinities. *Reinwardtia* 13 (3): 235–240.
- WIGHT, R. & WALKER-ARNOTT, G. A. 1834. *Prodromus Florae Peninsulae Indiae Orientalis* 1: 330. Parbury, Allen & Co.