



Banana Cultivars Microshoot Induction and Plantlet Formation using Cytokinin and Auxin

Sugiyono^{1*}, Prita Sari Dewi² and Rendie Prasetyo¹

¹Department of Botany, Faculty of Biology, Universitas Jenderal Soedirman, Purwokerto, Indonesia;

²Department of Agrotechnology, Faculty of Agriculture, Universitas Jenderal Soedirman, Purwokerto, Indonesia

*Corresponding author: sugiyono@unsoed.ac.id; gieks_sugiyono@hotmail.com

Abstract

Banana is a horticultural plant with very high potentials, which contains carbohydrates and vitamins that are useful in fulfilling people's food and nutritional needs. Hence, this study aims to produce superior banana seedlings and develop a protocol for their mass production using a plant *in vitro* culture technique. This was a two stage-experiment i.e. microshoot production and plantlet formation. The result showed that Gebyar cultivar produced more shoots than the Kepok Kuning cultivar, with an average of 4.25 microshoots explant⁻¹. However, Kepok Kuning produced more leaves than Gebyar, with an average of 4.64 leaves plantlet⁻¹. Banana shoots cultured on the media containing Indole-3-acetic acid (IAA) at a concentration of 2.5 μ M produced the highest leaves number. Meanwhile, those cultured on the media containing 1-Naphthalenesacetic acid (NAA) at a concentration of 7.5 μ M produced the highest roots number. A Murashige and Skoog (MS) medium supplemented with 6-Benzylaminopurine (BAP) up to 30 μ M and the one supplemented with 7.5 μ M of NAA are suitable for Kepok Kuning and Gebyar cultivars micropropagation with regard to microshoot induction and plantlet formation, respectively.

Keywords: BAP; Gebyar; *in vitro* culture; Kepok Kuning; NAA; sword sucker

Cite this as: Sugiyono, Dewi, P. S., & Prasetyo, R. (2021). Banana Cultivars Microshoot Induction and Plantlet Formation using Cytokinin and Auxin. *Caraka Tani: Journal of Sustainable Agriculture*, 36(2), 249-258. doi: <http://dx.doi.org/10.20961/carakatani.v36i2.50425>

INTRODUCTION

Banana is one of the horticultural plants with great potential due to its high nutritional value and being an income source for local farmers. This crop is also importantly cultivated by the people living in tropical countries and it is the world's fourth agricultural commodity after rice, wheat and maize (Satuhu and Supriyadi, 2007). According to Singh et al. (2014), almost all modern edible bananas originate from two wild diploid species, namely *Musa acuminata* designated as genotype AA and *Musa balbisiana* as BB. Some are found to have an AA or AB genotype but the vast majority is triploid. Many

domesticated bananas are proven to be triploid with a genotype of AAA, AAB or ABB. There are also seedless cultivated AA and AB diploid, AAAA, AAAB, AABB and ABBB tetraploids.

Indonesia's total banana production has been relatively stable in the last five years with an average of 7.202 million tons year⁻¹. This reached 7.280 million tons in 2019, with only 0.22% increase from 2015 (BPS - Statistics Indonesia, 2020). The Directorate of Horticultural Production Development and the Horticulture Research and Development Center prioritize banana as one of the fruits to be developed and studied. Production in the last 10 years has been influenced by plant pests and diseases

* Received for publication April 15, 2021

Accepted after corrections May 22, 2021

(Prasetyo et al., 2020), especially the outbreak of Panama disease (*Fusarium* wilt), Bacterium Blood disease caused by *Pseudomonas celebensis* and Moko disease (bacterial wilt) (Mondal et al., 2012; Ploetz, 2015). Furthermore, leaf wilt diseases such as Panama caused by the fungus *Fusarium oxysporum* Schlecht sp. *Cubans* and Moko caused by the bacterium *Raisotonia solanacearum* are the most dangerous to bananas. The two pathogens are soil-borne and easily transmitted, which enhance their spread to almost all banana plantations in Indonesia (Ploetz, 2015). They form chlamydospores in the soil which survive for a very long time (Ghag et al., 2015), therefore making the control of the diseases produced to be extremely difficult.

Most consumable bananas are triploid and sterile, making genetic improvement through conventional crossing difficult. Consequently, the integration of *in vitro* technique for genetic improvement is crucial. The provision of superior banana seedlings is an important strategy to increase production. High-quality seedlings can be developed using *in vitro* culture technique. The factors influencing this process include plant genetic composition and its expression, nutrition, physical growth factors and the utilization of Plant Growth Regulators (PGRs) and vitamins (Amoo and van Staden, 2013; Kadhimi et al., 2014).

The three banana cultivars used in this study were Ambon Nangka (*Musa acuminata* × *Musa balbisiana* AAB), Gebyar (*Musa acuminata* × *Musa balbisiana* AAB) and Kepok Kuning (*Musa acuminata* × *Musa balbisiana* ABB). They are widely cultivated in Indonesia, especially in Banyumas Regency, Central Java. Studies related to the microshoot induction of bananas with AAB and ABB genotypes were previously carried out by Hui et al. (2012), Kindimba and Msogoya (2014) and Yatim (2016). During the induction of axillary shoots growth, 6-Benzylaminopurine (BAP) and kinetin are commonly used due to being very active, long-lasting, easily translocated, stable and heat resistant. Meanwhile, cytokinins and auxins addition to the *in vitro* culture triggers cell division to form new cells, controls apical dominance and inhibits tissue aging (Schaller et al., 2014; Feng et al., 2017). The success of banana shoot multiplication using BAP has been reported by Ali et al. (2011); Vishnevetsky et al. (2011) and Govindaraju et al. (2012). In the plantlet formation phase, auxins such as

Indole-3-acetic acid (IAA), Indole-3-butyric acid (IBA) and 1-Naphthalenesacetic acid (NAA) are used. Banana shoots rooting in *in vitro* culture has been successfully carried out by adding auxins according to Muller and Leyser (2011); Müller et al. (2015) and Hossain et al. (2016).

This study aims to produce superior banana seedlings and develop a protocol for their mass production using *in vitro* plant culture technique, as well as to evaluate the influence of cytokinins and auxins types and concentrations on the three banana cultivars plantlet formation. The mass production of disease-free plantlets provides a means whereby plants including bananas (Daniells, 2007), are improved for sustainable cultivation (Viljoen et al., 2004).

MATERIALS AND METHOD

Plant material

The materials used in the microshoot induction phase were the sword suckers, i.e., vigorous shoots arising from the rootstock of banana plants, of three banana cultivars namely Gebyar, Ambon Nangka and Kepok Kuning. They were obtained from a local banana nursery in Purwokerto City, Banyumas Regency, Central Java, Indonesia. The use of sword sucker in banana micropropagation was also reported by Ali et al. (2011) and Govindaraju et al. (2012). During the plantlet formation phase, the Gebyar and Kepok Kuning cultivars microshoots obtained from the induction phase were used.

Explant preparation

Explant isolation was carried out by selecting banana sword suckers with an approximate size of 7 cm × 6 cm × 15 cm. The shoots were excised from the sword suckers and cut into 2 cm × 2 cm × 2 cm size and subsequently soaked in distilled water. Then, the explants were washed using sterile distilled water supplemented with few drops of Tween 20 to reduce surface tension. They were sterilized in a Laminar Air Flow cabinet (LAF) through immersion in 70% ethanol for 2 minutes, followed by double immersion in 0.2% HgCl₂ for five minutes. Sinha and Deka (2016) reported that HgCl₂ and Tween as effective chemicals for sterilizing Malbhog banana cultivar. Next, the explants were rinsed in sterile distilled water for 30 seconds and this was repeated three times. Each side of the sterilized explants was cut to remove residues and dead tissue which resulted in 1 cm × 1 cm × 1 cm explant

size as described by Govindaraju et al. (2012). Furthermore, they were inoculated and grown on a Murashige and Skoog (MS) medium supplemented with 15 μM BAP for axillary shoot induction, then the cultures were incubated at 24 °C with continuous light.

Microshoots induction and plantlet formation

MS basal medium (Sigma-Aldrich-M5519) supplemented with 20 g L⁻¹ sucrose, 2 μM IAA (Sigma-Aldrich-I2886), BAP and kinetin (Sigma-Aldrich-K0753) as treatments, solidified with 2.5 g L⁻¹ phytagel (Sigma-Aldrich-P8169), was used for microshoot induction. The treatments were arranged in a Split-Split Plot Design, with three replications. The main plot was banana cultivars consisting of Ambon Nangka, Gebyar and Kepok Kuning. The sub-plot was type of cytokinin consisting of BAP (Sigma-Aldrich-B3408) and kinetin (Sigma-Aldrich-K0753). The sub-sub-plot was concentration of cytokinin consisting of 15, 20, 25 and 30 μM . The cytokinin concentrations used are a modification of the results reported by Prayoga and Sugiyono (2010). The explants were sub-cultured into the same medium after six weeks of culturing.

For plantlet formation, microshoots were transferred to the root induction medium, namely an MS basal medium supplemented with 20 g L⁻¹ sucrose and solidified with 0.2% phytagel. The treatments were arranged in a Split-Split-Plot Design, with three replications. The main plot was banana cultivars which consisted of

Gebyar and Kepok Kuning, while the subplot was an auxin type consisting of IAA, IBA (Sigma-Aldrich-I5386) and NAA (Sigma-Aldrich-N0640). The sub-sub-plot was auxin concentration consisting of 2.5 μM , 5.0 μM , 7.5 μM and 10 μM . NAA and IBA had been reported as the best treatments for banana culture in rasthali cultivars (Govindaraju et al., 2012). The variables observed in this study were the formation of microshoots and plantlets. The parameters measured include the number of shoots, the number of leaves and the number of roots.

Data analysis

The data obtained were analyzed using an Analysis of Variance (ANOVA), followed by the Honestly Significant Difference (HSD) test with a confidence level of 95%.

RESULTS AND DISCUSSION

Microshoots induction

The *in vitro* culture of several banana cultivars has been successfully carried out, where explants were planted in the treatment medium (Figure 1). New shoots developed after six weeks of incubation (Figure 2) and few microshoots appeared after several sub-cultures (Figure 3). In microshoot induction, Gebyar and Kepok Kuning produced more microshoots than Ambon Nangka. Hence, Ambon Nangka did not proceed to the plantlet formation stage due to its low microshoot production.




Figure 1. The effect of different banana cultivars and cytokinin types and concentrations on microshoot formation. (A) The appearance of banana explant on treatment medium, (B) Gebyar microshoot development on MS medium supplemented with BAP

The small number of microshoots produced was probably due to the PGR concentration used (Bordoloi, 2016), the explants physiological

conditions such as apical dominance (Bhende and Kurien, 2015; Kebrom, 2017) and the culture period (Rahman et al., 2013). The cytokinin

concentration added to the media did not stimulate new shoots formation. According to Ahmed et al. (2014), the time required for explant shoot formation on a combined media (BAP and IAA) was slower than on the media containing only BAP or in combination with kinetin. This result is in agreement with the report by Buah et al. (2010) and Ashraf et al. (2014). BAP's better effect on shoot formation compared to kinetin may be attributed to its high stability within *in vitro* cultures, because it is not easily broken down and therefore persists in the medium. Possibly, the BAP amount conjugated in the medium was smaller than that of other growth regulators, leading to the presence of more BAP in their free or ionized forms, which were made readily available to plant tissues from the medium (Buah et al., 2010).

Furthermore, the explants in this study were taken at the dry season peak, hence it is suggested that they had high abscisic acid (ABA) content as a response to water stress. ABA is widely known as a stress hormone, and its level increases in the presence of stress, including water stress (Hu et al., 2016). The high concentration of endogenous ABA inhibits microshoots formation and growth (Feng et al., 2012). These conditions are thought to have influenced explants growth and microshoots development. To solve this problem, an extremely high cytokinin concentration have to be used to break the dormancy caused by ABA, which in turn stimulates microshoots development and growth (Goggin et al., 2015; Qiu et al., 2019). Nuraini et al. (2016) and Prasetyo et al. (2020) stated high cytokinin utilization breaks dormancy and initiates shoot growth.




Figure 2. The appearance of banana microshoot formation. (A) Axillary bud development in banana sword sucker explants. (B) Development of banana sword sucker with split open treatment in sub-culture media

The sword sucker explants used were grown intact and not split open, hence, the apical shoots might produce high endogenous auxin and inhibited lateral shoot growth. This inhibition of shoot growth by auxin was also reported by Kebrom (2017). Moreover, apical dominance breakage requires high exogenous cytokinin concentrations (Nuraini et al., 2016; Prasetyo et al., 2020). It is suspected that the exogenous cytokinin level added to the *in vitro* culture did not break the apical dominance as reported by Müller et al. (2015) and Ngomuo et al. (2014).

Plantlet formation

The shoot formation and root induction aimed to form plantlets that have both leaves and roots. The microshoots were separated individually from the original clump and then transferred to the treatment media. At the plantlet formation phase, shoot morphogenesis was influenced by banana cultivar (Table 1) and the interaction between the cultivar and auxin type used (Table 2). Gebyar cultivar produced more microshoots compared to Kepok Kuning and on average, it produced 4.25 axillary buds explant⁻¹. The difference in plants' genotypic

responses is observed in their ability to grow and regenerate. Each genotype has different abilities regarding axillary shoot growth, shoot number and rooting emergence (Basri, 2016).

Table 1. The average number of shoots formed

Types of cultivars	The number of axillary buds
Gebyar	4.25 ^a
Kepok Kuning	1.47 ^b

Note: Numbers followed by the same letters show a significant difference in DMRT (≤ 0.05)


Gebyar cultivars planted on a media containing NAA produced the greatest shoots number (5.7 shoots explant⁻¹), although this treatment was not significantly different from that of the media containing IBA (Table 2). PGR treatment is used to stimulate cell proliferation and differentiation through organogenesis or embryogenesis (Méndez-Hernández et al., 2019).

Not all cells in plant tissue respond to PGR, a cell only responds at a particular stage in the plant growth cycle (Tréhin et al., 1998; Schaller et al., 2014). Besides plant genotypes, explant's physiological conditions such as meristematic ability to grow and cells or tissues growth status determines bud regeneration's success. It is also related to cell metabolism, availability of endogenous PGR and the activity of genes controlling growth and development (Pillay and Tenkouano, 2011; Feng et al., 2012; Remakanthan et al., 2014).


Table 2. The average shoot number formed by two banana cultivars with different types of auxin

Type of cultivars	Type of auxin		
	NAA	IBA	IAA
Gebyar	5.75 ^a	4.58 ^a	2.42 ^b
Kepok Kuning	1.18 ^b	1.58 ^b	1.67 ^b

Note: Numbers followed by the same letters show a significant difference in DMRT (≤ 0.05)



A.



B.

Figure 3. The effect of auxin types and concentration with different banana cultivars on the number of shoots formation. (A) Kepok Kuning cultivar with an added 2.5 μM IAA. (B) Gebyar cultivar with an added 5.0 μM NAA

Leaf formation in the banana plantlet formation phase was controlled by the cultivar used and the interaction between the growth regulator type and PGR concentration used. Kepok Kuning produced more leaves than Gebyar, by producing 4.64 leaves plantlet⁻¹ on average (Table 3). IAA utilization in culture media showed a significantly different effect on the number of leaves formed. Banana shoots grown on the media supplemented with 2.5 μM IAA produced the greatest number of leaves, namely 5.5 leave plantlet⁻¹. Shoots number

is inversely related to the number of leaves formed. The greater the number of shoots produced, the smaller the number of leaves per shoot formed (Figure 4).

Table 3. The average number of leaves formed by two banana cultivars

Types of cultivars	Number of leaves
Gebyar	1.75 ^b
Kepok Kuning	4.64 ^a

Note: Numbers followed by the same letters show a significant difference in DMRT (≤ 0.05)




Figure 4. The effect of auxin types and concentration with different banana cultivars on leaf formation. (A) Kepok Kuning cultivar with added 2.5 μM IAA. (B) Gebyar cultivar with added 5.0 μM NAA

Auxin plays a very important role in leaf growth, by stimulating the development of prospective leaf meristem tissue (Xiong and Jiao, 2019). The leaves number is closely related to photosynthesis and plant metabolism, as well as nutrient absorption, while their increased growth is due to accelerated cell division and differentiation (Keller et al., 2011; Novak and Whitehouse, 2013). Also, their growth and development processes require PGR, such as auxin and cytokinin, as well as other nutrients contained in the growing media. Auxin and cytokinin work synergistically in stimulating plant growth, hence IAA which is a type of auxin, IAA functions in regulating cell division and stimulating growth to ensure an increase in leaves number (Scarpella et al., 2010; Skupa et al., 2014). Auxin addition affects leaf growth, especially the vascular tissue's length (Aloni, 2010; Zazimalova et al., 2010).

Leaves are essential plant organs, especially for photosynthesis, which facilitate organic materials production and optimum growth. The higher the number of leaves, length and width, the higher the leaf mass per unit area (LMA). LMA is an important leaf morphological trait affecting photosynthesis (Ren et al., 2019). The relationship between leaf number and individual size has metabolic and mechanical consequences that influence energy balances and carbon uptake at the whole plant level. In addition, it is also crucial to light interception and net carbon gain (Sun et al., 2019).

Root induction was carried out at the final stage of plant *in vitro* culture. When axillary shoots emerged, they were subsequently sub-cultured on rooting media. Banana root formation was influenced by the interaction between the type of auxins used and their concentration. Although the addition of the three auxin types, namely IBA, NAA and IAA, did not show a significantly different effect on roots number. Generally, NAA demonstrated quite better results and produced the highest roots number compared to the other two auxins (Table 4). Banana shoots grown on the media containing NAA at a concentration of 7.5 μM produced the highest roots number (13.3 roots plantlet⁻¹) (Figure 5).

Table 4. Interaction between auxin type and concentration on the plantlet formation

Type of auxin	Concentration of auxin (μM)	Number of leaves	Number of roots
IAA	2.5	5.50 ^a	4.50 ^{bc}
	5.0	2.17 ^{abc}	5.67 ^{bc}
	7.5	2.17 ^{abc}	3.83 ^{bc}
	10.0	3.83 ^{abc}	3.17 ^{bc}
	2.5	2.00 ^{abc}	4.67 ^{bc}
IBA	5.0	5.17 ^{ab}	7.00 ^{bc}
	7.5	3.33 ^{abc}	6.17 ^{bc}
	10.0	3.17 ^{abc}	4.17 ^{bc}
	2.5	1.50 ^{bc}	6.17 ^{bc}
NAA	5.0	1.17 ^c	2.33 ^c
	7.5	4.67 ^{abc}	13.33 ^a
	10.0	3.83 ^{abc}	9.67 ^{ab}

Note: Numbers followed by the same letters show a significant difference in DMRT (≤ 0.05)



Figure 5. The effect of auxin types and concentration with different banana cultivars on roots' formation after a 7.5 μ M NAA treatment

At low concentrations, IAA causes shoots and roots elongation, but at a higher concentration, it inhibits the elongation (Pamungkas, 2015; Lathifah and Dewi, 2016). According to Govindaraju et al. (2012) and Zhao (2014), NAA in low concentrations produced a higher roots number. Also, root's formation is related to the endogenous auxin and cytokinin contents in plant tissue. Therefore, lower auxin concentration is suggested to be used in inducing rooting on a nicely growing shoot.

CONCLUSIONS

Gebyar cultivar produced more shoots than Kepok Kuning, with an average of 4.25 microshoots explant⁻¹. However, Kepok Kuning produced more leaves than Gebyar, with an average of 4.64 leaves plantlet⁻¹. The media containing IAA with a concentration of 2.5 μ M produced the highest leaves number. Banana shoots that were cultured on a media containing NAA at a concentration of 7.5 μ M produced the highest roots number. Besides, good acclimatization to produce ready-to-plant seedlings and a study to optimize acclimatization conditions to provide a higher seedling survival rate are needed. The mass production of disease-free plantlets is expected to meet the increasing demand for banana seeds to increase the national production scale.

ACKNOWLEDGEMENT

The authors are grateful to the Rector of Jenderal Soedirman University and the Ministry of Research, Technology and Higher Education for the funding provided through Riset Unggulan Perguruan Tinggi (Grant No. 2193).

REFERENCES

- Ahmed, S., Sharma, A., Singh, A. K., Wali, V. K., & Kumari, P. (2014). *In vitro* multiplication of banana (*Musa* sp.) cv. Grand Naine. *African Journal of Biotechnology*, 13(27), 2696–2703. <https://doi.org/10.5897/ajb2014.13750>
- Ali, A., Sajid, A., Naveed, N. H., Majid, A., Saleem, A., Khan, U. A., Jafery, F. I., & Naz, S. (2011). Initiation, proliferation and development of micropropagation system for mass scale production of banana through meristem culture. *African Journal of Biotechnology*, 10(70), 15731–15738. Retrieved from <https://www.ajol.info/index.php/ajb/article/view/97536>
- Aloni, R. (2010). The induction of vascular tissues by auxin. In: Davies P.J. (eds) *Plant Hormones*. Dordrecht: Springer. https://doi.org/10.1007/978-1-4020-2686-7_22
- Amoo, S. O., & van Staden, J. (2013). Influence of plant growth regulators on shoot proliferation and secondary metabolite production in micropropagated *Huernia hystrix*. *Plant Cell, Tissue and Organ Culture*, 112, 249–256. <https://doi.org/10.1007/s11240-012-0230-x>
- Ashraf, M. F., Aziz, M. A., Kemat, N., & Ismail, I. (2014). Effect of cytokinin types, concentrations and their interactions on *in vitro* shoot regeneration of *Chlorophytum borivilianum* Sant. & Fernandez. *Electronic Journal of Biotechnology*, 17(6), 275–279. <https://doi.org/10.1016/j.ejbt.2014.08.004>
- Basri, A. H. (2016). Kajian pemanfaatan kultur jaringan dalam perbanyakan tanaman bebas virus. *Agrica Ekstensi*, 10(1), 64–73. Retrieved from <https://www.polbangtanmedan.ac.id/pdf/Jurnal%202016/Vol%2010%20No%201/08%20Arie.pdf>
- Bhende, S. S., & Kurien, S. (2015). Sucker production in banana. *Journal of Tropical Agriculture*, 53(2), 97–106. Retrieved from

- <http://jtropag.kau.in/index.php/ojs2/article/view/339>
- Bordoloi, N. D. (2016). Factors affecting *in vitro* shoot-tip culture of banana. *Acta Horticulturae*, 1113, 151–156. <https://doi.org/10.17660/ActaHortic.2016.1113.22>
- BPS - Statistics Indonesia. (2020). *SIMSPH Online, Survey Pertanian Hortikultura yang mencakup data produksi buah-buahan dan sayuran tahunan per kecamatan di seluruh wilayah Indonesia*. Badan Pusat Statistik. Retrieved from <https://www.bps.go.id/indikator/55/62/1/produksi-tanaman-buah-buahan.html>
- Buah, J. N., Danso, E., Taah, K. J., Abole, E. A., Bediako, E. A., Asiedu, J., & Baidoo, R. (2010). The effects of different concentrations cytokinins on the *in vitro* multiplication of plantain (*Musa* sp.). *Biotechnology*, 9(3), 343–347. <https://doi.org/10.3923/biotech.2010.343.347>
- Daniells, J. (2007). Global banana disease management - Getting serious with sustainability and food security. *Acta Horticulturae*, 828, 411–416. <https://doi.org/10.17660/ActaHortic.2009.828.43>
- Feng, J., Shi, Y., Yang, S., & Zuo, J. (2017). Cytokinins. In J. Li, C. Li, S.M. Smith (Ed.), *Hormone metabolism and signaling in plants*, pp. 77-106. Academic Press. <https://doi.org/10.1016/B978-0-12-811562-6.00003-7>
- Feng, J., Yuan, L., & Bao-zhong, H. (2012). Overview of plant shooting branch. *Journal of Northeast Agricultural University (English Edition)*, 19(2), 74–85. [https://doi.org/10.1016/s1006-8104\(13\)60042-2](https://doi.org/10.1016/s1006-8104(13)60042-2)
- Ghag, S. B., Shekhawat, U. K. S., & Ganapathi, T. R. (2015). Fusarium wilt of banana: biology, epidemiology and management. *International Journal of Pest Management*, 61(3), 250–263. <https://doi.org/10.1080/09670874.2015.1043972>
- Goggin, D. E., Emery, R. J. N., Kurepin, L. V., & Powles, S. B. (2015). A potential role for endogenous microflora in dormancy release, cytokinin metabolism and the response to fluridone in *Lolium rigidum* seeds. *Annals of Botany*, 115(2), 293–301. <https://doi.org/10.1093/aob/mcu231>
- Govindaraju, S., Saravanan, J., Jayanthi, B., Nancy, D., & Indra Arulselv, P. (2012). *In vitro* propagation of Banana (*Musa* sp.-Rasthali variety) from sword suckers for its commercial production. *Research in Plant Biology*, 2(5), 1–6. Retrieved from <https://updatepublishing.com/journal/index.php/ripb/article/view/2533>
- Hossain, M. A., Rubel, M. H., Nasiruddin, K. M., & Evamoni, F. Z. (2016). Influence of BAP and NAA on *in vitro* plantlet regeneration of local and exotic banana cultivars. *Journal of Bioscience and Agriculture Research*, 6(2), 553–564. <https://doi.org/10.18801/jbar.060216.66>
- Hu, B., Cao, J., Ge, K., & Li, L. (2016). The site of water stress governs the pattern of ABA synthesis and transport in peanut. *Scientific Reports*, 6, 1–11. <https://doi.org/10.1038/srep32143>
- Hui, A. V., Bhatt, A., & Keng, C. L. (2012). Micropropagation of *Musa acuminata* x *Musa balbisiana* cv. Pisang Awak (ABB genome) and three other cultivars. *Pakistan Journal of Botany*, 44(2), 777–780. Retrieved from [http://www.pakbs.org/pjbot/PDFs/44\(2\)/46.pdf](http://www.pakbs.org/pjbot/PDFs/44(2)/46.pdf)
- Kadhim, A. A., Alhasnawi, A. N., Mohamad, A., Yusoff, W. M. W., & Zain, C. R. C. M. (2014). Tissue culture and some of the factors affecting them and the micropropagation of strawberry. *Life Science Journal*, 11(8), 484–493. Retrieved from http://www.lifesciencesite.com/ljsj/life1108/063_23503life110814_484_493.pdf
- Kebrom, T. H. (2017). A growing stem inhibits bud outgrowth – The overlooked theory of apical dominance. *Frontiers in Plant Science*, 8, 1–7. <https://doi.org/10.3389/fpls.2017.01874>
- Keller, C. P., Grundstad, M. L., Evanoff, M. A., Keith, J. D., Lentz, D. S., Wagner, S. L., Culler, A. H., & Cohen, J. D. (2011). Auxin-induced leaf blade expansion in *Arabidopsis* requires both wounding and detachment. *Plant Signaling and Behavior*, 6(12), 1997–2007. <https://doi.org/10.4161/psb.6.12.18026>
- Kindimba, G., & Msogoya, T. (2014). Effect of benzylaminopurine on *in vivo* multiplication

- of French plantain (*Musa* spp. AAB) cv. 'Itoke sege.' *Journal of Applied Biosciences*, 74(1), 6086–6090. <https://doi.org/10.4314/jab.v74i1.1>
- Lathyfah, U., & Dewi, E. R. S. (2016). Pengaruh variasi konsentrasi Indole Acetic Acid (IAA) terhadap pertumbuhan tunas pisang barangan (*Musa acuminata* L. triploid AAA.) dalam kultur *in vitro*. *Bioma*, 5(1), 32–42. <https://doi.org/10.26877/bioma.v5i1.1492>
- Méndez-Hernández, H. A., Ledezma-Rodríguez, M., Avilez-Montalvo, R. N., Juárez-Gómez, Y. L., Skeete, A., Avilez-Montalvo, J., De-La-Peña, C., & Loyola-Vargas, V. M. (2019). Signaling overview of plant somatic embryogenesis. *Frontiers in Plant Science*, 10, 1–15. <https://doi.org/10.3389/fpls.2019.00077>
- Müller, D., & Leyser, O. (2011). Auxin, cytokinin and the control of shoot branching. *Annals of Botany*, 107(7), 1203–1212 <https://doi.org/10.1093/aob/mcr069>
- Mondal, B., Ray, S. K., Misra, D. K., & Khatua, D. C. (2012). Bacterial wilt of banana in West Bengal, India. *International Journal of Plant Protection*, 5(2), 227–231. Retrieved from [http://researchjournal.co.in/online/IJPP/IJPP%205\(2\)/5_A-227-231.pdf](http://researchjournal.co.in/online/IJPP/IJPP%205(2)/5_A-227-231.pdf)
- Müller, D., Waldie, T., Miyawaki, K., To, J. P. C., Melnyk, C. W., Kieber, J. J., Kakimoto, T., & Leyser, O. (2015). Cytokinin is required for escape but not release from auxin mediated apical dominance. *Plant Journal*, 82(5), 874–886. <https://doi.org/10.1111/tpj.12862>
- Ngomuo, M., Mneney, E., & Ndakidemi, P. (2014). The effect of bud splitting on suppression of apical dominance and inducing multiple buds development in banana shoot tip cultures of cv. 'Yangambi' (AAA) in Tanzania. *American Journal of Experimental Agriculture*, 4(12), 1853–1860. <https://doi.org/10.9734/ajea/2014/7290>
- Novak, S. D., & Whitehouse, G. A. (2013). Auxin regulates first leaf development and promotes the formation of protocorm trichomes and rhizome-like structures in developing seedlings of *Spathoglottis plicata* (Orchidaceae). *AoB PLANTS*, 5, 1–12. <https://doi.org/10.1093/aobpla/pls053>
- Nuraini, A., Sumadi, S., & Pratama, R. (2016). Aplikasi sitokinin untuk pematangan dormansi benih kentang G1 (*Solanum tuberosum* L.). *Kultivasi*, 15(3), 202–207. <https://doi.org/10.24198/kultivasi.v15i3.11765>
- Pamungkas, S. S. T. (2015). Pengaruh konsentrasi NAA dan BAP terhadap pertumbuhan tunas eksplan tanaman pisang cavendish (*Musa paradisiaca* L.) melalui kultur *in vitro*. *Gontor AGROTECH Science Journal*, 2(1), 31–45. <https://doi.org/10.21111/agrotech.v2i1.295>
- Pillay, M., & Tenkouano, A. (2011). *Banana breeding progress and challenges*. CRC Press.
- Ploetz, R. C. (2015). Management of Fusarium wilt of banana: A review with special reference to tropical race 4. *Crop Protection*, 73, 7–15. <https://doi.org/10.1016/j.cropro.2015.01.007>
- Prasetyo, R., Sugiyono, & Prayoga, L. (2020). Induksi tunas mikro pisang kultivar ambon nangka (*Musa* sp.) secara *in vitro*. *Vigor: Jurnal Ilmu Pertanian Dan Subtropika*, 5(2), 45–50. Retrieved from <https://jurnal.untidar.ac.id/index.php/vigor/article/view/3044>
- Prayoga, L., & Sugiyono, S. (2010). Uji perbedaan media dan konsentrasi BAP terhadap pertumbuhan tunas pisang raja secara kultur *in vitro*. *Agritech*, 12(2), 89–99. Retrieved from <http://jurnalnasional.ump.ac.id/index.php/AGRITECH/article/view/990>
- Qiu, Y., Guan, S. C., Wen, C., Li, P., Gao, Z., & Chen, X. (2019). Auxin and cytokinin coordinate the dormancy and outgrowth of axillary bud in strawberry runner. *BMC Plant Biology*, 19(1), 1–16. <https://doi.org/10.1186/s12870-019-2151-x>
- Rahman, S., Biswas, N., Hassan, M. M., & Ahmed, M. G., Mamun, A. N. K., Islam, M. R., Moniruzzaman, M., & Haque, M. E. (2013). Micro propagation of banana (*Musa* sp.) cv. Agnishwar by *in vitro* shoot tip culture. *International Research Journal of Biotechnology*, 4(4), 83–88. Retrieved from <https://www.interestjournals.org/articles/micro-propagation-of-banana-musa-sp-cv-agnishwar-by-in-vitro-shoot-tip-culture.pdf>
- Remakanthan, A., Menon, T. G., & Soniya, E. V. (2014). Somatic embryogenesis in banana (*Musa acuminata* AAA cv. Grand Naine): Effect of explant and culture conditions. *In vitro Cellular and Developmental Biology* -

- Plant*, 50(1), 127–136. <https://doi.org/10.1007/s11627-013-9546-4>
- Ren, T., Weraduwage, S. M., & Sharkey, T. D. (2019). Prospects for enhancing leaf photosynthetic capacity by manipulating mesophyll cell morphology. *Journal of Experimental Botany*, 70(4), 1153–1165. <https://doi.org/10.1093/jxb/ery448>
- Satuhu, S., & Supriyadi, A. (2007). *Pisang: budi daya, pengolahan, dan prospek pasar*. Jakarta: Penebar Swadaya.
- Scarpella, E., Barkoulas, M., & Tsiantis, M. (2010). Control of leaf and vein development by auxin. *Cold Spring Harbor Perspectives in Biology*, 2(1), a001511. <https://doi.org/10.1101/cshperspect.a001511>
- Schaller, G. E., Street, I. H., & Kieber, J. J. (2014). Cytokinin and the cell cycle. *Current Opinion in Plant Biology*, 21, 7–15. <https://doi.org/10.1016/j.pbi.2014.05.015>
- Singh, W. R., Singh, S. S., & Karuna, S. (2014). Analysis of banana genome groups of wild and cultivated cultivars of Manipur, India using sScore card method. *Advances in Applied Science Research*, 5(1), 35–38. Retrieved from <https://www.imedpub.com/articles/analysis-of-banana-genome-groups-of-wild-and-cultivate-d-cultivars-ofmanipur-india-using-sscore-card-method.pdf>
- Sinha, S. K., & Deka, A. C. (2016). Effect of osmotic stress on *in vitro* propagation of *Musa* sp. (Malbhog variety). *African Journal of Biotechnology*, 15(12), 465–471. <https://doi.org/10.5897/ajb2015.14446>
- Skupa, P., Opatrny, Z., & Petrasek, J. (2014). Auxin Biology: Applications and the Mechanisms Behind. In Nick, P., Opatrny Z. (Ed.) *Applied Plant Cell Biology: Cellular Tools and Approaches for Plant Biotechnology*. Berlin, Heidelberg: Springer. https://doi.org/10.1007/978-3-642-41787-0_3
- Sun, J., Wang, M., Lyu, M., Niklas, K. J., Zhong, Q., Li, M., & Cheng, D. (2019). Stem and leaf growth rates define the leaf size vs. number trade-off. *AoB PLANTS*, 11(6), plz063. <https://doi.org/10.1093/aobpla/plz063>
- Tréhin, C., Planchais, S., Glab, N., Perennes, C., Tregear, J., & Bergounioux, C. (1998). Cell cycle regulation by plant growth regulators: Involvement of auxin and cytokinin in the re-entry of *Petunia* protoplasts into the cell cycle. *Planta*, 206(2), 215–224. <https://doi.org/10.1007/s004250050393>
- Viljoen, A., Kunert, K., Kiggundu, A., Escalant, J. V., & Bornman, C.H. (2004). Biotechnology for sustainable banana and plantain production in Africa: The South African contribution. *South African Journal of Botany*, 70(1), 67–74. [https://doi.org/10.1016/S0254-6299\(15\)30308-2](https://doi.org/10.1016/S0254-6299(15)30308-2)
- Vishnevetsky, J., White, T. L., Palmateer, A. J., Flaishman, M., Cohen, Y., Elad, Y., Velcheva, M., Hanania, U., Sahar, N., Dgani, O., & Perl, A. (2011). Improved tolerance toward fungal diseases in transgenic Cavendish banana (*Musa* spp. AAA group) cv. Grand Nain. *Transgenic Research*, 20, 61–72. <https://doi.org/10.1007/s11248-010-9392-7>
- Xiong, Y., & Jiao, Y. (2019). The diverse roles of auxin in regulating leaf development. *Plants*, 8(7), 1–14. <https://doi.org/10.3390/plants8070243>
- Yatim, H. (2016). Multiplication of Raja Bulu banana (*Musa paradisiaca* L. AAB group) on several Benzyl Amino Purine (BAP) concentration by using *in vitro* method. *Agroekoteknologi*, 4(3), 1989–1995. Retrieved from <https://www.neliti.com/publications/107456/multiplication-of-raja-bulu-banana-musa-paradisiaca-l-aab-group-on-several-benzy>
- Zazimalova, E., Murphy, A. S., Yang, H., Hoyerova, K., & Hosek, P. (2010). Auxin transporters-why so many? *Cold Spring Harbor Laboratory Press*, 2(3), a001552. <https://dx.doi.org/10.1101/cshperspect.a001552>
- Zhao, Y. (2014). Auxin biosynthesis. *The Arabidopsis Book*, 12(12), e0173. <https://doi.org/10.1199/tab.0173>