

Current Issue



Vol. 23 No. 6 (2022)

[View All Issues](#)

Online biodiversitas.mipa.uns.ac.id, smujo.id/biodiv

ISSN: 1412-033X, E-ISSN: 2085-4722

Publisher: Society for Indonesian Biodiversity

Co-publisher: Department of Biology, FMNS, Universitas Sebelas Maret Surakarta

First Publication: 2000


Period of issuance: Starting on January 1, 2019, Biodiversitas issued monthly

Aims and Scope *Biodiversitas, Journal of Biological Diversity* or abbreviated as *Biodiversitas* encourages submission of manuscripts dealing with all biodiversity aspects of plants, animals and microbes at the level of the gene, species, and ecosystem as well as ethnobiology.

Article types The journal seeks original full-length research papers, reviews, and short communication.

Article types The journal seeks original full-length research papers, reviews, and short communication. Manuscript of original research should be written in no more than 8,000 words (including tables and figures), or proportional with articles in this publication number. Review articles will be accommodated, while, short communication should be written at least 2,000 words, except for pre-study.

Indexing The journal has been indexed/registered in SCOPUS, DOAJ, Google Scholar, Crossref, EBSCO, Microsoft Academic Search, etc.



Information

[For Readers](#)

[For Authors](#)

[For Librarians](#)

Journals List

[Biodiversitas Journal of Biological Diversity](#)

[Nusantara Bioscience](#)

[Prosiding Seminar Nasional Masyarakat Biodiversitas Indonesia](#)

[Asian Journal of Agriculture](#)

[Asian Journal of Ethnobiology](#)

[Asian Journal of Forestry](#)

[Asian Journal of Natural Product Biochemistry](#)

[Asian Journal of Tropical Biotechnology](#)

[International Journal of Bonorowo Wetlands](#)

[Cell Biology and Development](#)

[Indo-Pacific Journal of Ocean Life](#)

[International Journal of Tropical Drylands](#)

[International Journal of Tropical Drylands](#)

Reviewers List

[Reviewers](#)

Visitor Statistics

Statistics



Editorial Team

EDITOR-IN-CHIEF:

Sutarno

EDITORIAL MEMBERS:

English Editors: Graham Eagleton (grahameagleton@gmail.com)

English Editors: Suranto (surantouns@gmail.com)

Technical Editors: Solichatun (solichatun_s@yahoo.com)

Technical Editors: Artini Pangastuti (pangastuti_tutut@yahoo.co.id)

Distribution & Marketing: Rita Rakhmawati (oktia@yahoo.com)

Webmaster: Ari Pitoyo (aripitoyo@yahoo.co.id)

MANAGING EDITOR:

Ahmad Dwi Setyawan (unsjournals@gmail.com)

EDITORIAL BOARD:

Abd Fattah N. Abd Rabou, Islamic University of Gaza, Palestine

Agnieszka B. Najda, University of Life Sciences in Lublin, Lublin, Poland

Ajay Kumar Gautam, Abhilashi University Mandi, Himachal Pradesh, India

Alan J. Lymbery, Murdoch University, Perth, Australia

Bambang Hero Saharjo, Institut Pertanian Bogor, Bogor, Indonesia

- Daiane H. Nunes, State University of Londrina, Londrina, Brazil
- Darlina Md. Naim, University Sains Malaysia, Penang, Malaysia
- Ghulam Hassan Dar, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, Srinagar, India
- Hassan Pourbabaee, University of Guilan, Somehsara, Guilan, Iran
- Joko Ridho Witono, Center for Plant Conservation-Bogor Botanic Gardens, Indonesian Institute of Sciences, Bogor, Indonesia
- Kartika Dewi, Research Center for Biology, Indonesian Institute of Sciences, Cibinong, Bogor, Indonesia
- Katsuhiko Kondo, University of Missouri, Columbia, USA
- Kusumadewi Sri Yulita, Research Center for Biology, Indonesian Institute of Sciences, Cibinong, Bogor, Indonesia
- Livia Wanntorp, Naturhistoriska riksmuseet, Stockholm, Sweden
- M. Jayakara Bhandary, Government Arts and Science College, Karwar, Karnataka, India
- Mahdi Reyahi-Khoram, Islamic Azad University (Hamadan Branch), Hamadan, Iran
- Mahendra Kumar Rai, SGB Amravati University, Maharashtra, India
- Mahesh K. Adhikari, Adhikari Niwas, Kathmandu, Nepal
- Maria Panitsa, University of Patras, Agrinio, Greece
- Mochamad A. Soendjoto, Lambung Mangkurat University, Banjarbaru, Indonesia
- Mohamed M.M. Najim, University of Kelaniya, Kelaniya, Sri Lanka
- Mohib A. Shah, Nepean Telehealth Technology Centre, Sydney, Australia
- Nurhasanah, Universitas Mulawarman, Samarinda, Indonesia
- Praptiwi, Research Center for Biology, Indonesian Institute of Sciences, Cibinong, Bogor, Indonesia
- Rasool B.Tareen, University of Balochistan, Quetta, Pakistan
- Seyed Aliakbar Hedayati, Gorgan University of Agricultural Sciences and Natural Resources, Iran
- Seyed Mehdi Talebi, Arak University, Iran
- Shahabuddin, Universitas Tadulako, Palu, Indonesia
- Shahir Shamsir, Universiti Teknologi Malaysia, Skudai, Malaysia
- Shri Kant Tripathi, Mizoram University, Aizwal, India
- Sugeng Budiharta, Purwodadi Botanic Gardens, Indonesian Institute of Sciences, Pasuruan, Indonesia
- Subash C. Santra, University of Kalyani, India

Information

For Readers

For Authors

For Librarians

Journals List

Biodiversitas Journal of Biological Diversity

Nusantara Bioscience

Prosiding Seminar Nasional Masyarakat Biodiversitas Indonesia

Asian Journal of Agriculture

Asian Journal of Ethnobiology

Asian Journal of Forestry

Asian Journal of Natural Product Biochemist

Asian Journal of Tropical Biotechnology

International Journal of Bonorowo Wetlands

Cell Biology and Development

Indo-Pacific Journal of Ocean Life

International Journal of Tropical Drylands

Reviewers List

Reviewers

Visitor Statistics

Statistics



About the Society

Society for Indonesian Biodiversity (SIB) or Masyarakat Biodiversitas Indonesia (MBI) is an organization that engaged in the sphere of research, conservation and use of biodiversity in a sustainable manner at the level of genes, species, ecosystems, and cultural utilization (ethnobiology). SIB was founded in Surakarta, Indonesia on July 17, 2006. The nature of SIB membership is valid for lifetime. Prospective members must approve SIB Bylaw (AD/ART MBI) and pay a membership fee of IDR 200,000 (USD 15) per year. Full payment for the 5, 10 or 20 years (lifetime) get a rebate of 25%. Members get a sample of the journal "Biodiversitas" and "Nusantara Bioscience" print edition and/or online and get privileges and rebates for activities organized by the central Board of SIB.

[Download SIB Bylaw \(AD/ART MBI\) here.](#)

The following is a link SIB membership registration form.

[Download membership application form here.](#)

Submit a membership application form and photograph by email to the following address:
unsjournals@gmail.com

Or may also register online at the following link: <http://goo.gl/forms/Pb7B1vJy0P>

Information

[For Readers](#)

[For Authors](#)

[For Librarians](#)

Journals List

[Biodiversitas Journal of Biological Diversity](#)


[Nusantara Bioscience](#)

[Prosiding Seminar Nasional Masyarakat Biodiversitas Indonesia](#)

[Asian Journal of Agriculture](#)

[Asian Journal of Ethnobiology](#)

[Asian Journal of Forestry](#)



Vol. 22 No. 6 (2021)

Full Issue

[Front Cover](#)

Articles

Phenotypic characters and identification CYPs (Cyclophilin) gene in Cucumis melo L. cv. Gama Melon Parfum

WIKO ARIF WIBOWO, SIGIT DWI MARYANTO, BUDI SETIADI DARYONO

[PDF](#)

Rhizosphere Streptomycetes formulas as the biological control agent of

Information

[For Readers](#)

[For Authors](#)

[For Librarians](#)

Journals List

[Biodiversitas Journal of Biological Diversity](#)

[Nusantara Bioscience](#)

[Prosiding Seminar Nasional Masyarakat Biodiversitas Indonesia](#)

[Asian Journal of Agriculture](#)

[Asian Journal of Ethnobiology](#)

[Asian Journal of Forestry](#)

[Asian Journal of Natural Product Biochemistry](#)

[Asian Journal of Tropical Biotechnology](#)

[International Journal of Bonorowo Wetlands](#)

[Cell Biology and Development](#)

[Indo-Pacific Journal of Ocean Life](#)

[International Journal of Tropical Drylands](#)

Assessment of factual measurement times for chlorophyll-a fluorescence in rubber (*Hevea brasiliensis*) clones

ANDI NUR CAHYO, RUDI HARI MURTI, EKA T.S. PUTRA, TRI RINI NURINGTYAS, DENIS FABRE, PASCAL MONTORO

PDF

Antiplasmodial activity, biosynthetic gene clusters diversity, and secondary metabolite constituent of selected Indonesian *Streptomyces*

EMA DAMAYANTI, PUSPITA LISDIYANTI, ANDINI SUNDOWO, SHANTI RATNAKOMALA, ACHMAD DINOTO, JAKA WIDADA, MUSTOFA

PDF

Genetic diversity of *Pleurotus ostreatus* (Jacq.) P. Kumm. strains in Java based on random amplified polymorphic DNA markers

NURAENI EKOWATI, ARIS MUMPUNI, JUNI SAFITRI MULJOWATI, NUNIEK INA RATNANINGTYAS, ARDHINI RIN MAHARNING

PDF

Genetic diversity in several genotypes of Algerian lentil using biochemical markers

HAMIDA HAMMADI, DOUNIA HAMMOUDA-BOUSBIA, GHANIA CHAIB, ANFEL TIR

PDF

Diversity and potential contribution of wild edible plants to sustainable food security in North Wollo, Ethiopia


AHMED HASSEN

PDF

A case study of agroforestry practices and challenges in Mt. Merapi risk and hazard prone area of Indonesia

ZUHUD ROZAKI, NUR RAHMAWATI, OKI WIJAYA, ATIKAH FAJRIYAH MUBAROK, MASATERU SENGE, MOHD FAUZI KAMARUDIN

Genetic diversity of *Pleurotus ostreatus* (Jacq.) P. Kumm. strains in Java based on random amplified polymorphic DNA markers



PDF

DOI

<https://doi.org/10.13057/biodiv/d220658>

Issue

Vol. 22 No. 6 (2021)

Section

Section

Articles

NURAENI EKOWATI ♥

Mycology Laboratory, Faculty of Biology, Universitas Jenderal Soedirman. Jl. Dr. Soeparno 63, Purwokerto, Banyumas 53122, Central Java, Indonesia

<https://orcid.org/0000-0002-5927-0967>

ARIS MUMPUNI

Mycology Laboratory, Faculty of Biology, Universitas Jenderal Soedirman. Jl. Dr. Soeparno 63, Purwokerto, Banyumas 53122, Central Java, Indonesia

JUNI SAFITRI MULJOWATI

Mycology Laboratory, Faculty of Biology, Universitas Jenderal Soedirman. Jl. Dr. Soeparno 63, Purwokerto, Banyumas 53122, Central Java, Indonesia

NUNIEK INA RATNANINGTYAS

Mycology Laboratory, Faculty of Biology, Universitas Jenderal Soedirman. Jl. Dr. Soeparno 63, Purwokerto, Banyumas 53122, Central Java, Indonesia

ARDHINI RIN MAHARNING

Ecology Laboratory Faculty of Biology, Universitas Jenderal Soedirman. Jl. Dr. Soeparno 63, Purwokerto, Banyumas 53122, Central Java, Indonesia

Abstract

Abstract. Ekowati N, Mumpuni A, Muljowati JS, Ratnaningtyas NI, Maharning AR. 2021. Genetic diversity of *Pleurotus ostreatus* (Jacq.) P. Kumm. strains in Java based on Random Amplified Polymorphic DNA markers. *Biodiversitas* 22: 3488-3493. Genetic variation in a fungal population can occur due to mutation and recombination, resulting in changes in the nucleotides that encode specific DNA sequences. Strains with a high genetic distance and good production capabilities can be used to develop genetic breeding. This study aimed to investigate genetic relationship among *Pleurotus ostreatus* strains cultivated in Java (Bogor, Cianjur, Tasikmalaya, Purwokerto, Yogyakarta, Tawangmangu, Malang, and Madiun) based on random amplified polymorphic DNA (RAPD) markers. The research method consisted of DNA isolation and DNA amplification using six primers, i.e. OPA2, OPA3, OPA4, OPA7, OPA9, and OPA10. DNA band data were analyzed using NTSYSpc21 software to determine the level of genetic similarity, based on the Unweighted Pair Group Method with Arithmetic Average Algorithm (UPGMA). In all, 101 amplified DNA bands were obtained, with sizes ranging from 136 to 2320 bp and 96.0% of the bands were polymorphic. Based on cluster analysis, it shows that three clusters were formed. There were genetic variations and relationships among eight *P. ostreatus* strains in Java with a genetic similarity varying from 37-98%.

Information

For Readers

For Authors

For Librarians

Journals List

Biodiversitas Journal of Biological Diversity

Nusantara Bioscience

Prosiding Seminar Nasional Masyarakat Biodiversitas Indonesia

Asian Journal of Agriculture

Asian Journal of Ethnobiology

Asian Journal of Forestry

Asian Journal of Natural Product Biochemistry

Asian Journal of Tropical Biotechnology

International Journal of Bonorowo Wetlands

Cell Biology and Development

Indo-Pacific Journal of Ocean Life

International Journal of Tropical Drylands

Reviewers List

Reviewers

Reviewers

Visitor Statistics

Statistics


Visitors

 535,724	 15,685
 50,994	 11,750
 37,827	 9,609
 35,949	 8,421
 17,006	 7,459

FLAG counter

Most read articles by the same author(s)

- ARIS MUMPUNI, NURAENI EKOWATI, DANIEL JOKO WAHYONO, The existence of coprophilous macrofungi in Banyumas, Central Java, Indonesia, *Biodiversitas Journal of Biological Diversity*: Vol. 21 No. 1 (2020)
- NURAENI EKOWATI, ARIS MUMPUNI, NUNIEK INA RATNANINGTYAS, ARDHINI RIN MAHARNING, Compounds detection and inhibition activity of chloroform and ethyl acetate extracts of *Schizophyllum commune* on some cancer cell types, *Biodiversitas Journal of Biological Diversity*: Vol. 21 No. 12 (2020)
- ARIS MUMPUNI, ADI AMURWANTO, DANIEL JOKO WAHYONO, Molecular identification of coprophilous microfungi from Banyumas District, Central Java, Indonesia, *Biodiversitas Journal of Biological Diversity*: Vol. 22 No. 3 (2021)



Assessment of factual measurement times for chlorophyll-a fluorescence in rubber (*Hevea brasiliensis*) clones

ANDI NUR CAHYO, RUDI HARI MURTI, EKA T.S. PUTRA, TRI RINI NURINGTYAS, DENIS FABRE, PASCAL MONTORO

PDF

Antiplasmodial activity, biosynthetic gene clusters diversity, and secondary metabolite constituent of selected Indonesian *Streptomyces*

EMA DAMAYANTI, PUSPITA LISDIYANTI, ANDINI SUNDOWO, SHANTI RATNAKOMALA, ACHMAD DINOTO, JAKA WIDADA, MUSTOFA

PDF

Genetic diversity of *Pleurotus ostreatus* (Jacq.) P. Kumm. strains in Java based on random amplified polymorphic DNA markers

NURAENI EKOWATI, ARIS MUMPUNI, JUNI SAFITRI MULJOWATI, NUNIEK INA RATNANINGTYAS, ARDHINI RIN MAHARNING

PDF

Genetic diversity in several genotypes of Algerian lentil using biochemical markers

HAMIDA HAMMADI, DOUNIA HAMMOUDA-BOUSBIA, GHANIA CHAIB, ANFEL TIR

PDF

Diversity and potential contribution of wild edible plants to sustainable food security in North Wollo, Ethiopia

AHMED HASSEN

PDF

A case study of agroforestry practices and challenges in Mt. Merapi risk and hazard prone area of Indonesia

ZUHUD ROZAKI, NUR RAHMAWATI, OKI WIJAYA, ATIKAH FAJRIYAH MUBAROK, MASATERU SENGE, MOHD FAUZI KAMARUDIN

PDF

Biodiversitas 


COUNTRY Indonesia  Universities and research institutions in Indonesia	SUBJECT AREA AND CATEGORY Agricultural and Biological Sciences - Animal Science and Zoology - Plant Science Biochemistry, Genetics and Molecular Biology - Molecular Biology	PUBLISHER Biology department, Sebelas Maret University Surakarta  Universitas Negeri Sebelas Maret in Scimago Institutions Rankings	H-INDEX 19
PUBLICATION TYPE Journals	ISSN 1412033X, 20854722	COVERAGE 2014-2021	INFORMATION Homepage How to publish in this journal grahameagleton@gmail.com

SCOPE











"Biodiversitas, Journal of Biological Diversity" or Biodiversitas encourages submission of manuscripts dealing with all biodiversity aspects of plants, animals and microbes at the level of gene, species, and ecosystem.

Join the conversation about this journal

Quartiles



FIND SIMILAR JOURNALS



Genetic diversity of *Pleurotus ostreatus* (Jacq.) P. Kumm. strains in Java based on random amplified polymorphic DNA markers

NURAENI EKOWATI^{1,*}, ARIS MUMPUNI¹, JUNI SAFITRI MULJOWATI¹, NUNIEK INA RATNANINGTYAS¹,
ARDHINI RIN MAHARNING²

¹Mycology Laboratory, Faculty of Biology, Universitas Jenderal Soedirman. Jl. Dr. Soeparno 63, Purwokerto, Banyumas 53122, Central Java, Indonesia

²Ecology Laboratory Faculty of Biology, Universitas Jenderal Soedirman. Jl. Dr. Soeparno 63, Purwokerto, Banyumas 53122, Central Java, Indonesia.
Tel.: +62-281-638794, Fax.: +62-281-631700, *email: nuraeniekowati29@gmail.com

Manuscript received: 30 September 2020. Revision accepted: 30 May 2021.

Abstract. Ekowati N, Mumpuni A, Muljowati JS, Ratnaningtyas NI, Maharning AR. 2021. Genetic diversity of *Pleurotus ostreatus* (Jacq.) P. Kumm. strains in Java based on Random Amplified Polymorphic DNA markers. *Biodiversitas* 22: 3488-3493. Genetic variation in a fungal population can occur due to mutation and recombination, resulting in changes in the nucleotides that encode specific DNA sequences. Strains with a high genetic distance and good production capabilities can be used to develop genetic breeding. This study aimed to investigate genetic relationship among *Pleurotus ostreatus* strains cultivated in Java (Bogor, Cianjur, Tasikmalaya, Purwokerto, Yogyakarta, Tawangmangu, Malang, and Madiun) based on random amplified polymorphic DNA (RAPD) markers. The research method consisted of DNA isolation and DNA amplification using six primers, i.e. OPA2, OPA3, OPA4, OPA7, OPA9, and OPA10. DNA band data were analyzed using NTSYSpc21 software to determine the level of genetic similarity, based on the Unweighted Pair Group Method with Arithmetic Average Algorithm (UPGMA). In all, 101 amplified DNA bands were obtained, with sizes ranging from 136 to 2320 bp and 96.0% of the bands were polymorphic. Based on cluster analysis, it shows that three clusters were formed. There were genetic variations and relationships among eight *P. ostreatus* strains in Java with a genetic similarity varying from 37-98%.

Keywords: Genetic diversity, *Pleurotus ostreatus*, polymorphism, primers, RAPD

INTRODUCTION

Pleurotus ostreatus (Jacq.) P. Kumm. is a type of edible mushroom with high nutritive value and cultivated in several regions in Indonesia. This fungus is known as a white oyster mushroom because of its white color and shape cap. It can be grown on various agricultural wastes as media (Tsegaye and Tefera 2018). Among *P. ostreatus* cultivation centers in Indonesia, particularly Java including Bogor, Cianjur, Tasikmalaya, Purwokerto, Yogyakarta, Tawangmangu, Malang, and Madiun, it is suspected that cultivation of the fungus has resulted in several strains representing some genetic diversity. *P. ostreatus* has good nutritional value, resulting in increased interest in its cultivation. The determination of genotypic identity is also important for analyses of populations, structure, and evolution within and among species. An important issue is the identification of species beyond morphological and physiological criteria, as this affects fruiting bodies (Khan et al. 2011). This mushroom has low resource and space requirements and can be grown throughout the world and year-round from low-cost starting materials (Chakravarty 2011).

Information regarding genetic variation in *P. ostreatus* strains cultivated in Java is particularly important for selecting superior strains for further development. Inyod et al. (2017) reported that traditional morphological characterization must be supplemented with molecular

characterization to obtain more accurate results. Gupta et al. (2011) and Hasan et al. (2018) reported the development of molecular techniques that can genetically differentiate individuals from one another, in addition to fungal populations. Knowing genetic diversity in *P. ostreatus* will assist in the process of selecting parent strains (parental selection) that are genetically superior. One technique that can be used to determine genetic diversity is the random amplified polymorphic DNA (RAPD) technique. This method can be used to analyze whether there are significant genetic differences in isolates from different locations.

The RAPD technique can easily and quickly detect genetic changes throughout the genome and can be performed with small amounts of DNA. Therefore RAPD-PCR is becoming an extremely popular technique. It has numerous applications such as genetic mapping for fungi, DNA fingerprinting, population genetics studies, epidemiological studies, and the identification of fungal strains (Abdulmalk 2013). Genotyping is particularly important in analyses of population biology, involving the construction of genetic maps and recognizing evolutionary changes. Genetic variation can occur due to changes in the nucleotides that make up an organism's DNA molecules. These changes can in turn then affect phenotypes or reactions to a particular environment. Strains with a high genetic distance and good production ability can be used in genetic breeding (Khan et al. 2011).

The genetic diversity analysis on mushrooms has been using RAPD marker. This technique can determine the genetic similarity of *Pleurotus* strains (Ravash et al. 2010 and Khan et al. 2017). Analysis of the genetic diversity of *Pleurotus* have been conducted on wild or cultivated individual and showed high genetic diversity (Otieno and Onyango 2015). Khan et al. (2017) reported that genetic diversity on local and exotic of *P. ostreatus* using RAPD markers produced 70 bands that could be scored, of which 41 bands were polymorphic, with an average of 2.73 polymorphic bands per primer.

The *P. ostreatus* strains used in this study were commercially cultivated. However, to date, no studies have quantified the genetic diversity of isolates from these various cultivation centers. This study aimed to investigate genetic relationship among *P. ostreatus* strains cultivated in Java (Bogor, Cianjur, Tasikmalaya, Purwokerto, Yogyakarta, Tawangmangu, Malang, and Madiun) based on RAPD markers.

MATERIALS AND METHODS

Research materials

The study was conducted using a survey method, taking two samples of *P. ostreatus* mushrooms from each location. Eight strains were analyzed, one each originating from Bogor (PoB), Cianjur (PoC), Tasikmalaya (PoTs), Purwokerto (PoP), Yogyakarta (PoY), Tawangmangu (PoTw), Malang (PoMl), and Madiun (PoMd). All fungal strains were obtained in the form of pure mycelium culture. Then molecular research was done in Mycology Laboratory, Faculty of Biology, University of Jenderal Soedirman, Purwokerto Indonesia.

DNA extraction

The eight fungal strains were cultured on 50 mL potato dextrose broth (PDB) and incubated in a rotary shaker at 28°C for 15 days at a speed of 100 rpm. Mycelia were harvested using Whatman filter paper no. 41 in a Büchner funnel and used for DNA isolation (Ekowati et al. 2011).

The Nucleon Phytopure fungal DNA extraction RPN-8511/GE (Healthcare, UK) Kit was used to accomplish genomic DNA isolation. Fresh mycelia from 0.1 g to 0.3 g were crushed in liquid nitrogen and placed into 1.5 mL Eppendorf tubes. Then, 500 µL Phytopure I Reagent was added, and the mixture was homogenized by hand or vortex. Next, 200 µL Phytopure II Reagent was added. Samples were incubated at 65°C for 10 min in a water bath, then at 4°C for 20 min. Next, 400 µL cold chloroform that had been stored at 4°C was added to the samples and 20 µL Phytopure III reagent (resin) was added. Centrifugation was performed using a microcentrifuge at a speed of $1,300 \times g$ for 15 min at RT until the upper phase in each sample was clear. The supernatant portions containing DNA in a 200 to 250 µL volume were transferred to new 1.5 mL tubes. Cold isopropanol/propanol with a volume equal to that of the supernatant was added. Then the tubes were closed and inverted to combine the supernatant and isopropanol. Centrifugation was performed at $4,000 \times g$ for

5 min at room temperature. The supernatants were discarded and the precipitates (pellets) were washed by adding 100 µL 70% ethanol. Then the samples were centrifuged three times at $4,000 \times g$ for 5 min, and after each centrifugation, any remaining ethanol was removed by opening and inverting the tubes on tissue paper until the sediments (pellets) were dry. After drying, the pellets were each resuspended in 100 µL Tris-EDTA (TE) buffer. These aqueous DNA samples were stored at -20°C until further analyses. To assess the quality of genomic DNA extracted from each *P. ostreatus* strain, the samples were electrophoresed before RAPD analyses.

PCR reaction and electrophoresis

DNA amplification was performed with the PCR-RAPD technique, using six primers 10 nucleotides in length, i.e. OPA2, OPA3, OPA4, OPA7, OPA9, and OPA10. Each sample was prepared for PCR with the following reaction components: 13.0 µL kit Master mix (dNTP, Taq-DNA polymerase, 1X magnesium chloride buffer), 2.5 µL primer (30.0 pmol), and 3.0 µL the appropriate DNA sample (50.0 ng/mL). Nuclease-free water was added to a 25 µL total volume for each sample. The samples were placed in a thermocycler programmed as follows: one initial stage of denaturation at 92°C for 3 min, then a set of 45 cycles were carried out. Each cycle consisted of temperature settings of 92°C for 1 min to denature the DNA, 37°C for 1 min to anneal primers to template, and 72°C for 2 min for extension. After 45 cycles, extension at 72°C was carried out for 10 min. Then the samples were cooled to 4°C. The amplified samples were removed from the thermocycler and stored at -20°C for later gel electrophoresis analyses. Each reaction was repeated three times.

Amplified DNA samples were electrophoresed using 2% agarose gel in Tris-Borat-EDTA 1X (TBE). The gels were immersed in a 1X TBE running buffer. A total of 5 µL of each amplified sample was mixed with 1 µL loading dye, pipetted into a separate well in the gel, and then exposed to a 100V field for 30 min. Then the gels were immersed in ethidium bromide (EtBr) for 30 min. DNA bands were visualized with a UV transilluminator (Khan et al. 2017; FAMILONI et al. 2018).

Data analysis

The amplified DNA band data were analyzed for size and assessed based on the appearance of the bands. The data obtained were analyzed using the Numerical Taxonomy and Multivariate Analysis System Program, Version 2.1. (NTSYSpc21). Genetic similarity among accessions was calculated according to Jaccard's similarity coefficients using SIMQUAL (Similarity for Qualitative Data). The similarity coefficients were then used to construct dendrogram based on the unweighted pair group method with arithmetic average algorithm (UPGMA) (Rohlf 2000).

RESULTS AND DISCUSSION

RAPD profiles

DNA was isolated at concentrations between 210.20–2152.90 µg/mL with a DNA/RNA ratio between 1.86 and 2.10. The samples were of good quality, quantity, and purity (Figure 1). Template DNA purity is important for its use in PCR amplification. According to Sharon (2010), DNA isolation results are ideal if the ratio of DNA to RNA is between 1.8–2. Therefore, the samples possessed sufficient quality for further PCR analyses.

Genomic DNA samples were used for DNA amplification using six primers resulted in different amounts and sizes of amplified DNA band fragments. The results band sizes ranged from 136 to 2,320 base pairs (bp) (Figure 2). Our results are in line with previous studies, which have reported ranges of 200–2,000 bp in *P. nebrodensis* (Alam et al. 2009), 250–3,500 bp in *P. eryngii* (Ravash et al. 2010), and 646–779 bp in *P. sajor-caju* (Avin et al. 2013).




Figure 1. Genomic DNA band isolated from eight *Pleurotus ostreatus* strains. M: DNA marker 1 kb. Strains 1: Bogor, 2: Cianjur, 3: Tasikmalaya, 4: Purwokerto, 5: Yogyakarta, 6: Tawangmangu, 7: Malang, 8: Madiun, 9: Negative control (without DNA).




Figure 2. Pattern *Pleurotus ostreatus* DNA bands amplified using OPA2, OPA3, OPA4, OPA7, OPA9 and OPA10 primers. Note : M: 100 bp DNA marker of *P. ostreatus* strains 1: Bogor, 2: Cianjur, 3: Tasikmalaya, 4: Purwokerto, 5: Yogyakarta, 6: Tawangmangu, 7: Malang, 8: Madiun, 9: Negative control (without DNA).

Table 1. DNA size, numbers of amplified and polymorphic bands of eight *Pleurotus ostreatus* strains amplified using six RAPD primers

No.	Primer	Sequence (5'-3')	DNA band size (bp)	Numbers of amplified DNA bands	Numbers of polymorphic DNA bands
1	OPA 2	TGCCGAGCTG	136 - 2098	20	17
2	OPA 3	AGTCAGCCAC	170 - 1640	19	18
3	OPA 4	AATCGGGCTG	332 - 2056	17	17
4	OPA 7	GAAACGGGTG	330 - 2256	15	15
5	OPA 9	GGGTAACGCC	155 - 2288	17	17
6	OPA 10	GTGATCGCAG	276 - 2320	13	13
	Total			101	97
	Average			16.83	16.17

The results of polymorphism analyses indicated different levels of polymorphisms resulting from the use of each primer. Of the six primers, all exhibit the presence of polymorphisms among the eight strains of *P. ostreatus*. Thus, the six primers can be used to distinguish genotypes among these strains. The primer that could produce the most polymorphism patterns was OPA3 of 19 amplified DNA bands, and 18 (94.7%) were polymorphic. Based on the combined results of those six primers, 101 amplified DNA bands were observed, and 97 (96.0%) of the DNA bands were polymorphic. For each primer, the average yield was 16.83 DNA bands per primer (Table 1). Yadav et al. (2017) reported that 79.5% of RAPD bands of *Pleurotus* strains were polymorphic, and Pawlik et al. (2012) reported that 21 strains of *Pleurotus* obtained 90.4% polymorphism. The polymorphic band obtained from this current study is higher than those two previous studies which can be caused by the variety of primers used and the higher genetic variation of the strains used in this study.

Cluster analysis

To determine the genetic relationship of the eight *P. ostreatus* strains, the combined data from the six primers were analyzed based on the clustering method of UPGMA to obtain genetic similarity matrix (Table 2) and construct a dendrogram (Figure 3). Based on cluster analysis, it shows that three clusters were formed, the first cluster (I) being a group consisting of PoY, PoMl, PoTw, PoB, PoC, and PoP. Cluster I was divided into four sub-clusters; namely, the first sub-cluster consisted of PoY, PoMl and PoTw which were grouped with high genetic similarity between 0.96-0.98. The second sub-cluster consisted of one strain, PoB, the third sub-cluster consisted of one strain, PoC, and the fourth sub-cluster consisted of one strain, PoP. The second cluster consisted of one strain, PoMd, and the third cluster consisted of one strain, PoTs.


The different strains show variation in genetic similarity. However, the similarity of six of those strains was very high due to the number of amplicons and the percentage of polymorphic DNA bands determined by the number of selective nucleotides in the strain. In addition, it can be caused by the ancestor of the same strain so that there is only a slight genetic change at some locations. Khan et al. (2011) examined genetic variation in seven *Pleurotus* species using the RAPD technique and reported a similarity index of 0.76–0.86. Chandra et al. (2010) tested

eight strains of *P. ostreatus* in India and reported a similarity index between 0.111 and 0.727. Gupta et al. (2011) performed molecular characterization of *Pleurotus* spp. using the RAPD technique showed genetic similarity between 0.17-0.95. Vieira et al. (2013) carried out the molecular characterization of *P. ostreatus* in Brazil the reported a genetic similarity of 0.69. Windarsih et al. (2019) reported genetic similarity for four samples of *Gracilaria coronopifolia* ranged between 0.45-0.57.

In our study, *P. ostreatus* strains from Yogyakarta, Malang, and Tawangmangu showed very high genetic similarity while strain from Malang and Madiun showed closer genetic similarities as well as strain from Tasikmalaya and Cianjur (Figure 3). Geographically, *P. ostreatus* originating from Malang is closer to the strain from Madiun than to the strain from Yogyakarta. However, the Malang strain showed a closer genetic relationship with the Yogyakarta strain than the Madiun strain. Our UPGMA clustering results show that these genetic similarities are not only influenced by geographic distance but are based on genetic factors and other factors. Al-Tufaili et al. (2019) reported that the genetic convergence or divergence among studied strains may be due to many reasons, such as source of origin, different environmental conditions, and breeding or hybridization for these strains. The variation of cultivated strains depends on mutation and hybridization, range of spread, agricultural processes, domestication, or other factors, such as ecological, geographical, breeding system, and anthropogenic effects. Zhao et al. (2013) reported that the genetic distances among geographical populations of *P. eryngii* var. *tuoliensis* were positively correlated with geographic distance and longitudinal distances. This indicates that geographic isolation is an important factor for the observed genetic differentiation. Zhao et al. (2016) reported that adaptation to stress and a temporarily heterogeneous environment are two important factors that influence intraspecies genetic diversity of *P. ferulae* in China. Phylogenetic analyses have also shown that genetic variation in Chinese samples is not geographically linked to their origin. These results indicate that gene flow is not hindered by distance. The present study reveals the abundant diversity for *P. ferulae* in China although its habitat was restricted. Samples from China and Italy were separated according to their geographical origins, but they belonged to the same genetic group.

Table 2. Similarity matrix of eight *Pleurotus ostreatus* strains obtained through RAPD markers

Strains	PoB	PoC	PoTs	PoP	PoY	PoTw	PoMl	PoMd
PoB	1.00							
PoC	0.78	1.00						
PoTs	0.48	0.48	1.00					
PoP	0.70	0.72	0.41	1.00				
PoY	0.78	0.76	0.37	0.74	1.00			
PoTw	0.80	0.72	0.38	0.72	0.96	1.00		
PoMl	0.78	0.76	0.37	0.74	0.98	0.96	1.00	
PoMd	0.50	0.64	0.46	0.52	0.52	0.50	0.54	1.00

**Figure 3.** Dendrogram of genetic relationship among the eight strains of *Pleurotus ostreatus*

P. ostreatus strains from Madiun and Tasikmalaya showed quite high genetic differences versus the other six strains, which means that both strains can be used for genetic breeding development. Khan et al. (2011) stated that strains that have a high genetic distance and good production ability can be used in genetic breeding. Barh et al. (2019) reported that the genetic improvement in *Pleurotus* started with a simple selection technique, which later utilized hybridization (intraspecific, interspecific and intergeneric) and mutation breeding. The traits such as productivity, sporelessness, and quality improvement are important objectives on which most of the works have been done so far. However, new generation approaches such as molecular breeding, genetic transformation and genome editing techniques also added pace to the present improvement process. The improvement of *Pleurotus* mushroom primarily utilizes selection. Selection may be a direct selection from germplasm collected during fungal forays or from cultures made by multispore germination. Selection and mating of genetically diverse parents is another approach to exploit heterosis through hybridization. Avin et al. (2016) reported that the genetic variance components and heritability analyses indicated that the bulk of the *P. pulmonarius* genetic variations

observed for most traits were associated with genes that were dominant in their effects. The newly generated hybrids were identified by both morphological and molecular fingerprinting. Familoni et al. (2018) reported that the outcome of phylogenetic relationships based on the molecular analysis between different or similar populations of *Pleurotus* species provides a better resolution and understanding of their biogeography and speciation. RAPD is a valuable method in distinguishing the different genotypes in mushrooms and evaluating their genetic similarities.

In conclusion, there was variation in the genetic relationship among the eight strains of *P. ostreatus* in Java (Bogor, Cianjur, Tasikmalaya, Purwokerto, Yogyakarta, Tawangmangu, Malang, and Madiun) based on random amplified polymorphic DNA (RAPD) markers, with a genetic similarity varying from 37-98%. Based on cluster analysis, it shows that three clusters were formed.

ACKNOWLEDGEMENTS

We thank the Ministry of Research Technology and Higher Education for providing research funds through the

Strategic National Institution Scheme Project. Our gratitude also goes to LPPM Jenderal Soedirman University, Purwokerto, Banyumas, Indonesia for the sharing of information and collaboration; and the Dean of the Faculty of Biology, Jenderal Soedirman University, Purwokerto, for giving the research permission and supporting this research.

REFERENCES

- Abdulkalk HW. 2013. Rapid differentiation of *Pleurotus ostreatus* from *Pleurotus sapidus* using PCR technique. *Curr Res J Biol Sci* 5: 157-160. DOI: 10.19026/crjbs.5.5482
- Alam N, Shim MJ, Lee MW, Shin PG, Yoo YB, Lee TS. 2009. Phylogenetic relationship in different commercial strains of *Pleurotus nebrodensis* Based on ITS sequence and RAPD. *Mycobiol* 37: 183-188. DOI: 10.4489/MYCO.2009.37.3.183
- Al-Tufaili AKH, Aboohanah MA, Salman FA, Abdel-Hassan NM. 2019. Analysis of genetic distance and similarity in some radish cultivars (*Raphanus sativus* L.) by random amplified polymorphic DNA (RAPD) markers. *J Glob Pharm Technol* 11: 464-470.
- Avin FA, Bhassu S, Sabaratnam V. 2013. A simple and low-cost technique of DNA extraction from edible mushrooms examined by molecular phylogenetics. *Res Crops* 14:897-901.
- Avin FA, Bhassu S, Rameeh V, Tan YS, Vikineswary S. 2016. Genetics and hybrid breeding of *Pleurotus pulmonarius*: Heterosis, heritability and combining ability. *Euphytica* 209: 85-102. DOI: 10.1007/s10681-016-1638-x
- Barh A, Sharma VP, Annepu SK, Kamal S, Sharma S, Bhatt P. 2019. Genetic improvement in *Pleurotus* (oyster mushroom): A review. *Biotech* 9: 322. DOI: 10.1007/s13205-019-1854-x
- Chakravarty B. 2011. Trends in mushroom cultivation and breeding. *Aus J Agric Eng* 2: 102-109.
- Chandra S, Ghosh K, Acharya K. 2010. Comparative studies on the Indian cultivated *Pleurotus* species by RAPD fingerprinting. *Nat Sci* 8: 90-94.
- Ekowati N, Kasiandari RS, Pusposendjojo N, Soegihardjo CJ. 2011. Genetic variation base on random amplified polymorphic DNA molecular markers on shiitake mushrooms (*Lentinula edodes*). *Biota* 16: 178-186. DOI: 10.24002/biota.v16i2.97
- Familoni TV, Ogidi CO, Akinyele BJ, Onifade AK. 2018. Genetic diversity, microbiological study and composition of soil associated with wild *Pleurotus ostreatus* from different locations in Ondo and Ekiti States, Nigeria. *Chem Biol Technol Agric* 5: 7. DOI: 10.1186/s40538-018-0119-y
- Gupta B, Reddy PN, Kotasthane AS. 2011. Molecular characterization and mating type analysis of oyster mushroom (*Pleurotus* spp.) using single basidiospores for strain improvement. *World J Microbiol Biotechnol* 27: 1-9. DOI: 10.1007/s11274-010-0419-2
- Hasan HA, Almomany AM, Hasan S, Al-Abdallat AM. 2018. Assessment of genetic diversity among *Pleurotus* spp. isolates from Jordan. *J Fungi* 4: 52. DOI: 10.3390/jof4020052
- Inyod T, Sassanarakit S, Payapanon A, Keawsompong S. 2017. Morphological characteristics and molecular identification of a wild Thai isolate of the tropical mushroom Hed Taen Rad (*Macrocybe crassa*). *Biodiversitas* 18: 221-228. DOI: 10.13057/biodiv/d180128
- Khan SM, Nawaz A, Malik W, Javed N, Yasmin T, Rehman M, Qayyum A, Iqbal Q, Ahmad T, Khan AA. 2011. Morphological and molecular characterization of Oyster mushroom (*Pleurotus* spp.). *Afr J Biotechnol* 10: 2638-2643. DOI: 10.5897/AJB10.499
- Khan NA, Binyamin R, Awan FS, Khan AI, Waseem M. 2017. Genetic diversity of edible mushroom *Pleurotus* spp. revealed by randomly amplified polymorphic DNA fingerprinting. *Pak J Bot* 49: 1517-1521.
- Otieno OD, Onyango C. 2015. Genetic diversity of Kenyan native oyster mushroom (*Pleurotus*). *Mycologia* 107: 32-38. DOI: 10.3852/13-186.
- Pawlik A, Janusz G, Koszerny J, Malek W, Rogalski J. 2012. Genetic diversity of the edible mushroom *Pleurotus* sp. by amplified fragment length polymorphism. *Curr Microbiol* 65: 438-445. DOI: 10.1007/s00284-012-0175-7
- Ravash R, Shiran B, Alavi A, Bayat F, Rajae S, Zervakis GI. 2010. Genetic variability and molecular phylogeny of *Pleurotus eryngii* species-complex isolates from Iran, and notes on the systematics of Asiatic populations. *Mycol Progress* 9: 181-194. DOI: 10.1007/s11557-009-0624-2
- Rohlf FJ. 2000. NTSYS-pc: Numerical taxonomy and multivariate analysis system, version 2.1. Applied Biostatistics, New York.
- Sharon A. 2010. Molecular and Cell Biology Methods for Fungi. Department of Plant Sciences. Humana Press, New York.
- Tsegaye Z, Tefera G. 2018. Cultivation of oyster mushroom (*Pleurotus ostreatus* Kumm, 1871) using agro-industrial residues. *J Appl Microb Res* 1: 1-6.
- Vieira FR, Pereira DM, Nogueira de Andrade MC, Minihoni MT. 2013. Molecular characterization of *Pleurotus ostreatus* commercial strains by random amplified polymorphic DNA (RAPD). *Afr J Agric Res* 8: 3146-3150.
- Windarsih G, Utami DW, Yuriyah S. 2019. Genetic diversity and productivity of *Gracilaria coronopifolia* as alternative for food resource based on RAPD marker. *Biodiversitas* 20: 3758-3765. DOI: 10.13057/biodiv/d201239
- Zhao M, Huang C, Chen Q, Wu X, Qu J. 2013. Genetic variability and population structure of the mushroom *Pleurotus eryngii* var. *tuoliensis*. *PLoS One* 8: e83253. DOI: 10.1371/journal.pone.0083253
- Zhao M, Huang C, Wu X, Chen H, Qu J, Li Y, Gao W, Zhang J. 2016. Genetic variation and population structure of the mushroom *Pleurotus ferulae* in China inferred from nuclear DNA analysis. *J Integr Agr* 15: 2237-2246. DOI: 10.1016/S2095-3119(16)61383-7
- Yadav MK, Chandra R, Singh HB, Yadav SK, Yadav SK, Naik S, Dhakad PK. 2017. Genetic diversity characterization of *Pleurotus* strains by random amplified polymorphic DNA fingerprinting. *Intl J Curr Microbiol App Sci* 6: 1260-1267. DOI: 10.20546/ijcmas.2017.605.136.