

INTERNATIONAL FOOD RESEARCH JOURNAL

[Home](#)[Editorial Board](#)[Author's Guide](#)[Code of Ethics](#)[List of Issues](#)[Contact Us](#)[IFRJ Website](#)

EDITORIAL BOARD

Editor-in-Chief

Prof. Dr. Tan Chin Ping
Department of Food Technology,
Faculty of Food Science and Technology,
Universiti Putra Malaysia,
MALAYSIA

Associate Editors

Prof. Dr. Faridah Abas
Department of Food Science,
Faculty of Food Science and Technology,
Universiti Putra Malaysia,
MALAYSIA

Assoc. Prof. Dr. Chong Gun Hean
Department of Food Technology,
Faculty of Food Science and Technology,
Universiti Putra Malaysia,
MALAYSIA

Assoc. Prof. Dr. Anis Shobirin Meor Hussin
Department of Food Technology,
Faculty of Food Science and Technology,
Universiti Putra Malaysia,
MALAYSIA

Assoc. Prof. Dr. Ahmad Faizal Abdull Razis
Department of Food Science,
Faculty of Food Science and Technology,
Universiti Putra Malaysia,
MALAYSIA

Board Members

Prof. Dr. Nazamid Saari
Department of Food Science,
Faculty of Food Science and Technology,
Universiti Putra Malaysia,
MALAYSIA

Prof. Dr. Jinap Selamat
Department of Food Science,
Faculty of Food Science & Technology,
Universiti Putra Malaysia,
MALAYSIA

Prof. Dr. Wan Rosli Wan Ishak
School of Health Sciences,
Health Campus,
Universiti Sains Malaysia,
MALAYSIA

Prof. Dr. Karamatollah Rezaei
Department of Food Science and Engineering,
Faculty of Agricultural Engineering and Technology,
University of Tehran, Karaj,
IRAN

Prof. Dr. Olga Martin-Belloso
Food Technology Department,
University of Lleida,
SPAIN

Prof. Dr. Gulam Sumnu
Department of Food Engineering,
Middle East Technical University (METU),
TURKEY

Prof. Dr. Stefan Kasapis
Department of Biosciences and Food Technology,
School of Science,
RMIT University,
AUSTRALIA

Dr. Siree Chaiseri
Department of Food Science and Technology,
Faculty of Agro-Industry,
Kasetsart University,
THAILAND

Prof. Dr. John Gilbert
Orta Dogu Teknik Universitesi,

Prof. Dr. Steve Flint
Institute of Food Science and Technology,

FoodLife International Ltd,
TURKEY

School of Food and Nutrition,
Massey University,
NEW ZEALAND

Prof. Dr. Liu Yang

Institute of Agricultural Resources and Regional Planning,
Chinese Academy of Agricultural Sciences,
P.R. CHINA

Prof. Dr. Gregory R. Ziegler

Department of Food Science,
College of Agricultural Sciences,
The Pennsylvania State University,
USA

Prof. Dr. Yonghua Wang

School of Food Science and Engineering,
South China University of Technology,
CHINA

Prof. Dr. Van Viet Man LE

Department of Food Technology,
Ho Chi Minh City University of Technology,
VIETNAM

Prof. Dr. Yuanfa Liu

School of Food Science and Technology,
Jiangnan University,
P.R. CHINA

Prof. Dr. Yin Li

CAS Key Laboratory of Microbial Physiological and Metabolic
Engineering,
Institute of Microbiology,
Chinese Academy of Sciences,
CHINA

Dr. Franz Berthiller

Center for Analytical Chemistry,
Department of Agrobiotechnology (IFA-Tulln),
University of Natural Resources and Life Sciences,
AUSTRIA

Prof. Dr. Rekha S. Singhal

Food Engineering and Technology Department,
Institute of Chemical Technology,
INDIA

Prof. Dr. Yong Wang

Department of Food Science and Engineering,
College of Science and Engineering,
Jinan University,
CHINA

Assoc. Prof. Dr. Gan Chee Yuen

Analytical Biochemistry Research Centre (ABrC),
Universiti Sains Malaysia,
MALAYSIA

INTERNATIONAL FOOD RESEARCH JOURNAL

[Home](#)[Editorial Board](#)[Scope](#)[Subscription](#)[Guide](#)[List of Issues](#)[Contact Us](#)[FOOD Website](#)**VOLUME 21,2014****International Food Research Journal Volume 21 Issue 1, 2014****Mini Review**

1. **Recent research and development of *Monascus* fermentation products (1-12)**
 - Srinta, I., Ristiarini, S., Nugerahani, I., Sen, S. K., Zhang, B. B., Xu, G. R. and Blanc, P.J.
2. **Rice in health and nutrition (13-24)**
 - Rohman, A., Siti Helmiyati, Mirza Hapsari and Dwi Larasati Setyaningrum
3. **Rambutan seed fat as a potential source of cocoa butter substitute in confectionary product (25-31)**
 - Issara, U., Zzaman, W. and Yang, T.A.

Original Articles

4. **Microbiological quality analysis of shrimps collected from local market around Dhaka city (33-38)**
 - Samia, S., Galib, H. T., Tanvir, A. S., Basudeb, C. S., Md. Walliullah, Tasnia, A., Md. Sakil, M., Afsana, F. N., Sadia, K. P., Kamal, K. D., Mrityunjoy, A., Nusrat, J. U., Tasmina, R. and Rashed, N.
5. **Physico-chemical and sensory properties of musts and wines from *Melodorum fruticosum* Lour. (39-43)**
 - Tatdao, P., Norrasat, S. and Tiwawan, S.
6. **Isomerisation and degradation of lycopene during heat processing in simulated food system (45-50)**
 - Huawei, Z., Xiaowen, W., Elshareif, O., Hong, L., Qingrui, S. and Lianfu, Z.
7. **Effects of high pressure and thermal processing on phytochemical, color and microbiological qualities of herbal-plant infusion (51-57)**
 - Chaikham, P., Worametachanon, S. and Apichartsrangkoon, A.
8. **Biofilm assessment of *Vibrio parahaemolyticus* from seafood using Random Amplified Polymorphism DNA-PCR (59-65)**
 - Elexson, N., Yaya, R., Nor, A. M., Kantilal, H. K., Ubong, A., Yoshitsugu, N., Nishibuchi, M. and Son, R.
9. **Microwave drying kinetics of olive fruit (*Olea europaea* L.) (67-72)**
 - Mahdhaoui, B., Mechlouch, R. F., Mahjoubi, A. and Ben Brahim, A.
10. **Optimization of drying temperature and water extraction time of *Monascus*-fermented durian seed for the Monacolin K content using Response Surface Methodology (73-76)**
 - Srinta, I., Nugerahani, I., Sutedja, A. M. and Widharna, R. M.
11. **Effect of mash quantity and roasting duration on some physical and pasting properties of *gari* (77-82)**
 - Rahman Akinoso and Olatunde, S. J.
12. **Survivability and metabolic activity of *Lactobacillus casei* 01 incorporating lychee juice plus inulin under simulated gastrointestinal environment (83-89)**
 - Kingwatee, N., Apichartsrangkoon, A., Chaikham, P., Pankasemsuk, T. and Changrue, V.
13. **Effects of combined antioxidants and packing on lipid oxidation of salted dried snakehead fish (*Channa striata*) during refrigerated storage (91-99)**
 - Nitipong, J., Nongnuch, R., Kamonwan, R. and Teeraporn, K.
14. **Fate of the phenolic compounds during olive oil production with the traditional press method (101-109)**
 - Goldsmith, C. D., Stathopoulos, C. E., Golding, J. B. and Roach, P. D.
15. **The utilization of cocoa polyphenol extract to improve the shelf life of bulk frying oil used in open and deep frying (111-118)**
 - Misnawi, Teguh, W., Susijahadi, Eka, N. D. and Noor, A. F.
16. **Total phenolic content, reducing power, antioxidative and anti-amylase activities of five Bangladeshi fruits (119-124)**
 - Aklima, J., Mojumder, S. and Sikdar, D.
17. **Consumption pattern and consumers' opinion toward street food in Nakhon Si Thammarat province, Thailand (125-130)**
 - Khongtong, J., Ab Karim, S., Othman, M. and Bolong, J.

- acuminata* var. *Emas*) flour (2133-2139)
 ◦ Norhidayah, M., Noorlaila, A. and Nur Fatin Izzati, A.
12. Papaya seed flour (*Carica papaya*) affects the technological and sensory quality of hamburgers (2141-2145)
 ◦ Azevedo, L. A. and Campagnol, P. C. B.
 13. Comparison of protein, total fat, and omega-3 fatty acids content in yellowtail catfish (*Pangasius pangasius*) and long tail shad (*Hilsa (clupea) macrura*) in raw and pressurized fish (2147-2153)
 ◦ Asmah, R., Siti Sumaiyah, S. A. and Nurul, S. R.
 14. Effect of duck egg white addition on textural properties and microstructure of rice noodles (2155-2159)
 ◦ Sasipa Kerdnilek and Wunwidoon Garmjanagdonchorn
 15. Pasting properties mixtures of mangrove fruit flour (*Sonneratia caseolaris*) and starches (2161-2167)
 ◦ Jariyah, Widjanarko, S. B., Yumianta, Estiasih, T. and Sopade, P. A.
 16. Nutritional evaluation of different mango varieties available in Bangladesh (2169-2174)
 ◦ Ara, R., Motalab, M., Uddin, M. N., Fakhruddin, A. N. M. and Saha, B. K.
 17. Trans fatty acids in selected bakery products and its potential dietary exposure (2175-2181)
 ◦ Siti Nurshahbani, S. and Azrina, A.
 18. The stability of double fortification of salt with iodine and iron in different storage conditions (2183-2187)
 ◦ Helmyati, S., Narendra, Y. H., Septi, P., Rochyana, I. and Endri, Y.
 19. Effect of thermal and thermosonication on anthocyanin stability in jamun (*Eugenia jambolana*) fruit juice (2189-2194)
 ◦ Shaheer, C. A., Hafeeda, P., Kumar, R., Kathiravan, T., Dhananjay Kumar and Nadanasabapathi, S.
 20. Extraction and characterization of pectin from grapefruit (*Duncan cultivar*) and its utilization as gelling agent (2195-2199)
 ◦ Khan, A. A., Butt, M. S., Randhawa, M. A., Karim, R., Sultan, M. T. and Ahmed, W.
 21. Application of sonication and mixing for nanoencapsulation of the cyanobacterium *Spirulina platensis* in liposomes (2201-2206)
 ◦ Machado, A. R., Assis, L. M., Costa, J. A. V., Badiale-Furlong, E., Motta, A. S., Micheletto, Y. M. S. and Souza-Soares, L. A.
 22. Preparation of arrowroot starch nanoparticles by butanol-complex precipitation, and its application as bioactive encapsulation matrix (2207-2213)
 ◦ Winarti, C., Sunarti, T. C., Mangunwidjaja, D. and Richana, N.
 23. Effect of bitterns and constituent salts on the inosinate phosphatase activity in horse mackerel muscle (2215-2220)
 ◦ Hiroko Seki, Kazufumi Osako and Naoko Hamada-Sato
 24. Changes in chemical quality of cocoa butter during roasting of pulp pre-conditioned and fermented cocoa (*Theobroma cacao*) beans (2221-2227)
 ◦ Afoakwa, E. O., Ofosu-Ansah, E., Takrama, J. F., Budu, A. S. and Mensah-Brown, H.
 25. Functional properties of soy protein isolate of crude and enzymatically hydrolysed at different times (2229-2236)
 ◦ Kempka, A. P., Honaizer, T. C., Fagundes, E. and Prestes, R. C.
 26. Determination of major and minor capsaicinoids in different varieties of the *Capsicum* fruits using GC-MS and their inhibition effect of the chilli extract on α -amylase activity (2237-2243)
 ◦ Maokam, C., Techawongstien, S. and Chanthai, S.
 27. Extraction and enzymatic depolymerization of gum from *Artocarpus heterophyllus* Lam. seeds (2245-2251)
 ◦ Dasaesamoh, R. and Seechamnaturakit, V.
 28. Effect of cardamom leaves extract as antidiabetic, weight lost and hypocholesterolemic to alloxan-induced Sprague Dawley diabetic rats (2253-2261)
 ◦ Winarsi, H., Sasongko, N. D., Purwanto, A. and Nuraemi, I.
 29. Role of antioxidant enzymes in oxidative stress and immune response evaluation of aspartame in blood cells of wistar albino rats (2263-2272)
 ◦ Arbind, K. C., Sheela Devi, R. and Sundareswaran, L.
 30. Antidepressant activity of a polyherbal mixture in mice (2273-2277)
 ◦ El-Ashmawy, I. M., Bayad, A. E. and Omar, A. M. E.
 31. Effect of pre-slaughter stunning on the death of the poultry and myofiber apoptosis (2279-2284)
 ◦ Shahdan, I. A. and Rahman, M. T.
 32. Effect of sugar types on physical attributes and crystalline structure of sweet-dried chicken meat product (2285-2291)
 ◦ Wongwiwat, P. and Wattanachant, S.
 33. Effect of addition of rice bran oil extract on the stability of sunflower oil, sesame oil and their blends (2293-2298)
 ◦ Bardhan, J., Sahoo, B., Chakraborty, R. and Raychaudhuri, U.
 34. The optimization of extrusion condition of Phatthalung Sungyod rice extrudate: a preliminary study (2299-2304)
 ◦ Keawpeng, I., Charunuch, C., Roudaut, G. and Meunier, M.
 35. Essential oil composition of some spices treated with nitrogen in arid region (2305-2309)
 ◦ Khalid, A. K.

Effect of cardamom leaves extract as antidiabetic, weight lost and hypocholesterolemic to alloxan-induced *Sprague Dawley* diabetic rats

¹*Winarsi, H., ²Sasongko, N. D., ³Purwanto, A. and ¹Nuraeni, I.

¹Medical and Sciences Health Faculty, Nutrition Science Department of Jenderal Soedirman University, Purwokerto 53123, Indonesia

²Biology Faculty of Jenderal Soedirman University, Purwokerto, Indonesia ³General Hospital of Margono Soekarjo, Purwokerto. Indonesia

Article history

Received: 1 November 2013

Received in revised form:

8 April 2014

Accepted: 9 April 2014

Keywords

Cardamom-leaves
Blood-glucose and
cholesterol-level
Body-weight
Diabetic-rats

Abstract

Current study was proposed to know the potency of Cardamom leaves as antidiabetic, weight lost, and hypocholesterolemic on alloxan-induced Sprague-Dawley rats. The Cardamom leaves were prepared as extract using ethanol and its termed as ethanolic cardamom leaves extract (ECLE). Thirty of 45 rats were induced with alloxan at the dose of 120 mg/kg body weight, intraperitoneal. Blood glucose level of the rats were measured after 7 days of alloxan induction and forced to fast overnight. For this purpose, we split the rats into three different groups consisted of 15 rats each. Group I; the diabetic rats were given fed added by the ECLE, Group II; the diabetic rats were only given fed, and Group III; were non diabetic rats and given only fed. Intervention was observed for 14 days, and blood sample were taken 3 times i.e.: 0, 7 and 14 days after being intervened. One mL blood was taken from eyes sub-orbital using hematocrit capillary tube then transferred to eppendorf tube containing 10% EDTA. Samples were then centrifuged at 3,000 rpm, for 10 minutes. Blood glucose and cholesterol level were quantified after blood plasma was obtained. The rats body weight were measured simultaneously with that of blood sampling. Data were then analyzed by analysis of variance, and followed by Duncan test whenever there were a 5% significant different. Data showed that blood glucose level of rats group I decreased from 201.7 to 102.8 mg/dl ($P = 0.017$), cholesterol level from 77.6 to 56 mg/dl ($P = 0.025$), but not to the rats body weight which increased as also noted from the non-diabetic rat ($P = 0.92$). ECLE is potential to be used as functional food component for therapeutic of diabetic patient.

© All Rights Reserved

Introduction

Diabetes mellitus is caused by metabolic disorders. This diseases is characterized by hyperglycemia due to insulin secretion disorders, production or both (ADA, 2006). Diabetic is a progressive disease and strongly related to more complicated diseases. Moreover, diabetic has been known as the main cause of death in either human or animal (Avizeh *et al.*, 2010). There was no modern medicinal is safe to be used by patients, but, there are more than 800 plants/herbs are containing antidiabetic components (Alarcon-Aguilera *et al.*, 1998). The plant could be grouped as traditional antidiabetic when it has potential to cause hypoglycemia. Several medicines are chategorized capable to reduce hyperglycemic toward the diabetic patients, however, this group will cause side effects like diarrhoea lactic acidosis, and body weight. This situation is therefore, forcing us to search newer antidiabetic medicine or component with minimum risks to diabetic patients.

Diabetic patients in most of the cases will

also facing problems of hyperlipidemic and hypercholesterolemic. Alloxan-induced diabetic rats showed high level of lipid leads to becomes phospholipid and cholesterol. The high cholesterol level, the high risk for atherosclerosis, therefore, solving the problem of hypercholesterol of diabetic rats becomes a prerequisite.

Induction of alloxan to the guinea pigs causes also negatively to their weight due to their hyperglycemic (Szkudelski, 2001) and defects the protein and lipid tissues. The situation where body has a limit amount of energy, these two important components are then used by the body as energy sources, meanwhile the body has lots of glucose in its blood tissues (Shirwaikar *et al.*, 2004). This condition will then cause reduction of weight while it needs to be controlled.

The cardamom leaves are never being used by the growers or farmers, however, Winarsi *et al.* (2012) reported that the cardamom leaves contain 129.6 ± 6.9 mg/g flavonoid and vitamin C 19.22 ± 1.1 mg/g. It has been reported that flavonoid has antioxidant (Coskun *et al.*, 2005), antidiabetic (Dewanjee *et al.*, 2008),

*Corresponding author.

Email: winarsi12@gmail.com

hypocholesterolemic characteristic (Fuhrman *et al.*, 2000), and capable to control body weight (Yuji *et al.*, 2006). Winarsi *et al.* (2012) were also noted that the IC_{50} of cardamom leaves extract was smaller than that of ascorbic acids. In the current times, however, there is no available data showing the potential of ECLE as antidiabetic, hypocholesterolemic and controlling body weight *in vivo*. Currently, this study was done to overcome the potential characteristic of ECLE on reducing blood glucose and total cholesterol level, and therefore, controlling body weight of alloxan induced-diabetic rats.

Methods

Preparation of ECLE

Ethanol Cardamom Leaves Extract (ECLE) was prepared as in Winarsi *et al.* (2013). Cardamom leaves were cut into small pieces and grind to powder (cardamom leaves powder, CLP). The CLP was then immersed in ethanol 96% for several times until a clear filtrate was obtained, then evaporated to let get the extract.

Induction of alloxan to the rats

Fourty five males *Rattus norvegicus* L. strain Sprague Dawley rats, age of 2-3 months old, 220-250 gram weight, collection of the LPPT UGM Yogyakarta. During the acclimatisation, these rats were pellet fed of 10% from their body weight, the amount of water is ad libitum. After being forced to fast overnight, 30 of those 45 rats were induced with monohydrate alloxan in physiological NaCl sterile at the dose of 120 mg/kg body weight, intraperitoneal (Kim *et al.*, 2006; Winarsi *et al.*, 2013). Blood glucose level of the rats were measured with Glucotest meter (Nesco) after 7 days of alloxan induction and forced to fast overnight.

Grouping of treated rat and intervention

The diabetic rats were then grouped into 2 different groups consisting of 15 rats each. The rest 15 rats were grouped as control group. Total was three groups. Group I was diabetic rat given for feed plus ECLE, Group II was diabetic rat given for feed only, and Group III was non-diabetic rat given for feed only.

Intervention of ECLE to the diabetic rat

Intervention was done for a period of 14 days. The ECLE was given to the treated rat at the dose of 100 mg/kg body weight, pellet feed, however, given for only 10% of their body weight, and water at ad libitum (Winarsi *et al.*, 2013).

Preparation of blood samples

One ml of blood was taken from each rat for three times, i.e.: at the baseline, 7, and 14 days after having intervention. Blood was taken from eye sub orbital using the hematocrite tube then transferred into eppendorf tube containing of 10% EDTA. Part of the blood was separated for measuring the glucose level, while the rest was centrifuged to obtain the plasma for cholesterol level determination.

Determination of blood glucose level

Blood glucose level of the treated rat was determined using a Glucotest meter (Nesco). Before being used, the Glucotest meter was calibrated as written on the test-stripes code number. Every measurement need a single strip-test. Standard liquid for calibration is available in the kit. Strip-test was placed on the Glucotest meter and a sign of blood will appear on the screen to be ready for a further use. Blood was placed on to the strip-test and left for about 5 second until the screen shows the number of glucose level which was noted in mg/dl.

Determination of total cholesterol level

Blood plasma was determined for its cholesterol level using an end point method (Stanbio Cholesterol Liquicolor procedure no 1010). For this purpose, one ml reagent was added by one ml standard solution and one ml blood plasma are placed into cuvette-I and used as Blanko Reagent (BR). One ml of this reagent was then added by 0.01 ml standard solution and placed into the cuvette-II, this is called as standard solution (S). Another one ml of reagent was added by 0.01 ml blood plasma and placed in the cuvette-III and used as sample solution (U). All cuvettes containing different material were then incubated for 5 minutes at 37°C or alternatively incubation time of 10 minutes at the room temperature. Each sample was then read for the absorbance using spectrophotometer at 500 nm. The total cholesterol level can be calculated with the following equation:

$$\text{Total cholesterol (mg/dl)} = \frac{Au \times 200}{As}$$

Notes:

Au = the absorbance value of the plasma

As = the absorbance value of the standard solution

200 = standard for total cholesterol (mg/dl)

Determination of the treated rat body weight

The treated rat body weight was determined in the different times namely: baseline, 7, and 14 days after intervention using a weigh scale of Cook Master Kitchen Scale Model CMK6602, ISO 9001: 2008

Certified by SGS.

Statistical analysis

Data obtained were then analyzed by Analysis of Variance (Anova), and followed by Duncan test when there is significant different at the level of 5%.

Results

Rats which were induced by alloxan showed hyperglycemic, characterized by high level of glucose level in their blood 351.5 mg/dl and so called as diabetic rat, this number was higher than that of non induced rat which was only 110.3 mg/dl ($P = 6.18E-08$). With total cholesterol of 75.5 mg/dl, which was higher than the non diabetic ones 45.5 mg/dl. On the contrary, body weight of the diabetic rat were only 220.8 ± 23 g, a slightly lower than that of non-diabetic rat 247.6 ± 28 g ($P < 0.0005$) (Winarsi *et al.*, 2012).

Discussion

Blood glucose level of the diabetic rats

High blood glucose level in the treated rats were related with the induction of alloxan which is known as diabetogenic compound and toxic, selective to beta-pancreatic cells, leads to insulin deficiency. Pari *et al.* (2002) stated alloxan could decrease insulin secretion by beta-pancreatic cells and therefore, causing hyperglycemic. This situation will cause glucose can not be inserted in the metabolisms process but sticking in the blood.

Through a redox reaction, the high level of blood glucose will increase the formation of reactive oxygen species (ROS), leads to more NADH and FADH₂ enter to electron transport chain (Suarsana *et al.*, 2011). Increase of electron transport rate leads to the contribution of free radicals formation and cause a more severe diabetic.

Alloxan has an analog glucose character and therefore, this compound is known by the GLUT2, then entered to bilayer membrane of lipid plasma where beta-pancreatic cells are accumulated, and cause damage in granules those produce insulin (Lenzen, 2007). The cytotoxic activity of alloxan is mediated by free radicals formed during the redox reaction. Alloxan and its reduction product (dialuric acid) through redox cycles will form superoxide radicals and cause dismutation to become hydrogen peroxide. The more free radicals in the blood, the higher concentration of cytosol calcium leads to the faster damage of beta-pancreatic cells.

The beta-pancreatic cells are the glucose censorship, work to reduce blood glucose level by




Figure 1. Blood glucose level of diabetic rats during intervention. K, diabetic rats were given pellet + ECLE; P, diabetic rats were given pellet; N, non diabetic rats were given pellet.




Figure 2. Body weight of diabetic rats during intervention. K, diabetic rats were given pellet + ECLE; P, diabetic rats were given pellet; N, non diabetic rats were given pellet.




Figure 3. Cholesterol level of diabetic rats during intervention. K, diabetic rats were given pellet + ECLE; P, diabetic rats were given pellet; N, non diabetic rats were given pellet.

releasing insulin. When glucose passed the membrane plasma, it will be fosforilated by glucosidase and start to glycolysis activity. The ATP which was formed during the metabolism in the mitochondrion will force to close channel of KATP, leads to depolarization of plasma membrane. This situation will cause Ca²⁺ enter the blood through gradient concentration, and increase the Ca²⁺ concentration in the cytosol, leads to the exocytosis of insulin. The high concentration of Ca²⁺ in the cytosol, the high level of insulin secreted, and will damage beta-pancreatic cells (Maechler *et al.*, 2006). The chronic condition of hyperglycemic due to oxidative stress may cause damage, disfunction, and failure of several organs (Likidilid *et al.*, 2007), therefore, decreasing or minimalizing blood glucose level of diabetic patients becomes a prerequisite.

Current study showed that the diabetic rats given the ECLE for 7 consecutive days, decreased in their blood glucose level from 221.6 to 122.2 mg/dl ($P = 0,019$). The blood glucose level of this treated rats were continuously decrease when the ECLE was given to a longer time (14 days) of 201.7 to 102. 8

mg/dl ($P = 0,017$) in compare to those of diabetic group rat given feed only (Figure 1).

The ECLE contains flavonoid of 129.6 ± 6.9 mg/g (Winarsi *et al.*, 2012). Flavonoids are group of plant phenolic secondary metabolite which are abundantly present in nature. Eventhough, in their research did not define the type of flavonoids in the ECLE, but it is clear that this phenolic compound has similar structure to flavone, flavanone, flavonol, flavanonol, cetechin, anthocyanidin, and isoflavone. Several researchers stated that function of the flavonoid as antidiabetic (Brahmachari, 2008; Brahmachari, 2009). This compound reduced blood glucose level of diabetic rats. Using several kind of guinea pigs, they showed the positive hypoglycemic affect of flavonoid, and therefore, it can be used to reduce absorption of glucose or to increase glucose tolerance. A similar results have been reported by Bhathena and Velásquez (2002) who explore the use of isoflavone extract *in vitro*, they found this compound could stop glucose absorption on brush border colon of rabbit vesicula membrane. Naringenin was also reported to reduce uptake the rate of glucose within the vesicle membrane brush border of diabetic rat, to the same amount of non-diabetic rat (Li *et al.*, 2006). Several types of flavonoids like epicatechin gallate, myricetin, quercetin, apigenin, epigallocatechin gallate, and epigallocatechin were also can be used as competitive inhibitor on the glucose absorption against the sodium dependent glucose transporter-1. Further study by Johnston *et al.* (2005) and Zhao *et al.* (2004) noted that non-glycosylase of flavonoid clarify reduction of absorption activity of sodium-dependent glucose both *in vitro* or *in vivo*.

Besides reducing absorption of glucose, in order to control blood glucose level, flavonoid might also take another route by slowing down glucosidase-alpha activity in the colons. Effect on slowing down activity of the enzyme could be seen when luteolin, kaempferol, chrysin or galangin is taking place in the absorption and metabolism of carbohydrate (Matsui *et al.*, 2002). Kim *et al.* (2000) also proofed if activity of glucosidase-alpha was inhibited by flavonoid luteolin, amentoflavone, luteolin 7-o-glucoside and daidzein which runs as glycation inhibitors. Another possibility is by stimulating glucose absorption in the peripheral tissues and regulating enzymes of carbohydrate metabolism.

Hypoglycemic effects of flavonoid was also showed by pruning (naringenin 7-O- β -D-glucoside) which deliberately given intraperitoneally to diabetic rat. Moreover, it was also stated that flavonoid could not only to cause hypoglycemic but also potential to be used as anti-hyperglycemic especially of those

chrysin and its derivates, like silymarin, isoquercetin, and rutin (Hnatyszyn *et al.*, 2002). In a long period of intervention of this ECLE containing flavonoid, given to the diabetic rats per oral, showed a decrease of glucose in the blood plasma up to 70.7%, but in the diabetic rats were not given the compound reduced to 47%. Similar with this research that other types of flavonoids, hesperidin and naringin were also can reduce blood glucose level of the diabetic rats (Jung *et al.*, 2004). Those of flavonoid might take either cellular or molecular mechanism.

Flavonoids could also take a role in the secretion of insulin due to their insulin mimetic characteristic which might be related with pleiotropic mechanism (Winarsi and Purwanto, 2010). This action could help to suppress complication due to diabetic. This phenomenon showed that flavonoid is potential to be used as drugs to stimulate glucose absorption on peripheral tissues, and control enzymes activities within the carbohydrate metabolism pathway. In their study found that genistein was activated beta-pancreatic cells directly (Liu *et al.*, 2006), and cause the activation of cAMP/PKA signaling route leads to the insulinotropic effects. It might be concluded that flavonoid is a natural compound fit to be used as an alternative medicines in diabetic therapy.

Another study noted that flavonoid given routinely per oral to the diabetic rat caused reduction of glucose level in the blood plasma, increasing insulin, glycogen content, and activity of hexokinase enzyme. Prince and Kamalakkannan (2006) supported the previous study by saying that flavonoid which was given routinely reduced glucose 6-phosphatase and fructose 1,6-bisphosphatase significantly in the liver and tissues of diabetic rat.

The mechanism of antidiabetic effect of apigenin-6-C- β -fucopyranoside and L-apigenin-6-C-(2-O''- α -L-rhamnopyranosyl)- β -l-fucopyranoside by stimulating or inducing secretion of insulin in the hyperglycemic rat (Cazarolli *et al.*, 2009a; 2009b). Flavonoid was also reported that capable to stimulate synthesis of glycogen in the rat tissues through mechanism of transduction of insulin signals, then showed a double effect as anti-hyperglycemic or insulin secretion, and insulin mimetics or glycogen synthesis. Flavonoid is an antioxidant compound which can reduce cell damage due to diabetic. Quercetin, a strong type of flavonoid, was able to reduce blood glucose level normalize antioxidant level, superoxide dismutase, vitamin C, and vitamin E. The quercetin was reported to be more effective to control diabetic which induced by oxidative stress in a low dose (Mahesh and Menon, 2004).

The potential of alloxan, a diabetogenic

compound, will reduce in the presence of glutathione, especially when alloxan radicals are changed into dialuric acid. The oxidative radical compound formed can destroy beta-pancreatic cells in the langerhans. Quercetin is potential to minimize the damage of beta-pancreatic cells either as cells scavengers or taking role as chelating agent (Mira *et al.*, 2002; Anjaneyulu and Chopra 2004).

Intervention of ECLE orally showed a reduction in blood glucose level of the diabetic rat. Clinical study by Knekt *et al.* (2002) showed that improving condition on the diabetic rat might help to reduce risks of diabetic complications as retinopathy, nephropathy, neuropathy and foot diabetic, and also inhibit the possibility of atherosclerosis especially that of related with myocard infark and stroke. Several research reports also noted the significant role of flavonoid in reducing hyperglycemic (Anjaneyulu and Chopra, 2004; Dene *et al.*, 2005) and glycation of non-enzymatic protein on the guinea pigs.

Flavonoid is proven to be able to stimulate immune system. This statement was due to antioxidant character of flavonoid which was able to recover the beta-pancreatic cells damage caused by alloxan. The flavonoid glycosidic component takes place as suppressor of the hydroxyl radicals, and therefore, capable to block development of diabetic. It was also reported that flavonoid could interact with several protein transporter then it will inactivate some enzymes and causing concentration of hormones in the tissues. Quercetin capable to block the formation of free radicals and therefore, can be used as antidiabetic. Using ethanolic extract of ginger (*Zingiber officinale*) Elshater *et al.* (2009) reported this extract was able to reduce blood glucose level of diabetic rat. This extract was explored to contain phenolic compounds like gingerol, shogaol, zingerone, and paradol, which are capable to repair insulin secretion by beta-pancreatic cells. Raj *et al.* (2011) supported that ethanolic extraction of other rhizome (*Alpinia calcarata*) which contains flavonoid, alkaloid, and coumarin was capable to reduce blood glucose level through increasing secretion of insulin after repairing the beta-pancreatic cells, glycogenolysis and gluconeogenesis.

Hyperglycemic of the diabetic patient will reduce the system of antioxidant defence. Previous reports noted that this condition will be followed by low level of enzymatic antioxidants like glutathione peroxidase, superoxide dismutase, glutathione reductase, and vitamin C in both blood plasma and tissues (Kalaivani *et al.*, 2008; Middha *et al.*, 2011). Deficiency of vitamin C in blood plasma of diabetic patients might cause severe damage of the tissues due

to the presence of ROS. As a potential antioxidant, vitamin C is known capable to suppress the ROS. Dakhale *et al.* (2011) showed the combination of vitamin C and metformin could control blood glucose level of the diabetic patients. In this case, metformin repairs transport of glucose into the muscle cells, while vitamin C reduce the ROS.

Vitamin C (ascorbic acids) helps body to produce collagen and basic component for connective tissues. Collagen is an important element of blood vessels membrane, gum, and bone; and this material is strongly important to those who are in a recovery phase after having surgery. Though its mechanisms is not clear, but vitamin C is functioned as antioxidant by removing free radicals, improving immune system, protection of cancer, cataract, and slowing down degenerative diseases due to aging, macula retina, and other chronic diseases. Vitamin C is an important antioxidant in human and suppress the formation of oxygen radicals (McRae, 2007). Due to its similar structure to glucose, vitamin C could also take over the position of glucose in some reactions. Knekt *et al.* (2004) stated vitamin C was effective in blocking non enzymatic proteins. Besides that, in the guinea pigs, vitamin C is capable to regulate catabolism of cholesterol becomes acids, and even proofed to play significant role in lipid regulation. Several studies done previously, showed reduction of vitamin C level in the diabetic patients will increase oxidative stress (Després *et al.*, 2000; Franceschini, 2001).

Body weight of diabetic rats

The presence of alloxan in the blood, was still debatable to be the main cause of losing weight of the treated rat. Though Diniz *et al.* (2008) reported the presence of alloxan was not affecting the rat weight, but several reports stated differently. Iranloye *et al.* (2011) and Raj *et al.* (2011) noted alloxan which is potential diabetogenic compound could change the non-diabetic rat to become diabetic ones as showed by their blood glucose level of 351.5 mg/dl, and therefore, called hyperglycemic.

The diabetic rat will show a symptom of consume more feed than non-diabetic rat, but they showed a reduction in body weight. This might be because of insulin deficiency and glucose is not able to enter the cells, leads to the use of lipid and protein as energy sources instead of glucose itself. Losing of body weight, is due to severe damages in protein of the tissues. Kamalakkannan and Prince (2006) supported in previous reports by saying dehydration process is related with situations of polydipsia and polyuria, which promoted by high level of lipid and protein catabolism. Weight loss in the diabetic is similar to

gluconeogenesis, where in this pathway, the muscle size will be smaller than normal and therefore, lose of protein in the tissues. Gradual losing of body weight to the diabetic patients which related with chronic hyperglycemia is also severe to infection, maintaining body weight is therefore, very important to the diabetic.

Current study was capable to control weight loss of the treated rat, where 7 days after intervention of ECLE, their body weight was increase as in the non-diabetic rat ($P = 0.22$). Figure 2 showed the body weight of treated rat of the diabetic and non-diabetic ones was similar after 14 days of intervention with ECLE ($P = 0.92$). The increase of body weight did not automatically means as obesity or overweight, and therefore, it might be conclude that the ECLE might be used to control body weight of the diabetic rat due to alloxan induction.

Winarsi *et al.* (2012) noted ECLE contains 129 mg/g flavonoid, capable to control body weight. Flavonoid which is also contained in the green tea was reported to have biological function; consuming this compound in high dose will therefore control body weight, adipose tissues, and regulate of lipid metabolism (Ashida *et al.*, 2004). Lee *et al.* (2011) supported this statement by saying several natural compounds like catechins, capsaicin, conjugated linoleic acids, fucoxanthin, isoflavone of soy bean, glabridin, astaxanthin, and cyanidin-3-glucoside was effective to maintain body weight through varies mechanism.

Glabridin, the main flavonoid content of licorice (*Glycyrrhiza glabra*), was known capable to maintain body weight of both treated animals and human by spinning up differentiation of adipose tissues. This type of flavonoid was effective in reducing intra-abdominal fat through activation of peroxisome proliferator-activated receptor- γ (PPAR- γ) (Nakagawa *et al.*, 2004). Astaxanthin, a natural carotenoid showed pharmacological effects like: anti-tumor, anti-cancer, anti-diabetes, and anti-inflammation. Astaxanthin was also noted to reduce the weight of liver, triacylglycerol levels in liver and plasma, and total cholesterol (Ikeuchi *et al.*, 2007). Astaxanthin could stimulate the use of fatty acid of the treated rat, though its function in reducing body weight is still not clear yet.

The ECLE was not containing only flavonoid but also vitamin C (Winarsi *et al.*, 2012), which was reported to affect lipid oxidation. Blaak and Saris (2002) reported oxidized lipids takes a role in obesity and failure in weight loss. Deficiency of vitamin C, therefore, could cause metabolic disorders and potential to affect body weight. There was a significant

paradox effects between the presence of vitamin C and gaining body weight. Means the sufficient amount of vitamin C in the blood plasma will reduce the obesity level. Vitamin C is a cofactor in carnitine biosynthesis, a molecule to cause oxidation of fatty acid. The higher consumption of vitamin C the more fatty acids will be oxidized, leads the contribution of reduction of adipose cells and therefore, losing body weight. The current study, might then conclude that the ECLE could repair body weight of the treated rat through its contents which were flavonoids and vitamin C.

Cholesterol level of diabetic rats

The high level of lipid in the serum of diabetic patient is mainly due to blockade in sensitivity of lipase by insulin and leads to mobilization of fatty acid from lipid stock in the peripheral. Hyperlipidemic, which is also known to be one among the characteristics of diabetic, as a consequence of non blocking activity of lipolytic enzymes in the peripheral lipid. Over content of fatty acids in the alloxan-induced treated rats, promoted the conversion of lipid compound in the liver to become phospholipid and cholesterol. However, these two compounds, together with high level of triglycerides could be thrown as lipoprotein in the blood, to cause readable total cholesterol is low.

After 7 days of observation, total cholesterol levels of diabetic rats treated ECLE were not significantly different from diabetic rat group which was given feed only ($P = 0.74$), but higher than that of non-diabetic rat group ($P = 1.61E-05$). Observation done in 7 days following to the first observation, noted a reduce in total cholesterol from 77.6 to 56 mg/dl ($P = 0.025$) of the diabetic rat group with ECLE treatment, this level was still higher than that of non-diabetic rat group ($P = 0.009$) (Figure 3).

Decreasing of total cholesterol level in the alloxan-induced treated rat was due to the high level of flavonoid contained in the ECLE. Several studies showed that the use of flavonoids reduce risk of cardiovascular disease (Graf *et al.*, 2005; Scalbert *et al.*, 2005) and atherosclerosis (Aviram and Fuhrman, 1998). Mechanism flavonoids inhibit lipid lesion development by reducing the formation of oxidized LDL. Fuhrman *et al.* (2000) also reported that polyphenol of flavonoid, both *in vitro* and *in vivo*, can reduce the number of macrophages and oxidized LDL. The role of flavonoids *in vitro* in oxidized LDL modification proves that high levels of known compounds in the ECLE can prolong the lag time of LDL oxidation.

Apart from, blocking the production of oxidized

LDL, flavonoid is capable suppress proliferation of plain muscle cells which normally take part in atherosclerosis, and reduce total cholesterol. Flavonoid that comes from a natural compounds, has also antioxidant characteristic and it function as scavenger of free radicals, along the redox reaction it is also donating hydrogen atom to the free radicals. It might be concluded that flavonoid has more potential than tocopherol.

The current study, however, has not identified the particular flavonoid or types of it that effective in reducing blood glucose level, reducing total cholesterol, and repairing body weight. This might be because of all components which include in the ECLE which were not soludified in water. Nakagawa *et al.* (2004) reported that hydrophobic flavonoids could reduce total amount of abdominal fat and causing hypoglycemic through activation of PPAR- γ . And therefore, that flavonoid is not only potential to be used as antioxidant but also to activate PPAR- γ , leads to anti-atherosclerotic, because the PPAR- γ also taking place in reducing cholesterol level of the macrophage of spone cells.

In a general situation, the diabetic has a disorders in lipid metabolism charaterized with a low level of HDL-cholesterol but high level of triglycerides. It has already reported that vitamin C can protect the HDL-cholesterol from oxidation process and therefore, might include in the process of reverse transport (Hillstrom *et al.*, 2003). A reverse transport process of cholesterol includes in excluding cholesterol before esterification in the membrane of extrahepatic cells, and takes the cholesterol to the esterification of lecithin to become acyltransferase. Ester cholesterol in lipoprotein would then retransferred into liver to be processed further and excreted through bile. The oxidized HDL might also modify structure of apolipoprotein AI. The potential of HDL lipoprotein in activating lecithin as acyltransferase cholesterol will cause blocking in esterification and excluding the extrahepatic cholesterol.

Conclusion, ethanolic cardamom leaves extract was capable to reduce blood glucose level as well as total cholesterol, and also repair body weight of alloxan-induced diabetic rats. In the future, the cardamom leaves could be used as functional foods/drinks compound for diabetic patients.

Acknowledgement

We would like to thank you the Directorate General for Higher Education, Ministry of National Education of the Republic of Indonesia for funding the MP3EI Research in year 2013.

References

- Alarcon-Aguilera, F. J., Roman-Ramos, R., Perez-Gutierrez, S., Aguilar-Contreras, A., Contreras-Weber, C. C. and Flores-Saenz, J. L. 1998. Study of the antihyperglycemic effect of plants used as antidiabetes. *Journal of Intercultural Ethnopharmacology* 61(2): 101-110.
- American Diabetes Association (ADA). 2006. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 29: 43-48.
- Anjaneyulu, M. and Chopra, K. 2004. Quercetin, an antioxidant bioflavonoid, attenuates diabetic nephropathy in rats. *Clinical and Experimental Pharmacology and Physiology* 31(4): 244-248.
- Ashida, H., Furuyashiki, T., Nagayasu, H., Bessho, H., Sakakibara, H., Hashimoto, T. and Kanazawa, K. 2004. Anti-obesity actions of green tea: Possible involvements in modulation of the glucose uptake system and suppression of the adipogenesis-related transcription factors. *Biofactors* 22: 135-140.
- Aviram, M. and Fuhrman, B. 1998. Polyphenolic flavonoids inhibit macrophage-mediated oxidation of LDL and attenuate atherogenesis. *Atherosclerosis* 137 (supplement): S45-S50.
- Avizeh, R., Najafzadeh, H., Pourmahdi, M. and Mirzaee, M. 2010. Effect of glibenclamide and fruit extract of zizyphus spina-christi on alloxan-induced diabetic dogs. *The International Journal of Applied Research in Veterinary Medicine* 8(2): 109-113.
- Bhathena, S.J. and Velasquez, M.T. 2002. Beneficial role of dietary phytoestrogens in obesity and diabetes. *American Journal Clinical Nutrition*. 76(6): 1191-201.
- Blaak, E. E. and Saris, W. H. M. 2002. Substrate oxidation, obesity and exercise training. *Best Practice and Research Clinical Endocrinology & Metabolism* 6(4): 667-678.
- Brahmachari, G. 2008. Anti-diabetic agents of natural origin: a retrospective account of some promising chemotypes. *Natural Product Communications* 3(8): 1337-1354.
- Brahmachari, G. 2009. In: *Natural Products: Chemistry, Biochemistry and Pharmacology*, Brahmachari, G. (Eds). Narosa Publishing House Pvt. New Delhi.
- Cazarolli, L. H., Folador, P., Moresco, H. H., Brighente, I. M. C., Pizzolatti, M. G. and Silva, F. R. M. B. 2009a. Stimulatory effect of apigenin-6-C- β -L-fucopyranoside on insulin secretion and glycogen synthesis. *European Journal of Medicinal Chemistry* 44(11): 4668-4673.
- Cazarolli, L. H., Folador, P., Moresco, H. H., Brighente, I. M. C., Pizzolatti, M. G. and Silva, F. R. M. B. 2009b. Mechanism of action of the stimulatory effect of apigenin-6-C-(2''-O- α -L-rhamnopyranosyl)- β -fucopyranoside on 14C-glucose uptake. *Chemicobiological interactions* 179(2-3): 407-412.
- Coskun, O., Kanter, M., Korkmaz, A. and Oter, S. 2005. Quercetin, a flavonoid antioxidant, prevents and

- protects streptozotocin-induced oxidative stress and β -cell damage in rat pancreas. *Pharmacological Research* 51(2): 117-123.
- Dakhale, G. N., Chaudhari, H. V. and Shrivastava, M. 2011. Supplementation of vitamin C reduces blood glucose and improves glycosylated hemoglobin in type 2 diabetes mellitus: a randomized, double-blind study. *Advances in Pharmacological Sciences*. ID 195271: 1-5. <http://dx.doi.org/10.1155/2011/195271>
- Dene, B. A., Maritim, A. C., Sanders, R. A. and Watkins, J. B. 2005. Effects of antioxidant treatment on normal and diabetic rat retinal enzyme activities. *Journal of Ocular Pharmacology and Therapeutics* 21(1): 28-35.
- Després, J. P., Lemieux, I., Dagenais, G. R., Cantin, B. and Lamarche, B. 2000. HDL-cholesterol as a marker of coronary heart disease risk: the Québec cardiovascular study. *Atherosclerosis* 153(2): 263-272.
- Dewanjee, S., Bose, S. K., Sahu, R. and Mandal, S. C. 2008. Anti-diabetic effect of matured fruits of *Dispyros peregrina* in alloxan-induced diabetic rats. *International Journal of Green Pharmacy* 2: 95-99.
- Diniz, S. F., Amorim, F. P. L. G., Cavalcante-Neto, F. F., Bocca, A. L., Batista, A. C., Simm, G. E. P. M. and Silva, T. A. 2008. Alloxan-induced diabetes delays repair in a rat model of closed tibial fracture. *Brazilian Journal of Medical and Biological Research* 41(5): 373-379.
- Elshater, A. A., Muhammad, M. A. S. and Mahrous, M. A. M. 2009. Effect of ginger extract consumption on levels of blood glucose, lipid profile and kidney functions in Alloxan induced-diabetic rats. *Egyptian Academic Journal Biology Science* 2 (1): 153-162.
- Franceschini, G. 2001. Epidemiologic evidence for high-density lipoprotein cholesterol as a risk factor for coronary artery disease. *The American Journal of Cardiology* 88(12A): 9N-13N.
- Fuhrman, B., Rosenblat, M., Hayek, T., Coleman, R. and Aviram, M. 2000. Ginger extract consumption reduces plasma cholesterol, inhibits LDL oxidation and attenuates development of atherosclerosis in atherosclerotic, apolipoprotein E-deficient mice. *Journal of Nutrition* 130(5): 1124-1131.
- Graf, B. A., Milbury, P. E. and Blumberg, J. B. 2005. Flavonols, flavones, flavanones, and human health: epidemiological evidence. *Journal of Medicinal Food* 8(3): 281-290.
- Hillstrom, R. J., Yacapin-Ammons, A. K. and Lynch, S. M. 2003. Vitamin C inhibits lipid oxidation in human HDL. *Journal of Nutrition* 133(10): 3047-3051.
- Hnatyszyn, O., Miño, J., Ferraro, G. and Acevedo, C. 2002. The hypoglycemic effect of *Phyllanthus sellowianus* fractions in streptozotocin-induced diabetic mice. *Phytomedicine* 9(6): 556-559.
- Ikeuchi, M., Koyama, T., Takahashi, J. and Yazawa, K. 2007. Effects of astaxanthin in obese mice fed a high-fat diet. *Bioscience, Biotechnology, and Biochemistry* 71(4): 893-899.
- Iranloye, B. O., Arikawe, A. P., Rotimi, G. and Sogbade, A. O. 2011. Anti-diabetic and anti-oxidant effects of *Zingiber officinale* on alloxan-induced and insulin-resistant diabetic male rats. *Nigerian Journal of Physiological Sciences* 26(1): 89-96.
- Johnston, K., Sharp, P., Clifford, M. and Morgan, L. 2005. Dietary polyphenols decrease glucose uptake by human intestinal Caco-2 cells. *Febs Letters* 579(7): 1653-1657.
- Jung, U. J., Lee, M. K., Jeong, K. S. and Chol, M. S. 2004. The hypoglycemic effects of hesperidin and naringin are partly mediated by hepatic glucose regulating enzymes in C57 BL/KsJ-db/db Mice. *Journal of Nutrition* 134(10): 2499-2503.
- Kalaivani, A., Umamaheswari, A., Vinayagam, A. and Kalaivani, K. 2008. Anti-hyperglycemic and antioxidant properties of *Cassia auriculata* leaves and flowers on Alloxan induced diabetic rats. *Pharmacology online* 1: 204-217.
- Kamalakkannan, N. and Prince, P. S. M. 2006. Antihyperglycaemic and antioxidant effect of rutin, a polyphenolic flavonoid, in streptozotocin induced diabetic Wistar rats. *Basic and Clinical Pharmacology and Toxicology* 98(1): 97-103.
- Kim, S. J., Ju, B. J., Choi, W. C. and Kin, C. S. 2006. Hypoglycemic and antihyperlipidemic effect of four Korean medical plants in Alloxan induce diabetic rats. *American Journal Biochemistry Biotechnology* 2(4): 154-160.
- Kim, J. S., Kwon, C. S. and Son, K. H. 2000. Inhibition of alpha-glucosidase and amylase by luteolin, a flavonoid. *Bioscience, Biotechnology, and Biochemistry* 64(11): 2458-2461.
- Knekt, P., Kumpulainen, J., Järvinen, R., Rissanen, H., Heliövaara, M., Reunanen, A., Hakulinen, T. and Aromaa, A. 2002. Flavonoid intake and risk of chronic diseases. *American Journal of Clinical Nutrition* 76(3): 560-568.
- Knekt, P., Ritz, J., Pereira, M. A., O'Reilly, E. J., Augustsson, K. and Fraser, G. E. 2004. Antioxidant vitamins and coronary heart disease risk: a pooled analysis of 9 cohorts. *American Journal of Clinical Nutrition* 80(6): 1508-1520.
- Lee, J., Li, Y., Li, C. and Li, D. 2011. Natural products and body weight control. *American Journal of the Medical Sciences* 3(1): 13-19.
- Lenzen, S. 2007. Alloxan and Streptozotocin Diabetes. Download from http://www.saw-leipzig.de/forschung/projekte/zeitstrukturen-endokriner-systeme/endokrinologie-iii/endo_07-lenzen.pdf. on 24/1/2013.
- Li, J. M., Che, C. T., Lau, C. B., Leung, P. S. and Cheng, C. H. K. 2006. Inhibition of intestinal and renal Na⁺-glucose cotransporter by naringenin. *International Journal of Biochemistry and Cell Biology* 38(5-6): 985-995.
- Likidilid, A., Natchai, P., Saiphon, P. and Thavatchai, P. 2007. Glutathione and glutathione peroxidase in type 1 diabetic patients. *Journal Medical Association Thailand* 90(9): 1759-1767.
- Liu, D., Zhen, W., Yang, Z., Carter, J. D., Si, H. and Reynolds, K. A. 2006. Genistein acutely stimulates insulin secretion in pancreatic β -cells through a cAMP-dependent protein kinase pathway. *Diabetes*

- 55(4): 1043-1050.
- Maechler P., Carobbio, S. and Rubi, B. 2006. In beta-cells, mitochondria integrate and generate metabolic signals controlling insulin secretion. *The International Journal of Biochemistry and Cell Biology* 38(5-6): 696-709.
- Mahesh, T. and Menon, V. P. 2004. Quercetin alleviates oxidative stress in streptozotocin-induced diabetic rats. *Phytotherapy Research* 18(2): 123-127.
- Matsui, T., Kobayashi, M., Hayashida, S. and Matsumoto, K. 2002. Luteolin, a flavone, does not suppress postprandial glucose absorption through an inhibition of alpha-glucosidase action. *Bioscience, Biotechnology, and Biochemistry* 66(3): 689-692.
- McRae, M. P. 2007. Vitamin C supplementation for treating hypercholesterolemia: a meta-analysis of 16 randomized controlled trials. *The journal of the American Nutraceutical Association* 10(2): 21-28.
- Middha, S. K., Bhattacharjee, B., Saini, D., Baliga, M. S., Nagaveni, M. B. and Usha, T. 2011. Protective role of *Trigonella foenum graecum* extract against oxidative stress in hyperglycemic rats. *European Review for Medical and Pharmacological Sciences* 15: 427-335.
- Mira, L., Fernandez, M. T., Santos, M., Rocha, R., Florencio, M. H. and Jennings, K. R. 2002. Interactions of flavonoids with iron and copper ions: a mechanism for their antioxidant activity. *Free Radical Research* 36(11): 1199-1208.
- Nakagawa, K., Kishida, H., Arai, N., Nishiyama, T. and Mae, T. 2004. Licorice flavonoids suppress abdominal fat accumulation and increase in blood glucose level in obese diabetic KK-Ay mice. *Biological and Pharmaceutical Bulletin* 27(11): 1775-1778.
- Pari, L. and Venkateswaran, S. 2002. Hypoglycaemic activity of *Scoparia dulcis* L. extract in alloxan induced hyperglycaemic rats. *Phytotherapy Research* 16(7): 662-664.
- Prince, P. S. M. and Kamalakkannan, N. 2006. Rutin improves glucose homeostasis in streptozotocin diabetic tissues by altering glycolytic and gluconeogenic enzymes. *The Journal of Biochemical and Molecular Toxicology* 20(2): 96-102.
- Raj, N., Nadeem, S., Jain, S., Raj, C. and Nandi, K. C. P. 2011. Ameliorative effects of *Alpinia calcarata* in alloxan-induced diabetic rats. *Digest Journal of Nanomaterials and Biostructures* 6 (3): 991-997.
- Scalbert, A., Manach, C., Morand, C., Rémésy, C. and Jiménez, L. 2005. Dietary polyphenols and the prevention of diseases. *Critical Reviews in Food Science and Nutrition* 45(4): 287-306.
- Shirwaikar, A., Rajendran, K., Kumar, C. D. and Bodla, R. 2004. Antidiabetic activity of aqueous leaf extract of *Annona squamosa* in streptozotocin-nicotinamide type 2 diabetic rats. *Journal of Ethnopharmacology* 91(1): 171-175.
- Suarsana, I. N., Utama, I. H., Agung, I. G. and Suartini, A. 2011. Effect of hyperglycemia and vitamin E on malondialdehyde and antioxidant enzymes intracellular levels of rat pancreatic tissue. *Bandung Medical Journal* (43) 2: 72-26. (In Indonesia).
- Szkudelski, T. 2001. The mechanism of alloxan and streptozotocin action in B cells of the rat pancreas. *Physiological Research* 50: 536-546.
- Winarsi, H. and Purwanto, A. 2010. Soy germed protein plus Zn as an inducer insulin secretion on type-2 diabetes mellitus. *Hayati Journal of Biosciences* 17(3): 120-124.
- Winarsi, H., Sasongko, N. D., Purwanto, A., Arinton, I. G. and Nuraeni, I. 2012. *In Vitro* antioxidant activity of the stem and leaves *Amomum cardamomum* extracts. *Proceedings of International Conference on Medicinal Plants 2012*. Purwokerto: The 43rd National Meeting of National Working Group on Indonesia Medicinal Plant.
- Winarsi, H., Sasongko, N. D., Purwanto, A. and Nuraeni, I. 2013. Cardomum Leaves Extract Reduced Oxidative Stress Level in plasma alloxan-induced diabetic rats. *Proceedings of 13th Asean Food Conference*. Singapore: Meeting Future Food Demands, Security and Sustainability.
- Yuji, T., Tatsumasa, M., Mitsuaki, K., Yoshiro, S., Hideyuki, I. and Kaku, N. 2006. Licorice flavonoid oil effects body weight loss by reduction of body fat mass in overweight subjects. *Journal of Health Sciences* 52(6): 672-683 .
- Zhao, H., Yakar, S., Gavrilova, O., Sun, H., Zhang, Y., Kim, H., Setser, J., Jou, W. and Leroith, D. 2004. Phloridzin improves hyperglycemia but not hepatic insulin resistance in a transgenic mouse model of type 2 diabetes. *Diabetes* 53(11): 2901-2909.