

p-ISSN 2085-191X (print)  
e-ISSN 2338-7610 (online)

# Biosaintifika

Journal of Biology & Biology Education



Nationally Accredited by Ministry of Research and Technology 2019-2024

Biosaintifika	Volume 12	Number 3	Pages 288-487	Semarang December 2020	p-ISSN 2085-191X (print) e-ISSN 2338-7610 (online)
---------------	-----------	----------	---------------	---------------------------	---



## AIMS AND SCOPE

Biosaintifika [Nationally Accredited by Ministry of Research, Technology and Higher Education, Number 2/E/KPT/2015 periode 2015-2020], an electronic international journal, provides a forum for publishing the original research articles from contributors, and the novel technology news related to biological research and biological education.

Scientific articles dealing with botany, zoology, environmental, and biotechnology, etc. are particularly welcome. This journal encompasses original research articles including: botany; zoology; environmental; biotechnology; ethnobotany; ecotourism; biochemistry; biology education.

## PUBLICATION INFORMATION

Biosaintifika [p-ISSN 2085-191X | e-ISSN 2338-7610] has been published three a year, April, August and December by Department of Biology, Faculty of Mathematics and Sciences, Universitas Negeri Semarang, Indonesia collaborate with Perpustakaan Biologi Indonesia.

This journal has been a CrossRef Member [DOI 10.15294/biosaintifika] since 2014, so that all articles published by this journal have DOI unique numbers. This journal has been abstracted and distributed by EBSCO Publishing started from Volume 6 Number 2 Year 2014 to present. This journal has been a DOAJ since 2015, CABI since 2016 and ASEAN Citation Index since 2018.

## JOURNAL CITATIONS AND IMPACT FACTOR

Google Scholar  
h-index : 13  
i10-index : 31  
h5-index : 9  
h5-median : 11

Total articles in Google Scholar : 373 articles (since 2012)  
Total citations in Google Scholar : 1238 articles (since 2012)  
Total Citedness of Journal Articles in SCOPUS : 212 articles



For best and  
innovative research  
papers, click

**journal.unnes.ac.id**

## SUBSCRIPTION

Annual Subscription (3 issues: April, August & December). Combined Institutional Rate (print + electronic): Rp. 1.500.000,-. Please send your application to e-mail biosaintifika@mail.unnes.ac.id with :

Subject : Subscription

Content : Name, Address, Receipt of Payment

Payment to Bank Negara Indonesia (BNI)

Account Number 04 300 222 75 name Lutfia Nur Hadyanti

## JOURNAL HISTORY

Biosaintifika was established in 2009. The title with its subtitle was Biosaintifika: Berkala Ilmiah Biologi. Started from Volume 5, Number 2 Year 2013, the journal was only published by using one language namely English. The name of the journal with the sub-title was changed to Biosaintifika: Journal of Biology & Biology Education.





## Editorial Team

### Editor-in-Chief

1. Dyah Rini Indriyanti, [SCOPUS ID: 57194044439] Universitas Negeri Semarang, Indonesia

### International Editorial Board

1. Fahrul Huyop, [SCOPUS ID: 10240240100] Universiti Teknologi Malaysia, Malaysia
2. Mulyoto Pangestu, [SCOPUS ID: 35488897500] Monash University, Australia
3. Mukh Syaifudin, [SCOPUS ID: 6507831675] Badan Tenaga Nuklir Nasional, Indonesia
4. Yohanes Budi Widianarko, [SCOPUS ID: 6507055401] Soegijapranata Catholic University, Semarang, Indonesia
5. Amir Hamidy, [SCOPUS ID: 24341129500] Research Center For Biology-Cibinong Science Center, Indonesia
6. S. Sudarsono, [SCOPUS ID: 16450098400] Institut Pertanian Bogor, Indonesia
7. Iin Kurnia, [SCOPUS ID: 57193537585] Badan Tenaga Nuklir Nasional, Indonesia
8. Mohammad Irham, [SCOPUS ID: 55312221800] Research Center for Biology-Indonesian Institute of Sciences, Bogor, Indonesia

### Associate Editors

1. Talitha Widiatningrum, [SCOPUS ID: 56798544200] Universitas Negeri Semarang, Indonesia
2. Margareta Rahayuningsih, [SCOPUS ID: 57094066700] Universitas Negeri Semarang, Indonesia
3. Amin Retnoningsih, [SCOPUS ID: 51864647000] Universitas Negeri Semarang, Indonesia
4. Enni Suwarsi Rahayu, [SCOPUS ID: 56644635000] Universitas Negeri Semarang, Indonesia
5. Niken Subekti, [SCOPUS ID: 57195315756] Universitas Negeri Semarang, Indonesia



This work is licensed under a Creative Commons Attribution 4.0 International License.



### ABOUT THE JOURNAL

- [Aims and Scope](#)
- [Publication Ethics](#)
- [Indexing & Abstracting](#)
- [Editorial Team](#)
- [Reviewer Team](#)
- [Contact](#)

### COLLABORATE WITH



### FOR AUTHORS


- [Guidelines for Author](#)
- [Manuscript Template](#)
- [Mendeley Guidelines](#)
- [Online Submission](#)
- [Google Citation Analysis](#)
- [Scopus Citation Analysis](#)

### Readers

ID 228,630	GB 705
US 17,930	IR 676
IN 3,616	PK 641
PH 3,148	VN 596
MY 2,939	AU 575
TH 1,481	MX 521
CN 1,296	CA 477
BR 1,092	EG 470
JP 1,015	FR 463
SG 803	KR 449
TR 795	DE 433

Pageviews: 743,441

Flags Collected: 187



00680320 [View My Stats](#)

### USER

Username

Password

☐ Remember me

### JOURNAL CONTENT

Search

Search Scope




[Home](#) > **Indexing & Abstracting**

## Indexing & Abstracting

Biosaintifika: Journal of Biology & Biology Education has been covered by the following services:

1. Science and Technology Index (SINTA) 2
2. Google Scholar
3. DIMENSIONS
4. DOAJ
5. ASEAN Citation Index
6. Indonesian Publication Index
7. Indonesian Scientific Journal Database
8. Microsoft Academic
9. EBSCO Open Science Directory
10. EBSCO Academic Search Complete
11. National Library of Medicine (NLM)
12. CAB Abstract (CABI)
13. DOI Crossref
14. ROAD
15. BASE
16. Sherpa/Romeo
17. Ulrichsweb.com™ (Proquest)
18. WorldCat
19. Summon™ from Serials Solutions
20. Hinari
21. Mendeley
22. Citeulike
23. ResearchGate
24. Scilit



This work is licensed under a Creative Commons Attribution 4.0 International License.



### ABOUT THE JOURNAL

[Aims and Scope](#)
[Publication Ethics](#)
[Indexing & Abstracting](#)
[Editorial Team](#)
[Reviewer Team](#)
[Contact](#)

### COLLABORATE WITH



### FOR AUTHORS


[Guidelines for Author](#)
[Manuscript Template](#)
[Mendeley Guidelines](#)
[Online Submission](#)
[Google Citation Analysis](#)
[Scopus Citation Analysis](#)

### Readers

ID 228,630	GB 705
US 17,930	IR 676
IN 3,616	PK 641
PH 3,148	VN 596
MY 2,939	AU 575
TH 1,481	MX 521
CN 1,296	CA 477
BR 1,092	EG 470
JP 1,015	FR 463
SG 803	KR 449
TR 795	DE 433

Pageviews: 743,441

Flags Collected: 187



**00680321** [View My Stats](#)

### USER

Username

Password

☐ Remember me

### JOURNAL CONTENT

Search

Search Scope   
 All



## TABLE OF CONTENTS

Volume 12, Number 3, December 2020

- 288-296 **Stability and Antibacterial Property of Polyherbal Mouthwash Formulated Using Local Ingredients**  
Juman Nafea, Mohamed Edbeib, Kin Israel R. Notarte, Fahrul Huyop, Harisun Yaakub
- 297-303 **The Antioxidant Content and Activity of Various Plant Organs of Kitolod (*Isotoma longiflora*)**  
Grahita Rezhi Egarani, Sri Kasmiyati, Elizabeth Betty Elok Kristiani
- 304-310 **Identification of Gamma-Amino Butyric Acid Isolates Lactic Acid Bacteria Results from The Isolation of Rusip**  
S Feta Avila Prayitno, Agung Suprihadi, Endang Kusdiyantini
- 311-318 **Selection of Soybean (*Glycine max*) Germplasm Against Biotrophic Fungi Disease Based on Anatomical Resistance**  
Siti Samiyarsih, Ade Yuanita Putri Pratiwi, Juni Safitri Muljowati, Nur Fitrianto
- 319-328 **Carotenoid Production by *Rhodospiridium paludigenum* Using Orange Peel Extract as Substrate**  
Renna Eliana Warjoto, Jennifer Jennifer, Bibiana Widyawati Lay
- 329-334 **Nutritional Content of Bone Flour Made from Plecos Fish *Pterygoplichthys pardalis* from the Ciliwung River, Indonesia**  
Handhini Dwi Putri, Dewi Elfidasari, Haninah Haninah, Irawan Sugoro
- 335-342 **Potential of Ecology based Agrotourism for Agricultural Education in Besur Village, Lamongan Regency**  
Slamet Ifandi, Yustika Aulia Rahma
- 343-349 **Qualitative Tests of Secondary Metabolite Compounds in Ethanol Extract of *Spirulina platensis* from Karimun Jawa Sea, Indonesia**  
Renni Yuniati, Muhammad Zainuri, Hermin Kusumaningrum
- 350-355 **Antioxidant Activity of Jojoba (*Simmondsia chinensis*) Seed Residue Extract**  
Arnold Patogi Siahaan, Eti Rohaeti, Ali Mahmoud Muddathir, Irmanida Batubara
- 356-362 **Total Phenol Content of *Avicennia marina* Leaf and Its Relationship to the Environmental Quality**  
Endah Dwi Hastuti, Munifatul Izzati, Sri Darmanti
- 363-369 **Larvicidal Activity of *Brugmansia candida* against *Aedes aegypti* and *Culex quinquefasciatus* (Diptera: Culicidae)**  
Setiawan Khoirul Himmi, Didi Tarmadi, Dita Meisyara, Anugerah Fajar, Titik Kartika, Ikhsan Guswenrivo, Sulaeman Yusuf
- 370-377 **Histopathological of White Rats Aorta Induced by High-Fat Feed After Administered by Neem Leaf Ethanolic Extract**  
Baiq Miftahul Jannah, Sri Isdadiyanto, Agung Janika Sitasiwi



- 378-383 ***Bifidobacterium longum*, a Predominant Bifidobacterium in Early-life Infant Potentially Used as Probiotic**  
Dyah Fitri Kusharyati, Anwar Rovik, Dini Ryandini, Hendro Pramono
- 384-391 **Alkaline Phosphatase Expression From Mice Mesenchymal Stem Cells Induced By Flamboyant Flower (*Delonix regia*) Extract**  
Kartini Eriani, Deby Anggraini, Yudha Bintoro, Ichsan Ichsan, Al Azhar, Silmi Mariya
- 392-398 **Science Web-Module Integrated with Tannery Waste as Local Potential to Improve Students' Problem Solving**  
Atznain Iqma Miswami, Heru Nurcahyo
- 399-407 **Morphological Variation of Blue Panchax (*Aplocheilichthys panchax*) Lives in Different Habitat Assessed Using Truss Morphometric**  
Diah Mustikasari, Suhestri Suryaningsih, Agus Nuryanto
- 408-413 **Comparison of Nutritional Content of *Mangifera indica* L. var. Wirasangka with Three Other Varieties in Tegal Regency**  
Enni Suwarsi Rahayu, Amin Retnoningsih, Muhammad Abdullah, Priyanti Susanti
- 414-421 **Ethnobotanical Study of Early Childhood Medicinal Plants Used by the Local People in South Bangka Regency, Indonesia**  
Henri Henri, Vitriany Nababan, Luchman Hakim
- 422-430 **Anti-tumor Potential Ethnomedicinal Plants from Dayak Tribes in Central Kalimantan**  
Liswara Neneng, Asri Pujirahadju, Ferayani Ferayani, Tri Widayat
- 431-437 **Blood Hematological and Biochemical Parameters of *Osteochilus vittatus* with *Spirulina platensis* Supplementation in Biofloc System**  
Muhammad Azharul Rijal, Susanto Susanto, Sorta Basar Ida Simanjuntak, Hernayanti Hernayanti, Purnama Sukardi
- 438-445 **Biochemical and Histopathology Analysis of Liver Damage in Hypercholesterolemic Rats Induced by Tomato Extract**  
Retno Sri Iswari, Muchamad Dafip, Muhammad Rifa'i
- 446-452 **Characteristics of DNA Barcodes from Three *Thrixspermum* Orchids Based on ITS2 Regions**  
Siti Rohimah, Tri Ratnasari, Mukhamad Su'udi
- 453-458 **Plantlet Formation and Acclimatization of Sugarcane cv. PS 881 with Different Types and Concentration of Auxin**  
Rendie Prasetyo, Sugiyono Sugiyono, Elly Proklamasiningsih, Prita Sari Dewi
- 459-468 **Isolation of Potential Plant Growth Promoting Rhizobacteria (PGPR) from Cassava (*Manihot esculenta*) Rhizosphere Soil**  
Suri Raihan Safriani, Lenni Fitri, Yulia Sari Ismail
- 469-477 **In Situ Bioremediation Strategies for the Recovery of Mercury-contaminated Land in Abandoned Traditional Gold Mines in Indonesia**  
Winardi Winardi, Eko Haryono, Sudrajat Sudrajat, Endang Sutariningsih Soetarto
- 478-487 **Feed Resources Determination Based on Pollen Diversity in Trigona Bees (*Trigona* sp.) Colony**  
Muachiroh Abas, Eddy Tri Sucianto

# Morphological Variation of Blue Panchax (*Aplocheilichthys panchax*) Lives in Different Habitat Assessed Using Truss Morphometric

Diah Mustikasari, Suhestri Suryaningsih, Agus Nuryanto\*

Faculty of Biology, Universitas Jenderal Soedirman, Indonesia

\*Corresponding author: [agus.nuryanto@unsoed.ac.id](mailto:agus.nuryanto@unsoed.ac.id)

Submitted: 10 October 2020. Revised: 4 November 2020. Accepted: 21 December 2020

**Abstract.** Blue panchax (*Aplocheilichthys panchax* Hamilton, 1822) lives in broad ranges of habitat from open waters to closed waters, including at ex-tin mining pits in Bangka Island, Indonesia. Variable habitats might cause morphological variations due to different ecological factors. This study aimed to assess the morphological variation of blue panchax collected from different habitats using truss morphometric characters. Fish samples (70 individuals) were collected from abandoned ex-tin mining pits of different ages and a river in Bangka Island. Twenty-nine truss characters were analyzed using the Kruskal-Wallis test and post hoc with Dunn's test. The results showed that almost all of the body parts of blue panchax found in ex-tin mining pits and rivers were significantly different ( $p$ -value  $< 0.05$ ), except some truss characters of caudal peduncle. The post hoc of Dunn's test showed a positive correlation between habitats and truss characters (adj. sig  $< 0.05$ ). Some truss characters of killifish were different between ex-tin mining pits with different ages chronosequence, and some other characters were different between pits and river. This study provides the first data about the morphological variation of blue panchax in ex-tin mining pits of different ages. The data is valuable as a scientific basis of further utilization of ex-tin mining pits in the areas.

**Key words:** *Aplocheilichthys panchax*; morphometry; pits; river

**How to Cite:** Mustikasari, D., Suryaningsih, S., & Nuryanto, A. (2020). Morphological Variation of Blue Panchax (*Aplocheilichthys panchax*) Lives in Different Habitat Assessed Using Truss Morphometric. *Biosaintifika: Journal of Biology & Biology Education*, 12(3), 399-407.

**DOI:** <http://dx.doi.org/10.15294/biosaintifika.v12i3.26593>

## INTRODUCTION

Blue panchax (*Aplocheilichthys panchax*), locally known as ikan Kepala Timah, is one of the killifish species from the Order Cyprinodontiformes, Family Aplocheilichthyidae, and Genus *Aplocheilichthys*. Member of Genus *Aplocheilichthys* is widely distributed across the Indo-Malayan Islands, including Indonesia, the Indo-China region, and India (Vasil'eva et al., 2013; Dekar et al., 2018). *Aplocheilichthys panchax* is a species of the genus *Aplocheilichthys*. It is an endemic species to the Oriental Region (Costa, 2013; Sedlacek et al., 2014; Furness, 2015; Costa, 2016; Beck et al., 2017).

Blue panchax can live in a broad range of habitats (Manna et al., 2011). It can survive in open and closed waters such as lakes or pits of ex-tin mining, including newly formed and old pits. According to Kurniawan et al. (2019) and Irawan et al. (2014), the newly formed ex-tin mining pits are extreme ecosystems with very low pH values and dissolved oxygen (DO), but with high heavy metal content. Conversely, the older abandoned tin mining pits have a better water quality. Nevertheless, a previous study by Kurniawan (2019) had proved that blue panchax was

reported to live in newly abandoned tin mining pits in Bangka Island though have deplorable water quality conditions.

A study had shown that ecological characteristics have impacted fish genotype (Nguyen et al., 2017) and have a further effect on their morphology (Baillie et al., 2016; Endo & Watanabe, 2020). Other studies also proved that fish live in different habitats, showed variable morphologies, and, in extreme condition, might form different ecotypes (Rajeswari et al., 2017). Morphological variation among individual fish can be assessed using conventional and truss morphometric characters (Pazhayamadom et al., 2014; Mojekwu and Anumudu, 2015; Rawat et al., 2017). According to Ariyanto et al. (2011), truss morphometric provides a comprehensive, systematic, and fairly high-accuracy geometric picture of fish body shapes. So, this method can be used to distinguish between individual fish more precisely than standard morphometric. It has been proven that truss morphometric is an efficient technique to differentiate fish individuals than conventional morphometric (Ihya et al. 2020; Nabila et al. 2019; Pambudi et al. 2019)



It is assumed that different ecological factors among different ages of ex-tin mining pits and rivers in Bangka might cause morphological differences among blue panchax collected from such a diverse ecosystem. There is no study assessing the morphological variation of blue panchax inhabits different ages of abandoned tin mining pits and rivers in Bangka Province. The only research was about the existence and factors affecting blue panchax in the abandoned ex-tin mining pits (Mustikasari et al., 2020; Kurniawan et al., 2019). Therefore, this is the first research about the morphological variation of blue panchax live in different ages of ex-tin mining pits and river in Bangka Island. The data is vital for the management of the further utilization of ex-tin mining pits in the areas. This study aimed to assess the morphological variation of blue panchax collected from different habitats using truss morphometric characters.

## METHODS


### Study sites

The study was conducted in Pangkalpinang City and Bangka Regency of Bangka Belitung Archipelago Province, Indonesia (Figure 1). Fish samples were collected from ex-tin mining pits with chronosequences abandoned after mining activities. The pits were clustered into six different ages, i.e., Station A and Station B (< 5 years old), Station C and Station D (5-15 years), Station E and Station F (15-25 years), Station G (25 - 50 years), Station H (50 - 100 years), Station I and Station J (> 100 years), while Station K was Limbung River Stream.

Station I and Station J (> 100 years), and Limbung River Stream of Bangka Regency as Station K. The sampling site condition is shown in Figure 2.

### Sample collection and preservation

The 70 fish samples were collected at 09.00 am - 1.00 pm from ex-tin mining pits and a river using nets with mesh size of about 0.4 mm. Fresh individuals were placed in the labeled plastics bottle filled with 40% formalin. For permanent preservation, the samples were preserved with absolute ethanol. In the laboratory, the morphometric characters were measured by a ruler with an accuracy of 0.5 mm.



**Figure 1.** Map of research stations. Station A and Station B were pits with age < 5 years, Station C and Station D (5-15 years), Station E and Station F (15-25 years), Station G (25-50 years), Station H (50-100 years), Station I and Station J (> 100 years), while Station K was Limbung River Stream.




**Figure 2.** Waters condition of research stations, (a) Station A and (b) Station B were pits with age < 5 years; (c) Station C and (d) Station D (5-15 years); (e) Station E and (f) Station F (15-25 years); (g) Station G (25-50 years); (h) Station H (50-100 years); (i) Station I and (j) Station J (> 100 years); and (k) Station K was Limbung River Stream (private documentations).



### Morphometric measurement

Fish morphology was measured using truss network measurement. Truss morphometric was used to measure 29 diagonal distances among truss points and the truss characteristics encoded from A1 to D5 (Figure 3), while the description of each truss characteristics is presented in Table 2. The truss morphometric characters were analyzed using the Kruskal-Wallis test in SPSS Program version 25 to know significant differences of truss characters among individuals collected at different habitats.

Dunn's test was used for the post hoc of Kruskal-Wallis.



**Figure 3.** The truss network characteristics of Kepala Timah fish (*Apocheilus panchax*) (private documentations).

**Table 1.** Truss characteristics of Kepala Timah fish (*A. panchax*) and their descriptions

Part of Body	Code	Descriptions
Head	A1 (1 to 2)	distance between the snout or premaxilla and the pelvic maxilla (lower jaw)
	A2 (1 to 3)	distance between the snout and dorsal maxilla or anterior eye diameter (upper jaw)
	A3 (1 to 4)	distance between the snout and the pelvic operculum
	A4 (1 to 5)	distance between the snout and the dorsal operculum
	A5 (2 to 3)	distance between pelvic maxilla and dorsal maxilla or anterior eye diameter
	A6 (2 to 4)	distance between the pelvic maxilla and the pelvic operculum
	A7 (2 to 5)	distance between the pelvic maxilla and the dorsal operculum
	A8 (3 to 4)	distance between the dorsal maxilla or anterior eye diameter and the pelvic operculum
	A9 (3 to 5)	distance between the dorsal maxilla or anterior eye diameter and the dorsal operculum
	A10 (4 to 5)	distance between the pelvic operculum to the dorsal operculum
Anterior Body	B1 (4 to 6)	distance between the pelvic operculum and lower body-pectoral fin
	B2 (4 to 7)	distance between the pelvic operculum and the anterior dorsal fin
	B3 (5 to 6)	distance between the dorsal operculum and lower body-pectoral fin
	B4 (5 to 7)	distance between the dorsal operculum and the anterior dorsal fin
	B5 (5 to 8)	distance between the dorsal operculum and ventral or pelvic fin
	B6 (6 to 8)	distance between the lower body-pectoral fin and ventral or pelvic fin
	B7 (7 to 8)	distance between the anterior dorsal fin and ventral or pelvic fin
Posterior Body	C1 (7 to 9)	distance between the anterior dorsal fin and anterior anal fin
	C2 (7 to 10)	distance between the anterior and the posterior dorsal fin
	C3 (7 to 11)	distance between the anterior dorsal fin and posterior anal fin
	C4 (8 to 10)	distance between the ventral or pelvic fin and the posterior dorsal fin
	C5 (9 to 10)	distance between the anterior anal fin and the posterior dorsal fin
	C6 (9 to 11)	distance between anterior and posterior anal fin
	C7 (10 to 11)	distance between the posterior dorsal fin and rear anal fin
Caudal Peduncle	D1 (10 to 12)	distance between the posterior dorsal fin and pelvic-posterior caudal peduncle
	D2 (10 to 13)	distance between the posterior dorsal fin and dorsal-posterior caudal peduncle
	D3 (11 to 12)	distance between the posterior anal fin and pelvic-posterior caudal peduncle
	D4 (11 to 13)	distance between the posterior anal fin and dorsal-posterior caudal peduncle
	D5 (12 to 13)	the caudal peduncles' height


### RESULTS AND DISCUSSION

The general morphology of killifish collected from different aged ex-tin mining pits and river is shown in Figure 4.. It can be realized from Figure 4 that killifish individuals collected at the different ecosystems showed different colorations. Killifish individuals live in abandoned tin mining pits (Figure

4a-d) have brighter colors than fish in the river (Figure 4e). It can also be seen in Figure 4 that among different pits ages, killifish individual shows different body color. Nevertheless, body-color brightness does not correlate with pits ages. It is shown in Figure 4b that killifish collected from older than 5 to 15 years old pits has the brightest body color and more colorful than individuals collected in less than one to 5



years and older than 15 years pits (Figure 4a, 4c, and 4d).



**Figure 4.** General morphology of killifish collected from different habitats. (a) pits with age of < 5 years, (b) pits with age of 5-15 years, (c) pits with age of 15-25 years, (d) pits with age of 25-50 years, (e) Limbung River.

It is suggested that morphological differences among individuals of killifish live in different habitats is because of differences in ecological factors among the habitats. That argument rose based on the fact that sampling locations have distinct environmental parameters (Table 3). According to Nguyen et al. (2017), ecological factors might affect fish genotype, and according to Baillie et al. (2016) and Endo and Watanabe (2020), ecological characteristics have a further effect on fish morphology. Other studies also proved that fish live in different habitats show variable morphologies, and in extreme condition might form different ecotypes (Rajeswari et al., 2017).

Truss analysis was a measurement based on the ratio of truss character with head length or standard length. According to Paknejad et al. (2014), the truss morphometric network study effectively provides information about an organism's shape. It can describe the body shape of the fish more properly. Truss morphometric is a reliable method for morphological differentiation among fish samples. Therefore, it is expected that truss morphometric analysis could differentiate killifish collected from different habitats, such as among different aged ex-tin mining pits and between ex-tin mining pit habitats as closed waters with the river as open waters. The results of Dunn's test among truss morphometric characters of killifish from different habitats are presented in Table 2.

As can be seen in Table 2, that almost all of the truss characters of killifish showed significant differences ( $p$ -value < 0.05) among habitats, except on four characters (D1, D2, D4, and D5) that are parts of the caudal peduncle. These four characters indicated that the caudal peduncle could not differentiate among individuals collected at different habitats. The other

parts of the blue panchax fish's body showed significant differences based on the Kruskal-Wallis test. Dunn's test showed that some of these truss characters showed a positive correlation between different locations (adj. sig. < 0.05). The truss characters of A1, A5, B3, B7, C1, C2, C3, and C5 only showed the differences between ex-tin mining pits with different ages chronosequence. The truss characters of A2, A3, A4, A6, A7, A8, A9, A10, B1, B2, B4, B5, B6, C4, C6, C7, and D3 described the differences among pits with different ages and also between pits as closed water and the river as representative of open water.

The differences in morphometric among killifish can be caused by habitat characteristics where they live. They might differ in biotic and abiotic factors such as food availability, salinity, temperature, radiation, water depth, current flow, and other environmental factors that can affect the morphometric of the fish in each location (Sen et al., 2011; Kashefi et al., 2012; Muchlisin, 2013). Besides, the condition of ex-tin mining pits as closed water containing many elements such as metals and heavy metals (Kurniawan, 2017; Kurniawan et al., 2019) might contribute to the metabolism process impacted to morphometric characters of fish. It can happen in Bangka Island rivers too that the contamination of elements contributed to biological activity and morphometric characters of organisms in the waters. According to Lestari et al. (2018), human activities directly affecting rivers' water quality and freshwater fish diversity. According to Nuryanto et al. (2012) and Nuryanto et al. (2015), the existence of fish in rivers could be due to the different Physico-chemical characters of the rivers, especially its dissolved oxygen, carbon dioxide level, temperature, acidity (pH), substrates, organic and inorganic materials, water volume, the width of the river, and geology factor. These factors will naturally gradually change the microhabitat and impact of fish's diversity and characteristics in the river. According to Sari and Zakaria (2017), physical aspects (e.g., temperature), chemical factors (e.g., dissolved oxygen and acidity of the water), and biological factors (e.g., amount and type of food or food availability) can influence morphology and sexuality of fish.

Therefore, the differences in ecological characteristics can affect morphological characteristics (Baillie et al., 2016; Endo and Watanabe, 2020). Habitat variation affecting fish's community structures and previous studies had proven a positive correlation between fish morphology and their environment (Gebrekiros, 2016). The correlation between ecological parameters and morphological characters explains that the water quality, especially pH value in ex-tin mining pits or river, can influence the morphometric standard and truss characters where the low water



quality can inhibit the fish growth. These conditions can be seen in ex-tin mining pits that pits with age ranged between 1 and 5 years old was acidic, while pits with age between 25 and 50 years old was neu-

tral. The neutral pH can support biological life such as, plankton, and improves the physicochemical parameters, which in turn could promote fish's growth.

**Table 2.** The results of the Kruskal-Wallis test and Dunn's test of truss characteristics

Truss Parameters	Mean	Sig. of Kruskal-Wallis Test	Kruskal-Pairwise comparison from different research stations (adj. sig < 0.05 of Dunn's Test)***
A1*	0.35840.000		A-E (.009); A-F (.013); E-G (.016); F-G (.026)
A2*	0.30940.000		D-G (.038); E-F (.010); E-G (.038); E-K (.000); F-G (.005); G-K (.000)
A3*	0.68990.000		A-F (.001); A-K (.008); C-F (.002); C-K (.018); E-F (.000); E-K (.001); D-F (.025)
A4*	0.93930.000		A-D (.018); A-E (.005); A-G (.000); B-G (.010); E-K (.034); F-G (.002); G-K (.000)
A5*	0.30310.000		D-E (.001); D-F (.000)
A6*	0.34400.000		A-F (.036); C-F (.001); C-K (.006); E-F (.000); E-G (.035); E-K (.001); D-F (.040)
A7*	0.41990.000		A-F (.028); A-G (.000); B-G (.000); C-G (.012); D-G (.001); E-G (.001); G-K (.018)
A8*	0.53140.000		D-E (.024); D-F (.030); D-K (.003)
A9*	0.66870.000		A-F (.001); B-F (.025); D-F (.001); D-K (.020); E-F (.000); E-K (.008)
A10*	0.62960.000		D-G (.006); D-K (.001); E-F (.016); E-G (.000); E-K (.000)
B1*	0.13160.002		D-K (.042)
B2*	0.62570.000		A-E (.010); A-K (.000); B-E (.036); B-K (.000); C-K (.000); D-K (.046); G-K (.008)
B3*	0.18860.000		A-G (.024); D-G (.001); E-G (.008); F-G (.014)
B4*	0.56360.001		C-K (.021); F-K (.002)
B5*	0.29230.003		G-K (.044)
B6*	0.17910.000		A-D (.024); B-D (.027); C-G (.003); D-E (.002); D-F (.028); D-G (.000); G-K (.002)
B7*	0.40530.002		E-F (.000)
C1*	0.27760.000		A-C (.005); A-G (.002); C-D (.046); D-G (.014)
C2*	0.07010.000		A-G (.001); B-G (.001); D-G (.008)
C3*	0.29500.000		A-C (.021); A-G (.002); D-G (.009); E-G (.042)
C4*	0.44100.000		D-F (.042); E-F (.030); E-G (.036); F-K (.008); G-K (.010)
C5*	0.31200.000		A-G (.009); C-D (.043); C-E (.031); D-G (.001); E-G (.001)
C6*	0.24400.000		A-G (.001); A-K (.000); C-G (.037); C-K (.014); D-G (.013); D-K (.004); E-G (.049); E-K (.019)
C7*	0.14210.000		A-G (.022); B-G (.005); C-G (.009); D-G (.002); E-G (.016); G-K (.001)
D1**	0.17340.099		
D2**	0.12940.206		
D3*	0.15860.026		B-K (.004)
D4**	0.21110.557		
D5**	0.12640.160		

\*= sig. < 0.05 means significant differences of the parameters by Kruskal-Wallis test, and then they were continued to post hoc analysis with Dunn's test

\*\*=sig. > 0.05 means no significant differences of the parameters by Kruskal-Wallis test, and then they weren't continued to post hoc analysis with Dunn's test

\*\*\*=pairwise comparisons of pits with adj. sig. < 0.05 of Dunn's test means the significant differences of two pits for each truss parameter. The Bonferroni correction for multiple tests has adjusted the significance value.

Killifish body length differences among pits with chronosequence ages are positively correlated with the pits' water quality. The first indicator of water quality in ex-tin mining pits is pH value, where pit

with age of < 10 years has an acidic condition with a pH value of about 3 (Kurniawan et al., 2019). The acidification of habitat due to anthropogenic activities can impact the biological and ecological processes



(Kleinhappel et al., 2019). In acidic conditions, the biological activity such as plankton growth can be inhibited, causing plankton's appearance in the habitat to be minimum. The acidification in freshwaters reduces species' richness in general, including plankton, and it has negative consequences for aquatic organisms such as fish (Rychła et al., 2011; Hasler et al., 2018). The presence of plankton as nutrition for fish was significant for their growth, so directly and indirectly, the intensity of sunlight and pH value contributed to blue panchax fish's morphometric characters.

The low pH value directly can influence the metabolism of fish. Some study explained that the low pH value of acidic condition contribute to metabolism changes and inhibition of growth. According to Mota et al. (2018), fish exposure to acidic media experiencing various adverse effects, either on physiological and cytological conditions of fishes. Moreover, according to Srineetha et al. (2013) and Kwong et al. (2014), physiological impact of acidic environment includes ionocytes fluctuation, white blood cell production, and increases mucus production. Further impacts of low pH are increasing blood viscosity and affecting fish respiration through lowering total and rate of oxygen intake by fish.

The correlation between low pH and reducing growth and feed intake metabolism could impact growth, appetite, food conversion efficiency, a disruption to physiological homeostasis, blood acidosis, and blood plasma pH (Abbink et al., 2012; Kennedy and Picard, 2012). This condition impacted the morphometric characters of killifish in ex-tin mining pits, especially in pits with age < 5 years with low pH value. The acidic conditions of ex-tin mining pits, especially in pits with age < 5 years, caused this ecosystem not to have any biological life, organic substance, and nutrition to support the blue panchax fish maximum life and its growth. For a long time that

more than 15 years, the chronosequence effect in ex-tin mining pits caused the pH value to change to normal conditions (pH 7). The change of pH of chronosequence impacted the other changes such as DO, BOD, C-organic, total nitrogen, total phosphate, and others (Kurniawan et al., 2019). The consequence of changes can cause an increase in biological life, activity, and nutrition. The existence of biological substances can be the fundamental factor in supporting the blue panchax fish's life besides the chemical and physical aspects of water quality. Therefore, it is reasonable if this study observed that killifish inhabit less than 5 years old ex-tin mining pits have the shortest body length and body length tend to increase related to pits ages with the longest fish body was reach at pits ages between 25 and 50 years old.

Killifish were not found in ex-tin mining pits with age > 50 years. That condition contradicted the report by Raja et al. (2015) and Karuppaiah dan Ramesh (2016) that the members of Aplocheilidae could live in a broad range of water quality (Table 3). Simultaneously, ex-tin mining pits older than 50 years have some water quality parameters within ranges values reported by both groups of researchers. It is suggested that the other ecological factors in ex-tin mining pits with age > 50 years have influenced the existence of killifish in these habitats. It has been reported by Mustikasari et al. (2020) that the presence of killifish in ex-tin mining pits was strongly related to water quality.

Metal and heavy metal content in water could accumulate on fish tissue and organs through serial food chains (Rajeshkumar and Li, 2018; Kurniawan and Mustikasari, 2019). Heavy metal accumulation might disturb fish metabolism and inhibits fish growth. Under extreme conditions the accumulation of heavy metals in fishes' organs can cause mortality (Afshan et al., 2014).

**Table 3.** Water quality of habitat for Family Aplocheilidae

Parameters	Value						
	(a)	(b)	(c)	(d)	(e)	(f)	(g)
Temperature (°C)	21-25	25-28	26-30	28-30	25-32	30-32	19-32
pH	6.8-7.5	7.1-7.8	7.5-7.9	7.3-7.9	7.3-8.1	7.5-8.1	7.0-9.23
DO (mg.l <sup>-1</sup> )	4.8-11.6	4.1-6.4	3.0-3.8	2.0-3.8	0.2-0.3	0.2-0.3	0.02-14.4
COD (mg.l <sup>-1</sup> )	9-13	12-62	110-150	100-150	160-250	173-299	12.6-71.2
BOD (mg.l <sup>-1</sup> )	3.5-12.2	9.5-16.8	96-338	110-338	53-300	45-300	0.01-10.16
Hardness (mg.l <sup>-1</sup> )	33-98	-	-	-	-	-	34-356
Alkalinity (mg.l <sup>-1</sup> )	18-77	4.1-6.4	2.0-3.8	3.0-3.9	1.1-2	1.1-2.9	120-360
Conductivity (mhos.cm <sup>-1</sup> )	42-88	108-270	250-329	280-2000	850-2900	950-2900	240-1560
Turbidity (NTU)	-	14-22	16-25	21-30	26-32	29-32	-
TDS (ppm)	-	-	-	-	-	-	135-1451.6

Sources: (a) Reservoir Bhavanisagar, Tamil Nadu, India (Raja et al., 2015); (b) Dam Viagra, (c) Anaipatti River, (d) Solavandhan River, (e) Arapalaiyam River, (f) Anna Nagar River, India (Karuppaiah dan Ramesh, 2016); (g) Mysore Wetland, India (Prasad et al., 2009).

Previous studies have shown that fish morphology was affected by habitat characteristics (Baillie et al., 2016; Endo and Watanabe, 2020). Habitat characteristics might cause specific of body shape and coloration in fish (Rajeswari et al., 2017). However, there is no study assessing morphological variation in the abandoned ex-tin mining pits. Previous studies by Mustikasari et al. (2020) and Kurniawan et al. (2019) were only discussing blue panchax in the abandoned ex-tin mining pits and its ecological factors. Therefore, this study provides the first data about the morphological variation of blue panchax live in different aged ex-tin mining pits and river in Bangka Island. The data is vital for the management of the further utilization of ex-tin mining pits in the areas. In a broader sense, this study provides a scientific basis that variable sizes and colors of fish production can be obtained through ecological manipulations.

## CONCLUSION

It can be concluded that blue panchax fish collected from different ex-tin mining pits and Limbung River showed morphological variation. The variations are positively related to the ecological characteristics of each habitat.

## ACKNOWLEDGEMENT

The authors would like to thank the Directorate of Research and Society Services Ministry of Research and Technology, the Republic of Indonesia, for providing the funding through the Doctoral Dissertation Research grant. Our gratitude to the University of Jenderal Soedirman that facilitated the authors to conduct this research.

## REFERENCES

- Abbink, W., Garcia, A. B., Roques, J. A., Partridge, G. J., Kloet, K., & Schneider, O. (2012). The effect of temperature and pH on the growth and physiological response of juvenile yellowtail kingfish *Seriola lalandi* in recirculating aquaculture systems. *Aquaculture*, 330, 130-135.
- Afshan, S., Ali, S., Ameen, U. S., Farid, M., Bhargwana, S. A., Hannan, F., & Ahmad, R. (2014). Effect of different heavy metal pollution on fish. *Research Journal of Chemical and Environmental Sciences*, 2(1), 74-79.
- Ariyanto, D., Listiyowati, N., & Imron, I. (2011). Analisis truss morfometrik beberapa varietas ikan nila (*Oreochromis niloticus*). *Jurnal Riset Akuakultur*, 6(2), 187-196.
- Baillie, S. M., Muir, A. M., Hansen, M. J., Krueger, C. C., & Bentzen, P. (2016). Genetic and phenotypic variation along an ecological gradient in lake trout *Salvelinus namaycush*. *BMC Evolutionary Biology*, 16(1), 1-16.
- Beck, S. V., Carvalho, G. R., Barlow, A., Ruber, L., Tan, H. H., Nugroho, E., Wowor, D., Nor, S. A. M., Herder, F., Muchlisin, Z. A., & De Bruyn, M. (2017). Plio-Pleistocene phylogeography of the Southeast Asian blue panchax killifish, *Aplocheilichthys panchax*. *PLoS ONE*, 12(7), e0179557.
- Costa, W. J. E. M. (2013). Historical biogeography of aplocheiloid killifishes (Teleostei: Cyprinodontiformes). *Vertebrate Zoology*, 63(2), 139-154.
- Costa, W. J. E. M. (2016). Comparative morphology and classification of South American cynopoeciline killifishes (Cyprinodontiformes: Aplocheilichthys), with notes on family-group names used for aplocheiloids. *Vertebrate Zoology*, 66(2), 125-140.
- Dekar, M., Sarong, M. A., Batubara, A. S., & Muchlisin, Z. A. (2018). Ichthyofauna of Aceh River, Aceh Province, Indonesia. In *IOP Conference Series: Earth and Environmental Science*, 216(1), 012024. IOP Publishing.
- Endo, C., & Watanabe, K. (2020). Morphological variation associated with trophic niche expansion within a lake population of a benthic fish. *PLoS ONE*, 15(4), e0232114.
- Furness, A. I. (2015). The evolution of an annual life cycle in killifish: adaptation to ephemeral aquatic environments through embryonic diapause. *Biological Reviews*, 91(3), 796-812.
- Gebrekios, S. T. (2016). Factors affecting stream fish community composition and habitat suitability. *Journal of Aquaculture and Marine Biology*, 4(2), 00076.
- Hasler, C. T., Jeffrey, J. D., Schneider, E. V., Hannan, K. D., Tix, J. A., & Suski, C. D. (2018). Biological consequences of weak acidification caused by elevated carbon dioxide in freshwater ecosystems. *Hydrobiologia*, 806(1), 1-12.
- Ihya, S. S., Nuryanto, A., Prabowo, R. E., Bagawati, D., & Kusbiyanto. (2020). Phylogenetic relationships of ornamental Chaethodontidae in the South Coastal of West Java Indonesia. *Jurnal Biodjati*, 5(1), 82-89.
- Irawan, R. R., Sumarwan, U., Suhardjo, B., & Djohar, S. (2014). Strategic model of tin mining industry in Indonesia (case study of Bangka Belitung Province). *International Journal of Business and Management Review*, 2(3), 48-58.
- Karuppaiah, K. P., & Ramesh, U. (2016). Studies on the physicochemical and biological parameters of River Vaigai, Madurai Dist, Tamilnadu, India. *International Journal of Current Microbiology and Applied Sciences*, 5(1), 788-795.



- Kashefi, P., Bani, A., & Ebrahimi, E. (2012). Morphometric and meristic variations between non-reproductive and reproductive kutum females (*Rutilus frisii kutum*, Kamensky, 1901) in the south-west Caspian Sea. *Italian Journal of Zoology*, 79(3), 337-343.
- Kennedy, C. J., & Picard, C. (2012). Chronic low pH exposure affects the seawater readiness of juvenile Pacific sockeye salmon. *Fish Physiology and Biochemistry*, 38(4), 1131-1143.
- Kleinhappel, T. K., Burman, O. H., John, E. A., Wilkinson, A., & Pike, T. W. (2019). The impact of water pH on association preferences in fish. *Ethology*, 125(4), 195-202.
- Kurniawan, A. (2019). Diversitas metagenom bakteri di danau pascatambang timah dengan umur berbeda. *Disertasi*. Fakultas Biologi. Universitas Jenderal Soedirman. 103p.
- Kurniawan, A. (2017). Chronosequence effect of post tin mining ponds to metals residu and microecosystem change. *Omni-Akuatika*, 13(1), 60-65.
- Kurniawan, A., & Mustikasari, D. (2019). Review: Mekanisme akumulasi logam berat di ekosistem pascatambang timah. *Jurnal Ilmu Lingkungan*, 17(3), 408-415.
- Kurniawan, A., Oedjijono, Tamad, & Sulaeman, U. (2019). The pattern of heavy metals distribution in time chronosequence of ex-tin mining ponds in Bangka Regency, Indonesia. *Indonesian Journal of Chemistry*, 19(1), 254-261.
- Kwong, R. W., Kumai, Y., & Perry, S. F. (2014). The physiology of fish at low pH: the zebrafish as a model system. *Journal of Experimental Biology*, 217(5), 651-662.
- Lestari, W., Zayyana, D. A., Setyaningrum, N., & Amelia, T. (2018). The guild composition for modelling fish community in Banjaran River, Purwokerto. *Biosaintifika: Journal of Biology & Biology Education*, 10(3), 698-705.
- Manna, B., Aditya, G., & Banerjee, S. (2011). Habitat heterogeneity and prey selection of *Aplocheilichthys panchax*: an indigenous larvivorous fish. *Journal of Vector-Borne Diseases*, 48(3), 144-149.
- Mojekwu, T. O., & Anumudu, C. I. (2015). Advanced techniques for morphometric analysis in fish. *Journal of Aquaculture Research and Development*, 6: 354.
- Mota, V. C., Hop, J., Sampaio, L. A., Heinsbroek, L. T., Verdegem, M. C., Eding, E. H., & Verreth, J. A. (2018). The effect of low pH on physiology, stress status and growth performance of turbot (*Psetta maxima* L.) cultured in recirculating aquaculture systems. *Aquaculture Research*, 49(10), 3456-3467.
- Muchlisin, Z. A. (2013). Morphometric variations of *Rasbora* group (Pisces: Cyprinidae) in Lake Laut Tawar, Aceh Province, Indonesia, based on truss character analysis. *Hayati Journal of Biosciences*, 20(3), 138-143.
- Mustikasari, D., Nuryanto, A., & Suryaningsih, S. (2020). The presence of blue panchax (*Aplocheilichthys panchax*) in the waters, contaminated by heavy metals, of the abandoned tin mining pits of different age. *AACL Bioflux*, 13(5), 2538-2550.
- Nabila, A., Nuryanto, A., Prabowo, R. E., Bhagawati, D., & Kusbiyanto. (2019). Phylogenetic relationships among ornamental achanturid fish from Ujunggenteng and Taman Manalusu, West Java. *Biosaintifika*, 11(3), 325-331.
- Nguyen, N. H., Hamzah, A., & Thoa, N. P. (2017). Effects of genotype by environment interaction on genetic gain and genetic parameter estimates in Red Tilapia (*Oreochromis* spp.). *Frontiers in Genetics*, 8, 82.
- Nuryanto, A., Bhagawati, D., & Abulias, M. N. (2012). Fish diversity at Cileumeuh River in District of Majenang, Cilacap Regency, Central Java. *Jurnal Iktiologi Indonesia*, 12(2), 147-153.
- Nuryanto, A., Bhagawati, D., & Abulias, M. N. (2015). Ichthyofauna in the Cikawung River, Cilacap Regency, Central Java. *Jurnal Iktiologi Indonesia*, 15(1), 25-37.
- Paknejad, S., Heidari, A., & Mousavi-Sabet, H. (2014). Morphological variation of shad fish *Alosa brashnicowi* (Teleostei, Clupeidae) populations along the southern Caspian Sea coasts, using a truss system. *International Journal of Aquatic Biology*, 2(6), 330-336.
- Pambudi, D. S., Nuryanto, A., & Parabowo, R. E. (2019). Phylogeny of marine ornamental fish members of Labridae from the South Coast of West Java, Indonesia. *Biogenesis*, 7(2), 139-147.
- Pazhayamadom, D. G., Chakraborty, S. K., Jaiswar, A. K., Sudheesan, D., Sajina, A. M., & Jahaagerdar, S. (2014). Stock structure analysis of 'Bombay duck' (*Harpodon nehereus* Hamilton, 1822) along the Indian coast using truss network morphometrics. *Journal of Applied Ichthyology*, 31, 37-44.
- Prasad, A. G. D., Venkataramana, G. V., & Thomas, M. (2009). Fish diversity and its conservation in major wetlands of Mysore. *Journal of Environmental Biology*, 30(5), 713-718.
- Raja, M., Ramkumar, R., Kavitha, M., & Perumal, P. (2015). Distribution pattern, threatened status and conservation measures of fishes with relation to water quality and habitat characteristics of Bhavanisagar Reservoir, Tamil Nadu. *International Journal of Fisheries and Aquatic Studies*, 2(6), 11-18.
- Rajeshkumar, S., & Li, X. (2018). Bioaccumulation of heavy metals in fish species from the Meiliang

- Bay, Taihu Lake, China. *Toxicology Reports*, 5, 288-295.
- Rajeswari, M. V., Rajasree, S. R. R., & Balasubramanian, T. (2017). Effect of light levels on growth, survival and skin colour enhancement of marine angelfish, *Apothemichthys xanthurus* (Bennett, 1833). *Turkish Journal of Fisheries and Aquatic Sciences*, 17(6), 1083-1087.
- Rawat, S., Benakapa, S., Kumar, A.S.J., Naik, K. Pandey, G., & Pema, C.W. (2017). Identification of fish stocks based on truss morphometric: a review. *Journal of Fisheries and Life Sciences*, 2(1), 9-14.
- Rychła, A., Benndorf, J., & Buczyński, P. (2011). Impact of pH and conductivity on species richness and community structure of dragonflies (Odonata) in small mining lakes. *Fundamental and Applied Limnology/Archiv für Hydrobiologie*, 179(1), 41-50.
- Sari, T. S., & Zakaria, I. J. (2017). The relationship of body length and ratio pappilla with sex in gobi fish (*Sicyopterus macrostetholepis*). *Biosaintifika*, 9(2), 265-272.
- Sedlacek, O., Baciakova, B., & Kratochvil, K. (2014). Evolution of body colouration in killifishes (Cyprinodontiformes: Aplocheilidae, Nothobranchiidae, Rivulidae): is male ornamentation constrained by intersexual genetic correlation? *Zoologischer Anzeiger*, 253(3), 207-215.
- Sen, S., Jahageerda, S., Jaiswar, A. K., Chakraborty, S. K., Sajina, A. M., & Dash, G. R. (2011). Stock structure analysis of *Decapterus russelli* (Ruppell, 1830) from east and west coast of India using truss network analysis. *Fisheries Research*, 112(1), 38-43.
- Srineetha, U., Reddy, M. V., & Bhaskar, M. (2013). Effect of environmental acidic pH on oxygen consumption of fish, *Cyprinus carpio* (L). *Journal of Environmental Science, Toxicology and Food Technologsy*, 8(8), 17-21.
- Vasil'eva, E. D., Medvedev, D. A., Chi, T. T. L., Prazdnikov, D. V., Pavlov, D. S., Nga, N. T., & Vasil'ev, V. P. (2013). Species structure of the ichthyofauna of the inland waters of Phu Quoc Island, Gulf of Thailand, Vietnam. *Journal of Ichthyology*, 53(6), 380-396.