


RESEARCH NOTE

Open Access



# Foot self-care behavior and its predictors in diabetic patients in Indonesia

Yunita Sari<sup>1\*</sup> , Arif Setyo Upoyo<sup>1</sup>, Atyanti Isworo<sup>1</sup>, Agis Taufik<sup>1</sup>, Annas Sumeru<sup>1</sup>, Dian Anandari<sup>2</sup> and Eman Sutrisna<sup>3</sup>

## Abstract

**Objectives:** Previous studies have shown that diabetic foot ulcers are principally associated with improper foot self-care. Since appropriate foot self-care is essential to prevent diabetic foot-ulcers, any factors which can predict foot self-care behavior should be identified. However, until now, foot self-care behavior data and predictors of foot-care behavior in Indonesia remain unclear since such studies on Indonesian diabetic patients is very limited. Therefore, the purpose of this study was to investigate foot self-care behavior and to identify its predictors in Indonesia. The design of this study was cross-sectional. Cluster sampling was used, involving 546 type 2 diabetes mellitus patients registered in 22 primary healthcare centers. The questionnaires used in this study included the Diabetes Distress Scale, Beck Depression Inventory II, Family APGAR, Foot-Care Knowledge and Modified Diabetic Foot Care Behaviors.

**Results:** Foot self-care behavior and knowledge about foot care were poor. The predictors of foot self-care behavior were age, educational level, diabetes distress, family support, and knowledge. It needs the program to improve foot self-care knowledge and the program to reduce the diabetes distress in Indonesian diabetic patients. In performing of these programs, families should be involved to improve the support toward foot self-care behavior in patients.

**Keywords:** Behavior, Diabetes, Foot self-care, Predictor

## Introduction

Indonesia is one of the top ten countries with diabetes mellitus (DM) patients [1]. It is predicted there will be 14.1 million diabetic patients in Indonesia by 2035 [2]. One of the most feared and serious complications of DM in developing countries are diabetic foot ulcers [3]. Compared with US and worldwide prevalence, which ranges between 1.4% and 5.9%, the prevalence of diabetic foot ulcer in Indonesia is high, since it is 12% in hospitals and 24% in community settings [3–8]. The presence of a diabetic foot ulcer can affect both the physical and psychosocial life domains, resulting in a reduction in the quality of life, and even mortality [9, 10]. Considering the

significant impact of diabetic foot ulcers, strategies are needed to prevent their occurrence.

A previous study showed that the occurrence of diabetic foot ulcers was mainly associated with improper foot self-care [11, 12]. Previous studies showed that adequate foot self-care can reduce the total number of hospitalizations, and amputation by 50% [11, 13–16]. Therefore, assessment of predictors of foot self-care are required so that the best intervention can be designed and implemented. However, until now, the study about foot self-care behavior in Indonesian diabetic patients is very limited. Moreover, there is no data about in which skill of foot self-care practice that Indonesian diabetic patients is still lacking. Therefore, the first purpose of this study was to assess the foot self-care behavior in Indonesian diabetic patients.

Previous studies in other countries revealed that the factors which influence foot care behavior include

\*Correspondence: yunita.sari@unsoed.ac.id

<sup>1</sup> Department of Nursing, Faculty of Health Sciences, Universitas Jenderal Soedirman, Jl.Dr. Soeparno, Kampus Karangwangkal, Purwokerto, Indonesia

Full list of author information is available at the end of the article



© The Author(s) 2020. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

demographic variables like age, educational level, and occupation [17, 18]. Other factors such as peripheral neuropathy [19], diabetes distress [20], family support [21], depression [22], have also been identified as predictors of foot self-care behavior in diabetic patients. Until now, studies about the predictors of foot self-care behavior in Indonesian patients are very limited, and include only small numbers of participants. No studies have been conducted to investigate whether peripheral neuropathy, diabetes distress, family support, and depression, could serve as predictors of foot self-care behavior in diabetic patients in Indonesia. Until now, studies on predictors of foot self-care behavior in Indonesia have only focused on knowledge variable. However, studies which investigated knowledge as a predictor of foot self-care behavior have shown inconsistent findings in Indonesia. Therefore, the second purpose of this study was to investigate whether peripheral neuropathy, diabetes distress, family support, depression and knowledge affect foot self-care in Indonesian diabetic patients.

## Main text

### Methods

This is a descriptive cross-sectional study. The study was conducted between January 2, 2019 and June 29, 2019. Ethical approval was obtained from the Faculty of Medicine, Universitas Jenderal Soedirman, Purwokerto, Indonesia. Prior to collection of data, written informed consent was obtained from each participant. The purpose of the study, methods, risks and benefits of participating were explained to participants. Respondents were free to withdraw from the study at any time.

A cluster sampling method was used in this study, involving 22 clusters (primary health centers) in Banyumas Regency, Central Java, Indonesia. The sample size was calculated with 95% CI, an absolute estimated precision of 3%, a non-response rate of 5% and a design effect of 2. The final calculation of the sample size was 546 participants. The inclusion criteria were as follows: patients who were 18–80 years old with T2DM, with the ability to practice foot self-care and the ability to communicate. The exclusion criteria were as follows: patients with physical disabilities, immobile, with cognitive impairments, or dementia.

The presence of peripheral neuropathy was assessed with the Michigan Neuropathy Screening Instrument (MNSI). MNSI is a validated tool for screening for peripheral neuropathy in the distal extremity. The second part of the MNSI contains an objective physical assessment test. Previous studies have shown that the MNSI has high sensitivity (80%) and specificity (95%),

with a PPV of 97% and a NPV of 74% [23]. The total possible score was 8 points. A score  $\geq 2$  was considered abnormal [23].

Diabetes distress was assessed by means of the Diabetes Distress Scale [24]. This scale consists of 17 items that use the Likert scale. Items associated with distress experienced over the past month were scored from 1 (not a problem) to 6 (a very serious problem). The Diabetes Distress Scale has been validated in Indonesia. The Cronbach's alpha of the Indonesian version is satisfactory (0.78–0.83) [25].

Depression was assessed by means of Beck Depression Inventory II. This scale has been widely used to assess the severity of subjective depressive symptoms [26]. A total of 21 items were included in this questionnaire. The items were scored from 0 (no) to 3 (severe). Beck Depression Inventory II has been validated in Indonesia, with a Cronbach's alpha of 0.90 [27].

Family support was assessed based on the Family APGAR [28]. This scale consists of 5 items, scored from 0 (never) to 4 (always). The Cronbach's alpha of the subscale ranges between 0.63 and 0.83 [28, 29]. The Cronbach's alpha of the Indonesian version is satisfactory (0.83) [30].

Knowledge about foot self-care was measured by means of a validated foot care knowledge (FCK) questionnaire [19]. Previous studies have shown that the content validity index of this questionnaire is 0.91 [31]. The use of this scale in Indonesia has shown acceptable test–retest reliability, with a range of values of 0.67–1 [31]. The highest possible score was 11.

Foot self-care behavior was measured by means of the Modified Diabetic Foot Care Behaviors (MDFCB) questionnaire [32, 33]. The MDFCB has proven to be reliable, with a Cronbach's alpha coefficient of 0.81 [32]. The total number of items in MDFCB is 34. The formula used to calculate the score was as follows: standard score = (actual score/the highest possible score)  $\times$  100. If the score was less than 60, it was regarded as poor; if the score was between 60 and 80, it was considered as medium; and if the score was more than 80, it was considered good [34].

All data were analysed using SPSS version 23. The demographic variables were assessed using descriptive statistics (means, standard deviation, frequencies). Multivariate regression analysis was conducted to determine the predictors of foot self-care behavior.

## Results

### Socio demographic characteristics of the study participants

A total of 546 subjects participated in this study. The demographic characteristics are shown in Table 1.

**Table 1** Demography characteristics of the respondents (N = 546)

Demographic	n	%	Mean $\pm$ SD	Range
Age			60.14 $\pm$ 9.2	24–88 years
Gender				
Female	417	76.4		
Male	129	23.6		
Marital status				
Married	436	79.9		
Not married	9	1.6		
Divorced/widowed	101	18.5		
Education				
Less than high school graduate	535	97.9		
High school graduate	6	1.09		
Higher than high school graduate	5	0.91		
Occupation				
Farmer	42	7.7		
Non-farmer	137	25.1		
Unemployed/retired	367	67.2		
Income level				
Low income (less than USD\$138 a month)	463	84.8		
Middle income (USD\$138–USD\$177 USD a month)	78	14.3		
High income (higher than USD\$177 a month)	5	0.9		
Comorbidities				
Yes	355	65.1		
No	191	34.9		
Diabetes duration				
<1 years	53	9.7		
1–5 years	272	49.8		
6–10 years	130	23.8		
>10 years	91	16.7		
Peripheral neuropathy				
Yes	294	53.9		
No	252	46.1		
Current foot ulcer				
Yes	190	34.8		
No	356	65.2		

Mn mean, SD standard deviation

**Foot self-care behavior and knowledge about foot self-care**

Concerning foot self-care behavior, the average standard score was 47.4, indicating an overall poor level of foot-care behavior. Responses to questions regarding foot injury treatment and application of foot moisturizer got the lowest standard scores, with average standard scores of 26.3, and 16.3, respectively. Responses with average standard scores below 60 were: examining foot condition (50.7), appropriate footwear (46.3), and foot injury prevention (34.3).

The mean score for foot-care knowledge was  $5.33 \pm 2.2$ , out of a maximum possible score of 11, indicating an overall poor level of foot-care knowledge.

**Predictors of foot self-care behavior**

The results of multiple linear regression analysis are shown in Table 2. The predictors of foot self-care behavior were age, educational level, diabetes distress, family support, knowledge about foot self-care. These five

**Table 2** Result of multivariate linier regression (N = 546)

Variable	B	SE	$\beta$	T	p	R	R <sup>2</sup>	F
Constant term	7.910	5.969		1.325	0.000	0.444	0.197	10.806
Age	0.124	0.057	0.095	2.182	0.030*			
Gender	−2.092	1.296	−0.075	−1.614	0.107			
Marital status	−0.118	1.235	−0.004	−0.096	0.924			
Education	3.492	1.523	0.101	2.292	0.022*			
Occupation	−0.003	0.294	0.000	−0.010	0.992			
Income level	1.794	2.165	0.035	0.829	0.408			
Comorbidities	−1.195	1.055	−0.048	−1.132	0.258			
Diabetes duration	0.885	0.566	0.067	1.563	0.119			
Peripheral neuropathy	−0.122	0.245	−0.023	−0.501	0.617			
Current foot ulcer	3.218	1.706	0.082	1.886	0.060			
Diabetes distress	0.114	0.032	0.164	3.580	0.000**			
Depression	−0.110	0.088	−0.056	−1.250	0.212			
Foot-care knowledge	14.331	2.207	0.276	6.493	0.000**			
Family support	19.948	4.622	0.179	4.315	0.000**			

\*p &lt; 0.05, \*\*p &lt; 0.01

variables explained 19.7% of the variance ( $R = 0.444$ ,  $R^2 = 0.197$ ,  $F = 10.806$ ,  $p < 0.001$ ).

## Discussion

This is the first study to reveal that diabetes distress and family support are predictors of foot self-care behavior in Indonesia.

In this study, the average standard score for foot-care behavior was 47.4, which can be categorized as poor. More specifically, the categories which scored poorly were: use of appropriate footwear, examining foot condition, foot moisturizer use, foot injury prevention and treatment categories. The low score in these categories may be due to the fact that most respondents assume that foot examination, foot and skin injury prevention and treatment do not have to be conducted every day. The low score obtained for moisturizer use may be due to the fact that most patients do not know that moisturizer should not be applied between the toes. The low score regarding foot wear might due to the lack of knowledge about use of footwear and the climate of Indonesia. Indonesia is a hot country, and therefore many people prefer to use sandals instead of shoes.

Our study revealed that patients with an educational level higher than high school graduate obtained significantly higher foot care behavior scores than those with low educational backgrounds. Less educated people tend to have less health knowledge, leading to unhealthy behaviors than those with higher educational levels [35, 36]. Considering that most respondents have a low educational background, nurses who design the educational programs should focus on visual and

auditory demonstration methods, rather than on written instructions.

Our study found that age was one of the predictor of foot self-care behavior. Our study is not in accordance with a recent study which showed that foot self-care behavior in elderly patients were poor since they have physical difficulties and inadequate knowledge [37]. This difference might be due to that in our study we exclude elderly patients who have any physical difficulties such as patients who have loss of vision and who are immobile. Another reason is that elderly patients in this study might have more knowledge compared with adult patients since they have longer duration of having diabetes.

In this study, most of the patients' knowledge was categorized as poor. This lack of knowledge can be explained because most diabetic patients in Indonesia do not receive foot care management information. Based on the knowledge, attitude, and behavior theory, knowledge provides the basis for a positive attitude, and a positive attitude powers behavioral change [18, 38]. According to this theory, persons will do something if they believe that it will have a significant value for themselves. The implication of this theory is that nurses should teach persons with diabetes about the complications which can arise if they fail to properly perform foot care. The fear of the consequences of not treating their feet properly will become a motivation for diabetic patients to properly practice foot care.

Another finding from our study was that family support was a predictor of foot self-care behavior. Even though it is generally accepted that the family has an influence in the management of chronically ill patients [36, 39],

the role of family support on foot self-care behavior in Indonesia is still unknown. Families should be involved to improve family support toward foot self-care behavior in Indonesian diabetic patients. In Indonesia, family involvement can easily occur since the patient's extended family usually lives nearby.

In this study, we found that diabetes distress is one predictor of foot self-care practice. However, our study contradicts the conclusions of a recent study by Devarajoo and Chinna [40], which revealed that there was no relationship between diabetes distress and foot self-care. One possible difference between that study and our study is sociocultural differences. Cultural beliefs influence disease perceptions [41].

It is recommended for Indonesian nurses to design new prevention strategies based on the predictor factors found in this study. An educational program is required to improve the knowledge and skill of patients to perform foot self-care. Additional program which is required is a program of emotional/psychological intervention. This program could reduce diabetic distress in diabetic patients. In performing both of these programs, families should be involved to improve family support toward foot self-care behavior in patients.

## Conclusions

This study demonstrated that foot-care behavior in diabetic patients in Indonesia is poor. This study has shown that age, educational level, diabetes distress, family support and knowledge about foot self-care are predictors of foot self-care behavior in Indonesia.

## Limitation of the study

Firstly, the design of this study (cross-sectional) makes it difficult to establish causal relationships. Secondly, this study was conducted in Indonesia, therefore, the results cannot be generalized to other countries. However, our current study has distinctive strengths that are worth noting. It is the first study about foot self-care behavior that involve a large number of diabetic patients in Indonesia, and the first study to reveal that diabetes distress and family support are predictors of foot self-care behavior in Indonesia.

## Abbreviations

MNSI: Michigan Neuropathy Screening Instrument; Mn: mean; SD: standard deviation; T2DM: type 2 diabetes mellitus.

## Acknowledgements

The authors would like to thank the Ministry of Research, Technology and Higher Education of Indonesia for financially supporting this study.

## Authors' contributions

YS designed and carried out the study, organized and interpreted data, and drafted the manuscript. AI and AS conducted the interviews, and performed

statistical analysis. AT, AS, DA and ES performed statistical analysis and drafted the manuscript. All authors read and approved the final manuscript.

## Funding

This study was supported by a "Penelitian Terapan" Grant from the Ministry of Research, Technology and Higher Education, Indonesia (P/1786/UN23/14/PN/2019). The funding sponsor had no role in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

## Availability of data and materials

Datasets of this study are available from the corresponding author up-on request.

## Ethics approval and consent to participate

This study was approved by the Ethics Committee of the Faculty of Medicine, Jenderal Soedirman University. Written informed consents were obtained from all participants.

## Consent for publication

Not applicable.

## Competing interests

The authors declare that they have no competing interests.

## Author details

<sup>1</sup> Department of Nursing, Faculty of Health Sciences, Universitas Jenderal Soedirman, Jl.Dr. Soeparno, Kampus Karangwangkal, Purwokerto, Indonesia.

<sup>2</sup> Department of Public Health, Faculty of Health Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia. <sup>3</sup> Department of Pharmacology, Faculty of Medicine, Universitas Jenderal Soedirman, Purwokerto, Indonesia.

Received: 5 September 2019 Accepted: 10 January 2020

Published online: 01 February 2020

## References

- Shaw JE, Sicree RA, Zimmet PZ. Global estimates of the prevalence of diabetes for 2010 and 2030. *Diab Res Clin Pract.* 2010;87(1):4–14.
- Guariguata L, Whiting DR, Hambleton I, Beagley J, Linnenkamp U, Shaw JE. Global estimates of diabetes prevalence for 2013 and projections for 2035. *Diab Res Clin Pract.* 2014;103(2):137–49.
- Mariam TG, Alemayehu A, Tesfaye E, Mequannt W, Temesgen K, Yetwale F, et al. Prevalence of diabetic foot ulcer and associated factors among adult diabetic patients who attend the diabetic follow-up clinic at the University of Gondar Referral Hospital, North West Ethiopia, 2016: Institutional-Based cross-sectional study. *J Diab Res.* 2017. <https://doi.org/10.1155/2017/2879249>.
- Yusuf S, Okuwa M, Irwan M, Rassa S, Laitung B, Thalib A, et al. Prevalence and risk factor of diabetic foot ulcers in a Regional Hospital Eastern Indonesia. *Open J Nurs.* 2016. <https://doi.org/10.4236/ojn.2016.61001>.
- Abbott CA, Carrington AL, Ashe H, Bath S, Every LC, Griffiths J, et al. The North-West Diabetes Foot Care Study: incidence of, and risk factors for, new diabetic foot ulceration in a community-based patient cohort. *Diab Med.* 2002;19(5):377–84.
- Margolis DJ, Malay DS, Hoffstad OJ, Leonard CE, MaCurdy T, de Nava KL, et al. Incidence of diabetic foot ulcer and lower extremity amputation among Medicare beneficiaries, 2006 to 2008: Data Points #2. In *Rockville (MD)*; 2011.
- Lauterbach S, Kostev K, Kohlmann T. Prevalence of diabetic foot syndrome and its risk factors in the UK. *J Wound Care.* 2010;19(8):333–7.
- Boulton AJM. The diabetic foot: grand overview, epidemiology and pathogenesis. *Diab Metab Res Rev.* 2008;24(S1):S3–6. <https://doi.org/10.1002/dmrr.833>.
- de Jesus Pereira MT, Magela Salome G, Guimaraes Openheimer D, Cunha Esposito VH, Aguinaldo de Almeida S, Masako Ferreira L. Feelings of powerlessness in patients with diabetic foot ulcers. *Wounds.* 2014;26(6):172–7.
- Rice JB, Desai U, Cummings AKG, Birnbaum HG, Skornicki M, Parsons NB. Burden of diabetic foot ulcers for medicare and private insurers. *Diab Care.* 2014;37(3):651–8.



11. Chiwanga FS, Njelekela MA. Diabetic foot: prevalence, knowledge, and foot self-care practices among diabetic patients in Dar es Salaam, Tanzania—a cross-sectional study. *J Foot Ankle Res*. 2015;8:20.
12. Adib-Hajbaghery M, Alinaqipoor T. Comparing the effects of two teaching methods on healing of diabetic foot ulcer. *J Caring Sci*. 2012;1(1):17–24.
13. Abbas ZG, Lutale JK, Bakker K, Baker N, Archibald LK. The “Step by Step” Diabetic Foot Project in Tanzania: a model for improving patient outcomes in less-developed countries. *Int Wound J*. 2011;8(2):169–75.
14. Canavan RJ, Unwin NC, Kelly WF, Connolly VM. Diabetes- and nondiabetes-related lower extremity amputation incidence before and after the introduction of better organized diabetes foot care: continuous longitudinal monitoring using a standard method. *Diab Care*. 2008;31(3):459–63.
15. Anichini R, Zecchini F, Cerretini I, Meucci G, Fusilli D, Alviggi L, et al. Improvement of diabetic foot care after the implementation of the International Consensus on the Diabetic Foot (ICDF): results of a 5-year prospective study. *Diab Res Clin Pract*. 2007;75(2):153–8.
16. Krishnan S, Nash F, Baker N, Fowler D, Rayman G. Reduction in diabetic amputations over 11 years in a defined U.K. population: benefits of multidisciplinary team work and continuous prospective audit. *Diab Care*. 2008;31(1):99–101.
17. Seid A, Tsige Y. Knowledge, practice, and barriers of foot care among diabetic patients attending Felege Hiwot Referral Hospital, Bahir Dar, Northwest Ethiopia. *Adv Nurs*. 2015. <https://doi.org/10.1155/2015/934623>.
18. Li R, Yuan L, Guo X-H, Lou Q-Q, Zhao F, Shen L, et al. The current status of foot self-care knowledge, behaviours, and analysis of influencing factors in patients with type 2 diabetes mellitus in China. *Int J Nurs Sci*. 2014;1(3):266–71.
19. Pollock RD, Unwin NC, Connolly V. Knowledge and practice of foot care in people with diabetes. *Diab Res Clin Pract*. 2004;64(2):117–22.
20. Tol A, Shojaezadeh D, Sharifirad G, Eslami A, Mohajeritehrani M, Baghbani A. Evaluation of self-care practices and relative components among type 2 diabetic patients. *J Educ Health Promot*. 2012;1(1):19.
21. Ravi S, Kumar S, Gopichandran V. Do supportive family behaviors promote diabetes self-management in resource limited urban settings? A cross sectional study. *BMC Public Health*. 2018;18(1):826.
22. Egede LE, Ellis C, Grubaugh AL. The effect of depression on self-care behaviors and quality of care in a national sample of adults with diabetes. *Gen Hosp Psychiatry*. 2009;31(5):422–7.
23. Feldman EL, Stevens MJ, Thomas PK, Brown MB, Canal N, Greene DA. A practical two-step quantitative clinical and electrophysiological assessment for the diagnosis and staging of diabetic neuropathy. *Diab Care*. 1994;17(11):1281–9.
24. Polonsky WH, Fisher L, Earles J, Dudl RJ, Lees J, Mullan J, et al. Assessing psychosocial distress in diabetes. *Diab Care*. 2005;28(3):626–31.
25. Farm BAS, Perwitasari DA, Thobari JA, Cao Q, Krabbe PFM, Postma MJ. Translation, revision, and validation of the diabetes distress scale for Indonesian type 2 diabetic outpatients with various types of complications. *Value Heal Reg Issues*. 2017;12:63–73.
26. Beck AT, Steer RA, Ball R, Ranieri WF. Comparison of beck depression inventories-IA and-II in psychiatric outpatients. *J Pers Assess*. 1996;67(3):588–97. [https://doi.org/10.1207/s15327752jpa6703\\_13](https://doi.org/10.1207/s15327752jpa6703_13).
27. Ginting H, Näring G, van der Veld WM, Srisayekti W, Becker ES. Validating the beck depression inventory-II in Indonesia's general population and coronary heart disease patients. *Int J Clin Heal Psychol*. 2013;13(3):235–42.
28. Neabel B, Fothergill-Bourbonnais F, Dunning J. Family assessment tools: a review of the literature from 1978–1997. *Heart Lung*. 2000;29(3):196–209.
29. Chin YF, Huang TT, Hsu BR. Impact of action cues, self-efficacy and perceived barriers on daily foot exam practice in type 2 diabetes mellitus patients with peripheral neuropathy. *J Clin Nurs*. 2013;22(1–2):61–8. <https://doi.org/10.1111/j.1365-2702.2012.04291.x>.
30. Mutmainah A, Kusnanto H, Hilman O. The effect of family APGAR score on depression rate in type 2 Diabetes mellitus patients at first-level health facilities. *Rev Prim Care Pract Educ*. 2018;1(2):62.
31. Indrayana S, Guo S-E, Lin C-L, Fang S-Y. Illness perception as a predictor of foot care behavior among people with type 2 diabetes mellitus in Indonesia. *J Transcult Nurs*. 2019;30(1):17–25.
32. Hadi Sulistyoo AA, Sae Sia W, Maneewat K. The effect of a foot care camp on diabetic foot care knowledge and the behaviours of individuals with diabetes mellitus. *J Res Nurs*. 2018;23(5):416–25. <https://doi.org/10.1177/1744987118765903>.
33. Kurniawan T, Yudianto K. Diabetes self-management and its related factors. *J Keperawatan Padjadjaran*. 2017; 4(3). <http://jkp.fkep.unpad.ac.id/index.php/jkp/article/view/289>.
34. Sulistyoo AA H, W S-S, K M. Diabetic foot care knowledge and behaviors of individuals with diabetes mellitus in Indonesia. *GSTF J Nurs Heal Care (JNHCC)*. 2018. 5(1). <http://dl6.globalstf.org/index.php/jnhcc/article/view/1234>.
35. Yuan F, Qian D, Huang C, Tian M, Xiang Y, He Z, et al. Analysis of awareness of health knowledge among rural residents in Western China. *BMC Public Health*. 2015;15:55.
36. Dalstra JAA, Kunst AE, Mackenbach JP. A comparative appraisal of the relationship of education, income and housing tenure with less than good health among the elderly in Europe. *Soc Sci Med*. 2006;62(8):2046–60.
37. Miikkola M, Lantta T, Suhonen R, Stolt M. Challenges of foot self-care in older people: a qualitative focus-group study. *J Foot Ankle Res*. 2019;12:5.
38. Knowles S, Lam LT, McInnes E, Elliott D, Hardy J, Middleton S. Knowledge, attitudes, beliefs and behaviour intentions for three bowel management practices in intensive care: effects of a targeted protocol implementation for nursing and medical staff. *BMC Nurs*. 2015;14:6.
39. Harwell TS, Helgeson SD, Gohdes D, McInerney MJ, Roumagoux LP, Smilie JG. Foot care practices, services and perceptions of risk among medicare beneficiaries with diabetes at high and low risk for future foot complications. *Foot Ankle Int*. 2001;22(9):734–8.
40. Devarajoo C, Chinna K. Depression, distress and self-efficacy: the impact on diabetes self-care practices. *PLoS ONE*. 2017;12(3):e0175096.
41. Kahissay MH, Fenta TG, Boon H. Beliefs and perception of ill-health causation: a socio-cultural qualitative study in rural North-Eastern Ethiopia. *BMC Public Health*. 2017;17(1):124.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)



[Search](#)[Explore journals](#)  
[Menu](#)[Get published](#)[About BMC](#)[Login](#)

---

# BMC Research Notes

---

[Home](#) [About](#) [Articles](#) [Submission Guidelines](#)

---

## About ▾

[Contact](#)[Editorial Board](#)

### ▾ Article types

[Journal News](#)[In Review](#)

## About

- [Content we consider](#)
  - [Open access](#)
  - [Data availability policy](#)
  - [Article-processing charges](#)
  - [Indexing services](#)
  - [Peer-review policy](#)
  - [Editorial policies](#)
  - [Appeals and complaints](#)
  - [Citing articles in BMC Research Notes](#)
  - [Benefits of publishing with BMC](#)
- 

## Content we consider

### Aims and scope

*BMC Research Notes* is an open access journal publishing peer-reviewed contributions from across all scientific and clinical disciplines, including intriguing initial observations, updates to previous work and established methods, valid negative results, and scientific data sets and descriptions. We are committed to making it easier to find, cite and share your short reports by providing an inclusive forum for valuable data and research observations.

---

*BMC Research Notes* does not make editorial decisions on the basis of the interest of a study or its likely impact. Studies must be scientifically valid; for research articles this includes a scientifically sound research question, the use of suitable methods and analysis, and following community-agreed standards relevant to the research field.

<b>Health sciences and medicine</b>	Biology and life sciences	Physical sciences and engineering	Scientific community
-------------------------------------	---------------------------	-----------------------------------	----------------------

Health science contributions can cover

- Allergy and immunology
  - Brain and nervous system
  - Health services research
  - Infectious disease
  - Internal medicine
  - [Oncology](#)
  - [Mental health](#)
  - Nutrition
  - [Public health](#)
  - Surgery
  - and more
- 

Find out more about our unique article types



publication, plus VAI or local taxes where applicable.

If the corresponding author's institution participates in our open access membership program, some or all of the publication cost may be covered (more details available on the [membership page](#)). We routinely waive charges for authors from [low-income countries](#). For other countries, article-processing charge waivers or discounts are granted on a case-by-case basis to authors with insufficient funds. Authors can request a waiver or discount during the submission process. For further details, see our [article-processing charge page](#).

Visit Springer Nature's [open access funding & support services](#) for information about research funders and institutions that provide funding for APCs.

Springer Nature offers agreements that enable institutions to cover open access publishing costs. Learn more about our [open access agreements](#) to check your eligibility and discover whether this journal is included.

For more information on APCs please see our [Journal Pricing FAQs](#)

## Indexing services

[Back to top](#)

All articles published in *BMC Research Notes* are included in:

- ANVUR
- BFI List
- CAB Abstracts
- CLOCKSS
- CNKI
- Chemical Abstracts Service (CAS)
- Chinese Academy of Sciences (CAS) - GoOA
- DOAJ
- Dimensions
- EBSCO Academic Search

- EBSCO Biomedical Reference Collection
- EBSCO Discovery Service
- EBSCO STM Source
- EMBiology
- Emerging Sources Citation Index
- Gale
- Google Scholar
- Medline
- Meta
- Naver
- Norwegian Register for Scientific Journals and Series
- OCLC WorldCat Discovery Service
- Pathway Studio
- Portico
- ProQuest-ExLibris Primo
- ProQuest-ExLibris Summon
- PubMedCentral
- SCImago
- SCOPUS
- TD Net Discovery Service
- UGC-CARE List (India)
- Zoological Record

The full text of all articles is deposited in [digital archives](#) around the world to guarantee long-term digital preservation. You can also access all articles published by BioMed Central on [SpringerLink](#).

We are working closely with relevant indexing services including Web of Science (Clarivate Analytics) to ensure that articles published in *BMC Research Notes* will be available in their databases when appropriate.

---

## Peer-review policy

## BMC Research Notes

[Home](#) [About](#) [Articles](#) [Submission Guidelines](#) [Join The Editorial Board](#)

Search articles within this journal



Submit manuscript

### Reproducibility and research integrity



Following a call from the UK House of Commons Science and Technology Committee for evidence on reproducibility and research integrity, and the roles different institutions play in this, *BMC Research Notes* partnered with the UK Reproducibility Network to provide a platform to share feedback on the topic with the wider scientific community.

Browse collection articles here

### Aims and scope

*BMC Research Notes* is an open access journal publishing peer-reviewed contributions from across all scientific and clinical disciplines, including intriguing initial observations, updates to previous work and established methods, valid negative results, and scientific data sets and descriptions. We are committed to making it easier to find, cite and share your short reports by providing an inclusive forum for valuable data and research observations.

More information about our scope

### **BMC** Series

#### Important information

[Editorial board](#)

[About our article types](#)

[For authors](#)

[For editorial board members](#)

[For reviewers](#)

Activate Windows  
Go to Settings to activate Windows.

[Search](#)[Explore journals](#)  
[Menu](#)[Get published](#)[About BMC](#)[Login](#)

# BMC Research Notes

[Home](#) [About](#) [Articles](#) [Submission Guidelines](#)

## Articles ▾

[Data notes](#)

### ▾ Collections

[Supplements](#)[Reviewer acknowledgements](#)

## Articles

[Search by keyword](#)[Search by citation](#)[Volume 13 \(2020\) ▾](#)[Search](#)

## 545 result(s)

within Volume 13 of BMC Research Notes

Page 10 of 11

Sort by

[Newest first ▾](#)

### **A comparison of erythrocyte sedimentation rates of bloods anticoagulated with trisodium citrate and EDTA among TB presumptive patients at the University of Gondar comprehensive specialized hospital, northwest Ethiopia**

The purpose of this study was comparing the erythrocyte sedimentation rate (ESR) results of trisodium citrate (TSC) and ethylene diamine tetra-acetic acid (EDTA) anticoagulants. A comparative cross-sectional s...

Research note | Published on: 4 February 2020

› [Full Text](#) › [PDF](#)

---

### **R-spondin 2 mediates neutrophil egress into the alveolar space through increased lung permeability**

R-spondin 2 (RSPO2) is required for lung morphogenesis, activates Wnt signaling, and is upregulated in idiopathic lung fibrosis. Our objective was to investigate whether RSPO2 is similarly important in homeost...

S. R. Jackson, M. F. D. M. Costa, C. F. Pastore, G. Zhao, A. I. Weiner, S. Adams, G. Palashikar, K. Quansah, K. Hankenson, D. R. Herbert and A. E. Vaughan

*BMC Research Notes* 2020 13:54

Research note | Published on: 4 February 2020

› [Full Text](#) › [PDF](#)

---

### **High-throughput sequencing of virus-infected *Cucurbita pepo* samples revealed the presence of Zucchini shoestring virus in Zimbabwe**

Plant-infecting viruses remain a serious challenge towards achieving food security worldwide. Cucurbit virus surveys were conducted in Zimbabwe during the 2014 and 2015 growing seasons. Leaf samples displaying...

Charles Karavina, Jacques Davy Ibaba and Augustine Gubba

*BMC Research Notes* 2020 13:53

Research note | Published on: 3 February 2020

› [Full Text](#) › [PDF](#)

---

### **Foot self-care behavior and its predictors in diabetic patients in Indonesia**

Previous studies have shown that diabetic foot ulcers are principally associated with improper foot self-care. Since appropriate foot self-care is essential to prevent diabetic foot-ulcers, any factors which c...

Yunita Sari, Arif Setyo Upoyo, Atyanti Isworo, Agis Taufik, Annas Sumeru, Dian Anandari and Eman Sutrisna

*BMC Research Notes* 2020 13:38

Research note | Published on: 1 February 2020

➤ [Full Text](#) ➤ [PDF](#)

---

### **Correlation between detergent activity and anti-herpes simplex virus-2 activity of commercially available vaginal gels**

Herpes simplex virus-2 (HSV-2) infections are almost exclusively sexually transmitted. The presence of vaginal gels during sexual activity may have a significant positive or negative impact on viral transmissi...

Andrea Szöllősi, Tímea Raffai, Anita Bogdanov, Valéria Endrész, László Párducz, Ferenc Somogyvári, László Janovák, Katalin Burián and Dezső P. Virok

*BMC Research Notes* 2020 13:52

Research note | Published on: 31 January 2020

➤ [Full Text](#) ➤ [PDF](#)

---

### **Outer membrane protein A (OmpA) of extraintestinal pathogenic *Escherichia coli***

Extraintestinal Pathogenic *E. coli* (ExPEC), are responsible for host diseases such as Neonatal Meningitis *Escherichia coli* (NMEC), the second-leading cause of neonatal bacterial meningitis, Avian Pathogenic *E. co...*

Daniel W. Nielsen, Nicole Ricker, Nicolle L. Barbieri, Heather K. Allen, Lisa K. Nolan and Catherine M. Logue

*BMC Research Notes* 2020 13:51

Research note | Published on: 31 January 2020

➤ [Full Text](#) ➤ [PDF](#)





# Source details

## BMC Research Notes

Open Access

Scopus coverage years: from 2008 to Present

Publisher: Springer Nature

E-ISSN: 1756-0500

Subject area: Biochemistry, Genetics and Molecular Biology: General Biochemistry, Genetics and Molecular Biology

Source type: Journal

CiteScore 2021

3.5

SJR 2021

0.527

SNIP 2021

0.950

[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Source Homepage](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

### Improved CiteScore methodology

CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore 2021

$$3.5 = \frac{9,405 \text{ Citations 2018 - 2021}}{2,712 \text{ Documents 2018 - 2021}}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022

$$3.5 = \frac{7,031 \text{ Citations to date}}{2,032 \text{ Documents to date}}$$

Last updated on 05 September, 2022 • Updated monthly

## CiteScore rank 2021

Category	Rank	Percentile
Biochemistry, Genetics and Molecular Biology	#89/204	56th
General Biochemistry, Genetics and Molecular Biology		

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)

## About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

## Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

## Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

---

**ELSEVIER**

[Terms and conditions](#) ↗ [Privacy policy](#) ↗

Copyright © [Elsevier B.V](#) ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the [use of cookies](#) ↗.





Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

- Home
- Journal Rankings
- Country Rankings
- Viz Tools
- Help
- About Us



Latest Research Findings


Online & Print Weekly Journal

Groundbreaking Global Science Research Delivered to You & Online Archive Access From 1997.

nature.com

OPEN

BMC Research Notes

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
United Kingdom	Biochemistry, Genetics and Molecular Biology	BioMed Central Ltd.	79
 Universities and research institutions in United Kingdom	Biochemistry, Genetics and Molecular Biology (miscellaneous)		
	Medicine		
	Medicine (miscellaneous)		
PUBLICATION TYPE	ISSN	COVERAGE	INFORMATION
Journals	17560500	2008-2021	<a href="#">Homepage</a>
			<a href="#">How to publish in this journal</a>

## Online & Print Weekly Jour

**Reviews, News, & Analysis**


Groundbreaking Global Science Research  
Get Your Online & Print Subscription Now!

nature.com

OPEN

### SCOPE

BMC Research Notes publishes scientifically valid research outputs that cannot be considered as full research or methodology articles. We support the research community across all scientific and clinical disciplines by providing an open access forum for sharing data and useful information; this includes, but is not limited to, updates to previous work, additions to established methods, short publications, null results, research proposals and data management plans.

 Join the conversation about this journal

## 100% Asian Telcos

See you in In-Person in Bangkok, 2 – 3 November 2022

Telecoms World Asia 2022



Quartiles



# 100% Asian Telcos

See you in In-Person in Bangkok, 2 – 3 November 2022







Telecoms World Asia 2022



FIND SIMILAR JOURNALS ?

1 PLoS ONE USA	2 Pan African Medical Journal NGA	3 Clinical Epidemiology and Global Health NLD	4 Frontiers in Public Health CHE	5 Biomedical and Environmental Science GBR
73% similarity	62% similarity	60% similarity	60% similarity	59% similarity

SJR	Total Documents	Citations per document



← Show this widget in your own website

Just copy the code below and paste within your html code:

`<a href="https://www.scima`

**SCImago Graphica**

Explore, visually communicate and make sense of data with our **new data visualization tool**.

