



Home About Login Register Search Current Archives Announcements Statistics Indexing Site

Home > Archives > Vol 5, No 3 (2020)

Vol 5, No 3 (2020)

December

TABLE OF CONTENTS

Ilmu Pertanian (Agricultural Science)	
Volume 5 no. 3 December 2020	
ARTICLES	
Morphological characteristics and yield performance of mutant soybean [Glycine max (L.) Merr.] M6 genotypes in Jatihangor, Sumedang, West Java Andrew Yool and Meddy Rachmad	117
Physiological characteristics of two accessions of moringa (<i>Moringa oleifera</i> Lam.) at different planting site in Madura Catur Watonowati, Endang Sulistyanyingsih, Didi Indradewa, and Budiatuti Kumasih	123
Strength assessment of rice hills from different planting distance by loading simulation Edi Santosa, Herdiana Agustia, Dwi Guntono, and Sofyan Zaman	131
Similarity level of <i>Nepenthes</i> spp. based on the qualitative characters Putri Lukmanasari, Adis Purwaningsih, Rudi Hari Murti, and Zukhrif	140
Effects of reducing rates of NPK, ZA, and KCl fertilizers on the growth and yield of shallot in multiple cropping system in Bantul Kasidi, Endang Sulistyanyingsih, Didi Indradewa, and Budiatuti Kumasih	150
Response of upland rice (<i>Oryza sativa</i> L.) cultivars to different shade levels in sandy soil Irwani Sukir Padang, Tohari, and Jaka Widada	158
Branch pruning and chicken manure application to improve growth and yield of broccoli in lowland area Dita Herliana and Fagil A. Alkhatib	166

Recent Issues

Upcoming Issues

Recent issues :

Ilmu Pertanian (Agricultural Science)	
Volume 5 no. 3 December 2020	
ARTICLES	
Morphological characteristics and yield performance of mutant soybean [Glycine max (L.) Merr.] M6 genotypes in Jatihangor, Sumedang, West Java Andrew Yool and Meddy Rachmad	117
Physiological characteristics of two accessions of moringa (<i>Moringa oleifera</i> Lam.) at different planting site in Madura Catur Watonowati, Endang Sulistyanyingsih, Didi Indradewa, and Budiatuti Kumasih	123
Strength assessment of rice hills from different planting distance by loading simulation Edi Santosa, Herdiana Agustia, Dwi Guntono, and Sofyan Zaman	131
Similarity level of <i>Nepenthes</i> spp. based on the qualitative characters Putri Lukmanasari, Adis Purwaningsih, Rudi Hari Murti, and Zukhrif	140
Effects of reducing rates of NPK, ZA, and KCl fertilizers on the growth and yield of shallot in multiple cropping system in Bantul Kasidi, Endang Sulistyanyingsih, Didi Indradewa, and Budiatuti Kumasih	150
Response of upland rice (<i>Oryza sativa</i> L.) cultivars to different shade levels in sandy soil Irwani Sukir Padang, Tohari, and Jaka Widada	158
Branch pruning and chicken manure application to improve growth and yield of broccoli in lowland area Dita Herliana and Fagil A. Alkhatib	166

Vol 5, no 3 (2020)

Ilmu Pertanian (Agricultural Science)	
Volume 5 no. 2 August 2020	
ARTICLES	
Morphological characters of rice and yield of three accessions (<i>Oryza sativa</i> L.) in the field with degraded trench Dagang Hari Nugroho and Dwi Tarnawa Satrio Putra	58
Evaluation of drought tolerance in some tropical wheat genotypes (<i>Triticum aestivum</i> L.) at different nutrient stress level Muhammad Rabbil, Kamuladin Komaludin, Yuni Mulya, Rudi Belandjaja, and Anisa Nur	65
NPK uptake and growth of maize on endogenous pest as affected by the application of mycorrhizal fungi multi species and commercial fungicides Mugjo Ade Sapitro, Dwi Kusnadi, Gajip Wulandari, and Sri Murni Rahmah Darmi	76
Effects of acids and alkaline involves the systems of an layering in two plant doses of 5000 P and 5000 P using both chemical, inorganic and amino acids No Betty Wahidatul, Mudo Yudianto, and Dwi Tarnawa Satrio Putra	86
Growth and yield of five prospective shallot selected accessions from true shallot seeds in lowland areas Endang Sulistyanyingsih, Anisa Purwaningsih, and Didi Indradewa	93
Effects of water flow rate and surface cover plant density on the growth of clambered (<i>Ipomoea pes-caprae</i> L.) Anisa Purwaningsih, Anisa Purwaningsih, and Didi Indradewa	98
The influence of disease and phytochemical inhibitor on loss of N and all palm harvest in year Mito Daniel Riyadi, Dwi Tarnawa Satrio Putra, and the Kusnadi	130

Vol 5, No 2 (2020)

Ilmu Pertanian (Agricultural Science)	
Volume 5 no. 1 April 2020	
ARTICLES	
A Review: Challenges and opportunities of implementing water productivity index (WPI) at data farms in Malaysia Samsul Hani, Anis Faridat Komaludin, Sri Nurana Mahdi Daud, Zaini Azzah, Dwi Tarnawa Satrio Putra, Rudi Belandjaja, and Yuni Mulya	1
Effects of varied fish waste and raw manure on NPK availability and uptake of lowland rice on peat soil in Palauwang River Dwi Tarnawa Satrio Putra, Rudi Belandjaja, and Yuni Mulya	11
Effect of saline irrigation method on the growth of rice (<i>Oryza sativa</i> L.) Adhik, Budiatuti Kumasih, and Didi Indradewa	18
Effect of humic acid on the growth and yield of rice maize (<i>Oryza sativa</i> L.) submersed rice Fagil A. Alkhatib, Endang Sulistyanyingsih, and Didi Indradewa	25
The role of iron oxidizing bacteria to the quality of biochar on and soilborne soil Kusnadi, Rabbil, Anisa Nur, Sri Murni Rahmah Darmi, and Yuni Mulya	29
The development of smart screening technology for (SST) for screening cassava cultivars based on cytosolic analysis Judi Hattoro, Herdiana Agustia, Didi Indradewa, Rudi Hari Murti, Anisa Nur, Anisa Nur, Anisa Nur, and Didi Indradewa	46
Transformation of biomass (T) of oil palm seeds at different storage periods Bismillah, Tarnawa, and Priya Fatma	51

Vol 5, No 1 (2020)

Ilmu Pertanian (Agricultural Science)



[Home](#) [About](#) [Login](#) [Register](#) [Search](#) [Current](#) [Archives](#) [Announcements](#) [Statistics](#) [Indexing Site](#)

[Home](#) > [About the Journal](#) > **Editorial Team**

Editorial Team

Editor in Chief

Dr. Rudi Hari Murti, Department of Agronomy, Faculty of Agriculture, Universitas Gadjah Mada, Indonesia

Editorial Board

Prof. Dr. Irwandi Jaswir, Bio Process & Molecular Engineering Research Unit (BPMERU), International Islamic University Malaysia, Malaysia

Dr. Endang Septiningsih, Department of Soil and Crop Sciences, College of Agriculture and Life Science, Texas A&M University, United States

Dr. Chee How Teo, Centre for Research in Biotechnology for Agriculture, University of Malaya, Malaysia

Dr. Sastia Prama Putri, Department of Biotechnology, Graduate School of Engineering, Osaka University, Japan

Dr. Kim Kyung-Min, Division of Plant Biosciences, School of Applied Biosciences, College of Agriculture and Life Science, Kyungpook National University, Korea, Republic of

Dr. Bunyamin Tar'an, University of Saskatchewan - Canada

Dr. Agung Wahyu Susilo, Indonesian Coffee and Cocoa Research Institute, Indonesia

Dr. Sintho Wahyuning Ardie, SP, M.Si, Department of Agronomy and Horticulture, Institut Pertanian Bogor, Bogor, Indonesia

Dr. Ir. Benito Heru Purwanto, M.P., M.Agr.Sc., Soil Science Department, Faculty of Agriculture, Universitas Gadjah Mada, Indonesia

Associate (Handling) Editors

Widhi Dyah Sawitri, Ph.D, Faculty of Agriculture, Universitas Gadjah Mada, Indonesia

Nur Ainun Harlin Jennie Pulungan, Ph.D, Department of Soil, Faculty of Agriculture, Universitas Gadjah Mada, Indonesia

Dr. Dyah Weny Respatie, Department of Agronomy, Faculty of Agriculture, Universitas Gadjah Mada, Indonesia

Enik Nurlaili Afifah, Department of Agronomy, Faculty of Agriculture, Universitas Gadjah Mada, Indonesia

Editorial Assistants

Eka Candra Wardani, Faculty of Agriculture, Universitas Gadjah Mada, Indonesia

Sri Junandi, Perpustakaan Universitas Gadjah Mada, Indonesia

Laili Hidayah S.Sos., M.A., Faculty of Agriculture Library Gadjah Mada University, Indonesia

Palupi Jatiasri, Department of Plant Protection, Faculty of Agriculture, Universitas Gadjah Mada, Indonesia

[Recent Issues](#)

[Upcoming Issues](#)

Recent issues :





Vol 5, No 3 (2020)

December

Table of Contents

Research Articles



Morphological characteristics and yield performance of mutant soybean [*Glycine max* (L.) Merr.] M6 genotypes in Jatinangor, Sumedang, West Java

Andrew Yoel, Meddy Rachmadi

[doi: 10.22146/ipas.54104](#) Abstract views : 619 | views : 468

117-122




Physiological characteristics of two accessions of moringa (*Moringa oleifera* Lam.) at different planting sites in Madura

Catur Wasonowati, Endang Sulistyaningsih, Didik Indradewa, Budiastuti Kurniasih

[doi: 10.22146/ipas.47245](#) Abstract views : 490 | views : 551

123-130




Strength assessment of rice hills from different planting distance by loading simulation

Edi Santosa, Herdhata Agusta, Dwi Guntoro, Sofyan Zaman

[doi: 10.22146/ipas.31895](#) Abstract views : 537 | views : 550

131-139



Similarity level of *Nepenthes* spp. based on the qualitative characters

Putri Lukmanasari, Aziz Purwantoro, Rudi Hari Murti, Zulkifli Zulkifli

[doi: 10.22146/ipas.55728](#) Abstract views : 555 | views : 747

140-149




Effect of reducing rates of NPK, ZA, and KCl fertilizers on the growth and yield of shallot in multiple cropping system in Bantul

Karsidi Karsidi, Endang Sulistyaningsih, Didik Indradewa, Budiastuti Kurniasih

[doi: 10.22146/ipas.36816](#) Abstract views : 472 | views : 432

150-157



Response of upland rice (*Oryza sativa* L.) cultivars to different shade levels in sandy soil

Irwan Suluk Padang, Tohari Tohari, Jaka Widada

[doi: 10.22146/ipas.30643](#) Abstract views : 488 | views : 411

158-165



Branch pruning and chicken manure application to improve growth and yield of brocolli in lowland area

Okti Herliana, Yugi R Ahadiyat

doi:10.22146/ipas.43101  Abstract views : 408 |  views : 341

Recent Issues

Upcoming Issues

Recent issues :

[illegible]

Ilmu Pertanian (Agricultural Science) **ISSN 0126-4214** (print), **ISSN 2527-7162** (online) is published by Faculty of Agriculture Universitas Gadjah Mada collaboration with Perhimpunan Sarjana Pertanian Indonesia (PISPI) and licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](#).

00303280 View My Stats

Focus & Scope

Editorial Board

Author Guidelines

Reviewers Guidelines

Author Fees

Publication Ethics

Peer Review Process

Online Submission

Visitor Statistics

Journal History

DOAJ DIRECTORY OF
OPEN ACCESS JOURNALS

Ilmu Pertanian (Agricultural Science)

 (<https://creativecommons.org/licenses/by-sa/4.0/>)



Full-text formats available: PDF

<https://doaj.org/toc/2527-7162?source=%7B%22query%22%3A%7B%22filtered%22%3A%7B%22filter%22%3A%7B%22bool%22%3A%7B%22must%22%3A%5B%7B%22terms%22%3A...> 1/16

Ilmu Pertanian (Agricultural Science)

eISSN : 2527-7162 | pISSN : 0126-4214

Agriculture

Universitas Gadjah Mada



S2

Sinta Score



Indexed by GARUDA

18

H-Index

17

H5-Index

1327

Citations

1004

5 Year Citations



Branch pruning and chicken manure application to improve growth and yield of broccoli in lowland area

Okti Herliana* and Yugi R. Ahadiyat

Department of Agrotechnology, Faculty of Agriculture, Universitas Jenderal Soedirman

Jl. Dr. Soeparno no. 61, Purwokerto 53123, Indonesia

*Corresponding author: okti.herliana@unsoed.ac.id

Received: 29th January 2019; **Revised:** 27th August 2020; **Accepted:** 02nd September 2020

Keywords:

Broccoli, branch pruning, chicken manure, lowland

ABSTRACT

The objective of this study was to determine the effect of branch pruning and chicken manure application on improving the growth and yield of broccoli in the lowland. The research was conducted in the dry land of Kutasari Village, Baturraden District, Banyumas Regency at 138 m above sea level. The research was arranged in a Factorial Randomized Complete Block Design (RCBD) with three replications consisting of two factors of chicken manure application (without chicken manure and with chicken manure at doses of 10 ton.ha⁻¹ and 20 ton.ha⁻¹) and branch pruning (without and with branch pruning of 25% and 50%). The pruning was performed when the plants initiated flowers. The observed data were analyzed with the F test and followed with Duncan's Multiple Range Test (DMRT) at $\alpha = 5\%$. The branch pruning resulted in a significant effect on the number of leaves, leaf width, head weight, and head diameter. Meanwhile, the chicken manure application showed a significant effect on all of the observed variables. There was an interaction effect of branch pruning and chicken manure application on the leaf width, head weight, and head diameter. The best result was obtained by the combined treatment of 50% branch pruning and 20 ton.ha⁻¹ chicken manure, resulting in the head weight of 1048.33 g.plant⁻¹ and leaf width of 1705.41 cm².

INTRODUCTION

Broccoli is a vegetable crop with a high economic value, containing essential minerals of vitamins of A, B complex, and E, as well as calcium, magnesium, zinc, and iron as reported by Gad and El-Moez (2011), which could fulfill the nutritional need as well as high anti-oxidant and beta-carotene, making its role even higher in preventing cancer within the human body (Wasonowati, 2009). Indonesia's broccoli production is around 183,816 ton.ha⁻¹ (Badan Pusat Statistik, 2019), hence, it cannot meet the local market and international market demands that keep on increasing 20% to 30% annually. Besides, Indonesia's broccoli contains a lot of chemical residues with limited planting areas and expensive

production costs for its fertilization. Most varieties of this plant can only be produced well at highland. Therefore, it is necessary to find technology to develop this plant in lowland area using organic fertilizer to decrease the use of synthetic fertilizers as well as to support sustainable broccoli production. For this purpose, it is also important to improve the broccoli quality by reducing the use of synthetic chemical substances (Budiastuti et al., 2009). Ouda and Mahadeen (2008) reported that recently, consumers are demanding higher quality and safer food, and they are highly interested in organic products.

Broccoli cultivation in lowland area has agro-climatic obstacles because broccoli flowering requires a relatively low temperature (vernalization) at the end of its vegetative phase (Booij and Struik,

1990; Grevsen, 1998). If this low temperature is not met, then broccoli will continue its vegetative phase. Broccoli heads are also reportedly to experience a decrease in when the temperature in its growth environment is less optimal during the flowering process (Heather et al., 1992; Grevsen, 1998; Björkman and Pearson, 1998). Another problem of broccoli cultivation in tropical lowland is the high number of branches. Jaya et al. (2006) reported that the number of branches of broccoli planted in the lowland of Lombok Island (± 125 m above sea level) could reach up to 12 branches. This high number of branches do not present any profit economically because they could decrease the main head size and increase the harvesting costs because they do not have different harvesting periods (Wien and Wurr, 1997).

The need for macronutrients becomes an important factor in broccoli cultivation for generating optimum production. Nitrogen element needs to be added to produce green leaves and big flower, yet the excessive use of urea will reduce the natural soil properties. One of the solutions could be applied to reduce the use of urea fertilizer is by using organic fertilizer. According to Simanungkalit et al. (2006), organic fertilizer consists of organic matters deriving from plants and or animal, which has undergone an engineering process and a change into solid or liquid. In addition to having macro-elements, organic fertilizers have microelements as their strength, which are not found in synthetic chemical fertilizers. Organic fertilizers are also eco-friendly, and they can easily be found in the land cultivation area. Chicken manure has greater N and P elements compared to cow or goat manure. It has N element three times greater than cow or goat manure because chicken only has one discharge hole, making its liquid and solid excretion mixed (Musnamar, 2004). Pinem et al. (2015) mentioned that the addition of chicken manure to plants has some effects on increasing plant growth performance. Thus, this study was conducted to determine the effects of branch pruning and chicken manure application on improving plant growth and yield of broccoli in lowland area.

MATERIALS AND METHODS

The study was carried out in Kutasari Village, Baturraden District, Banyumas Regency, from April to July 2015. The site is 138 m above sea level with Inceptisol soil type. The materials used consisted of

broccoli seeds, organic fertilizer in the form of chicken manure, synthetic chemical fertilizer consisting of Urea, SP36, and KCl at a dose of 50% of recommendation doses and pesticides. Meanwhile, the tools used were hoe, scythe, electric scale, ruler, measuring tools (thermometer, lux meter, and measuring cup) and soil analysis tools.

This research was arranged in a Factorial Randomized Complete Block Design (RCBD) consisted of two factors, which were chicken manure doses (A0: 0 ton.ha⁻¹, A1: 10 ton.ha⁻¹ and A2: 20 ton.ha⁻¹) and branch pruning (P0: without pruning, P1: 25% pruning and P2: 50%), with three replications, resulting in nine combined treatments and 27 plots with a size of 3 m × 2 m each and a plant spacing of 60 cm × 50 cm. There were 20 plants per plot.

The broccoli seeds were germinated for 15 days and then transplanted. Chicken manure application was carried out a week before transplanting as basic fertilizer. Urea, SP36, and KCl fertilizers were applied at a dose of 50% of recommendation dose at 10 days after planting. Branch pruning was done when the plant started flower initiation by cutting the base of the branch that was attached to the stem using a knife sterilized with alcohol.

The variables for observation included growth and yield components. The growth variables observed were plant height (cm), number of leaves, and leaf width before cut and leaf width after harvest (cm²), while the yields components observed were canopy diameter (cm), head weight (g), and head diameter (cm). The observed data were analyzed with the F test and followed with Duncan's Multiple Range Test (DMRT) at $\alpha=5\%$ (Gomez and Gomez, 1983).

RESULTS AND DISCUSSION

General condition of the research site

Based on the climatology data in the research site obtained from the nearest meteorology station during the experiment, the lowest temperature was 17.4°C and the highest was 34.3°C with the average daily temperature of 25.3°C. The lowest temperature occurred in July 2015 during the flower development phase. As reported earlier, broccoli requires relatively low temperatures to initiate flowering (Grevsen and Olesen, 1999) and to produce a good quality flower (Heather et al., 1992). The soil analysis conducted

in the Laboratory of Soil Resources, Department of Agrotechnology, Universitas Jenderal Soedirman, indicated that the soil pH in the experiment site was 6.33, which was slightly acidic at a medium-fertility level, as can be seen from the elements contained in it. The nitrogen content was 188.916 ppm (medium), available phosphorus content was 8.17 (low), potassium content was 0.625 % (medium), and carbon content was 3.616 (medium).

The statistical analysis results indicated that there were varied responses in the observed variables. The matrix of Duncan test results is shown in Table 1.

Effects of branch pruning on broccoli growth and yield

The analysis of variance indicated that branch pruning significantly affected the number of leaves, initial leaf width, canopy diameter, stem diameter, and head diameter. The effect of branch pruning was not significant on the plant height. The data of the observed variables are shown in Table 2.

Branch pruning significantly affected the number of leaves. The treatment of 50 % branch pruning (P2) produced 19.07 leaves, which was 8.43 % higher than that in the plants treated with 25 % branch pruning (P1) and without pruning (P0). The leaf width after pruning was significantly different between each treatment. This result was because branch pruning caused some leaves were also disposed, hence the higher the percentage of disposed branches the smaller the leaf width. Canopy diameter was also significantly between each treatment, with the best result being shown in P2 (75.67 cm). Similar

result was also observed in the stem diameter with the best result being shown in P2 (3.35 cm). The treatment of P2 also resulted in the best values of yield component, including head diameter (24.56 cm) (Table 2).

Branch pruning treatment tended to decrease the plant's leaf width but increase the head diameter (Table 2). This result indicated that the light use efficiency was higher in those plants that have not so many leaves covering each other. The trimmed leaf surface did not overlap, thereby optimizing sunlight interception that will be used in the photosynthesis process (Niinemets et al., 2006), as what happened in the plants treated with 50 % branch pruning. Pasaribu et al. (2015) stated that leaves covering each other will inhibit the light penetration and photosynthetic processes, causing lower assimilation rate.

The high rate of light interception resulted in the high photosynthetic process and photosynthate products allocated to the increasing dry plant biomass. This is because light is an important factor in plant metabolism (Lakitan, 1996). Earlier, Van den Boogaard et al. (2001) reported an increase in the relative growth rate of the cauliflower plant after cutting. The research conducted by Jaya (2009) also indicated that broccoli branch pruning could accelerate harvest period, increase head weight and head diameter. An increase of head weight of around 23.7 % occurred when the branch was cut 100 % compared to the plant whose branch was not cut. Results of another study showed that the cutting of two and three main branches in eggplant gave the

Table 1. Variance analysis (F test) on branch pruning and chicken manure application

Observed variable	P	A	P×A
Plant height (cm)	ns	ns	ns
Number of leaves	*	*	ns
Initial leaf width (cm ²)	*	*	ns
Pruning leaf width (cm ²)	*	*	*
Canopy diameter (cm)	*	*	ns
Stem diameter (cm)	*	*	ns
Head weight (g.plant ⁻¹)	*	*	*
Head diameter (cm)	*	*	ns

Remarks: P=Pruning, A=Chicken manure application, P×A= Interaction of pruning and chicken manure; (*) = significantly different, (ns)= non-significant

Table 2. Effects of branch pruning on broccoli growth and yield

Treatment	PH (cm)	NL	ILW (cm ²)	CD (cm)	SD (cm)	HD (cm)
P0 (Without pruning)	45.11	16.29 c	1146.70 c	61.11 c	2.75 c	17.06 c
P1 (25% pruning)	49.01	17.59 b	1355.07 b	71.70 b	3.05 b	19.81 b
P2 (50% pruning)	47.89	19.08 a	1507.09 a	75.67 a	3.35 a	24.56 a
CV (%)	13.48	17.13 *	24.82 *	24.55 *	24.17 *	36.11 *

Remarks: Means in the same column followed by the same letters were not significantly different based on DMRT at $\alpha=5\%$. PH: Plant Height, NL: Number of Leaves, ILW: Initial Leaf Width, PLW: after Pruning Leaf Width, CD: Canopy Diameter, SD: Stem diameter, HW: Head Weight, HD: Head Diameter

Table 3. Broccoli growth and yield as effect of chicken manure fertilization

Treatment	PH (cm)	NL	ILW (cm ²)	CD (cm)	SD (cm)	HD (cm)
A0 (Without chicken manure)	38.48	15.12 c	1164.88 c	63.70 c	2.79 b	18.10 c
A1 (10 ton.ha ⁻¹)	49.59	17.78 b	1317.13 b	70.19 b	3.09 a	19.89 b
A2 (20 ton.ha ⁻¹)	53.93	20.07 a	1526.85 a	74.59 a	3.27 a	23.44 a
CV (%)	13.48	17.13 *	24.82 *	24.55 *	24.17 *	36.11 *

Remark: Means in the same column followed by the same letters were not significantly different based on DMRT at $\alpha=5\%$. PH: Plant Height, NL: Number of Leaf, ILW: Initial Leaf Width, PLW: after Pruning Leaf Width, CD: Canopy Diameter, SD: Stem diameter, HW: Head Weight, HD: Head Diameter

best result compared to without cutting. This was because too many leaves growing from the branches will act as sinks because they were covered by the leaves of the main stem, making it hard for the sunlight as the source in a photosynthetic process to be received optimally. According to Raden et al. (2009), the number of eggplants increased through controlling the number of the branch in the main stem by keeping the productive branch by which improving the allocation of the photosynthetic product. Reducing the branches could increase the intercepted light to the leaves, thereby improving the photosynthetic rate and directly increasing the weight and number of eggplants. The cutting of two and three main branches resulted in the higher weight of the eggplants (Maghfoer, 2014).

Effects of chicken manure application on broccoli growth and yield

Data analysis indicated that chicken manure had a significant effect on the number of leaves, initial leaf width, canopy diameter, stem diameter, and head diameter. The data of the observed variables

were shown in Table 3.

The results indicated that chicken manure application could improve the growth of broccoli. The application of 20 ton.ha⁻¹ chicken manure (A2) resulted in the highest number of leaves number (20.07). The similar results were also observed in leaf width before branch pruning, in which the highest values were resulted by the application of 20 ton.ha⁻¹ chicken manure (A2), which were 1526.85 cm². Meanwhile, the highest canopy and stem diameter were found in the plants treated with 10 ton.ha⁻¹ chicken manure (A1), which were 74.59 cm and 3.217 cm, consecutively. The application of 20 ton.ha⁻¹ chicken manure (A2) was also found to improve the yield component, resulting in the highest head diameter (23.44 cm) (Table 3).

The organic fertilization in the form of chicken manure could improve soil's fertility by applying 20 ton.ha⁻¹ chicken manure, producing the highest average in each observed variable than other treatments. This result was because the nutrient in chicken manure could be quickly absorbed by the plant, enhancing the growth rate. Nutrient contents

of chicken manure were water content of 57%, organic matter content of 19%, N content of 1.2%, P_2O_5 content of 1.1%, K_2O content of 0.6%, and C/N ratio of 7 (Laboratory of Soil Resources of Agriculture Faculty, Universitas Jenderal Soedirman). The ability of organic fertilizer to be absorbed by plants depends on its C/N ratio. The lower the C/N ratio, the quicker the fertilizer can be absorbed by the plants. When the C/N ratio is low, the mineralization process of N will be more dominant than N immobilization, enabling the organic matters to be the source of N for the plant. Besides, the use of organic fertilizer can also improve soil's biological quality, which could increase the microorganism activity and, thus could be expected to improve soil fertility (Mujiyati and Supriyadi, 2009).

An increased microorganism activity is an indicator of soil fertility that can improve soil structure (Dauda et al., 2008). Microorganisms are capable of turning hard soil into a more friable one, making the soil aeration better. As reported by Djazuli and Pitono's (2009), the application of chicken manure increased the weight of leaf, root, and entire plant of Purwoceng's (*Pimpinella pruatjan*) compared to the application of goat manure, cow manure, and compost.

Organic fertilizer has micronutrients content that are not contained in synthetic chemical fertilizers. Organic manure is also reported to be able to serve as alternative practice to mineral fertilizers (Naeem et al., 2006). According to Amannullah et al. (2010), chicken manure contains several macronutrients (N,

P, and K) highly needed by plant and some micro-nutrients (mangan (Mn), calcium (Ca), and iron (Fe)) and several other nutrients that could enhance plant production. To dissolve the nutrients needed by plant, a sufficient amount of water is needed to be the nutrient solvent to enable its absorption by the plant. The research by Jigme's et al. (2015) indicated that the application of 200 g.polybag⁻¹ chicken manure showed the best result in broccoli compared to other treatments. The MPHP⁺ of chicken manure treatment could increase the consumption weight by 159% to 165% higher than the treatment with no mulch in all types of fertilizers and increase the consumption weight by 144% to 206% higher than the straw mulch treatment in all types of fertilizers (Multazam, 2014).

The increase in available nutrients such as ammonium and nitrate as the effect of the chicken manure application resulted in the high total of tea yield (Gross et al., 2008). Thus, the application of chicken manure enhances soil aggregation, soil aeration, and water holding capacity. The favorable soil properties resulted in healthy plants with a large root system and vegetative growth by which gained the high yield and head diameter of broccoli plants. However, the highest yield was obtained by the application of inorganic treatment because of the high quantity of available nutrients.

Branch pruning significantly decreased the leaf width as the cutting percentage got higher. This phenomenon indicates that a high leaf width as a result of the many leaves growing on the broccoli




Figure 1. Effects of branch pruning and chicken manure application on the after pruning leaf width of broccoli; (P0)= without pruning; (P1)= 25% pruning; (P2)= 50% pruning; (A0)= without chicken manure; (A1)= 10 ton.ha⁻¹ chicken manure; (A2)= 20 ton.ha⁻¹ chicken manure.




Figure 2. Effects of branch pruning and chicken manure application on head weight of broccoli; (P0)= without pruning; (P1)= 25% pruning; (P2)= 50% pruning; (A0)= without chicken manure; (A1)= 10 ton.ha⁻¹ chicken manure; (A2)= 20 ton.ha⁻¹ chicken manure.

branches is less effective in forming biomass. This happened because many leaves growing from the branches acts as sinks since they are covered by the leaves from the main stem and/or many leaves have not developed perfectly. To avoid the reduction of plant growth and yield, pruning is used to control the distribution of photosynthetic apparatus, as well as to enhance the capacity of source in developing leaves to support the growth of stems and roots (Iqbal et al., 2012). However, the optimal result depends on the plant growth and development, such as defoliation process, growth stages, and age of leaf tissues (Stagnari et al., 2018). The highest leaf width was generated in a combined treatment of 50% branch pruning and 20 ton.ha⁻¹ chicken manure (P2A2), resulting in a leaf width of 1705.41 cm² (Figure 1). Similarly, the highest head weight was also produced in a combined treatment of 50% branch pruning and 20 ton.ha⁻¹ Chicken manure (P2A2), producing a head weight of 1048.33 g.plant⁻¹ (Figure 2). Some similar results were observed by Van Den Boogaard et al. (2001) and Li et al. (2012), reporting that pruning reduce internal shading to increase light interception per unit leaf area and directly enhances photosynthetic rate.

The large area of leaves could result in higher photosynthetic rate to translocate photosynthate to sink and obtain high yield. The high head weight is also supported by the increase in the leaf and head diameter (Singh et al., 2018).

CONCLUSIONS

The treatment of 50% branch pruning could improve the growth and yield of broccoli in lowland area. The head weight of plants treated with 50% branch pruning increased by 26.90% compared to that of plants treated with 25% pruning and without pruning. The application of 20 ton.ha⁻¹ chicken manure produced the highest head compared to the application of 20 ton.ha⁻¹ chicken manure and without chicken manure. The combined treatment of 50 % branch pruning and 20 ton.ha⁻¹ chicken manure produced the highest yield of broccoli of 1048.33 g.plant⁻¹.

ACKNOWLEDGMENT



The authors would like to thanks LPPM UNSOED that have facilitated this research through the “Penelitian Dosen Pemula” Research program funded from the budget of BLU UNSOED of 2015.

REFERENCES

- Amanullah, M.A., S. Sekar, and P. Muthukrishnan. 2010. Prospects and potential of poultry manure. *Asian Journal of Plant Sciences*, 3: 641–652.
- Badan Pusat Statistik. 2019. Produksi sayuran Indonesia. <https://www.bps.go.id/indicator/55/61/1/produksi-tanaman-sayuran.html>

- Björkman, T. and K. J. Pearson. 1998. High temperature arrest of inflorescence development in broccoli (*Brassica oleracea* var. *italica* L.). *Journal of Experimental Botany*, 49: 101–106.
- Booij, R. and P. C. Struik. 1990. Effect of temperature on leaf and curd initiation in relation to juvenility in cauliflower. *Scientia Horticulturae*, 44: 201–214.
- Budiastuti, S., D. Harjoko, and G. Shelti. 2009. Peningkatan potensi dan kualitas brokoli kopeng di Semarang Jawa Tengah melalui budidaya organik. *Agrivita*, 31 : 158–165.
- Dauda, S.N., F.A. Ajayi, and E. Ndor. 2008. Growth and yield of water melon (*Citrullus lanatus*) as affected by poultry manure application. *Journal of Agriculture and Social Science*, 4: 121–124
- Djazuli, M. and J. Pitono. 2009. Pengaruh jenis dan taraf pupuk organik terhadap produksi dan mutu purwoceng. *Jurnal Penelitian Tanaman Industri*, 15 : 40–45
- Gad, N. and M. R. A. El-Moez. 2011. Broccoli growth, yield quantity and quality as affected by cobalt nutrition. *Agriculture and Biology Journal of North America*, 2 : 226–231.
- Gomez, K. A. and A.A Gomez. 1983. *Statistical procedures for agriculture research*. 2nd ed., New York: John Wiley & Sons.
- Grevsen, K. 1998. Effect of temperature on head growth of broccoli (*Brassica oleracea* L. var. *italica*): parameter estimates for a predictive model. *Journal of Horticultural Science & Biotechnology*, 73: 235–244.
- Grevsen, K. and J. E. Olesen. 1999. Modelling development of broccoli (*Brassica oleracea* L. var *italica*) from transplanting to head initiation. *Journal of Horticultural Science & Biotechnology*, 74: 698–705.
- Gross, A., R. Arusi, and A. Nejidat. 2008. Assessment of extraction methods with fowl manure for the production of liquid organic fertilizers. *Bioresource Technology*, 99: 327–334.
- Hammad, H.S., A. A. M. Al-Mandalawi and G.J Hamdi. 2019. Effect of manure on growth and yield of broccoli. *International Journal of Vegetable Science*, 25, 400. (Abstr.).
- Heather, D. W., J. B. Sieczka, M. H. Dickson, and D. W. Wolfe. 1992. Heat tolerance and holding ability in broccoli. *Journal of American Society for Horticultural Science*, 117: 887–892.
- Iqbal, N., A. Masood, and N.A. Khan. 2012. Analyzing the significance of defoliation in growth, photosynthetic compensation and source-sink relations. *Photosynthetica*, 50: 161–170.
- Jaya, I. K. D. 2009. Pengaruh pemangkasan cabang terhadap hasil tanaman brokoli (*Brassica oleracea* var *italica*) di dataran rendah. *Crop Agro*, 2: 15–21
- Jaya, I. K. D., N. Novianthy, and M. Martajaya, 2006. Pertumbuhan, perkembangan dan hasil tanaman brokoli (*Brassica oleracea* L. var *italica*) di dataran rendah. *Prosiding Seminar Nasional Hortikultura Perhimpunan Hortikultura Indonesia*. 91–96.
- Jigme, N. Jayamangkala, P. Sutigoolabud, J. Inthasan and S. Sakhonwasee. 2015. The effect of organic fertilizers on growth and yield of broccoli (*Brassica oleracea* L. var. *italica* Plenck cv. Top Green). *Journal of Organic Systems*, 10: 9–14
- Lakitan, B. 1996. *Fisiologi pertumbuhan dan perkembangan tanaman*. 1st ed., Jakarta: Grafindo Persada.
- Li, W., J. Luo, X. Tian, C. Peng, and X. Zhou. 2012. Patterns of defoliation and their effect on the plant growth and photosynthetic characteristics of *Ipomoea cairica*. *Weed Biology and Management*, 12: 40–46.
- Maghfoer, M.D., R. Soelistiyono, and N. Herlina. 2014. Response eggplant (*Solanum melongena*) to combination of inorganic-organic N and EM4. *Agrivita*, 35: 296–303
- Mujiyati and Supriyadi. 2009. Pengaruh pupuk kandang dan NPK terhadap populasi bakteri *Azotobacter* dan *Azospirillum* dalam tanah pada budidaya cabai (*Capsicum annum*). *Bioteknologi*, 6 : 63–69.
- Multazam, M.A., A. Suryanto, and N. Herlina. 2014. Pengaruh macam pupuk organik dan mulsa pada tanaman brokoli (*Brassica oleracea* L. Var. *Italica*). *Jurnal Produksi Tanaman*, 2: 154–161.
- Musnamar, E.I. 2004. *Pupuk organik : cair dan padat, pembuatan, aplikasi*. 10th ed., Jakarta: Penebar Swadaya.
- Naeem, M., J. Iqbal and M.A.A. Bakhsh, 2006. Comparative study of inorganic fertilizers and organic manures on yield and yield components of mungbean (*Vigna radiata* L.). *Journal of Agriculture and Social Science*, 2: 227–229.
- Niinemets, U., A. Portsmouth, and M. Tobias, 2006. Leaf size modifies support biomass distribution among stems, petioles and mid-ribs in temperate plants. *New Phytologist*, 171: 91–104.

- Ouda, B.A. and A.Y. Mahadeen, 2008. Effect of fertilizers on growth, yield, yield components, quality and certain nutrient contents in broccoli (*Brassica oleracea*). *International Journal of Agriculture and Biology*, 10: 627–32
- Pasaribu, R.P., H. Yetti, and Nurbaiti. 2015. Pengaruh pemangkasan cabang utama dan pemberian pupuk pelengkap cair organik terhadap pertumbuhan dan produksi tanaman tomat (*Lycopersicum esculentum* Mill.). JOM Faperta, 2.
- Pinem, D.Y.F., T. Irmansyah, and F. E. T. Sitepu. 2015. Respons pertumbuhan dan produksi brokoli terhadap pemberian pupuk kandang ayam dan jamur pelarut fosfat. *Jurnal Online Agroekoteknologi*, 3: 198–205
- Raden, I., B.S. Purwoko, Hariadi, M. Ghulamahdi, and E. Santosa. 2009. Pengaruh tinggi pangkasan batang utama dan jumlah cabang primer yang dipelihara terhadap produksi minyak jarak. *Jurnal Agronomi Indonesia*, 37: 159–166.
- Simanungkalit, R.D.M.; D.A. Suriadikarta; R. Saraswati; D. Setryorini; and W. Hartatik. 2006. *Pupuk organik dan pupuk hayati*. 1st ed., Bogor: Balai Besar Litbang Sumberdaya Lahan Pertanian. Badan Penelitian dan Pengembangan Pertanian. Balai Besar Penelitian dan Pengembangan Sumberdaya Lahan Pertanian.
- Singh, G., S. Sarvanan, A. Kerketta, and J. Rajesh. 2018. Effect of organic manures and inorganic fertilizers on plant growth, yield and flower bud quality of broccoli (*Brassica oleracea* var. Italica) cv-Green Magic. *International Journal of Pure & Applied Bioscience*, 6: 1338–1342.
- Stagnari, F., A. Galieni, S. D’egidio, G. Pagnani, N. Ficcadenti, and M. Pisant. 2018. Defoliation and S nutrition on radish: growth, polyphenols and antiradical activity. *Horticultura Brasileira*, 36: 313–319.
- Van den Boogaard, R., K. Grevsen and K. Thorup-Kristensen, 2001. Effects of defoliation on growth of cauliflower. *Scientia Horticulturae*, 91: 1–16.
- Wasnowati, C. 2009. Kajian saat pemberian pupuk dasar nitrogen dan umbi bibit pada tanaman brokoli (*Brassica oleraceae* L.). *Agrovigor*, 2: 14–22.
- Wien, H. C. and D. C. E. Wurr. 1997. *Cauliflower, broccoli, cabbage and brussels sprouts*, p.662. In: H.C. Wien (ed.). *The Physiology of Vegetable Crops*. Wallingford UK: CAB International.



Internet Hotspot x [ipad] Editor Deci x www.google.com x 77620074 (1).pdf x Berisikan Teknis x article.php x www.google.com x

mail.google.com/mail/u/0/#search/JIP+UGM/RMfcgwXfprKzpDGzQitMnWkLUSjWT

Anda kehabisan ruang penyimpanan. Coba [kosongkan ruang](#) atau [beli penyimpanan tambahan](#).

Gmail JIP UGM

Tulis

Kotak Masuk 538

Berbintang

Ditunda

Penting

Terakhir

Draf 139

Muat

Mulai rapat

Gabung ke rapat

Hangout

Old -

ane wieshia

[ipad] Editor Decision

Enik Nurlaili Afifah - enik.nurlaili.aj@mail.ugm.ac.id

Rab, 2 Sep 09 24

Old Heriana

We have reached a decision regarding your submission to Ilmu Pertanian (Agricultural Science), "Branch Pruning and Chicken Manure Application to Improve Growth and Yield Broccoli in Lowland".

Our decision is to: Accept Submission

To facilitate prompt publication, please pay your author publication charges 1000000.00 (IDR) in the account (BNB) 9588812014100653 with name UGM FPN KAF PUBLIKASI JURNAL ILMIAH

If you have already paid, please send the form transactions in this email within 2020/09/09. If you have any questions about these charges, please contact us by replying to this email.

Thank you

Erik Nurlaili Afifah
Department of Agronomy, Faculty of Agriculture, Universitas Gadjah Mada
Phone 081215952747

PASCAPANEN KEL...pdf 77620074 (1).pdf DPT SEMINAR_D...pptx registrasi video eks...jpg Banner-Semnas-20...jpg Show all X

Internet Hotspot x [ipad] Editor Revisi x www.google.com x 77620074 (1).pdf x Berisikan Teknis x article.php x www.google.com x

mail.google.com/mail/u/0/#search/JIP+UGM/RMfcgwXfprKzpDGzQitMnWkLUSjWT

Anda kehabisan ruang penyimpanan. Coba [kosongkan ruang](#) atau [beli penyimpanan tambahan](#).

Gmail JIP UGM

Tulis

Kotak Masuk 538

Berbintang

Ditunda

Penting

Terakhir

Draf 139

Muat

Mulai rapat

Gabung ke rapat

Hangout

Old -

ane wieshia

[ipad] Editor Revision

Admin JIP - jip.faperta@ugm.ac.id

Sab, 1 Sep 08 30



Old Heriana

Kami informasikan bahwa tim editor telah memeriksa perbaikan naskah anda. Mohon untuk memperbaiki kalimat di Bab Hasil dan Pembahasan menjadi Past Tense dan melengkapi edisi buku dari referensi Musnandar 2004. Selengkapnya dapat dicek di file terlampir.

Demikian informasi ini kami sampaikan. Atas perhatian Ibu, kami ucapkan terimakasih.

Admin JIP
Faculty of Agriculture, Universitas Gadjah Mada
Phone +628112632106
jip.faperta@ugm.ac.id
Ilmu Pertanian (Agricultural Science) ISSN 0126-4214 (print), ISSN 2527-7162 (online)
Faculty of Agriculture, Gadjah Mada University
Jalan Flora, Bulaksumur, Yogyakarta 55281
<http://jurnal.ugm.ac.id/jip>

PASCAPANEN KEL...pdf 77620074 (1).pdf DPT SEMINAR_D...pptx registrasi video eks...jpg Banner-Semnas-20...jpg Show all X



Internet Hotspot x [psd] Editor Deci x www.google.com x 77620074 (1).pdf x Berisikan Teknis x article.php x www.google.com x

mail.google.com/mail/u/0/#search/JIP+UGM/RMfcgw/XVPx2enptklpXMBTQTvqmQkQv

Anda kehabisan ruang penyimpanan. Coba [kosongkan ruang](#) atau [beli penyimpanan tambahan](#).

Gmail JIP UGM

Tulis

Kotak Masuk 538

Berbintang

Ditunda

Penting

Terakhir

Draf 139

Muat

Mulai rapat

Gabung ke rapat

Hangout

Okt -

aria wiedzha

Admin JIP <jip.faperta@ugm.ac.id>
Kepada saya -
Okt Herliana

Sen, 24 Agu 07:42

Your article has been reviewed by all reviewers. Please revise based on the reviewer's comments and editor's comments on this previous email.
Please revise your article by 2020/08/31

Admin JIP
Faculty of Agriculture, Universitas Gadjah Mada
Phone +628112632106
jip.faperta@ugm.ac.id
Ilmu Pertanian (Agricultural Science) ISSN 0126-4214 (print), ISSN 2527-7162 (online)
Faculty of Agriculture, Gadjah Mada University
Jalan Flora, Bulaksumur, Yogyakarta 55281
<http://jurnal.ugm.ac.id/jip>

Okti Herliana <o.herliana@gmail.com>
Kepada Admin -
Dear Editor of JIP

Kam, 27 Agu 13:25

PASCAPANEN KEL...pdf 77620074 (1).pdf DPT SEMINAR_D...pptx registrasi video eks...jpg Banner-Semnas-20...jpg Show all X

Internet Hotspot x [psd] Editor Deci x www.google.com x 77620074 (1).pdf x Berisikan Teknis x article.php x www.google.com x

mail.google.com/mail/u/0/#search/JIP+UGM/RMfcgw/XVPx2enptklpXMBTQTvqmQkQv

Anda kehabisan ruang penyimpanan. Coba [kosongkan ruang](#) atau [beli penyimpanan tambahan](#).

Gmail JIP UGM

Tulis

Kotak Masuk 538

Berbintang

Ditunda

Penting

Terakhir

Draf 139

Muat

Mulai rapat

Gabung ke rapat

Hangout

Okt -

aria wiedzha

Okti Herliana <o.herliana@gmail.com>
Kepada Admin -
Dear Editor of JIP

Kam, 27 Agu 13:25

Thank you for give me information about my revised paper from both of reviewer. I have sent my revised paper to OJS and waiting for final decision. I wish my paper can be accept and published on Journal Ilmu Pertanian (JIP UGM).
Thank you for your attention.

Regards

Okti Herliana
Laboratorium Agroekologi Fakultas Pertanian
Universitas Jenderal Soedman
xxx

43101-161036-1-R...

PASCAPANEN KEL...pdf 77620074 (1).pdf DPT SEMINAR_D...pptx registrasi video eks...jpg Banner-Semnas-20...jpg Show all X