



Source details

IOP Conference Series: Earth and Environmental Science

Scopus coverage years: from 2010 to Present

Publisher: Institute of Physics Publishing

ISSN: 1755-1307 E-ISSN: 1755-1315

Subject area: [Earth and Planetary Sciences: General Earth and Planetary Sciences](#)

[Environmental Science: General Environmental Science](#)

Source type: Conference Proceeding

CiteScore 2021
0.6

SJR 2021
0.202

SNIP 2021
0.409

[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Source Homepage](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

Improved CiteScore methodology

CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore 2021

$$0.6 = \frac{45.063 \text{ Citations 2018 - 2021}}{74.324 \text{ Documents 2018 - 2021}}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022

$$0.6 = \frac{39.465 \text{ Citations to date}}{68.948 \text{ Documents to date}}$$

Last updated on 06 June, 2022 • Updated monthly

CiteScore rank 2021

Category	Rank	Percentile
Earth and Planetary Sciences	#153/191	20th
General Earth and Planetary Sciences		
Environmental Science	#191/228	16th
General Environmental Science		

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)

About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

ELSEVIER

[Terms and conditions](#) ↗ [Privacy policy](#) ↗

Copyright © [Elsevier B.V](#) ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the [use of cookies](#) ↗.



PAPER • OPEN ACCESS

The 3rd International Conference on Life and Applied Sciences *for Sustainable Rural Development* (ICLAS-SURE)

<http://iclas.conference.unsoed.ac.id/>

To cite this article: 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **746** 011001

View the [article online](#) for updates and enhancements.

You may also like

- [Analysis of Blue Swimmer Crab \(*Portunus Pelagicus*\) Processing Efficiency In The Sort Stage In Pt. Blue Star Anugrah Cold Storage Company, Pemalang](#)
T Junaidi, U F Arafah, A Margiwyatno et al.
- [Optimization of wavelet- and curvelet-based denoising algorithms by multivariate SURE and GCV](#)
R Mortezaejad and A Gholami
- [11th Joint Conference on Chemistry in Conjunction with the 4th Regional Biomaterials Scientific Meeting](#)



The Electrochemical Society
Advancing solid state & electrochemical science & technology

242nd ECS Meeting

Oct 9 – 13, 2022 • Atlanta, GA, US

Abstract submission deadline: **April 8, 2022**

Connect. Engage. Champion. Empower. Accelerate.

MOVE SCIENCE FORWARD



Submit your abstract



PREFACE

International Conference on Life and Applied Sciences for Sustainable Rural Development (ICLAS-SURE) is an annual international event organized by Institute of Research and Community Service, Universitas Jenderal Soedirman (Unsoed), Indonesia. Universitas Jenderal Soedirman (Unsoed) is one of the outstanding National University in Indonesia, which is located in Purwokerto, Central Java, Indonesia. This university was established by Minister of Higher Education and Science, Republic Indonesia, based on Presidential Decree No. 195 dated September 23, 1963. Since 1963, Universitas Jenderal Soedirman has been experiencing on rural resource development as well as community services.

Following the success of the 1st and 2nd **ICLAS-SURE**, this year, the **Institute of Research and Community Service, Universitas Jenderal Soedirman**, organize **The 3rd ICLAS-SURE**. The vision of Jenderal Soedirman University is to be **globally recognized as a university that focuses on sustainable rural and local wisdom development**. Hopefully, this core competence in sustainable rural development shall initiate the university to be nationally and internationally renowned as the center of rural community empowerment. To achieve this vision and cope with the COVID 19 pandemic, this year, we bring the particular theme, "Interdisciplinary approaches and applied technologies for sustainable rural-environmental resources based on local wisdom before and during COVID-19 pandemic". COVID-19 has led to a significant loss of output, employment, and income, affecting rural development. To develop a sustainable rural development, we must fulfil three basic needs, i.e. people welfare improvement, protection of natural, landscape, and cultural resources, and food security through a sustainable farming production.



COMMITTEE

Steering Committee

- Prof. Rifda Naufalin
<https://www.scopus.com/authid/detail.uri?authorId=57193676582>
- Prof. II Soo Moon - Dongguk University, South Korea
<https://www.scopus.com/authid/detail.uri?authorId=7101610481>
- Prof. B Mohan Kumar - Kerala Agricultural University, India
<https://www.scopus.com/authid/detail.uri?authorId=55435104500>
- Prof Peter Idowu - PennState Univ, USA
<https://www.scopus.com/authid/detail.uri?authorId=6603845249>
- Dr. Yeong Sheng Tey, - Universiti Putra Malaysia
<https://www.scopus.com/authid/detail.uri?authorId=26026139600>
- Prof. Oceandy Delvac - Manchester Universit, England
<https://www.scopus.com/authid/detail.uri?authorId=6506557120>

Organizing Committee

- Assoc. Prof. Maria Dyah Nur Meinita
<https://www.scopus.com/authid/detail.uri?authorId=37061363700>
- Prof. Retno Supriyanti
<https://www.scopus.com/authid/detail.uri?authorId=24332824200>
- Assoc. Prof. Poppy Arsil
<https://www.scopus.com/authid/detail.uri?authorId=56131258600>

Editorial Board

- Eko Murdyantoro, MT
<https://www.scopus.com/authid/detail.uri?authorId=56028724800>
- Amin Fatoni, Ph.D.
<https://www.scopus.com/authid/detail.uri?authorId=55488648900>
- Professor Jae-Suk Choi
Silla University, South Korea
<https://www.scopus.com/authid/detail.uri?authorId=35104615200>
- Professor Md. Abdul Hannan, PhD
Bangladesh Agricultural University, Bangladesh
<https://www.scopus.com/authid/detail.uri?authorId=55122778800>
- Assistant Professor Dr. Eng. Paulos Getachew
Addis Ababa University, Ethiopia
<https://www.scopus.com/authid/detail.uri?authorId=55820965900>

PAPER • OPEN ACCESS

Peer Review Declaration

To cite this article: 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **746** 011002

View the [article online](#) for updates and enhancements.

You may also like

- [Peer review declaration](#)
- [Peer review declaration](#)
- [Peer review declaration](#)



The Electrochemical Society
Advancing solid state & electrochemical science & technology

242nd ECS Meeting

Oct 9 – 13, 2022 • Atlanta, GA, US

Abstract submission deadline: **April 8, 2022**

Connect. Engage. Champion. Empower. Accelerate.

MOVE SCIENCE FORWARD



Submit your abstract



PEER REVIEW DECLARATION

All papers published in this volume of IOP Conference Series: Earth and Environmental Science have been peer reviewed through processes administered by the Editors. Reviews were conducted by expert referees to the professional and scientific standards expected of a proceedings journal published by IOP Publishing.

The ICLAS-SURE committee is committed to maintaining the highest level of integrity in the content published paper in The 3rd ICLAS-SURE. All the papers we received have been peer-reviewed. There are two steps of the review process we conducted

1. Preliminary Review

All papers submitted to ICLAS-SURE must fit the scope of The 3rd ICLAS-SURE. The scope was checked by the Editorial board. Then all of the papers had undergone a plagiarism check, English grammar check, and double-blind review by two reviewers. Based on the reviewer comments and the result of plagiarism and grammar check, we decided the paper which can be processed to the next step

- **Type of peer review:** Double-blind, author and reviewer identities are hidden to each other
- **Scope:** Biosciences, Agriculture, Engineering and applied sciences for rural development.
- **Plagiarism and grammar checking:** Turnitin & Grammarly
- **Conference submission management system:** Open Conference Systems (OCS).

2. Content review

The papers passed the first review were reviewed to the next step, review by Scientific Committee and reviewers. The from the following aspects: Originality, Methodology, Novelty and Scientific Structure.

- Type of peer review: Double-blind by Scientific Committee and reviewers
- Number of abstract presented : 182 titles
- Number of submissions paper sent for review : 113 papers
- Number of submissions accepted: 45 papers
- Acceptance Rate (Number of Submissions Accepted / Number of Submissions Received X 100): 39,8 %
- Average number of reviews per paper: 2 round
- Total number of reviewers involved: 22 reviewers

Contact person for queries:

Eko Murdyantoro AM

Jenderal Soedirman University, Jawa Tengah, Indonesia

Email : icmeasure.lppm@unsoed.ac.id



Table of contents

[Volume 746](#)

2021

[Previous issue](#)[Next issue](#)

3rd International Conference on Life and Applied Sciences for Sustainable Rural Development (ICLAS-SURE 2020) November 18-19, 2020 Central Java, Indonesia

Accepted papers received: 09 April 2021

Published online: 13 May 2021

[Open all abstracts](#), in this issue

Preface

011001

THE FOLLOWING ARTICLE IS OPEN ACCESS

[The 3rd International Conference on Life and Applied Sciences for Sustainable Rural Development \(ICLAS-SURE\)](#)<http://iclas.conference.unsoed.ac.id/>

[Open abstract](#), The 3rd International Conference on Life and Applied Sciences for Sustainable Rural Development (ICLAS-SURE) <http://iclas.conference.unsoed.ac.id/> [View article](#), The 3rd International Conference on Life and Applied Sciences for Sustainable Rural Development (ICLAS-SURE) <http://iclas.conference.unsoed.ac.id/> [PDF](#), The 3rd International Conference on Life and Applied Sciences for Sustainable Rural Development (ICLAS-SURE) <http://iclas.conference.unsoed.ac.id/>

011002

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Peer Review Declaration](#)

[Open abstract](#), [Peer Review Declaration](#) [View article](#), [Peer Review Declaration](#) [PDF](#), [Peer Review Declaration](#)

Agriculture for Rural Development

012001

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Comparative study between cow and goat milk yogurt based on composition and sensory evaluation](#)

A Ibrahim, R Naufalin, E Muryatmo and H Dwiyantri

[Open abstract](#), [Comparative study between cow and goat milk yogurt based on composition and sensory evaluation](#) [View article](#), [Comparative study between cow and goat milk yogurt based on composition and sensory evaluation](#) [PDF](#), [Comparative study between cow and goat milk yogurt based on composition and sensory evaluation](#)

012002

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Homegardening for food and nutritional security and for biodiversity conservation during the pandemic times](#)

B. Mohan Kumar

[Open abstract](#), Homegardening for food and nutritional security and for biodiversity conservation during the pandemic times [View article](#), Homegardening for food and nutritional security and for biodiversity conservation during the pandemic times [PDF](#), Homegardening for food and nutritional security and for biodiversity conservation during the pandemic times

012003

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Induce resistance of rice plants against bacterial leaf blight by using salicylic acid application](#)

W S Suharti and N W A Leana

[Open abstract](#), Induce resistance of rice plants against bacterial leaf blight by using salicylic acid application [View article](#), Induce resistance of rice plants against bacterial leaf blight by using salicylic acid application [PDF](#), Induce resistance of rice plants against bacterial leaf blight by using salicylic acid application

012004

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Effect of edible coating application by spraying method on the quality of red chili during storage](#)

C Wibowo, P Haryanti and R Wicaksono

[Open abstract](#), Effect of edible coating application by spraying method on the quality of red chili during storage [View article](#), Effect of edible coating application by spraying method on the quality of red chili during storage [PDF](#), Effect of edible coating application by spraying method on the quality of red chili during storage

012005

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Key drivers of organic rice productivity in Sleman and Magelang Regencies](#)

Laksmi Yustika Devi, Irham, Subejo, Esti Anatasari, Azizatun Nurhayati and Arif Wahyu Widada

[Open abstract](#), Key drivers of organic rice productivity in Sleman and Magelang Regencies [View article](#), Key drivers of organic rice productivity in Sleman and Magelang Regencies [PDF](#), Key drivers of organic rice productivity in Sleman and Magelang Regencies

012006

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Nutrient digestibility, intake rate, and performance of Indonesian native cattle breeds fed rice straw ammoniation and concentrate](#)

Muhamad Bata, Sri Rahayu and Efka Aris Rimbawanto

[Open abstract](#), Nutrient digestibility, intake rate, and performance of Indonesian native cattle breeds fed rice straw ammoniation and concentrate [View article](#), Nutrient digestibility, intake rate, and performance of Indonesian native cattle breeds fed rice straw ammoniation and concentrate [PDF](#), Nutrient digestibility, intake rate, and performance of Indonesian native cattle breeds fed rice straw ammoniation and concentrate

012007

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Functional properties of hydrothermally modified lesser yam \(*Dioscorea esculenta*\) starch](#)

Laksmi Putri Ayuningtyas, Ashri Mukti Benita and Desy Triastuti

[Open abstract](#), Functional properties of hydrothermally modified lesser yam (*Dioscorea esculenta*) starch [View article](#), Functional properties of hydrothermally modified lesser yam (*Dioscorea esculenta*) starch [PDF](#), Functional properties of hydrothermally modified lesser yam (*Dioscorea esculenta*) starch

012008

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Extraction time optimization of antibacterial activities of kecombrang flower extract with microwave assisted extraction \(MAE\) method](#)

R Naufalin, Erminawati, N Herliya and N Latifasari

[Open abstract](#), Extraction time optimization of antibacterial activities of kecombrang flower extract with microwave assisted extraction (MAE) method [View article](#), Extraction time optimization of antibacterial activities of kecombrang flower extract with microwave assisted extraction (MAE) method [PDF](#), Extraction time optimization of antibacterial activities of kecombrang flower extract with microwave assisted extraction (MAE) method

012009

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Volatile compounds profile of some Indonesian shallot varieties](#)

Siti D Indrasari, Desi Arofah, Kristamtini, Sudarmaji and Dody D Handoko

[Open abstract](#), Volatile compounds profile of some Indonesian shallot varieties [View article](#), Volatile compounds profile of some Indonesian shallot varieties [PDF](#), Volatile compounds profile of some Indonesian shallot varieties

012010

THE FOLLOWING ARTICLE IS OPEN ACCESS

[The implementation of rice's Good Agricultural Practices \(GAP\) in Panarukan-Situbondo](#)

G. I. A. Yekti and Y. Suryaningsih

[Open abstract](#), The implementation of rice's Good Agricultural Practices (GAP) in Panarukan-Situbondo [View article](#), The implementation of rice's Good Agricultural Practices (GAP) in Panarukan-Situbondo [PDF](#), The implementation of rice's Good Agricultural Practices (GAP) in Panarukan-Situbondo

012011

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Triple helix as an empowerment strategy for labor fishermen: a proposed model through action research study](#)

Laeli Budiarti, Christina Tri Setyorini, Dewi Susilowati, Warsidi, Purnama Sukardi and Miftahul Jannah

[Open abstract](#), Triple helix as an empowerment strategy for labor fishermen: a proposed model through action research study [View article](#), Triple helix as an empowerment strategy for labor fishermen: a proposed model through action research study [PDF](#), Triple helix as an empowerment strategy for labor fishermen: a proposed model through action research study

012012

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Farmers' term of trade in Indonesia: an overview during pandemic COVID-19](#)

M Pinilih, D Rakhmawati and R Rosyidi

[Open abstract](#), Farmers' term of trade in Indonesia: an overview during pandemic COVID-19 [View article](#), Farmers' term of trade in Indonesia: an overview during pandemic COVID-19 [PDF](#), Farmers' term of trade in Indonesia: an overview during pandemic COVID-19

012013

THE FOLLOWING ARTICLE IS OPEN ACCESS

[The factors contributing to the sustainability of agribusiness MSMEs in Sukoharjo Regency during the Covid-19 pandemic](#)

I Khomah, N Setyowati, M Harisudin, R K Adi and A Qonita

[Open abstract](#), The factors contributing to the sustainability of agribusiness MSMEs in Sukoharjo Regency during the Covid-19 pandemic [View article](#), The factors contributing to the sustainability of

agribusiness MSMEs in Sukoharjo Regency during the Covid-19 pandemic [PDF](#), The factors contributing to the sustainability of agribusiness MSMEs in Sukoharjo Regency during the Covid-19 pandemic

012014

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Marker identification and phylogenetic analysis of saline tolerant rice varieties](#)

Suprayogi, P S Dewi, E Oktaviani, A W Aisya and R G N Prasetya

[Open abstract](#), Marker identification and phylogenetic analysis of saline tolerant rice varieties [View article](#), Marker identification and phylogenetic analysis of saline tolerant rice varieties [PDF](#), Marker identification and phylogenetic analysis of saline tolerant rice varieties

012015

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Perspectives on the development of local food policy using the Analytical Hierarchy Process](#)

P Arsil, K E Sularso, A Mulyani and N F Hardana

[Open abstract](#), Perspectives on the development of local food policy using the Analytical Hierarchy Process [View article](#), Perspectives on the development of local food policy using the Analytical Hierarchy Process [PDF](#), Perspectives on the development of local food policy using the Analytical Hierarchy Process

012016

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Adaption of local rice cultivars Banten to drought environment](#)

Rusmana, S Ritawati, I Rohmawati and E P Ningsih

[Open abstract](#), Adaption of local rice cultivars Banten to drought environment [View article](#), Adaption of local rice cultivars Banten to drought environment [PDF](#), Adaption of local rice cultivars Banten to drought environment

012017

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Antioxidant activity of kecombrang preserving powder using foam mat drying method](#)

R Naufalin, Erminawati, R Wicaksono, A T Febryani and N Latifasari

[Open abstract](#), Antioxidant activity of kecombrang preserving powder using foam mat drying method [View article](#), Antioxidant activity of kecombrang preserving powder using foam mat drying method [PDF](#), Antioxidant activity of kecombrang preserving powder using foam mat drying method

Biosciences for Rural Development

012018

THE FOLLOWING ARTICLE IS OPEN ACCESS

[A comparison of the effectiveness banana stem sap and virgin coconut oil on diabetic wound healing](#)

Yunita Sari, Atyanti Isworo, Arif Setyo Upoyo, Annas Sumeru, Dhadhang Wahyu

Kurniawan and Eman Sutrisna

[Open abstract](#), A comparison of the effectiveness banana stem sap and virgin coconut oil on diabetic wound healing [View article](#), A comparison of the effectiveness banana stem sap and virgin coconut oil on diabetic wound healing [PDF](#), A comparison of the effectiveness banana stem sap and virgin coconut oil on diabetic wound healing

012019

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Diversity of Introduced Species of Fishes in Penjalin Reservoir Central Java Indonesia](#)

N Setyaningrum, Sugiharto and P Susatyo

[Open abstract](#), [Diversity of Introduced Species of Fishes in Penjalin Reservoir Central Java Indonesia](#) [View article](#), [Diversity of Introduced Species of Fishes in Penjalin Reservoir Central Java Indonesia](#) [PDF](#), [Diversity of Introduced Species of Fishes in Penjalin Reservoir Central Java Indonesia](#)

012020

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Investigation of condition factor of wild spiny lobster juvenile *Panulirus* spp. inhabit in Cilacap waters, Indonesia](#)

F E D Haryono, T Winanto, Amron, M Trenggono, R T Harisam and D Wisudyanti

[Open abstract](#), [Investigation of condition factor of wild spiny lobster juvenile *Panulirus* spp. inhabit in Cilacap waters, Indonesia](#) [View article](#), [Investigation of condition factor of wild spiny lobster juvenile *Panulirus* spp. inhabit in Cilacap waters, Indonesia](#) [PDF](#), [Investigation of condition factor of wild spiny lobster juvenile *Panulirus* spp. inhabit in Cilacap waters, Indonesia](#)

012021

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Species diversity and conservation status of marine ornamental fish traded at three market spots in the southern coast of West Java](#)

A Nuryanto, D Bhagawati and Kusbiyanto

[Open abstract](#), [Species diversity and conservation status of marine ornamental fish traded at three market spots in the southern coast of West Java](#) [View article](#), [Species diversity and conservation status of marine ornamental fish traded at three market spots in the southern coast of West Java](#) [PDF](#), [Species diversity and conservation status of marine ornamental fish traded at three market spots in the southern coast of West Java](#)

012022

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Mangrove cluster as adaptation pattern of mangrove ecosystem in Segara Anakan Lagoon](#)

Endang Hilmi, Lilik Kartika Sari, Amron, Tri Nur Cahyo and Asrul Sahri Siregar

[Open abstract](#), [Mangrove cluster as adaptation pattern of mangrove ecosystem in Segara Anakan Lagoon](#) [View article](#), [Mangrove cluster as adaptation pattern of mangrove ecosystem in Segara Anakan Lagoon](#) [PDF](#), [Mangrove cluster as adaptation pattern of mangrove ecosystem in Segara Anakan Lagoon](#)

012023

THE FOLLOWING ARTICLE IS OPEN ACCESS

[The Primary culture of caudal fin, gill lamella, hepatopancreas and spleen of *Osteochilus vittatus*](#)

Gratiana E. Wijayanti and Atang

[Open abstract](#), [The Primary culture of caudal fin, gill lamella, hepatopancreas and spleen of *Osteochilus vittatus*](#) [View article](#), [The Primary culture of caudal fin, gill lamella, hepatopancreas and spleen of *Osteochilus vittatus*](#) [PDF](#), [The Primary culture of caudal fin, gill lamella, hepatopancreas and spleen of *Osteochilus vittatus*](#)

012024

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Tofu wastewater industry with urea fertilizer as a cultivation medium for the microalga *Spirulina plantensis*](#)

P HT Soedibya, T B Pramono, P Sukardi, B Kusuma, S Marnani, R Fitriadi and T Aditama

[Open abstract](#), [Tofu wastewater industry with urea fertilizer as a cultivation medium for the microalga *Spirulina plantensis*](#) [View article](#), [Tofu wastewater industry with urea fertilizer as a](#)

cultivation medium for the microalga *Spirulina plantensis* [PDF](#), Tofu wastewater industry with urea fertilizer as a cultivation medium for the microalga *Spirulina plantensis*

012025

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Composition and diversity of macroalgae community in the coast of Karang Bolong, Nusakambangan Island](#)

Dwi Sunu Widyartini, Hernayanti and Romanus Edy Prabowo

[Open abstract](#), Composition and diversity of macroalgae community in the coast of Karang Bolong, Nusakambangan Island [View article](#), Composition and diversity of macroalgae community in the coast of Karang Bolong, Nusakambangan Island [PDF](#), Composition and diversity of macroalgae community in the coast of Karang Bolong, Nusakambangan Island

012026

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Distribution and accumulation of heavy metals from waters and sediments to *Scylla serrata* in Segara Anakan, Cilacap](#)

M H Sastranegara, W Lestari, E Sudiana, Oedjijono and E K Nasution

[Open abstract](#), Distribution and accumulation of heavy metals from waters and sediments to *Scylla serrata* in Segara Anakan, Cilacap [View article](#), Distribution and accumulation of heavy metals from waters and sediments to *Scylla serrata* in Segara Anakan, Cilacap [PDF](#), Distribution and accumulation of heavy metals from waters and sediments to *Scylla serrata* in Segara Anakan, Cilacap

012027

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Nutritional information access and dietary behavior among people with diabetes during Covid-19 pandemic](#)

Yovita Puri Subardjo, Gumintang Ratna Ramadhan, Dika Betaditya, Muflihatus

Syarifah and Nurafifah Fauziana Abidin

[Open abstract](#), Nutritional information access and dietary behavior among people with diabetes during Covid-19 pandemic [View article](#), Nutritional information access and dietary behavior among people with diabetes during Covid-19 pandemic [PDF](#), Nutritional information access and dietary behavior among people with diabetes during Covid-19 pandemic

012028

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Screening of microfungi from spent mushroom for decolorizing and removing heavy metals from batik effluent and its toxicity](#)

Ratna Stia Dewi and Hana

[Open abstract](#), Screening of microfungi from spent mushroom for decolorizing and removing heavy metals from batik effluent and its toxicity [View article](#), Screening of microfungi from spent mushroom for decolorizing and removing heavy metals from batik effluent and its toxicity [PDF](#), Screening of microfungi from spent mushroom for decolorizing and removing heavy metals from batik effluent and its toxicity

012029

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Several ecological factors that determine the survival of temperature resistant *Phytoseius amba*](#)

Bambang Heru Budianto, Rokhmani and Edi Basuki

[Open abstract](#), Several ecological factors that determine the survival of temperature resistant *Phytoseius amba* [View article](#), Several ecological factors that determine the survival of temperature resistant *Phytoseius amba* [PDF](#), Several ecological factors that determine the survival of temperature resistant *Phytoseius amba*

012030

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Investigation of total organic matter \[TOM\] content during high and low water in inter-tidal zone sediment at Teluk Penyu Coast, Cilacap, Indonesia](#)

F E D Haryono, Z Y Illahi and R Dewi

[Open abstract](#), Investigation of total organic matter [TOM] content during high and low water in inter-tidal zone sediment at Teluk Penyu Coast, Cilacap, Indonesia [View article](#), Investigation of total organic matter [TOM] content during high and low water in inter-tidal zone sediment at Teluk Penyu Coast, Cilacap, Indonesia [PDF](#), Investigation of total organic matter [TOM] content during high and low water in inter-tidal zone sediment at Teluk Penyu Coast, Cilacap, Indonesia

Engineering & Applied Sciences for Rural Development

012031

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Degradation of phenol in batik industry wastewater using thin layer TiO₂ photocatalyst](#)

K Riyani, T Setyaningtyas and A Riapanitra

[Open abstract](#), Degradation of phenol in batik industry wastewater using thin layer TiO₂ photocatalyst [View article](#), Degradation of phenol in batik industry wastewater using thin layer TiO₂ photocatalyst [PDF](#), Degradation of phenol in batik industry wastewater using thin layer TiO₂ photocatalyst

012032

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Energy Efficiency Calculation and Air Handling Unit Design Based on Cooling Load Capacity at MASTEK Mosque](#)

Catur Harsito, Ariyo Nurachman Satiya Permana and Finda Sihta

[Open abstract](#), Energy Efficiency Calculation and Air Handling Unit Design Based on Cooling Load Capacity at MASTEK Mosque [View article](#), Energy Efficiency Calculation and Air Handling Unit Design Based on Cooling Load Capacity at MASTEK Mosque [PDF](#), Energy Efficiency Calculation and Air Handling Unit Design Based on Cooling Load Capacity at MASTEK Mosque

012033

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Factors affecting electricity demand in Cambodia](#)

Virak Dy and Naraphorn Paoprasert

[Open abstract](#), Factors affecting electricity demand in Cambodia [View article](#), Factors affecting electricity demand in Cambodia [PDF](#), Factors affecting electricity demand in Cambodia

012034

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Factors affecting electricity consumption of residential consumers in Cambodia](#)

V Noeurn

[Open abstract](#), Factors affecting electricity consumption of residential consumers in Cambodia [View article](#), Factors affecting electricity consumption of residential consumers in Cambodia [PDF](#), Factors affecting electricity consumption of residential consumers in Cambodia

012035

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Forecasting the amount of rainfall in West Kalimantan using Generalized Space-time Autoregressive model](#)

R Utami, N Nurhayati and S Maryani

[Open abstract](#), Forecasting the amount of rainfall in West Kalimantan using Generalized Space-time Autoregressive model [View article](#), Forecasting the amount of rainfall in West Kalimantan using Generalized Space-time Autoregressive model [PDF](#), Forecasting the amount of rainfall in West Kalimantan using Generalized Space-time Autoregressive model

012036

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Implementation of autoregressive integrated moving average model to predict total electron content from GPS satellite receiver in Bandung](#)

Marifatul Nur Yuniati and Agus Sugandha

[Open abstract](#), Implementation of autoregressive integrated moving average model to predict total electron content from GPS satellite receiver in Bandung [View article](#), Implementation of autoregressive integrated moving average model to predict total electron content from GPS satellite receiver in Bandung [PDF](#), Implementation of autoregressive integrated moving average model to predict total electron content from GPS satellite receiver in Bandung

012037

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Hardware-based microgrid testbed to facilitate development of Distributed Energy Resource \(DER\) systems for sustainable growth](#)

Peter B. Idowu and Raja Suryadevara

[Open abstract](#), Hardware-based microgrid testbed to facilitate development of Distributed Energy Resource (DER) systems for sustainable growth [View article](#), Hardware-based microgrid testbed to facilitate development of Distributed Energy Resource (DER) systems for sustainable growth [PDF](#), Hardware-based microgrid testbed to facilitate development of Distributed Energy Resource (DER) systems for sustainable growth

012038

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Physical and mechanical properties of coffee waste composites and viselin fabrics as alternative base materials for manufacturing products in the interior field](#)

Purwanto

[Open abstract](#), Physical and mechanical properties of coffee waste composites and viselin fabrics as alternative base materials for manufacturing products in the interior field [View article](#), Physical and mechanical properties of coffee waste composites and viselin fabrics as alternative base materials for manufacturing products in the interior field [PDF](#), Physical and mechanical properties of coffee waste composites and viselin fabrics as alternative base materials for manufacturing products in the interior field

012039

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Portable wastewater treatment plant using banana stem filter media in small scale motor vehicle washing services](#)

Y Kusumawardani, S Subekti, W Astuti and S Soehartono

[Open abstract](#), Portable wastewater treatment plant using banana stem filter media in small scale motor vehicle washing services [View article](#), Portable wastewater treatment plant using banana stem filter media in small scale motor vehicle washing services [PDF](#), Portable wastewater treatment plant using banana stem filter media in small scale motor vehicle washing services

012040

THE FOLLOWING ARTICLE IS OPEN ACCESS

[The synthesis of \$\text{Ag}_3\text{PO}_4\$ under graphene oxide and hydroxyapatite aqueous dispersion for enhanced photocatalytic activity](#)

U Sulaeman, R D Permadi and H Diastuti

[Open abstract](#), The synthesis of Ag₃PO₄ under graphene oxide and hydroxyapatite aqueous dispersion for enhanced photocatalytic activity [View article](#), The synthesis of Ag₃PO₄ under graphene oxide and hydroxyapatite aqueous dispersion for enhanced photocatalytic activity [PDF](#), The synthesis of Ag₃PO₄ under graphene oxide and hydroxyapatite aqueous dispersion for enhanced photocatalytic activity

012041

THE FOLLOWING ARTICLE IS OPEN ACCESS

[An assessment Indonesia's Ocean Thermal Energy Conversion \(OTEC\) as an electrical energy resource](#)

M Trenggono, R R Hidayat, T N Cahyo, M D Mahardiono and A D Destrianty

[Open abstract](#), An assessment Indonesia's Ocean Thermal Energy Conversion (OTEC) as an electrical energy resource [View article](#), An assessment Indonesia's Ocean Thermal Energy Conversion (OTEC) as an electrical energy resource [PDF](#), An assessment Indonesia's Ocean Thermal Energy Conversion (OTEC) as an electrical energy resource

012042

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Design improvement for safety risks using hazard and operability method](#)

Retna Kristiana and Kushardiono

[Open abstract](#), Design improvement for safety risks using hazard and operability method [View article](#), Design improvement for safety risks using hazard and operability method [PDF](#), Design improvement for safety risks using hazard and operability method

012043

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Increased reliability over current relay \(ocr\) as a transformer protection with non-cascade coordination patterns](#)

Hari Prasetyo, Ari Fadli, Prisantono and Widhiatmoko Herry Purnomo

[Open abstract](#), Increased reliability over current relay (ocr) as a transformer protection with non-cascade coordination patterns [View article](#), Increased reliability over current relay (ocr) as a transformer protection with non-cascade coordination patterns [PDF](#), Increased reliability over current relay (ocr) as a transformer protection with non-cascade coordination patterns

012044

THE FOLLOWING ARTICLE IS OPEN ACCESS

[IOT Based Climate Monitoring System](#)

Muhammad Aziz Muslim, Raden Arief Setyawan, Achmad Basuki, Angger Abdul Razak, Fakhriy P Hario and Edward Fernando

[Open abstract](#), IOT Based Climate Monitoring System [View article](#), IOT Based Climate Monitoring System [PDF](#), IOT Based Climate Monitoring System

012045

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Geochemical of Volcanic Rock in Southern Part of Slamet Volcano, Indonesia](#)

Adi Candra, Januar Aziz Zaenurrohman, Siswandi and Aprian Wahyu Nugroho

[Open abstract](#), Geochemical of Volcanic Rock in Southern Part of Slamet Volcano, Indonesia [View article](#), Geochemical of Volcanic Rock in Southern Part of Slamet Volcano, Indonesia [PDF](#), Geochemical of Volcanic Rock in Southern Part of Slamet Volcano, Indonesia

JOURNAL LINKS

[IOP Science home](#)

- [Journals](#)
- [Books](#)
- [About IOPscience](#)
- [Contact us](#)
- [Developing countries access](#)
- [IOP Publishing open access policy](#)

IOP Publishing home

PAPER • OPEN ACCESS

Antioxidant activity of kecombrang preserving powder using *foam mat drying* method

To cite this article: R Naufalin *et al* 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **746** 012017

View the [article online](#) for updates and enhancements.

You may also like

- [Development of modified starch technology \(*maltodextrin*\) from commercial tapioca on semi production scale using oil heater dextrinator](#)
Agus Triyono, Raden Cecep Erwan Andriansyah, Rohmah Luthfiyanti et al.
- [Utilization of -amylase enzyme from *Bacillus stearothermophilus* RSAll₁₈ for maltodextrin production from sago starch](#)
R A Arfah, A Ahmad, S Dali et al.
- [Antibacterial And Antioxidant Activities Of Ethanol Extracts Of Cocoa Husk \(*Theobroma cacao* L.\) With Maltodextrine In Various Concentration](#)
A Hasanuddin, K Anwar, M Mappatoba et al.



The Electrochemical Society
Advancing solid state & electrochemical science & technology

241st ECS Meeting

May 29 – June 2, 2022 Vancouver • BC • Canada

Abstract submission deadline: Dec 3, 2021

Connect. Engage. Champion. Empower. Accelerate.
We move science forward



Submit your abstract



Antioxidant activity of kecombrang preserving powder using foam mat drying method

R Naufalin*, Erminawati, R Wicaksono, A T Febryani, and N Latifasari

Food Technology Study Program, Faculty of Agriculture, Jenderal Soedirman University, Purwokerto 53123

*Email: rifda.naufalin@unsoed.ac.id

Abstract. Food production is one of the easily damaged products, both in processing and during product storage. Therefore, to prevent damage made preservative powders from natural ingredients. One of them is obtained from the flowers of the kecombrang (*Etlingera elatior*). Kecombrang plant flowers contain bioactive components, namely, alkaloids, polyphenols, flavonoids, and essential oils. This study used a reasonably practical drying, namely foam-mat drying technique. This study aims to determine the best formulation in making kecombrang flower preservative powder using the foam mat drying method. This study used a randomized block design (RBD). The factors studied included maltodextrin concentration and drying temperature. The concentration of maltodextrin consists of three levels, namely 5%, 10%, and 15%, and the temperature consists of three levels, namely 50°C, 60°C, and 70°C. The variables observed were qualitative and quantitative analyses. The best treatment results qualitatively and quantitatively, namely at the drying temperature treatment of 50°C and 5% maltodextrin concentration with a water content value of 5.92%, antioxidant activity value 61.77%, total phenol levels 2.12 mg TAE / g, total flavonoid levels 0.13 mg TAE/g. The qualitative test shows that kecombrang flower preservative powder contains phenolic compounds, flavonoids, steroids, tannins, saponins, and alkaloids.

1. Introduction

The food product is one of the easily damaged products, both in processing and during product storage. Food that is damaged will experience changes caused by physical damage, chemical reactions, or the activity of organisms such as rats, parasites, insects, microbes, and others. Therefore, additional materials such as preservatives are needed to maintain food products' quality, so they are not easily damaged [1].

One of the natural ingredients used as a preservative, antioxidant, and antimicrobial is the kecombrang (*Etlingera elatior*). Kecombrang has been widely used in medicine and food additives, especially flowers, that effectively reduce microbiological damage. Kecombrang flower is an alternative natural preservative because it contains bioactive components: alkaloids, phenols, flavonoids, and essential oils [2]. The manufacture of natural preservatives from the kecombrang plant's flowers is quite useful because it contains suitable antimicrobials [3] and antioxidants [4] in preventing the damage of a food product.

The preservatives are made into powder form because the powder form has the advantages of being more durable, lighter, and has a smaller volume to make packaging more comfortable. Powdered preservatives are quite useful and can be stored for a long time compared to preservatives in



extract or liquid form [5]. The manufacture of preservative powder uses the foam-mat drying method (foam drying technique). In this preservative powder formulation, a filler is needed, namely, maltodextrin. The use of this maltodextrin concentration is around 5-15%. This is consistent with the statement regarding the addition of maltodextrin in the range of 6-20% in the manufacture of apple powder products using the foam mat drying method, which is proven to reduce hygroscopic properties, stabilize empty cavities in the foam and increase granulation in powders [6]. The use of concentrations above 15% can reduce the quality of preservative powders. The addition of more than 20% maltodextrin can reduce polyphenol levels in kombucha tea instant drinks, where polyphenols act as antioxidants [7].

The foam drying technique's effectivity is determined by the drying speed, which can be done with the correct temperature setting and filler concentration. Too high a temperature will cause the loss of volatile compounds such as aroma and accelerate the material's browning reaction. Simultaneously, a too low temperature will cause the drying process to be less efficient and encourage damage during the process.

According to other studies, foam's presence will accelerate the evaporation process of water even without the temperature being too high; products that are dried using foam at a temperature of 50°C - 80°C can produce 2-3% water content. The powder from the foam mat drying method has a low density (light) and is crumbly [8]. Therefore, it is necessary to study the heating temperature and maltodextrin concentration as an effective filler in making preservative powder formulations in this study. Based on these descriptions, this study's objectives are to determine the best treatment on the kecombrang flower preservative powder formulation using the foam mat drying method.

2. Research Methodology

2.1. Experimental design

This research is an experimental study with the experimental design used was a Randomized Block Design. The factors observed were maltodextrin concentration, which consists of three levels, namely 5%, 10%, and 15%, and the temperature, which consists of three levels, namely 50, 60, and 70°C. From these treatments obtained nine treatment combinations and carried out three times repetition to get 27 experimental units.

2.2. Preparation and Manufacture of Flower Simplicia Powder

The red kecombrang flowers are sorted into good ones, then washed first, then chopped into small pieces and evenly distributed with a knife. Then spread it out on a baking sheet and then dry it with a *cabinet dryer* at 50°C for 4 hours. The dried flowers are then ground with a disk mill grinder until a homogeneous powder is obtained.

2.3. Kecombrang Flower Extract

Kecombrang flower powder and water are put into the extractor with a ratio of 1:14. Kecombrang flower extraction process was carried out for 3 hours 42 minutes at a temperature of 60°C using an extractor.

2.4. Drying Preservative Powder with Foam-mat drying

The liquid extract of kecombrang flower was added with maltodextrin with each different concentration (5%, 10%, 15%) and 1% tween 80, then stirred using a mixer for 15 minutes foam was formed. Kecombrang flower liquid extract foam is poured into a glass plate and flattened; after that, it is put into a *cabinet dryer* for the drying process with a temperature of 50°C, 60°C, 70°C, then the dry product is mashed with mortar to obtain a homogeneous preservative powder.

2.5. Qualitative Analyses

After obtaining a homogeneous preservative powder, a qualitative analysis was carried out, namely the phytochemical test including the alkaloid, phenol hydroquinone, flavonoid, tannin, saponin, and steroid/triterpenoid test. Furthermore, the preservative powder will be analyzed

quantitatively, including moisture content, total phenol content, determination of flavonoid levels, antioxidant activity, and antibacterial activity.

2.6. Quantitative Analyses

2.6.1. Grading Total Phenol. A total of 0.4 ml of sample was added to 1.5 ml of Folin-Cicalteu 10% and stood for 5 minutes at 25°C. Then Sodium bicarbonate (NaHCO₃) 1.5 ml was shaken and left in a dark room for 90 minutes. Next step, the absorbance was measured using a spectrophotometer at a wavelength of 725 nm.

2.6.2. Grading Total Flavonoid. A total of 0.1 ml of sample supernatant was added to 1 ml AlCl₃ 2% (2 g AlCl₃ in 100 ml acetate acid glacial solid 5%) and 1 ml calcium acetate solid 120 mm (1.176 g calcium acetate in 100 ml distilled water). After that, it was incubated for 1 hour at 25°C and measured the absorbance using a spectrophotometer UV Vis of 435 nm.

2.6.3. Antioxidant Activity. 0.2 ml of sample solution for each dilution was taken and then mixed with 2.8 ml of 2,2-diphenil-1- pikrilhidrazil (DPPH) solution in a dark test tube. The solution was then vortexed and then incubated for 30 minutes; then, measured the absorbance using a spectrophotometer UV Vis of 517 nm.

Furthermore, the IC₅₀ price is determined, namely the concentration of the sample, which has a DPPH absorbance inhibition of 50%, using regression soft wear with a value of $y = 50\%$ so that the IC₅₀ inhibitory concentration value is obtained. The lower the IC₅₀ value, the higher the free anti-radical activity.

2.7. Analysis Data.

The variable data obtained were analysed using ANOVA with a level of 95%, if the treatment had a significant effect, then continued with the Duncan Multiple Range Test with a level of 5%. The best formulation is done by testing the effectiveness index.

3. Result and Discussions

3.1 Qualitative Analyses

The bioactive content found in kecombrang is saponins, flavonoid, phenol, triterpenoid, tannin, and alkaloids which activities as antimicrobials and antioxidants for natural preservatives [9].

Table 1. Phytochemical analysis of kecombrang flower preservative powder

Treatment	Saponin	Flavonoid	Phenol	Triterpenoid	Tannin	Alkaloid
50°C, 5%	++	+++	++++	+++	++++	++++
50°C, 10%	++	+++	+++	++	+++	+++
50°C, 15%	++	++	+++	++	++	++
60°C, 5%	++	+++	+++	+++	+++	++++
60°C, 10%	++	++	++	++	++	+++
60°C, 15%	+++	+	++	++	-	+++
70°C, 5%	++	+++	+++	++++	++	+++
70°C, 10%	+++	++	+++	++	++	+++
70°C, 15%	+++	+	++	+	-	+++

Note: - (negative), + (weak), ++ (positive), +++ (strong), ++++ (very strong)

3.1.1. Alkaloid. In alkaloid testing, sample was added to HCl and then reagent [10]. A positive result in the Meyer test is indicated by the formation of a white precipitate. In contrast, a positive result for

alkaloids in the Wagner test is shown by forming a light brown to red precipitate. Positive alkaloid formation.

In Table 1, it can be seen that the formation of the most potent alkaloid compounds at treatment temperatures of 50°C and 60°C with a concentration of 5% maltodextrin. The appearance of an orange precipitate indicates a positive result for alkaloids in the Dragendorff analyses. These residues are metal complexes with alkaloids. In the Dragendorff reagent reaction, bismuth nitrate reacts with potassium iodide to form bismuth (III) iodide deposits, which dissolves in excess iodide potassium tetra iodo bismutate. In the alkaloid test with the Dragendorff reaction, the lone pair on nitrogen is used to form a covalent bond with bismuth to give an orange to red precipitate [11].

3.1.2. Phenol. Table 1 shows that the phenolic compounds produced from kecombrang flower preservative powder are the strongest at the lowest temperature (50°C) and a low concentration of 5% maltodextrin. The presence of phenolics is indicated by a blue to green color formed when the sample is exposed to FeCl₃. The more Phenol is contained, the darker the color will be. This is consistent with research that explains that phenol testing is carried out by reacting a sample of 1 mL with 5% FeCl₃. The presence of phenolic compounds is characterized by forming a bluish-green to blackish color [12].

3.1.3. Flavonoid. The table shows that the formed flavonoid compounds are more potent with temperature treatment and low maltodextrin concentration at 50°C. In contrast, the formed flavonoid compounds are weak if the temperature and the maltodextrin concentration are higher, 70°C. Flavonoids in kecombrang flowers contain phenolic compounds with carbonyl groups, flavones with 3-OH groups, and flavones with free ortho-dihydroxy and ortho-hydroxy carbonyls [13]. The flavonoid test with H₂SO₄ reagent shows a positive sample when the solution has a striking color change to yellow, red, or brown [10].

3.1.4. Saponin. In the table, it is known that the higher the temperature and the concentration of maltodextrin, the stronger the saponin content formed. Saponin testing is characterized by the formation of foam when the sample is shaken. Saponin test results yield a solution with the appearance of stable foam as high as ± 15 mm, which indicates a positive effect [14]. In general, if the result is positive, the addition of HCl 2 N is intended to increase polarity so that the hydrophilic groups will bind more stably and the formed foam becomes stable. When shaken, saponins form because of the hydrophilic groups that bind to water, while hydrophobes will attach to air. In the micellar structure, the polar groups face outwards while the non-polar groups face inwards; this forms the foam [15].

3.1.5. Steroid/Triterpenoid. In table 1, it is known that the formation of the most potent steroid content at 70°C treatment and 5% maltodextrin concentration. Test for terpenoids and steroids using Lieberman Burchard's reagent [16]. Steroid and triterpenoid tests using the Liebermann-Bouchard method, the extract was dissolved in chloroform and then added with Liebermann-Bouchard reagent (acetic acid anhydrous-H₂SO₄) showed positive results in the presence of a brownish-red color change for steroids and brown-purple for triterpenoids. The reaction of triterpenoids with Liebermann's reagent produces a red-purple color while steroids give a green-blue color. This is based on triterpenoid and steroid compounds' ability to form color by H₂SO₄ in an anhydrous acetic acid solvent. The difference in color produced by triterpenoids and steroids is due to differences in groups on the C-4 atom [17].

3.1.6. Tanin. In table 1, it can be seen that the tannin content is the strongest at a treatment temperature of 50°C and a concentration of 5% maltodextrin. The higher temperature and the maltodextrin concentration used, the lower the color density. In the treatment of maltodextrin concentration of 15%, temperature 60 and 70°C showed no tannin content in the preservative powder.

While kecombrang do not contain tannins, in Table 1, the tannin test results green to blackish blue colors were formed in several treatment samples. In testing, the FeCl₃ used can react with other compounds with the same color results. The main properties of tannins depend on the phenolic group contained in the tannins. Color reactions occur when combined with FeCl₃. FeCl₃ will give a green or

blue-black color when reacted with tannins. But this test is not good because, in addition to tannins which can give color reactions, other substances can also provide the same color reaction [18].

3.2. Analysis Result of Quantitative Data

3.2.1. Total Phenol. The results of the analysis of variance showed that the drying temperature had a very significant effect on the total phenol content and the maltodextrin concentration also had a very substantial impact on the total phenol content of kecombrang flower preservative powder. Meanwhile, the interaction of drying temperature and maltodextrin concentration did not significantly affect the total phenol content of kecombrang flower preservative powder. The principle of measuring total Phenol is by measuring the Folin-Ciocalteu reagent's ability to reduce phenol compounds that form blue complexes. Analysis of total phenol levels is necessary because there is a link between antioxidant and antimicrobial activity, which is influenced by phenolic compounds' composition [19].




Figure 1. Effect of drying temperature




Figure 2. Effect of maltodextrin concentration

Total phenol levels against total phenol levels From Figure 1, it can be seen that the degree of drying temperature greatly affects the total phenol content, and the results are significantly different. The highest average total phenol value was obtained from the treatment at 50°C, namely 1.8248 mg TAE / g, at 60°C at 1.4948 mg TAE / g, and the lowest at 70°C at 1.1998 mg TAE / g. The decrease in the total amount of Phenol is due to the oxidation process due to heat treatment so that the drying process can reduce the content of phenol compounds [20]. The decrease may be due to chemical changes, decomposition of phenol compounds, or phenol-protein complexes' formation due to temperature and pressure [21]. The literature states that there is a relationship between temperature and phenolic content. Increasing the temperature causes an increase in phenolic content to a specific temperature, decreasing with higher temperature [22].

The phenol component in the extraction of kecombrang plant parts is thought to have a polarity close to ethanol's polarity. The use of ethanol solvents is more effective in dissolving phenol compounds [9][23]. The difference in total phenol content can be caused due to environmental factors for different kecombrang plants, such as differences in ultraviolet radiation, plant growth, soil composition, temperature, and rainfall [24].

Figure 2 also shows that maltodextrin concentration has a very significant effect on the total phenol value. The highest average total phenol value was obtained from the treatment of 5% maltodextrin concentration, namely 1.7544 mg TAE / g, at 10% maltodextrin concentration valued at 1.5358 mg TAE / g, and the lowest at 15% maltodextrin concentration worth 1.2291 mg. TAE / g. The higher the addition of maltodextrin causes a decrease in the total phenol content. This is due to the increasing number of total solids in the material, namely maltodextrin as a filler. The total Phenol measured is less, where the maltodextrin is white. Simultaneously, the complex color is blue, so when measured with a spectrophotometer, the blue color intensity decreases so that the total content phenol tends to decrease [25].

3.2.2. Total Flavonoid. The total flavonoid levels produced ranged from 0.06 to 0.13 mg TAE / g. From Figure 3, it can be seen that the degree of drying temperature has an effect on the total flavonoid levels with significantly different results. The highest mean value of total flavonoids was obtained from treatment at 50°C, namely 0.1254 mg TAE / g, at 60°C at 0.0978 mg TAE / g, and the lowest at 70°C at 0.0697 mg TAE / g. According to the literature, a temperature of 50°C is relatively safe and prevents damage to specific secondary metabolites, especially flavonoids. Flavonoids are phenolic compounds that have a conjugated aromatic system that is easily damaged on relatively long heating with high temperatures. Several groups of flavonoids have glycoside bonds with sugar molecules. Glycoside bonds will easily break or break on high-temperature heating [26].




Figure 3. Effect of drying temperature on concentration total flavonoid levels




Figure 4. Effect of maltodextrin on total flavonoid levels

High temperature affects the increase in total flavonoids. This is because, at higher temperatures, contact between the materials and the solvent works well. As a result, the solvent's dissolve ability is higher when the solvent is hot and lower when it is cold. According to the literature, optimal resistance to flavonoid compounds has a temperature range of 0°C – 100°C [27].

Figure 4 also shows that the concentration of maltodextrin has a significant effect on the total flavonoid value. The highest mean value of total flavonoids was obtained from the treatment of 5% maltodextrin concentration, namely 0.1043 mg TAE / g, at 10% maltodextrin concentration worth 0.0980 mg TAE / g, and the lowest at 15% maltodextrin concentration worth 0.0906 mg TAE / g. This is directly proportional to the total phenol content and antioxidant activity, where flavonoids are phenolic compounds that act as anti-free radicals. It is related to the value of the sample's antioxidant activity. From the resulting data, it can be seen that the total flavonoid value is lower than the total phenol value. Increasing temperature and extraction time need to be considered; too high extraction temperature and long extraction time and exceeding the optimum time limit can cause the loss of compounds in the solution due to evaporation. Vice versa, if the extraction temperature is too low, it will cause not all active compounds to be extracted from ingredients, resulting in a low percentage of the active compounds obtained [28].

3.2.3. Antioxidant Activity. The analysis results showed that the treatment of drying temperature and maltodextrin concentration significantly affected kecombrang flower preservative powder's antioxidant activity. Simultaneously, the interaction of drying temperature and maltodextrin concentration has no significant effect on kecombrang flower preservative powder's antioxidant activity. The IC₅₀ values obtained ranged from 49.49 - 204.44 ppm.




Figure 5. Effect of drying temperature on Antioxidant Activity




Figure 6. Effect of maltodextrin concentration on Antioxidant Activity

From Figure 5, it can be seen that the higher the temperature can cause the antioxidant activity to decrease. The most increased antioxidant activity was at 50°C with an IC_{50} value of 73.204%; at 60°C, it became 98.954%, and the lowest at 70°C was around 151.974%. The antioxidant activity obtained was calculated the IC_{50} value with a linear regression equation. The IC_{50} value is inversely proportional to the antioxidant ability of a compound contained in the test material. The smaller the IC_{50} value, the greater the antioxidant capacity. When the electrons become paired by a free radical scavenger's presence, the absorbance decreases stoichiometry according to the number of electrons taken. The presence of antioxidant compounds can change the color of the DPPH solution from purple to yellow. The absorbance changes resulting from this reaction have been used extensively to test the ability of some molecules to scavenge free radicals [29].

Other studies suggest that the antioxidant activity decreases when the drying temperature is high [30]. This is because the higher heating temperature causes the secondary metabolites that act as antioxidants (flavonoid compounds) to become damaged. The literature states that the higher the drying temperature, the lower the antioxidant activity and impair the sample's antioxidant activity [31].

Based on Figure 6, it can be seen that as well as temperature, the effect of maltodextrin concentration also has a very significant impact on antioxidant activity, so that the higher the maltodextrin concentration causes the IC_{50} value to increase, causing the ability to capture free radicals from bioactive compounds of kecombrang flowers to decrease. The moderate antioxidant activity against the highest concentration of maltodextrin was at a concentration of 5% worth 93.658%, while the lowest was at a concentration of 15%, namely 122.499%. A substance has antioxidant properties when the IC_{50} value is less than 200 ppm. If the IC_{50} value obtained is between 200-1000 ppm, the meaning is less active but still has potential as an antioxidant [32].

The addition of higher maltodextrin concentration causes a decrease in antioxidant activity levels. This is thought to be due to the increasing number of total solids contained in the material, namely maltodextrin as a filler so that the measured antioxidant activity is less so that with the increase in the total solids in a material, the measured levels of antioxidant activity will be smaller [33]. Also, it is thought that it is also caused by changes in antioxidant compounds due to the heating process, namely vitamin C and other phenol compounds that are oxidized. It is possible that heating causes phenol compounds to decompose so that their ability as an antioxidant decreases. The antioxidant activity in noni leaf instant drinks is closely related to total Phenol and vitamin C. So that by decreasing the concentration of phenol and vitamin C, the flavonoid content will decrease and cause the antioxidant activity to decrease [25].

4. Conclusion

Kecombrang flower preservative powder formulations positively contain bioactive compounds, including phenolic compounds, flavonoids, saponins, steroids, and alkaloids. Except for negative tannins at treatment temperatures of 60 and 70°C with a concentration of 15% maltodextrin. The best formulation was obtained at a temperature treatment of 50°C with a concentration of 5% maltodextrin. It produces the highest results at an antioxidant activity value of 61.77%, total phenol content of 2.12 mg TAE / g, and total levels of flavonoids 0.13 mg TAE / g.

Acknowledgments

The authors wish to thanks to Institute for Research and Community Service of Jenderal Soedirman University, which has provided research funding through Institution Research Scheme 2020.

References

- [1] Naufalin R and Rukmini H S 2018 Antibacterial activity of kecombrang flower extract (*Nicolaia speciosa*) microencapsulation with food additive materials formulation *IOP Conf. Ser. Earth Environ. Sci.* **102**
- [2] Naufalin R 2019 Natural preservation opportunities and challenges in improving food safety *AIP Conf. Proc.* **2094**
- [3] Putri F A, Naufalin R and Wicaksono R 2019 Antimicrobial edible coating application of Kecombrang flower concentrate to reduce microbial growth on gourami fish sausage *IOP Conf. Ser. Earth Environ. Sci.* **250**
- [4] Naufalin R, Wicaksono R, Erminawati, Arsil P and Gulo K I K 2019 Application of Concentrates Flower Kecombrang on Edible Coating as Antioxidant to Suppress Damage on Gourami Sausage *IOP Conf. Ser. Earth Environ. Sci.* **255**
- [5] Kamsiati E 2006 Pembuatan bubuk sari buah tomat (*Licopersicon esculentum* Mill.) dengan metode "foammat drying" *J. Teknol. Pertan.* **7** 113–9
- [6] Jakubczyka E, Gondeka E, Tamborb K, Jakubczyk E, Gondek E and Tambor K 2011 Characteristics of selected functional properties of apple powders obtained by the foam-mat drying method *ICEF 11 International Congress on Engineering and Food* (International Association of Engineering and Food Athens, Greece)
- [7] Fiana R M, Murtius W S and ASben A 2016 Pengaruh konsentrasi maltodekstrin terhadap mutu minuman instan dari teh kombucha *J. Teknol. Pertan. Andalas* **20** 1–8
- [8] Ramadhia M, Kumalaningsih S and Santoso I 2012 Pembuatan tepung lidah buaya (aloe vera l.) dengan metode foam-mat drying *J. Teknol. Pertan.* **13** 125–37
- [9] Naufalin R and Herastuti S R 2017 Antibacterial activity of *Nicolaia speciosa* fruit extract *Int. Food Res. J.* **24** 379–85
- [10] Harborne J B 1984 *Metode Fitokimia: Penuntun Cara Modern Menganalisis Tumbuhan* (Bandung: ITB Bandung)
- [11] Marliana S D and Suryanti V 2005 Suyono. 2005. Skrining fitokimia dan analisis kromatografi lapis tipis komponen kimia buah labu siam (*Sechium edule* Jacq. Swartz.) dalam ekstrak etanol *Biofarmasi* **3** 26–31
- [12] Syafitri N E, Bintang M and Falah S 2014 Kandungan fitokimia, total fenol, dan total flavonoid ekstrak buah harendong (*Melastoma affine* D. Don) *Curr. Biochem.* **1**
- [13] Farida S and Maruzy A 2016 Kecombrang (*Etlingera elatior*): Sebuah Tinjauan Penggunaan Secara Tradisional, Fitokimia Dan Aktivitas Farmakologinya *Indones. J. Plant Med.* **9** 19–28
- [14] Madduluri S, Rao K B and Sitaram B 2013 In vitro evaluation of antibacterial activity of five indigenous plants extract against five bacterial pathogens of human *Int. J. Pharm. Pharm. Sci.* **5** 679–84
- [15] Simaremare E S 2014 Skrining fitokimia ekstrak etanol daun gatal (*Laportea decumana* (Roxb.) Wedd) *Pharm. J. Farm. Indones. (Pharmaceutical J. Indones.)* **11**

- [16] Ningsih D R, Zufahair Z and Kartika D 2016 Identification of secondary metabolites compounds and antibacterial activities on the extract of soursop leaf *Molekul* **11** 101–11
- [17] Marliana E and Saleh C 2011 Uji Fitokimia dan Aktivitas Antibakteri Ekstrak Kasar Etanol Fraksi n-Heksana, Etil Asetat dan Metanol dari Buah Labu Air (*Lagenaria siceraria* (Molina) Standl) *J. Kim. Mulawarman* **8** 63–9
- [18] Susanti M C E 2000 Autokondensat tanin dan penggunaan sebagai perekat kayu lamina
- [19] Bahri S R, Fredj R Ben, Boughalleb N, Shriaa J, Saguem S, Hilbert J-L, Troitin F, Ammar S, Bouzid S and Harzallah-Skhiri F 2014 Phenolic Composition and Antioxidant and Antimicrobial Activities of Extracts Obtained from *Crataegus azarolus* L. var. *aronia* (Willd.) Batt. Ovaries Calli. *J. Bot.*
- [20] Naufalin R, Jenie B S L, Kusnandar F, Sudarwanto M and Rukmini H 2005 Antibacterial Activity of Kecombrang Flower Extract Toward Pathogenic and Food Spoilage Bacteria *J. Teknol. dan Ind. Pangan* **16** 119–25
- [21] Estiasih T and Sofia E 2009 Stabilitas antioksidan bubuk keluwak (*Pangium edule* Reinw) selama pengeringan dan pemasakan *J. Teknol. Pertan.* **10** 115–22
- [22] Liyana P C and Shahidi F 2005 Optimization of extraction of phenolic compounds from wheat using response surface methodology *Food Chem.* **93** 47–56
- [23] Pertiwi D I, Naufalin R, Arsil P, Erminawati, Wicaksono R and Auliya T 2019 Quality of simplician bioactive components and liquid extract of kecombrang flower powder from temperature and time optimization results *IOP Conf. Ser. Earth Environ. Sci.* **406**
- [24] Borges L L, Alves S F, Sampaio B L, Conceição E C, Bara M T F and Paula J R 2013 Environmental factors affecting the concentration of phenolic compounds in *Myrcia tomentosa* leaves *Brazilian J. Pharmacogn.* **23** 230–8
- [25] Yuliwaty S T and Susanto W H 2014 Pengaruh Lama Pengeringan dan Konsentrasi Maltodekstrin terhadap Karakteristik Fisik Kimia dan Organoleptik Minuman Instan Daun Mengkudu (*Morinda citrifolia* L)[In Press Januari 2015] *J. Pangan dan Agroindustri* **3** 41–52
- [26] Oktavia J D 2011 Pengoptimuman Ekstraksi Flavonoid Daun Salam (*Syzygium polyanthum*) dan Analisis Sidik Jari Dengan Kromatografi Lapis Tipis
- [27] Putri D D and Nurmagustina D E 2014 Kandungan Total Fenol dan Aktivitas Antibakteri Kelopak Buah Rosela Merah dan Ungu Sebagai Kandidat Feed Additive Alami Pada Broiler *J. Penelit. Pertan. Terap.* **14** 174–80
- [28] Ibrahim A M, Yunianta Y and Sriherfyna F H 2015 Pengaruh Suhu dan Lama Waktu Ekstraksi terhadap Sifat Kimia dan Fisik pada Pembuatan Minuman Sari Jahe Merah (*Zingiber officinale* var. *Rubrum*) dengan Kombinasi Penambahan Madu sebagai Pemanis [in press April 2015] *J. Pangan dan Agroindustri* **3** 530–41
- [29] Tiaraswara R A and Leni Herliani Afrianti L H A 2016 Optimalisasi Formulasi Hard Candy Ekstrak Daun Mulberry (*Morus* sp.) dengan Menggunakan Design Expert Metode D-Optimal
- [30] Dewi F K 2010 Aktivitas antibakteri ekstrak etanol buah mengkudu (*Morinda citrifolia*, Linnaeus) terhadap bakteri pembusuk daging segar
- [31] Sayekti E D, Asngad A and Chalimah S 2016 Aktivitas antioksidan teh kombinasi daun katuk dan daun kelor dengan variasi suhu pengeringan
- [32] Molyneux P 2004 The Use of the Stable Free Radical Diphenylpicryl-hydrazyl (DPPH) for Estimating Antioxidant Activity *Songklanakarin J. Sci. Technol.* **26** 211–9
- [33] Naufalin R and Rukmini H S 2013 Microcapsule Application of Kecombrang Flower Extract : Effects of Concentration, Types of Fraction, pH of Medium, and NaCl on Microbiological Properties of Minced Beef *Anim. Prod.* **15** 8–14

CERTIFICATE OF PRESENTATION

This is to certify that

Rifda Naufalin*, Agung Triska Febryani, Erminawati Wuryatmo, Rumpoko Wicaksono

Has presented a paper entitled:

Antioxidant Activity Of Kecombrang Preserving Powder Using Foam Mat Drying Method

in the 3rd International Conference on Multidisciplinary Approaches
for Sustainable Rural Development 2020.

3rd ICMA-SURE 2020

“Interdisciplinary approaches and applied technologies for sustainable rural-environmental resources based on local
wisdoms before and during COVID-19 pandemic”



Prof. Dr. Rifda Naufalin S.P., M.Si.

Head of Institute of Research and Community Service
Jenderal Soedirman University



Assoc. Prof. Maria Dyah Nur Meinita

Conference Chair

Registered



3rd International Conference on Multidisciplinary Approaches for Sustainable Rural Development (ICMA-SURE)

organized by LPPM Universitas Jenderal Soedirman



Letter of Acceptance and Invitation

Date : October 12th 2020
No : 162/LOA/ICMA-2020

Dear Rifda Naufalin

Thank you for submitting your abstract for presentation at the 3rd International Conference on Multidisciplinary Approaches for Sustainable Rural Development (ICMA-SURE) 2020. After reviewing your abstract, we are pleased to inform you that your Abstract entitled:

ANTIOXIDANT ACTIVITY OF KECOMBRANG PRESERVING POWDER USING FOAM MAT DRYING METHOD

ID Paper: 1649

meets preliminary acceptance requirements set forth by our Scientific Committee to be presented as Oral Presentation at the conference. The conference will be held online on 18-19 November 2020 using the Zoom application. Please send your presentation material by 15 November 2020. The Oral/Poster Presentation Guidelines can be found at the following link:

<https://conference.unsoed.ac.id/index.php/icma/index/pages/view/presentationguidelines>

Regarding the payment, you also need to re-register to get a bill number. Please make bill payments before 20 October 2020. The payment and re-registration steps are explained in the ICMA SURE 2020 Payment Guideline.

If you require any further information, please do not hesitate to contact us or visit our website. We look forward to seeing you at the conference.

Yours sincerely,

Assoc. Prof. Maria Dyah Nur Meinita
Chairperson of ICMA SURE 2020

