

COVERING LETTER

Dear **Editor-in-Chief**,

I herewith enclosed a research article,

Title:

Isolation and Characterization of The Endophytic Bacteria, And Their Potential As Maize Diseases Control

Author(s) name:

Endang Mugiastuti, Suprayogi, Nur Prihatiningsih and Loekas Soesanto

Address

(Fill in your institution's name and address, your personal cellular phone and email)

Faculty of Agriculture, Jenderal Soedirman University, Purwokerto, HP. 08122826706,
email: endangmugiastuti@gmail.com

For possibility publication on the journal:

(fill in *Biodiversitas* or *Nusantara Bioscience* or *mention the others*)

Biodiversitas

Novelty:

(state your claimed novelty of the findings versus current knowledge)

Research to get biological control agents that can control 2 main diseases of maize. Biological agents are obtained from 2 altitudes, so it is expected that biological agents can be applied in various maize growing areas

Statements:

This manuscript has not been published and is not under consideration for publication to any other journal or any other type of publication (including web hosting) either by me or any of my co-authors.

Author(s) has been read and agree to the Ethical Guidelines.

List of five potential reviewers

(Fill in names of five potential reviewers **that agree to review your manuscript** and their **email** addresses. He/she should have Scopus ID and come from different institution with the authors; and from at least three different countries)

Place and date:

Purwokerto, august 2019

Sincerely yours,

(fill in your name, no need scanned autograph)

Endang Mugiastuti

ISOLATION AND CHARACTERIZATION OF THE ENDOPHYTIC BACTERIA, AND THEIR POTENTIAL AS MAIZE DISEASES CONTROL

ENDANG MUGIASTUTI^{1*}, SUPRAYOGI¹, NUR PRIHATININGSIH¹ AND LOEKAS SOESANTO¹

Faculty of Agriculture, Jenderal Soedirman University, Purwokerto. Jl. Dr. Suparno, Karangwangkal, Purwokerto 53123, Central Java, Indonesia.
Tel./Fax. +62-281-638791, *email: endangmugiastuti@gmail.com.

Manuscript received: 1411 2019 (Date of abstract/manuscript submission). Revision accepted: 20.

Abstract. The research aimed to isolate and characterize morphologically and biochemically the endophytic bacteria, and their potential to control maize diseases, especially sheath blight and bacterial wilt. The study was conducted at the Plant Protection Laboratory from April to August 2019. The study consisted of four stages: isolation and characterization of endophytic bacteria, the antagonism test of the endophytic bacterial to *R. solani*, the antagonism test of the endophytic bacteria to *Pantoea* sp., and the mechanism test of the endophytic bacteria as biological control agents and plant growth-promoting bacteria. Based on the research, it has been successfully isolated, and characterized morphologically and biochemically characterized four endophytic bacteria isolates that have the potential to be developed as biopesticides to control maize disease, especially *R. solani* and *Pantoea* sp. They can suppress the growth of *R. solani* by more than 50%, have a strong antagonistic index against *Pantoea* sp (> 4), and can produce protease and lipase enzyme, and phosphate solubilization.

Key words: *Bacillus* sp., Fluorescents *Pseudomonads*, *Pantoea* sp, *Rhizoctonia solani*

Running title: Isolation and characterization of the endophytic

INTRODUCTION

Maize is a strategic food commodity in the world. In Indonesia, the government seeks to achieve self-sufficiency in maize through increasing production of sustainable maize. However, these efforts have faced several obstacles, one of them is the presence of plant diseases such as sheath blight, caused by *Rhizoctonia solani* Kuhn, and bacterial wilt caused by (*Pantoea* sp.). *R. solani* can infect up to the midrib of the cob (Djaenuddin et al. 2017), resulting in a decrease in the yield of up to 100%. (Muis 2007). *Pantoea* sp. can attack all stages of the plant causing wilting and leaf blight, and is known as Stewart's wilt (Pataky 2004; Ammar et al. 2014). The pathogens can cause 40-100% yield loss.

Over the past 3 decades, the concept of sustainable and environmentally friendly agriculture have been carried out by minimizing the use of chemicals, both synthetic fertilizers and synthetic pesticides. In the management of pests and plant diseases, biological control is developed by applying biological control agents including the endophytic bacteria (Shanti and Vittal 2013). Many endophytic bacteria can pass the endodermic barrier across from the root cortex to the vascular system, and subsequently develop as endophytes in stems, leaves, tubers, and other organs (Compant et al. 2005). The use of endophytic bacteria as biological agents has an advantage compared to rhizosphere bacteria because endophytic bacteria live and survive in the plant tissue during plant development, thus protecting the plants.

Bacillus sp. and fluorescents *Pseudomonads* are reported to be able to live as endophytes and are widely used as biological control agents for soil-borne and air-borne diseases. The endophytic bacteria could control plant diseases through several mechanisms including competition, hyperparasite, producing microbial inhibiting compounds (antibiotics, lysis enzymes, other physical or chemical disorders), enhancing plant resistance, and promoting plant growth (Compant et al 2005, Pal and McSpadden 2006; Rosenblueth and Martinez Romero 2006;).

Based on the mechanisms, the use of endophytic bacteria isolated from maize, both upland and lowland, suggested potentially alternative control for sheath blight (*R. solani*) and bacterial wilt (*Pantoea* sp). The research aimed to isolate and characterize morphologically and biochemically the endophytic bacterial as well as their potential to control pathogens that cause disease in maize especially *R. solani* and *Pantoea* sp.

MATERIALS AND METHODS

This research was conducted at the Laboratory of Plant Protection, Faculty of Agriculture, Jenderal Soedirman University, Purwokerto, Central Java, Indonesia, from April to August 2019

Isolation *R. solani*

R. solani was isolated from maize with sheath blight symptoms and there was sclerotium as a resistant structure from the pathogenic fungi in Banyumas. Samples were isolated on PDA medium to obtain pure *R. solani* isolates.

Isolation *Pantoea* sp.

Pantoea sp. isolated from diseased maize, which was taken from the maize growing area in Banyumas Regency. *Pantoea* sp. was isolated according to Coplin et al. 2012; Aini et al. 2013 and Desi et al. 2014. Diseased leaves or stems were washed with running water, then dried with a tissue. Diseased samples were cut 1.5 x 5 cm, then sterilized with ethanol 70% and rinsed with sterile water 3 times. Furthermore, the sample was crushed with 5 ml of sterile distilled water using a sterile mortar. The bacterial suspension was streaked on nutrient agar and incubated 3-5 days. Bacterial colonies that exhibit the character of *Pantoea* sp. are yellow, shiny, slimy, flat or convex, then separated as pure cultures of *stewartii* candidates. The culture was then tested by Gram Reaction (KOH test), Hugh-Leifson test, pigment production in YDC medium, oxidase test, hypersensitivity test, and pathogenicity test on maize.

Isolation and characterization of endophytic bacteria

Sampling for the isolation of endophytic bacteria was carried out in Banyumas and Purbalingga, Central Java, Indonesia, with purposive stratified random sampling. Samples were taken from two areas of altitude, i.e., low-moderate lands (0-500 m above sea level), and highlands (> 500 m above sea level) (Nuryanto et al., 2014). In each district, 2 locations were selected for the low-medium lands, and 1 location for the highlands. Age of maize plants was 20-30 days after planting, when the number of endophytic microbial populations that can be cultured is in the highest population (Cavaglieri et al. 2009).

The endophytic bacteria are isolated from the roots and stems of healthy maize plants. Roots and stems are washed, sterilized with 70% ethanol (1 minute), 20% sodium hypochlorite (5 minutes) and Ringer's thiosulfate solution (5 minutes). Separately, the roots and stems of 10 g each were crushed with 90 ml PBS on a sterile mortar. Subsequently, samples were plated on NA and Kings B media (Cavaglieri et al. 2009). To isolate *Bacillus* sp., the suspension is heated for 10 minutes at 80 ° C, before plating on NA. Bacterial isolates were further purified and characterized, such as morphological characteristics, gram properties, catalase tests, and hypersensitivity tests

The antagonism test of endophytic bacterial to *R. solani*

The antagonism test of endophytic bacteria on *R. solani* was carried out using the dual culture method. The level of inhibition of antagonist is calculated using the formula (Abidin et al., 2015).

$$I = \frac{C - T}{C} \times 100\%$$

I = The level of inhibition of antagonist (%)

C = The radius of pathogen colonies opposite antagonist

T = The radius of the colony of pathogens towards antagonist

The antagonism test of endophytic bacterial to bacterial pathogens

Antagonism testing was carried out using the double-layer test method (Santiago et al. 2015). Endophytic bacteria to be tested were grown on the NA medium, incubated at 28 °C for 48 hours. In the upside-down position, 1 ml of chloroform was added to the cup lid and left for 2 hours. Next, add 5 mL so that 0.6% water containing 0.5 mL of *P. stewartii* bacterial suspension. The culture was re-incubated for 24 hours, and there were clear zones around the antagonistic bacterial colony. The antibiotic activity was assessed based on the diameter of the clear zone compared to the diameter of the colony. Characterization of the type of antibiosis can be divided into bactericidal and bacteriostatic types according to the method of Djatmiko et al. 2017.

The mechanism test of endophytic bacteria as controlling agents biological and plant growth-promoting microbial

The testing mechanism of endophytic bacteria is carried out for bacteria that have the potential in testing the antagonism of the fungus *R. solani* and *Pantoea* sp.

1). Protease Test

The activity of the ability of antagonistic bacteria to produce extracellular protease enzymes was tested using Skim Milk Agar (SMA) medium. Each bacterium to be tested was grown in a medium SMA and incubated at 28 °C for 24-48 hours. The presence of clear zones around the colony shows that positive bacteria produce protease enzymes (Abed et al. 2016). The protease activity index is assessed based on the diameter of the clear zone compared to the diameter of the colony..

$$\text{Protease index} = \frac{(\text{clear zone diameter} - \text{colony diameter})}{\text{colony diameter}}$$

2.) Lipase test

Detection of the ability of bacteria to produce the enzyme lipase was done by growing the antagonistic bacteria on a medium containing 1% Tween 80. The presence of lipase enzyme activity was demonstrated by milky white sediment around the bacterial colony, after incubating at 28 °C for 4-7 days. The lipolytic index was measured using a formula Djuric et al. (201).

$$\text{Lipolytic index} = \frac{(\text{milky white diameter} - \text{colony diameter})}{\text{colony diameter}}$$

3.) Uji fosfatase

Detection of the ability of bacteria to produce the enzyme phosphatase was done by growing bacterial isolates on Pikovkaya medium. After incubating for 7 days at 28 C, the presence of a clear zone around the bacterial colony shows that the bacteria has the ability to produce the phosphatase enzyme to dissolve phosphates. The solubility index is measured using a formula (Farooq and Bano 2013)

$$\text{Phosphatase Index} = \frac{\text{clear zone diameter} - \text{colony diameter}}{\text{colony diameter}}$$

RESULTS AND DISCUSSION

Isolation and characterization of endophytic bacteria

The results of the exploration, isolation, and characterization of endophytic bacteria obtained 23 isolates of endophytic bacteria, consisted of 9 isolates of the *fluorescent Pseudomonads* and 14 isolates of *Bacillus* sp. (Table 1). *Fluorescent Pseudomonads* colony on King's B is round, with a flat edge, and yellowish-white, to greenish-yellow, gram-negative, rod-shaped, non-spore and fluorescent under ultraviolet light. According to Arwiyanto et al. 2007), *P. fluorescens* have round, flat-edged, fluidal and release greenish-yellow colony in the King's B. Individual rod-shaped bacteria with a size (0.5-1.0) - (1.5-4.0) µm. The *P. fluorescens* isolates is gram-negative, which can form catalase, a positive oxidase, needed to grow aerobes.

Bacillus sp. has a spherical colony, cell rod-shaped, gram-positive, and endospores within cells. *Bacillus* sp. has the characteristics of a circular colony and punctiform (small round), variations in the entire margin and lobate, white dull, non-slimy, gram-positive, has endospores, flagellum and some are motile.

Based on its distribution, *fluorescent Pseudomonads* and *Bacillus* sp. found in all sampling locations, high or low-medium lands. This shows that *fluorescent Pseudomonads* and *Bacillus* sp spread and can live in various altitudes, both high and low-medium land. According to Bacon and Hilton 2002 and Ganeshan and Kumar 2005, *P. fluorescens* and *Bacillus* sp, are species of bacteria with a wide range of life and are very adaptive in various environments. Both types of bacteria are also found in the roots or corn stalks. According to Ganeshan and Kumar 2005; Orole and Adejumo 2011; Costa et al. 2013, *Bacillus* sp, and *Pseudomonas* sp. including a group of endophytic bacteria have a wide range of life and more isolated in maize.

Antagonism test between the endophytic bacteria against *R. solani*

Based on the results of in vitro tests (Table 2), 24 isolates of the endophytic bacteria were able to inhibit the growth of *R. solani*, with varying degrees of inhibition. The endophytic bacteria that have inhibition rates above 50%, i.e *Pseudomonas* Pf BK.A1 (51%), *Bacillus* sp. B.K.A1 (55.39%), *Bacillus* sp BK.A3 (51.52%), PP.A5 (50.66%), and PPD.B2 (50.8%). The effect of the endophytic bacteria in inhibiting the growth of *R. solani* is inversely proportional to the dry weight mycelium. The greater the percentage of inhibition of endophytic bacteria to the growth of *R. solani*, the smaller the dry weight mycelium (Table 2.)

The endophytic bacteria can inhibit the growth of *R solani* shown by the inhibition zone around the bacterial colony (Fig. 1). The endophytic bacteria have anti-pathogenic properties and can produce antibiotic compounds. The ability of the endophytic bacteria to control plant pathogens occurs through the mechanism of antibiosis, competition, lysis, inducing resistance and producing growth substances. Bacteria capable of producing secondary metabolites that can inhibit growth or damage pathogens (Hastuti et al. 2014). These compounds, including alkaloids, phenols, flavonoids, glycosides, and phytoalexin (Soesanto et al. 2010).



Fig.1. Antagonism test between the endophytic bacteria against *R. solani* (a) and *Pantoea* sp..

Table 1. Isolation and characterization of endophytic bacteria

Land	Sampling location	sample	Gram test	Catalase test	oxidase test	Colony morphology	colony pigment	Fluorescence on KB Medium	Cell morphology	Endospores	Isolat
Highland	1. Purbalingga, Pratin 7.13'33" LS, 109.17'21" BT, 1.190 m dpl	Root	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PP.A1
		Root	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. PP.A3
		Root	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. PP.A5
		Stem	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PP. B4
Medium-Lowland	2. Banyumas, Baturaden 7.19'1" LS, 109.14'29" BT, TT 520 m dpl	Root	-	+	+	round	Greenish yellow	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> BB.A2
		Root	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. BB.A3
		Stem	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BB.B4
	1. Banyumas, Sumbang 7.21'54" LS, 109.17'33" BT, TT 200 m dpl	Root	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> BS.A2
		Root	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BS.A1
		Root	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. BS.A3
		Stem	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. BS. B1
	2. Purbalingga, Bojongsari, 7.20'12" LS, 109.20'22" BT, TT 190 m dpl	Root	-	+	+	round	Greenish yellow	+	Small rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PB. A 4
		Stem	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. PB. B1
		Stem	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BB. B3
		Root	-	+	+	round	Greenish yellow	+	Small rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PPD A1
	3. Purbalingga, Padamara, 7.22'28" LS, 109.13'24" BT, TT 180 m dpl	Stem	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PPD. B1
		Stem	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PPD. B5
		Stem	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. PPD. B2
		Stem	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. PPD. B4
	4. Banyumas, Kembaran 7.23'47" LS, 109.17'9" BT, TT 110 m dpl	Root	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> BK. A1
		Root	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BK.A1
		Root	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BK.A3
		Stem	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BK.B3

152
153

Table 2. Inhibition of endophytic bacteria against *R. solani*

No	Isolate	Inhibition rate (%)	Dry weights Mycelium
1	Control	0	0,093
Endophytic bacteria from the root			
2	<i>Pseudomonas</i> Pf BB.A2	49,00	0,038
3	<i>Pseudomonas</i> Pf BS.A 2	45,00	0,027
4	<i>Pseudomonas</i> Pf BK.A1	51,00	0,017
5	<i>Pseudomonas</i> Pf PPD.A1	10,33	0,059
6	<i>Pseudomonas</i> Pf PP.A1	38,33	0,017
7	<i>Pseudomonas</i> Pf PB.A4	18,00	0,037
8	<i>Bacillus</i> sp.BB.A3	40,42	0,030
9	<i>Bacillus</i> sp.BS.A1	48,73	0,016
10	<i>Bacillus</i> sp. BSA3	37,42	0,039
11	<i>Bacillus</i> sp. B.K.A1	55,39	0,002
12	<i>Bacillus</i> sp. B.K.A3	51,52	0,003
13	<i>Bacillus</i> sp.PP.A3	46,65	0,019
14	<i>Bacillus</i> sp.PP.A5	50,66	0,009
Endophytic bacteria from the stem			
15	<i>Pseudomonas</i> Pf PPD.B1	27,00	0,020
16	<i>Pseudomonas</i> Pf PPD.B5	49,33	0,013
17	<i>Pseudomonas</i> Pf PP.B4	65,67	0,004
18	<i>Bacillus</i> sp. BB.B4	44,44	0,026
19	<i>Bacillus</i> sp. BS.B1	49,74	0,012
20	<i>Bacillus</i> sp. BK. B3	40,36	0,031
21	<i>Bacillus</i> sp.PPD.B2	50,8	0,007
22	<i>Bacillus</i> sp. PPD.B4	39,44	0,036
23	<i>Bacillus</i> sp. PB.B1	37,29	0,047
24	<i>Bacillus</i> sp. PB.B3	44,9	0,022

154

The endophytic bacterial antagonism test against *Pantoea* sp.

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

The results of antagonism between the endophytic bacteria and *Pantoea* sp. show varied results. The endophytic bacteria that can inhibit bacterial growth are indicated by the presence of clear zones around the endophytic bacterial colonies (Fig.1). From the nine isolate *Pseudomonas* sp. were tested, only three isolates were able to inhibit the growth of the *Pantoea* sp., i.e Pf BS.A2, Pf BK.A1 Pf PPD.B5. While the isolates Pf BB.A2, Pf PPD.A1, Pf PP.A1, Pf PPD.B1, and Pf PP.B4 are not able to inhibit the growth of pathogenic bacteria. Meanwhile, all isolate *Bacillus* sp tested (thirteen isolates) were able to inhibit the growth of *Pantoea* sp. (Table 3).

The presence of clear zones around endophytic bacterial colonies shows the ability of endophytic bacteria to produce antibiotics to inhibit the growth of *Pantoea* sp. *P. fluorescens* P60 can produce antibiotics that inhibit the growth of pathogens (Soesanto 2011). *Pseudomonas fluorescens* is reported to produce phenazine-1-carboxylic acid (PCA) and other derivatives, 2,4 diacetyl phloroglucinol (DAPG), pyrrolnitrin (PRN) and or pyoluteorin (Plt) (Heydari and Pessarakli 2010). Nasrun and Burhanudin (2016) mention that *P. fluorescens* produce secondary metabolites, i.e. antimicrobial, cyanide acid and 2,4-diacetyl phloroglucinol phenazine, pyrrolnitrin, pyoluteorin antibiotics

The level of bacteria's ability to inhibit growth can be shown by the large diameter of the clear zone. The results showed that the antagonism index ranged from 1.67 - 8.17. Based on this index, most endophytic bacteria have a strong antagonism (index of antagonism > 4) (Davis and Stout 1971). Furthermore, bacterial isolates that showed antagonistic activity were tested for types of antagonism based on Djatmiko (2007). Based on the type of antagonistic activity, ten isolates the endophytic bacteria were bacteriostatic and nine isolates the endophytic bacteria were bactericidal. Bacteriostatic activity, growth inhibition is temporary, it is shown that regrowth of bacteria after being transferred to a new medium, which is free from the influence of antagonistic bacteria. Bactericidal activity, inhibition is permanent. Bacteria were unable to grow even though they are transferred to new medium

177 Table 3. Inhibition of endophytic bacteria against *Pantoea* sp.

No	Isolate	Antagonism	Antagonism index	Antagonism category*	Antagonism activity
Endophytic bacteria from the root					
1	<i>Pseudomonas</i> Pf BB.A2	-	0	-	-
2	<i>Pseudomonas</i> Pf BS.A 2	+	4,91	strong	bacteriostatic
3	<i>Pseudomonas</i> Pf BK.A1	+	4,42	strong	bacteriostatic
4	<i>Pseudomonas</i> Pf PPD.A1	-	0	-	-
5	<i>Pseudomonas</i> Pf PP.A1	-	0	-	-
6	<i>Pseudomonas</i> Pf PB.A4	+	5,29	strong	bactericidal
7	<i>Bacillus</i> sp.BB.A3	+	8,17	strong	bacteriostatic
8	<i>Bacillus</i> sp.BS.A1	+	4,00	strong	bacteriostatic
9	<i>Bacillus</i> sp. BSA3	+	5,07	strong	bactericidal
10	<i>Bacillus</i> sp. B.K.A1	+	4,01	strong	bakteriostatik
11	<i>Bacillus</i> sp. B.K.A3	+	4,91	strong	bacteriostatic
12	<i>Bacillus</i> sp.PP.A3	+	6,63	strong	bactericidal
13	<i>Bacillus</i> sp.PP.A5	+	6,56	strong	bactericidal
Endophytic bacteria from the stem					
14	<i>Pseudomonas</i> PPD.B1	-	0	-	-
15	<i>Pseudomonas</i> PPD.B5	+	5,86	strong	bactericidal
16	<i>Pseudomonas</i> PP.B4	-	0	-	-
17	<i>Bacillus</i> sp. BB.B4	+	7,80	strong	bactericidal
18	<i>Bacillus</i> sp. BS.B1	+	6,22	strong	bacteriostatic
19	<i>Bacillus</i> sp. BK. B3	+	5,33	strong	bacteriostatic
20	<i>Bacillus</i> sp.PPD.B2	+	5,00	strong	bacteriostatic
21	<i>Bacillus</i> sp. PPD.B4	+	8,75	strong	bacteriostatic
22	<i>Bacillus</i> sp. PB.B1	+	1,67	weak	bacteriostatic
23	<i>Bacillus</i> sp. PB.B3	+	5,67	strong	bactericidal

178 • Based on Davis and Stout, 1971

179 Test the mechanism of endophytic bacteria as biological control agents and plant growth-promoting microbes

180 The mechanism test is carried out on endophytic bacteria that have the potential to control the fungus *R. solani* and
 181 *Pantoea* sp., i.e. *Bacillus* sp. B.K.A1, *Bacillus* sp. B.K.A3, *Bacillus* sp. PP.A5, *Bacillus* sp. PPD.B2. The results of enzyme
 182 activity tests are as shown in Table 4. The production of compounds related to biocontrol of pathogens and/or promotion
 183 of plant growth in bacterial isolates was evaluated by measuring the production of antimicrobial compounds and hydrolytic
 184 enzymes (amylases, lipases, proteases, and chitinases) and phosphate solubilization. the isolates that have high protein and
 185 fat hydrolysis enzymes have the potential as biological control agents because proteins and fats are constituents of
 186 pathogen cells (Mota et al 2016).

187 The four isolates *Bacillus* sp. tested were able to produce protease, lipase and phosphatase enzymes, with varied
 188 activity indexes. All isolates of *Bacillus* sp. those tested had a high index of protease and lipase enzymes (> 3) (Table 4.,
 189 Fig 2.). Protease and lipase enzymes, related to the ability of the endophytic bacteria as biological control agents.
 190 According to Anderson et al. 2014, the extracellular protease enzyme produced by *P. fluorescens* can inactivate antibiotic
 191 compounds produced by *Pantoea agglomerans*. The phosphate solubilization is related to the ability of endophytic bacteria
 192 as a plant growth promoter, providing phosphates for plants.

193 Table 4. Test results of proteases, lipases and phosphate solubilization.

No	Isolate	Protease Test		Lipase Test		Phosphate solubilization	
		activity	index	activity	index	activity	index
Endophytic bacteria from the root							
1	<i>Bacillus</i> sp. B.K.A1	+	3.75	+	3.23	+	1.17
2	<i>Bacillus</i> sp. B.K.A3	+	3.20	+	3.73	+	1.27

3	<i>Bacillus</i> sp.PP.A5	+	5.00	+	4.40	+	1.46
Endophytic bacteria from the stem							
4	<i>Bacillus</i> sp.PPD.B2	+	3.00	+	3.90	+	2.60



Fig. 2. Hydrolysis enzyme activity, (a) protease , (b) lipase and (c) phosphate solubilization.

CONCLUSION

Based on research carried out, it has been successfully isolated, morphologically and biochemically characterized four the endophytic bacteria that have the potential to be developed as biopesticides to control maize disease, especially *R. solani* and *Pantoea* sp. They can suppress the growth of *R.solani* by more than 50%, have a strong antagonistic index against *Pantoea* sp (> 4), and can produce protease and lipase enzyme, and phosphate solubilization.

ACKNOWLEDGEMENTS

This research was a part of the main study funded by Doctoral Dissertation Grant from the Indonesian Ministry of Research, Technology and Higher Education; for that, I deeply thank for the financial support

REFERENCES

207 Abed HN, Rouag.,Mouatasssem D, Rouabhi A. 2016. Screening for *Pseudomonas* and *Bacillus* antagonistic rhizobacteria strains for the biocontrol of
208 Fusarium wilt of chickpea. Eurasian Journal of Soil Science 5(3):182–191

209 Abidin Z, Aini LQ, Abadi AL. 2015. Effect of *Bacillus* sp. and *Pseudomonas* sp. on the growth of the pathogenic fungus *Sclerotium rolfsii* Sacc. causes
210 of seedling diseases in soybean plants. Jurnal HPT 3(1): 1-10.

211 Aini,LQ, Suryani L, Sugiharto AN,. Abadi A. 2013. Identification of bacterial wilt and leaf blight disease on maize (*Zea Mays*) found in Kediri,
212 Indonesia. Agrivita 35 (1): 1-7.

213 Ammar, E, V.R. Correa, S.A. Hogenhout, and M.G. Redinbaugh. 2014. Immunofluorescence localization and ultrastructure of Stewart’s wilt disease
214 bacterium *Pantoea stewartii* in maize leaves and in its flea beetle vector *Chaetocnema pulicaria* (Coleoptera: Chrysomelidae). Journal of Microscopy
215 and Ultrastructure 2: 28–33

216 Anderson LM, Stockwell VO, Loper JE. 2004. An extracellular protease of *Pseudomonas fluorescens* inactivates antibiotics of *Pantoea agglomerans*.
217 Phytopathology 94:1228-1234

218 Arwiyanto T, Maryudani. YMS, Nurul N, Azizah. 2007. The phenotypic properties of *Pseudomonas fluorescens*, biological control agents for lincat
219 disease in temanggung tobacco. Biodiversitas 8(2) : 147-151.

220 Cavaglieri L, Orlando J, Etcheverry M. 2009. Rhizosphere microbial community structure at different maize plant growth stages and root locations.
221 Microbiological

222 Compant SB, Duffy, Nowak J. Clement C, Barka EA. 2005. Use of Plant Growth- Promotng Bacteria for Biocontrol of Plant Diseases: principles,
223 Mechanisme of Action, and Future Prospects. Applied and Enviromental Microbiology 71(9): 4951-4959.

224 Coplin DL, Redinbaugh M G. 2012. The Bacterium *Pantoea stewartii* Uses Two Different Type III Secretion Systems To Colonize Its Plant Host and
225 Insect Vector. Applied and Enviromental Microbiology 78(17): 6327-6336.

226 Costa FG, Zucchi TD, de Melo IS. 2013. Biological control of Phypathogenic Fungi by Endophytic Actinomycetes Isolated from Maize (*Zea Mays* L.).
227 Braz. Arch.Biol.Technol 56(6):948-955.

228 Davis WW, Stout TR. 1971. Disc plate methods of microbiological antibiotic assay. Applied Microbiology 22(4):659-665.

229 Djaenuddin N, Nonci N, Muis A. 2017. Effectiveness of the formula *Bacillus subtilis* TM4 for disease control in maize plants. Jurnal Fitopatologi
230 Indonesia 13(4): 113-118

231 Djatmiko HA, Arwiyanto T, Hadisutrisno B, Sunarminto BH. 2007. Potential of three bacterial genera from three plant rhizosphere as biological agents
232 controlling lincat disease. Jurnal ilmu-ilmu Pertanian 9(1):40-47.

233 Djuric SA, Pavic, Jarak M, Pavlovic S, Starovic M, Pivic R, Josic D. 2011. Selection of indigenous fluorescent pseudomonad isolates from maize
234 rhizospheric soil in Vojvodina as possible PGPR. Romanian Biotechnological Letters 16(5): 6580–6590.DOI:10.1094/PHI-I-2004-0113-01.

235 Farooq U, Bano A. 2013. Screening of indigenous bacteria from rhizosphere of maize (*Zea mays* L.) for their plant growth promotion ability and
236 antagonism against fungal and bacterial pathogens. Journal of Animal and Plant Sciences, 23(6), pp.1642– 1652.

237 Ganeshan G, Kumar AM. 2005. *Pseudomonas fluorescens*, a potential bacterial antagonist to control plant diseases. Journal of Plant Interactions 1(3):
238 123_134.

239 Hastuti RD, Saraswati R, Sari AP. 2014. The effectiveness of the endophytic microbes in promoting plant growth and controlling leaf blight disease in
240 the lowland rice. Jurnal Tanah dan Iklim 38(2) : 109-118.

241 Heydari A, Pessarakhi M. 2010. A review on biological control of fungal plant pathogens using microbial antagonists. Journal of Biological Science
242 10(4): 273-290

243 Motaa MS, Gomes CB, Júniora ITS, Moura AB. 2017. Bacterial selection for biological control of plant disease: criterion determination and validation.
244 Brazilian Journal of Microbiology 48 : 62–70

245 Muis A. 2007. Mmanagement of sheath blight disease (*Rhizoctonia solani* Kuhn.) in maize. Jurnal Litbang Pertanian 26(3) : 100-103Desi 2014

246 Nasrun, Burhanudin. 2016. Evaluation of the efficacy of the formula *Pseudomonas fluorescens* for controlling bacterial wilt (*Ralstonia solanacearum*)
247 patchouli. Buletin Penelitian Tanaman Rempah dan Obat, 27(1): 67-76.

248 Nuryanto B, Priyatmojo A, Hadisutrisno B. 2014. Pengaruh Tinggi Tempat dan Tipe Tanaman Padi terhadap Keparahana Penyakit Hawar Pelepah.
249 Penelitian Pertanian Tanaman Pangan 33(1):1–8.

250 Orole OO, Adejumo TO. 2011. Bacterial and fungal endophytes associated with grains and roots of maize. Journal of Ecology and the natural
251 Enviroment 3(9):298-303.

252 Pal KK, McSpadden Gardener B. 2006. Biological Control of Plant Pathogens.

253 Pataky JK. 2004. Stewart’s wilt of corn. The Plant Health Instructor.

254 Rosenblueth M, Martinez-Romero E. 2006. Bacterial endophytes and ther interaction with hosts (Review). MPMI 19 (8): 827-837.

255 Santiago TR, Grabowski C, Rossato M, Romeiroa RS., Mizubuti ESG. 2015. Biological control of eucalyptus bacterial wilt with rhizobacteria. Biological
256 Control 80:14–22. <http://dx.doi.org/10.1016/j.biocontrol.2014.09.007>.

257 Shanti AT, Vittal RR. 2013. Biocontrol potenciales of Plant Growth promoting Rhizobacteria Against Fuusarium Wilt Disease of Cucurbit. Esci J. Plant
258 Pathol 2(3): 155-161.

259 Soesanto L, Mugiastuti E, Rahayuniati RF. 2010. Study of the antagonistic mechanism of *Pseudomonas fluorescens* P60 against *Fusarium oxysporum*
260 f.sp. *lycopersici* in tomatoes in vivo. Jurnal HPT Tropika 10(2) : 108-115

261 Soesanto L, Mugiastuti E, Rahayuniati RF. 2011. Utilization of some animal broths as a liquid formula for *Pseudomonas fluorescens* P60 to control
262 *Sclerotium rolfsii* in cucumber plants. Jurnal Perlindungan Tanaman Indonesia, 17(1): 7–17.

263 SUBMISSION CHECKLIST

264
265
266 Ensure that the following items are present:
267

The first corresponding author must be accompanied with contact details:		Give mark (X)
• E-mail address		x
• Full postal address (incl street name and number (location), city, postal code, state/province, country)		x
• Phone and facsimile numbers (incl country phone code)		x

All necessary files have been uploaded, and contain:

• Keywords	x
• Running titles	x

• All figure captions	x
• All tables (incl title and note/description)	x

Further considerations

• Manuscript has been “spell & grammar-checked” Better, if it is revised by a professional science editor or a native English speaker	x
• References are in the correct format for this journal	x
• All references mentioned in the Reference list are cited in the text, and vice versa	x
• Colored figures are only used if the information in the text may be losing without those images	x
• Charts (graphs and diagrams) are drawn in black and white images; use shading to differentiate	x

268

269

potential as maize diseases control

Commented [LN1]: Where?

Commented [LN2]: Concluding statement is required here

INTRODUCTION

MATERIALS AND METHODS

This research was conducted at the Laboratory of Plant Protection, Faculty of Agriculture, Jenderal Soedirman University, Purwokerto, Central Java, Indonesia, from April to August 2019

44 **Isolation *R. solani***
45 *R. solani* was isolated from maize with sheath blight symptoms and there was sclerotium as a resistant structure from
46 the pathogenic fungi in Banyumas. Samples were isolated on PDA medium to obtain pure *R. solani* isolates.

47 **Isolation *Pantoea* sp.**
48 *Pantoea* sp. isolated from diseased maize, which was taken from the maize growing area in Banyumas Regency.
49 *Pantoea* sp. was isolated according to Coplin et al. 2012; Aini et al. 2013 and Desi et al. 2014. Diseased leaves or stems
50 were washed with running water, then dried with a tissue. Diseased samples were cut 1.5 x 5 cm, then sterilized with
51 ethanol 70% and rinsed with sterile water 3 times. Furthermore, the sample was crushed with 5 ml of sterile distilled water
52 using a sterile mortar. The bacterial suspension was streaked on nutrient agar and incubated 3-5 days. Bacterial colonies
53 that exhibit the character of *Pantoea* sp. are yellow, shiny, slimy, flat or convex, then separated as pure cultures of *P.*
54 *stewartii* candidates. The culture was then tested by Gram Reaction (KOH test), Hugh-Leifson test, pigment production in
55 YDC medium, oxidase test, hypersensitivity test, and pathogenicity test on maize.

56 **Isolation and characterization of endophytic bacteria**
57 Sampling for the isolation of endophytic bacteria was carried out in Banyumas and Purbalingga, Central Java,
58 Indonesia, with purposive stratified random sampling. Samples were taken from two areas of altitude, i.e., low-moderate
59 lands (0-500 m above sea level), and highlands (> 500 m above sea level) (Nuryanto et al., 2014). In each district, 2
60 locations were selected for the low-medium lands, and 1 location for the highlands. Age of maize plants was 20-30 days
61 after planting, when the number of endophytic microbial populations that can be cultured is in the highest population
62 (Cavaglieri et al. 2009).

63 The endophytic bacteria are isolated from the roots and stems of healthy maize plants. Roots and stems are washed,
64 sterilized with 70% ethanol (1 minute), 20% natrium hypochlorite (5 minutes) and Ringer's thiosulfate solution (5
65 minutes). Separately, the roots and stems of 10 g each were crushed with 90 ml PBS on a sterile mortar. Subsequently,
66 samples were plated on NA and Kings B media (Cavaglieri et al. 2009). To isolate *Bacillus* sp., the suspension is heated
67 for 10 minutes at 80 ° C, before plating on NA. Bacterial isolates were further purified and characterized, such as
68 morphological characteristics, gram properties, catalase tests, and hypersensitivity tests

69 **The antagonism test of endophytic bacterial to *R. solani***
70 The antagonism test of endophytic bacteria on *R. solani* was carried out using the dual culture method. The level of
71 inhibition of antagonist is calculated using the formula (Abidin et al., 2015).

72
73
$$I = \frac{C - T}{C} \times 100\%$$

74
75 I = The level of inhibition of antagonist (%)
76 C = The radius of pathogen colonies opposite antagonist
77 T = The radius of the colony of pathogens towards antagonist

78 **The antagonism test of endophytic bacterial to bacterial pathogens**
79 Antagonism testing was carried out using the double-layer test method (Santiago et al. 2015). Endophytic bacteria to be
80 tested were grown on the NA medium, incubated at 28 C for 48 hours. In the upside-down position, 1 ml of chloroform
81 was added to the cup lid and left for 2 hours. Next, add 5 mL so that 0.6% water containing 0.5 mL of *P. stewartii* bacterial
82 suspension. The culture was re-incubated for 24 hours, and there were clear zones around the antagonistic bacterial colony.
83 The antibiotic activity was assessed based on the diameter of the clear zone compared to the diameter of the colony.
84 Characterization of the type of antibiosis can be divided into bactericidal and bacteriostatic types according to the method
85 of Djatmiko et al. (2017).

86 The mechanism test of endophytic bacteria as controlling agents biological and plant growth-promoting microbial
87 The testing mechanism of endophytic bacteria is carried out for bacteria that have the potential in testing the
88 antagonism of the fungus *R. solani* and *Pantoea* sp.

89 **Protease Test**
90 The activity of the ability of antagonistic bacteria to produce extracellular protease enzymes was tested using Skim
91 Milk Agar (SMA) medium. Each bacterium to be tested was grown in a medium SMA and incubated at 28 C for 24-48
92 hours. The presence of clear zones around the colony shows that positive bacteria produce protease enzymes (Abed et al.
93 2016). The protease activity index is assessed based on the diameter of the clear zone compared to the diameter of the
94 colony..

95 Protease index =
$$\frac{\text{clear zone diameter} - \text{colony diameter}}{\text{colony diameter}}$$

96

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Lipase test

Detection of the ability of bacteria to produce the enzyme lipase was done by growing the antagonistic bacteria on a medium containing 1% Tween 80. The presence of lipase enzyme activity was demonstrated by milky white sediment around the bacterial colony, after incubating at 28 C for 4-7 days. The lipolytic index was measured using a formula Djuric et al. (201).

$$\text{Lipolytic index} = \frac{(\text{milky white diameter} - \text{colony diameter})}{\text{colony diameter}}$$

Uji fosfatase

Detection of the ability of bacteria to produce the enzyme phosphatase was done by growing bacterial isolates on Pikovkaya medium. After incubating for 7 days at 28 C, the presence of a clear zone around the bacterial colony shows that the bacteria has the ability to produce the phosphatase enzyme to dissolve phosphates. The solubility index is measured using a formula (Farooq and Bano 2013)

$$\text{Phosphatase Index} = \frac{\text{clear zone diameter} - \text{colony diameter}}{\text{colony diameter}}$$

RESULTS AND DISCUSSION

Isolation and characterization of endophytic bacteria

The results of the exploration, isolation, and characterization of endophytic bacteria obtained 23 isolates of endophytic bacteria, consisted of 9 isolates of the *fluorescent Pseudomonads* and 14 isolates of *Bacillus* sp. (Table 1). *Fluorescent Pseudomonads* colony on King's B is round, with a flat edge, and yellowish-white, to greenish-yellow, gram-negative, rod-shaped, non-spore and fluorescent under ultraviolet light. According to Arwiyanto et al. (2007), *P. fluorescens* have round, flat-edged, fluidal and release greenish-yellow colony in the King's B. Individual rod-shaped bacteria with a size (0.5-1.0) - (1.5-4.0) µm. The *P. fluorescens* isolates is gram-negative, which can form catalase, a positive oxidase, needed to grow aerobes.

Bacillus sp. has a spherical colony, cell rod-shaped, gram-positive, and endospores within cells. *Bacillus* sp. has the characteristics of a circular colony and punctiform (small round), variations in the entire margin and lobate, white dull, non-slimy, gram-positive, has endospores, flagellum and some are motile.

Based on its distribution, *fluorescent Pseudomonads* and *Bacillus* sp. found in all sampling locations, high or low-medium lands. This shows that *fluorescent Pseudomonads* and *Bacillus* sp spread and can live in various altitudes, both high and low-medium land. According to Bacon and Hilton 2002 and Ganeshan and Kumar 2005, *P. fluoresscens* and *Bacillus* sp. are species of bacteria with a wide range of life and are very adaptive in various environments. Both types of bacteria are also found in the roots or corn stalks. According to Ganeshan and Kumar 2005; Orole and Adejumo 2011; Costa et al. 2013, *Bacillus* sp. and *Pseudomonas* sp. including a group of endophytic bacteria have a wide range of life and more isolated in maize.

Antagonism test between the endophytic bacteria against *R. solani*

Based on the results of in vitro tests (Table 2), 24 isolates of the endophytic bacteria were able to inhibit the growth of *R. solani*, with varying degrees of inhibition. The endophytic bacteria that have inhibition rates above 50%, i.e *Pseudomonas* Pf BK.A1 (51%), *Bacillus* sp. B.K.A1 (55.39%), *Bacillus* sp BK.A3 (51.52%), PP.A5 (50.66%), and PPD.B2 (50.8%). The effect of the endophytic bacteria in inhibiting the growth of *R. solani* is inversely proportional to the dry weight mycelium. The greater the percentage of inhibition of endophytic bacteria to the growth of *R. solani*, the smaller the dry weight mycelium (Table 2.)

The endophytic bacteria can inhibit the growth of *R solani* shown by the inhibition zone around the bacterial colony (Fig. 1). The endophytic bacteria have anti-pathogenic properties and can produce antibiotic compounds. The ability of the endophytic bacteria to control plant pathogens occurs through the mechanism of antibiosis, competition, lysis, inducing resistance and producing growth substances. Bacteria capable of producing secondary metabolites that can inhibit growth or damage pathogens (Hastuti et al. 2014). These compounds, including alkaloids, phenols, flavonoids, glycosides, and phytoalexin (Soesanto et al. 2010).

Formatted: Font: Italic, Highlight

150 **Table 1.** Isolation and characterization of endophytic bacteria
151

Land	Sampling location	sample	Gram test	Catalase test	oxidase test	Colony morphology	colony pigment	Fluorescence on KB Medium	Cell morphology	Endospores	Isolat
Highland	1. Purbalingga, Pratin 7.13'33" LS, 109.17'21" BT, TT 1.190 m dpl	Root	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PP.A1
		Root	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. PP.A3
		Root	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. PP.A5
		Stem	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PP. B4
Medium-Lowland	2.Banyumas, Baturaden 7.19"1" LS, 109.14'29" BT, TT 520 m dpl	Root	-	+	+	round	Greenish yellow	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> BB.A2
		Root	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. BB.A3
		Stem	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BB.B4
	1.Banyumas, Sumbang7.21'54" LS, 109.17'33"BT, TT 200 m dpl	Root	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> BS.A2
		Root	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BS.A1
		Root	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. BS.A3
		Stem	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. BS. B1
	2. Purbalingga, Bojongsari, 7.20'12" LS, 109.20'22" BT, TT 190 m dpl	Root	-	+	+	round	Greenish yellow	+	Small rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PB. A 4
		Stem	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. PB. B1
		Stem	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BB. B3
		Root	-	+	+	round	Greenish yellow	+	Small rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PPD A1
	3.Purbalingga, Padamara, 7.22'28" LS, 109.13'24" BT, TT 180 m dpl	Stem	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PPD. B1
		Stem	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PPD. B5
		Stem	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. PPD. B2
		Stem	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. PPD. B4
	4. Banyumas, Kembaran 7.23'47" LS, 109.17'9" BT, TT 110 m dpl	Root	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> BK. A1
		Root	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BK.A1
		Root	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BK.A3
		Stem	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BK.B3



Figure 1. Antagonism test between the endophytic bacteria against *R. solani* (a) and *Pantoea* sp..

Table 2. Inhibition of endophytic bacteria against *R. solani*

No	Isolate	Inhibition rate (%)	Dry weights Mycelium
1	Control	0	0,093
Endophytic bacteria from the root			
2	<i>Pseudomonas</i> Pf BB.A2	49,00	0,038
3	<i>Pseudomonas</i> Pf BS.A 2	45,00	0,027
4	<i>Pseudomonas</i> Pf BK.A1	51,00	0,017
5	<i>Pseudomonas</i> Pf PPD.A1	10,33	0,059
6	<i>Pseudomonas</i> Pf PP.A1	38,33	0,017
7	<i>Pseudomonas</i> Pf PB.A4	18,00	0,037
8	<i>Bacillus</i> sp.BB.A3	40,42	0,030
9	<i>Bacillus</i> sp.BS.A1	48,73	0,016
10	<i>Bacillus</i> sp. BSA3	37,42	0,039
11	<i>Bacillus</i> sp. B.K.A1	55,39	0,002
12	<i>Bacillus</i> sp. B.K.A3	51,52	0,003
13	<i>Bacillus</i> sp.PP.A3	46,65	0,019
14	<i>Bacillus</i> sp.PP.A5	50,66	0,009
Endophytic bacteria from the stem			
15	<i>Pseudomonas</i> Pf PPD.B1	27,00	0,020
16	<i>Pseudomonas</i> Pf PPD.B5	49,33	0,013
17	<i>Pseudomonas</i> Pf PP.B4	65,67	0,004
18	<i>Bacillus</i> sp. BB.B4	44,44	0,026
19	<i>Bacillus</i> sp. BS.B1	49,74	0,012
20	<i>Bacillus</i> sp. BK. B3	40,36	0,031
21	<i>Bacillus</i> sp.PPD.B2	50,8	0,007
22	<i>Bacillus</i> sp. PPD.B4	39,44	0,036
23	<i>Bacillus</i> sp. PB.B1	37,29	0,047
24	<i>Bacillus</i> sp. PB.B3	44,9	0,022

The endophytic bacterial antagonism test against *Pantoea* sp.

The results of antagonism between the endophytic bacteria and *Pantoea* sp. show varied results. The endophytic bacteria that can inhibit bacterial growth are indicated by the presence of clear zones around the endophytic bacterial colonies (Fig.1). From the nine isolate *Pseudomonas* sp. were tested, only three isolates were able to inhibit the growth of the *Pantoea* sp., i.e Pf BS.A2, Pf BK.A1 Pf PPD.B5. While the isolates Pf BB.A2, Pf PPD.A1, Pf PP.A1, Pf PPD.B1, and Pf PP.B4 are not able to inhibit the growth of pathogenic bacteria. Meanwhile, all isolate *Bacillus* sp tested (thirteen isolates) were able to inhibit the growth of *Pantoea* sp. (Table 3).

The presence of clear zones around endophytic bacterial colonies shows the ability of endophytic bacteria to produce antibiotics to inhibit the growth of *Pantoea* sp. *P. fluorescens* P60 can produce antibiotics that inhibit the growth of pathogens (Soesanto 2011). *Pseudomonas fluorescens* is reported to produce phenazine-1-carboxylic acid (PCA) and other derivatives, 2,4 diacetyl phloroglucinol (DAPG), pyrrolnitrin (PRN) and or pyoluteorin (Plt) (Heydari and Pessarakli 2010). Nasrun and Burhanudin (2016) mention that *P. fluorescens* produce secondary metabolites, i.e. antimicrobial, cyanide acid and 2,4-diacetyl phloroglucinol phenazine, pyrrolnitrin, pyoluteorin antibiotics.

The level of bacteria's ability to inhibit growth can be shown by the large diameter of the clear zone. The results showed that the antagonism index ranged from 1.67 - 8.17. Based on this index, most endophytic bacteria have a strong antagonism (index of antagonism > 4) (Davis and Stout 1971). Furthermore, bacterial isolates that showed antagonistic activity were tested for types of antagonism based on Djatmiko (2007). Based on the type of antagonistic activity, ten isolates the endophytic bacteria were bacteriostatic and nine isolates the endophytic bacteria were bactericidal. Bacteriostatic activity, growth inhibition is temporary, it is shown that regrowth of bacteria after being transferred to a new medium, which is free from the influence of antagonistic bacteria. Bactericidal activity, inhibition is permanent. Bacteria were unable to grow even though they are transferred to new medium.

Table 3. Inhibition of endophytic bacteria against *Pantoea* sp.

No	Isolate	Antagonism	Antagonism index	Antagonism catagory*	Antagonism activity
Endophytic bacteria from the root					
1	<i>Pseudomonas</i> Pf BB.A2	-	0	-	-
2	<i>Pseudomonas</i> Pf BS.A 2	+	4,91	strong	bacteriostatic
3	<i>Pseudomonas</i> Pf BK.A1	+	4,42	strong	bacteriostatic
4	<i>Pseudomonas</i> Pf PPD.A1	-	0	-	-
5	<i>Pseudomonas</i> Pf PP.A1	-	0	-	-
6	<i>Pseudomonas</i> Pf PB.A4	+	5,29	strong	bactericidal
7	<i>Bacillus</i> sp.BB.A3	+	8,17	strong	bacteriostatic
8	<i>Bacillus</i> sp.BS.A1	+	4,00	strong	bacteriostatic
9	<i>Bacillus</i> sp. BSA3	+	5,07	strong	bactericidal
10	<i>Bacillus</i> sp. B.K.A1	+	4,01	strong	bakteriostatik
11	<i>Bacillus</i> sp. B.K.A3	+	4,91	strong	bacteriostatic
12	<i>Bacillus</i> sp.PP.A3	+	6,63	strong	bactericidal
13	<i>Bacillus</i> sp.PP.A5	+	6,56	strong	bactericidal
Endophytic bacteria from the stem					
14	<i>Pseudomonas</i> PPD.B1	-	0	-	-
15	<i>Pseudomonas</i> PPD.B5	+	5,86	strong	bactericidal
16	<i>Pseudomonas</i> PP.B4	-	0	-	-
17	<i>Bacillus</i> sp. BB.B4	+	7,80	strong	bactericidal
18	<i>Bacillus</i> sp. BS.B1	+	6,22	strong	bacteriostatic
19	<i>Bacillus</i> sp. BK. B3	+	5,33	strong	bacteriostatic
20	<i>Bacillus</i> sp.PPD.B2	+	5,00	strong	bacteriostatic
21	<i>Bacillus</i> sp. PPD.B4	+	8,75	strong	bacteriostatic
22	<i>Bacillus</i> sp. PB.B1	+	1,67	weak	bacteriostatic
23	<i>Bacillus</i> sp. PB.B3	+	5,67	strong	bactericidal

*Based on Davis and Stout, 1971

Test the mechanism of endophytic bacteria as biological control agents and plant growth-promoting microbes

The mechanism test is carried out on endophytic bacteria that have the potential to control the fungus *R. solani* and *Pantoea* sp., i.e. *Bacillus* sp. B.K.A1, *Bacillus* sp. B.K.A3, *Bacillus* sp. PP.A5, *Bacillus* sp. PPD.B2. The results of enzyme activity tests are as shown in Table 4. The production of compounds related to biocontrol of pathogens and/or promotion of plant growth in bacterial isolates was evaluated by measuring the production of antimicrobial compounds and hydrolytic enzymes (amylases, lipases, proteases, and chitinases) and phosphate solubilization. the isolates that have high protein and fat hydrolysis enzymes have the potential as biological control agents because proteins and fats are constituents of pathogen cells (Mota et al 2016).

The four isolates *Bacillus* sp. tested were able to produce protease, lipase and phosphatase enzymes, with varied activity indexes. All isolates of *Bacillus* sp. those tested had a high index of protease and lipase enzymes (> 3) (Table 4.,

Fig 2.). Protease and lipase enzymes, related to the ability of the endophytic bacteria as biological control agents. According to Anderson et al. (2014), the extracellular protease enzyme produced by *P. fluorescens* can inactivate antibiotic compounds produced by *Pantoea agglomerans*. The phosphate solubilization is related to the ability of endophytic bacteria as a plant growth promoter, providing phosphates for plants.

Table 4. Test results of proteases, lipases and phosphate solubilization.

No	Isolate	Protease Test activity	index	Lipase Test activity	index	Phosphate solubilization activity	index
Endophytic bacteria from the root							
1	<i>Bacillus</i> sp. B.K.A1	+	3.75	+	3.23	+	1.17
2	<i>Bacillus</i> sp. B.K.A3	+	3.20	+	3.73	+	1.27
3	<i>Bacillus</i> sp.PP.A5	+	5.00	+	4.40	+	1.46
Endophytic bacteria from the stem							
4	<i>Bacillus</i> sp.PPD.B2	+	3.00	+	3.90	+	2.60



Figure 2. Hydrolysis enzyme activity, (a) protease , (b) lipase and (c) phosphate solubilization.

The author should expand the discussion by looking at previous published studies and compare with current findings.

Formatted: English (South Africa)

CONCLUSION

Based on research carried out, it has been successfully isolated, morphologically and biochemically characterized four the endophytic bacteria that have the potential to be developed as biopesticides to control maize disease, especially *R. solani* and *Pantoea* sp. They can suppress the growth of *R.solani* by more than 50%, have a strong antagonistic index against *Pantoea* sp (> 4), and can produce protease and lipase enzyme, and phosphate solubilization.

ACKNOWLEDGEMENTS

This research was a part of the main study funded by Doctoral Dissertation Grant from the Indonesian Ministry of Research, Technology and Higher Education; for that, I deeply thank for the financial support

REFERENCES

Abed HN, Rouag.,Mouatassem D, Rouabhi A. 2016. Screening for *Pseudomonas* and *Bacillus* antagonistic rhizobacteria strains for the biocontrol of Fusarium wilt of chickpea. Eurasian Journal of Soil Science 5(3):182–191

Abidin Z, Aini LQ, Abadi AL. 2015. Effect of *Bacillus* sp. and *Pseudomonas* sp. on the growth of the pathogenic fungus *Sclerotium rolfsii* Sacc. causes of seedling diseases in soybean plants. Jurnal HPT 3(1): 1-10.

Aini,LQ, Suryani L, Sugiharto AN., Abadi A. 2013. Identification of bacterial wilt and leaf blight disease on maize (*Zea Mays*) found in Kediri, Indonesia. Agrivita 35 (1): 1-7.

Ammar, E, V.R. Correa, S.A. Hogenhout, and M.G. Redinbaugh. 2014. Immunofluorescence localization and ultrastructure of Stewart's wilt disease bacterium *Pantoea stewartii* in maize leaves and in its flea beetle vector *Chaetocnema pulicaria* (Coleoptera: Chrysomelidae). Journal of Microscopy and Ultrastructure 2: 28–33

Anderson LM, Stockwell VO, Loper JE. 2004. An extracellular protease of *Pseudomonas fluorescens* inactivates antibiotics of *Pantoea agglomerans*. Phytopathology 94:1228-1234

Arwiyanto T, Maryudani. YMS, Nurul N, Azizah. 2007. The phenotypic properties of *Pseudomonas fluorescens*, biological control agents for lincat disease in temanggung tobacco. Biodiversitas 8(2) : 147-151.

Cavaglieri L, Orlando J, Etcheverry M. 2009. Rhizosphere microbial community structure at different maize plant growth stages and root locations. Microbiological

- Compant SB, Duffy, Nowak J, Clement C, Barka EA. 2005. Use of Plant Growth- Promoting Bacteria for Biocontrol of Plant Diseases: principles, Mechanisms of Action, and Future Prospects. *Applied and Environmental Microbiology* 71(9): 4951-4959.
- Coplin DL, Redinbaugh M G. 2012. The Bacterium *Pantoea stewartii* Uses Two Different Type III Secretion Systems To Colonize Its Plant Host and Insect Vector. *Applied and Environmental Microbiology* 78(17): 6327-6336.
- Costa FG, Zucchini TD, de Melo IS. 2013. Biological control of Phytopathogenic Fungi by Endophytic Actinomycetes Isolated from Maize (*Zea Mays* L.). *Braz. Arch.Biol.Technol* 56(6):948-955.
- Davis WW, Stout TR. 1971. Disc plate methods of microbiological antibiotic assay. *Applied Microbiology* 22(4):659-665.
- Djaenuddin N, Nonci N, Muis A. 2017. Effectiveness of the formula *Bacillus subtilis* TM4 for disease control in maize plants. *Jurnal Fitopatologi Indonesia* 13(4): 113-118
- Djarmiko HA, Arwiyanto T, Hadisutrisno B, Sunarminto BH. 2007. Potential of three bacterial genera from three plant rhizosphere as biological agents controlling lincat disease. *Jurnal ilmu-ilmu Pertanian* 9(1):40-47.
- Djuric SA, Pavic, Jarak M, Pavlovic S, Starovic M, Pivic R, Josic D.. 2011. Selection of indigenous fluorescent pseudomonad isolates from maize rhizospheric soil in Vojvodina as possible PGPR. *Romanian Biotechnological Letters* 16(5): 6580–6590.DOI:10.1094/PHI-I-2004-0113-01.
- Farooq U, Bano A. 2013. Screening of indigenous bacteria from rhizosphere of maize (*Zea mays* L.) for their plant growth promotion ability and antagonism against fungal and bacterial pathogens. *Journal of Animal and Plant Sciences*, 23(6), pp.1642– 1652.
- Ganeshan G, Kumar AM. 2005. *Pseudomonas fluorescens*, a potential bacterial antagonist to control plant diseases. *Journal of Plant Interactions* 1(3): 123-134.
- Hastuti RD, Saraswati R, Sari AP. 2014. The effectiveness of the endophytic microbes in promoting plant growth and controlling leaf blight disease in the lowland rice. *Jurnal Tanah dan Iklim* 38(2) : 109-118.
- Heydari A, Pessarakhi M. 2010. A review on biological control of fungal plant pathogens using microbial antagonists. *Journal of Biological Science* 10(4): 273-290
- Motaa MS, Gomes CB, Júnior ITS, Moura AB. 2017. Bacterial selection for biological control of plant disease: criterion determination and validation. *Brazilian Journal of Microbiology* 48 : 62–70
- Muis A. 2007. Management of sheath blight disease (*Rhizoctonia solani* Kuhn.) in maize. *Jurnal Litbang Pertanian* 26(3) : 100-103
- Nasrun, Burhanudin. 2016. Evaluation of the efficacy of the formula *Pseudomonas fluorescens* for controlling bacterial wilt (*Ralstonia solanacearum*) patchouli. *Buletin Penelitian Tanaman Rempah dan Obat*, 27(1): 67-76.
- Nuryanto B, Priyatomojo A, Hadisutrisno B.. 2014. Pengaruh Tinggi Tempat dan Tipe Tanaman Padi terhadap Keparahan Penyakit Hawar Pelepah. *Penelitian Pertanian Tanaman Pangan* 33(1):1–8.
- Orole OO, Adejumo TO. 2011. Bacterial and fungal endophytes associated with grains and roots of maize. *Journal of Ecology and the natural Environment* 3(9):298-303.
- Pal KK, McSpadden Gardener B. 2006. Biological Control of Plant Pathogens.
- Pataky JK. 2004. Stewart's wilt of corn. The Plant Health Instructor.
- Rosenblueth M, Martinez-Romero E. 2006. Bacterial endophytes and their interaction with hosts (Review). *MPMI* 19 (8): 827-837.
- Santiago TR, Grabowski C, Rossato M, Romeiroa RS., Mizubuti ESG. 2015. Biological control of eucalyptus bacterial wilt with rhizobacteria. *Biological Control* 80:14–22. <http://dx.doi.org/10.1016/j.biocontrol.2014.09.007>.
- Shanti AT, Vittal RR. 2013. Biocontrol potentials of Plant Growth promoting Rhizobacteria Against Fusarium Wilt Disease of Cucurbit. *Esci J. Plant Pathol* 2(3): 155-161.
- Soesanto L, Mugiastuti E, Rahayuniati RF. 2010. Study of the antagonistic mechanism of *Pseudomonas fluorescens* P60 against *Fusarium oxysporum* f.sp. *lycopersici* in tomatoes in vivo. *Jurnal HPT Tropika* 10(2) : 108-115
- Soesanto L, Mugiastuti E, Rahayuniati RF. 2011. Utilization of some animal broths as a liquid formula for *Pseudomonas fluorescens* P60 to control *Sclerotium rolfsii* in cucumber plants. *Jurnal Perlingungan Tanaman Indonesia*, 17(1): 7–17.

Isolation and characterization of the endophytic bacteria, and their potential as maize diseases control

Abstract. The research aimed to isolate and characterize morphologically and biochemically the endophytic bacteria, and their potential to control maize diseases, especially sheath blight and bacterial wilt. The study was conducted at the Plant Protection Laboratory, Faculty of Agriculture, Jenderal Soedirman University, from April to August 2019. The study consisted of four stages: isolation and characterization of endophytic bacteria, the antagonism test of the endophytic bacterial to *R. solani*, the antagonism test of the endophytic bacteria to *Pantoea* sp., and the mechanism test of the endophytic bacteria as biological control agents and plant growth-promoting bacteria. Based on the research, it has been successfully isolated, and characterized morphologically and biochemically characterized four endophytic bacteria isolates that have the potential to be developed as biopesticides to control maize disease, especially *R. solani* and *Pantoea* sp. They can suppress the growth of *R. solani* by more than 50%, have a strong antagonistic index against *Pantoea* sp (> 4), and can produce protease and lipase enzyme, and phosphate solubilization.

Key words: *Bacillus* sp., Fluorescents *Pseudomonads*, *Pantoea* sp., *Rhizoctoni solani*

Running title: Isolation and characterization of the endophytic

INTRODUCTION

Maize is a strategic food commodity in the world. In Indonesia, the government seeks to achieve self-sufficiency in maize through increasing production of sustainable maize. However, these efforts have faced several obstacles, one of them is the presence of plant diseases such as sheath blight, caused by *Rhizoctonia solani* Kuhn, and bacterial wilt caused by (*Pantoea* sp.). *R. solani* can infect up to the midrib of the cob (Djaenuddin et al. 2017), resulting in a decrease in the yield of up to 100%. (Muis 2007). *Pantoea* sp. can attack all stages of the plant causing wilting and leaf blight, and is known as Stewart's wilt (Pataky 2004; Ammar et al. 2014). The pathogens can cause 40-100% yield loss.

Over the past 3 decades, the concept of sustainable and environmentally friendly agriculture have been carried out by minimizing the use of chemicals, both synthetic fertilizers and synthetic pesticides. In the management of pests and plant diseases, biological control is developed by applying biological control agents including the endophytic bacteria (Shanti and Vittal 2013). Many endophytic bacteria can pass the endodermic barrier across from the root cortex to the vascular system, and subsequently develop as endophytes in stems, leaves, tubers, and other organs (Compant et al. 2005). The use of endophytic bacteria as biological agents has an advantage compared to rhizosphere bacteria because endophytic bacteria live and survive in the plant tissue during plant development, thus protecting the plants.

Bacillus sp. and fluorescent *Pseudomonads* are reported to be able to live as endophytes and are widely used as biological control agents for soil-borne and air-borne diseases. The endophytic bacteria could control plant diseases through several mechanisms including competition, hyperparasite, producing microbial inhibiting compounds (antibiotics, lysis enzymes, other physical or chemical disorders), enhancing plant resistance, and promoting plant growth (Compant et al 2005, Pal and McSpadden 2006; Rosenblueth and Martinez Romero 2006;).

Based on the mechanisms, the use of endophytic bacteria isolated from maize, both upland and lowland, suggested potentially alternative control for sheath blight (*R. solani*) and bacterial wilt (*Pantoea* sp). The research aimed to isolate and characterize morphologically and biochemically the endophytic bacterial as well as their potential to control pathogens that cause disease in maize especially *R. solani* and *Pantoea* sp.

MATERIALS AND METHODS

This research was conducted at the Laboratory of Plant Protection, Faculty of Agriculture, Jenderal Soedirman University, Purwokerto, Central Java, Indonesia, from April to August 2019

Commented [LN1]: Concluding statement is required here

44 **Isolation *R. solani***

45 *R. solani* was isolated from maize with sheath blight symptoms and there was sclerotium as a resistant structure from
46 the pathogenic fungi in Banyumas. Samples were isolated on PDA medium to obtain pure *R. solani* isolates.

47 **Isolation *Pantoea* sp.**

48 *Pantoea* sp. isolated from diseased maize, which was taken from the maize growing area in Banyumas Regency.
49 *Pantoea* sp. was isolated according to Coplin et al. 2012; Aini et al. 2013 and Desi et al. 2014. Diseased leaves or stems
50 were washed with running water, then dried with a tissue. Diseased samples were cut 1.5 x 5 cm, then sterilized with
51 ethanol 70% and rinsed with sterile water 3 times. Furthermore, the sample was crushed with 5 ml of sterile distilled water
52 using a sterile mortar. The bacterial suspension was streaked on nutrient agar and incubated 3-5 days. Bacterial colonies
53 that exhibit the character of *Pantoea* sp. are yellow, shiny, slimy, flat or convex, then separated as pure cultures of *P.*
54 *stewartii* candidates. The culture was then tested by Gram Reaction (KOH test), Hugh-Leifson test, pigment production in
55 YDC medium, oxidase test, hypersensitivity test, and pathogenicity test on maize.

56 **Isolation and characterization of endophytic bacteria**

57 Sampling for the isolation of endophytic bacteria was carried out in Banyumas and Purbalingga, Central Java,
58 Indonesia, with purposive stratified random sampling. Samples were taken from two areas of altitude, i.e., low-moderate
59 lands (0-500 m above sea level), and highlands (> 500 m above sea level) (Nuryanto et al., 2014). In each district, 2
60 locations were selected for the low-medium lands, and 1 location for the highlands. Age of maize plants was 20-30 days
61 after planting, when the number of endophytic microbial populations that can be cultured is in the highest population
62 (Cavaglieri et al. 2009).

63 The endophytic bacteria are isolated from the roots and stems of healthy maize plants. Roots and stems are washed,
64 sterilized with 70% ethanol (1 minute), 20% sodium hypochlorite (5 minutes) and Ringer's thiosulfate solution (5
65 minutes). Separately, the roots and stems of 10 g each were crushed with 90 ml PBS on a sterile mortar. Subsequently,
66 samples were plated on NA and Kings B media (Cavaglieri et al. 2009). To isolate *Bacillus* sp., the suspension is heated
67 for 10 minutes at 80 ° C, before plating on NA. Bacterial isolates were further purified and characterized, such as
68 morphological characteristics, gram properties, catalase tests, and hypersensitivity tests

69 **The antagonism test of endophytic bacterial to *R. solani***

70 The antagonism test of endophytic bacteria on *R. solani* was carried out using the dual culture method. The level of
71 inhibition of antagonist is calculated using the formula (Abidin et al., 2015).

72

$$73 \quad I = \frac{C - T}{C} \times 100\%$$

74

75 I = The level of inhibition of antagonist (%)

76 C = The radius of pathogen colonies opposite antagonist

77 T = The radius of the colony of pathogens towards antagonist

78 **The antagonism test of endophytic bacterial to bacterial pathogens**

79 Antagonism testing was carried out using the double-layer test method (Santiago et al. 2015). Endophytic bacteria to be
80 tested were grown on the NA medium, incubated at 28 C for 48 hours. In the upside-down position, 1 ml of chloroform
81 was added to the cup lid and left for 2 hours. Next, add 5 mL so that 0.6% water containing 0.5 mL of *P. stewartii* bacterial
82 suspension. The culture was re-incubated for 24 hours, and there were clear zones around the antagonistic bacterial colony.
83 The antibiotic activity was assessed based on the diameter of the clear zone compared to the diameter of the colony.
84 Characterization of the type of antibiosis can be divided into bactericidal and bacteriostatic types according to the method
85 of Djatmiko et al. (2017).

86 The mechanism test of endophytic bacteria as controlling agents biological and plant growth-promoting microbial

87 The testing mechanism of endophytic bacteria is carried out for bacteria that have the potential in testing the
88 antagonism of the fungus *R. solani* and *Pantoea* sp.

89 **Protease Test**

90 The activity of the ability of antagonistic bacteria to produce extracellular protease enzymes was tested using Skim
91 Milk Agar (SMA) medium. Each bacterium to be tested was grown in a medium SMA and incubated at 28 C for 24-48
92 hours. The presence of clear zones around the colony shows that positive bacteria produce protease enzymes (Abed et al.
93 2016). The protease activity index is assessed based on the diameter of the clear zone compared to the diameter of the
94 colony..

95 Protease index = $\frac{\text{clear zone diameter} - \text{colony diameter}}{\text{colony diameter}}$

96

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Lipase test

Detection of the ability of bacteria to produce the enzyme lipase was done by growing the antagonistic bacteria on a medium containing 1% Tween 80. The presence of lipase enzyme activity was demonstrated by milky white sediment around the bacterial colony, after incubating at 28 C for 4-7 days. The lipolytic index was measured using a formula Djuric et al. (201).

$$\text{Lipolytic index} = \frac{(\text{milky white diameter} - \text{colony diameter})}{\text{colony diameter}}$$

Uji fosfatase

Detection of the ability of bacteria to produce the enzyme phosphatase was done by growing bacterial isolates on Pikovkaya medium. After incubating for 7 days at 28 C, the presence of a clear zone around the bacterial colony shows that the bacteria has the ability to produce the phosphatase enzyme to dissolve phosphates. The solubility index is measured using a formula (Farooq and Bano 2013)

$$\text{Phosphatase Index} = \frac{\text{clear zone diameter} - \text{colony diameter}}{\text{colony diameter}}$$

RESULTS AND DISCUSSION

Isolation and characterization of endophytic bacteria

The results of the exploration, isolation, and characterization of endophytic bacteria obtained 23 isolates of endophytic bacteria, consisted of 9 isolates of the *fluorescent Pseudomonads* and 14 isolates of *Bacillus* sp. (Table 1). *Fluorescent Pseudomonads* colony on King's B is round, with a flat edge, and yellowish-white, to greenish-yellow, gram-negative, rod-shaped, non-spore and fluorescent under ultraviolet light. According to Arwiyanto et al. (2007), *P. fluorescens* have round, flat-edged, fluidal and release greenish-yellow colony in the King's B. Individual rod-shaped bacteria with a size (0.5-1.0) - (1.5-4.0) µm. The *P. fluorescens* isolates is gram-negative, which can form catalase, a positive oxidase, needed to grow aerobes.

Bacillus sp. has a spherical colony, cell rod-shaped, gram-positive, and endospores within cells (Table 1.). *Bacillus* sp. has the characteristics of a circular colony and punctiform (small round), variations in the entire margin and lobate, white dull, non-slimy, gram-positive, has endospores, flagellum and some are motile.

Based on its distribution, *fluorescent Pseudomonads* and *Bacillus* sp. found in all sampling locations, high or low-medium lands. This shows that *fluorescent Pseudomonads* and *Bacillus* sp spread and can live in various altitudes, both high and low-medium land. According to Bacon and Hilton 2002 and Ganeshan and Kumar 2005, *P. fluorescens* and *Bacillus* sp. are species of bacteria with a wide range of life and are very adaptive in various environments. Both types of bacteria are also found in the roots or corn stalks. According to Ganeshan and Kumar 2005; Orole and Adejumo 2011; Costa et al. 2013, *Bacillus* sp. and *Pseudomonas* sp. including a group of endophytic bacteria have a wide range of life and more isolated in maize.

Antagonism test between the endophytic bacteria against *R. solani*

Based on the results of in vitro tests (Table 2), 24 isolates of the endophytic bacteria were able to inhibit the growth of *R. solani*, with varying degrees of inhibition. The endophytic bacteria that have inhibition rates above 50%, i.e *Pseudomonas* Pf BK.A1 (51%), *Bacillus* sp. B.K.A1 (55.39%), *Bacillus* sp BK.A3 (51.52%), PP.A5 (50.66%), and PPD.B2 (50.8%). The effect of the endophytic bacteria in inhibiting the growth of *R. solani* is inversely proportional to the dry weight mycelium. The greater the percentage of inhibition of endophytic bacteria to the growth of *R. solani*, the smaller the dry weight mycelium (Table 2.)

The endophytic bacteria can inhibit the growth of *R solani* shown by the inhibition zone around the bacterial colony (Fig. 1). The endophytic bacteria have anti-pathogenic properties and can produce antibiotic compounds. The ability of the endophytic bacteria to control plant pathogens occurs through the mechanism of antibiosis, competition, lysis, inducing resistance and producing growth substances. Bacteria capable of producing secondary metabolites that can inhibit growth or damage pathogens (Hastuti et al. 2014). These compounds, including alkaloids, phenols, flavonoids, glycosides, and phytoalexin (Soesanto et al. 2010).

Formatted: Font: Italic, Highlight

150 **Table 1.** Isolation and characterization of endophytic bacteria
151

Land	Sampling location	sample	Gram test	Catalase test	oxidase test	Colony morphology	colony pigment	Fluorescence on KB Medium	Cell morphology	Endospores	Isolat
Highland	1. Purbalingga, Pratin LS, 7.13'33" BT, 109.17'21" TT 1.190 m dpl	Root	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PP.A1
		Root	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. PP.A3
		Root	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. PP.A5
		Stem	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PP. B4
Medium-Lowland	2.Banyumas, Baturaden 7.19"1" LS, 109.14'29" BT, TT 520 m dpl	Root	-	+	+	round	Greenish yellow	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> BB.A2
		Root	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. BB.A3
		Stem	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BB.B4
	1.Banyumas, Sumbang7.21'54" LS, 109.17'33"BT, TT 200 m dpl	Root	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> BS.A2
		Root	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BS.A1
		Root	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. BS.A3
		Stem	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. BS. B1
	2.Purbalingga, Bojongsari, 7.20'12" LS, 109.20'22" BT, TT 190 m dpl	Root	-	+	+	round	Greenish yellow	+	Small rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PB. A 4
		Stem	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. PB. B1
		Stem	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BB. B3
		Root	-	+	+	round	Greenish yellow	+	Small rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PPD A1
	3.Purbalingga, Padamara, 7.22'28" LS, 109.13'24" BT, TT 180 m dpl	Stem	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PPD. B1
		Stem	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PPD. B5
		Stem	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. PPD. B2
		Stem	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. PPD. B4
	4.Banyumas, Kembaran 7.23'47" LS, 109.17'9" BT, TT 110 m dpl	Root	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> BK. A1
		Root	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BK.A1
		Root	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BK.A3
		Stem	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BK.B3



Figure 1. Antagonism test between the endophytic bacteria against *R. solani* (a) and *Pantoea* sp..

Table 2. Inhibition of endophytic bacteria against *R. solani*

No	Isolate	Inhibition rate (%)	Dry weights Mycelium
1	Control	0	0,093
Endophytic bacteria from the root			
2	<i>Pseudomonas</i> Pf BB.A2	49,00	0,038
3	<i>Pseudomonas</i> Pf BS.A 2	45,00	0,027
4	<i>Pseudomonas</i> Pf BK.A1	51,00	0,017
5	<i>Pseudomonas</i> Pf PPD.A1	10,33	0,059
6	<i>Pseudomonas</i> Pf PP.A1	38,33	0,017
7	<i>Pseudomonas</i> Pf PB.A4	18,00	0,037
8	<i>Bacillus</i> sp.BB.A3	40,42	0,030
9	<i>Bacillus</i> sp.BS.A1	48,73	0,016
10	<i>Bacillus</i> sp. BSA3	37,42	0,039
11	<i>Bacillus</i> sp. B.K.A1	55,39	0,002
12	<i>Bacillus</i> sp. B.K.A3	51,52	0,003
13	<i>Bacillus</i> sp.PP.A3	46,65	0,019
14	<i>Bacillus</i> sp.PP.A5	50,66	0,009
Endophytic bacteria from the stem			
15	<i>Pseudomonas</i> Pf PPD.B1	27,00	0,020
16	<i>Pseudomonas</i> Pf PPD.B5	49,33	0,013
17	<i>Pseudomonas</i> Pf PP.B4	65,67	0,004
18	<i>Bacillus</i> sp. BB.B4	44,44	0,026
19	<i>Bacillus</i> sp. BS.B1	49,74	0,012
20	<i>Bacillus</i> sp. BK. B3	40,36	0,031
21	<i>Bacillus</i> sp.PPD.B2	50,8	0,007
22	<i>Bacillus</i> sp. PPD.B4	39,44	0,036
23	<i>Bacillus</i> sp. PB.B1	37,29	0,047
24	<i>Bacillus</i> sp. PB.B3	44,9	0,022

The endophytic bacterial antagonism test against *Pantoea* sp.

The results of antagonism between the endophytic bacteria and *Pantoea* sp. show varied results. The endophytic bacteria that can inhibit bacterial growth are indicated by the presence of clear zones around the endophytic bacterial colonies (Fig.1). From the nine isolate *Pseudomonas* sp. were tested, only three isolates were able to inhibit the growth of the *Pantoea* sp., i.e Pf BS.A2, Pf BK.A1 Pf PPD.B5. While the isolates Pf BB.A2, Pf PPD.A1, Pf PP.A1, Pf PPD.B1, and Pf PP.B4 are not able to inhibit the growth of pathogenic bacteria. Meanwhile, all isolate *Bacillus* sp tested (thirteen isolates) were able to inhibit the growth of *Pantoea* sp. (Table 3).

The presence of clear zones around endophytic bacterial colonies shows the ability of endophytic bacteria to produce antibiotics to inhibit the growth of *Pantoea* sp. *P. fluorescens* P60 can produce antibiotics that inhibit the growth of pathogens (Soesanto 2011). *Pseudomonas fluorescens* is reported to produce phenazine-1-carboxylic acid (PCA) and other derivatives, 2,4 diacetyl phloroglucinol (DAPG), pyrrolnitrin (PRN) and or pyoluteorin (Plt) (Heydari and Pessarakli 2010). Nasrun and Burhanudin (2016) mention that *P. fluorescens* produce secondary metabolites, i.e. antimicrobial, cyanide acid and 2,4-diacetyl phloroglucinol phenazine, pyrrolnitrin, pyoluteorin antibiotics.

The level of bacteria's ability to inhibit growth can be shown by the large diameter of the clear zone. The results showed that the antagonism index ranged from 1.67 - 8.17. Based on this index, most endophytic bacteria have a strong antagonism (index of antagonism > 4) (Davis and Stout 1971). Furthermore, bacterial isolates that showed antagonistic activity were tested for types of antagonism based on Djatmiko (2007). Based on the type of antagonistic activity, ten isolates the endophytic bacteria were bacteriostatic and nine isolates the endophytic bacteria were bactericidal. Bacteriostatic activity, growth inhibition is temporary, it is shown that regrowth of bacteria after being transferred to a new medium, which is free from the influence of antagonistic bacteria. Bactericidal activity, inhibition is permanent. Bacteria were unable to grow even though they are transferred to new medium.

Table 3. Inhibition of endophytic bacteria against *Pantoea* sp.

No	Isolate	Antagonism	Antagonism index	Antagonism catagory*	Antagonism activity
Endophytic bacteria from the root					
1	<i>Pseudomonas</i> Pf BB.A2	-	0	-	-
2	<i>Pseudomonas</i> Pf BS.A 2	+	4,91	strong	bacteriostatic
3	<i>Pseudomonas</i> Pf BK.A1	+	4,42	strong	bacteriostatic
4	<i>Pseudomonas</i> Pf PPD.A1	-	0	-	-
5	<i>Pseudomonas</i> Pf PP.A1	-	0	-	-
6	<i>Pseudomonas</i> Pf PB.A4	+	5,29	strong	bactericidal
7	<i>Bacillus</i> sp.BB.A3	+	8,17	strong	bacteriostatic
8	<i>Bacillus</i> sp.BS.A1	+	4,00	strong	bacteriostatic
9	<i>Bacillus</i> sp. BSA3	+	5,07	strong	bactericidal
10	<i>Bacillus</i> sp. B.K.A1	+	4,01	strong	bakteriostatik
11	<i>Bacillus</i> sp. B.K.A3	+	4,91	strong	bacteriostatic
12	<i>Bacillus</i> sp.PP.A3	+	6,63	strong	bactericidal
13	<i>Bacillus</i> sp.PP.A5	+	6,56	strong	bactericidal
Endophytic bacteria from the stem					
14	<i>Pseudomonas</i> PPD.B1	-	0	-	-
15	<i>Pseudomonas</i> PPD.B5	+	5,86	strong	bactericidal
16	<i>Pseudomonas</i> PP.B4	-	0	-	-
17	<i>Bacillus</i> sp. BB.B4	+	7,80	strong	bactericidal
18	<i>Bacillus</i> sp. BS.B1	+	6,22	strong	bacteriostatic
19	<i>Bacillus</i> sp. BK. B3	+	5,33	strong	bacteriostatic
20	<i>Bacillus</i> sp.PPD.B2	+	5,00	strong	bacteriostatic
21	<i>Bacillus</i> sp. PPD.B4	+	8,75	strong	bacteriostatic
22	<i>Bacillus</i> sp. PB.B1	+	1,67	weak	bacteriostatic
23	<i>Bacillus</i> sp. PB.B3	+	5,67	strong	bactericidal

*Based on Davis and Stout, 1971

Test the mechanism of endophytic bacteria as biological control agents and plant growth-promoting microbes

The mechanism test is carried out on endophytic bacteria that have the potential to control the fungus *R. solani* and *Pantoea* sp., i.e. *Bacillus* sp. B.K.A1, *Bacillus* sp. B.K.A3, *Bacillus* sp. PP.A5, *Bacillus* sp. PPD.B2. The results of enzyme activity tests are as shown in Table 4. The production of compounds related to biocontrol of pathogens and/or promotion of plant growth in bacterial isolates was evaluated by measuring the production of antimicrobial compounds and hydrolytic enzymes (amylases, lipases, proteases, and chitinases) and phosphate solubilization. the isolates that have high protein and fat hydrolysis enzymes have the potential as biological control agents because proteins and fats are constituents of pathogen cells (Mota et al 2016).

The four isolates *Bacillus* sp. tested were able to produce protease, lipase and phosphatase enzymes, with varied activity indexes. All isolates of *Bacillus* sp. those tested had a high index of protease and lipase enzymes (> 3) (Table 4.,

Fig 2.). Protease and lipase enzymes, related to the ability of the endophytic bacteria as biological control agents. According to Anderson et al. (2014), the extracellular protease enzyme produced by *P. fluorescens* can inactivate antibiotic compounds produced by *Pantoea agglomerans*. The phosphate solubilization is related to the ability of endophytic bacteria as a plant growth promoter, providing phosphates for plants.

Table 4. Test results of proteases, lipases and phosphate solubilization.

No	Isolate	Protease Test activity	index	Lipase Test activity	index	Phosphate solubilization activity	index
Endophytic bacteria from the root							
1	<i>Bacillus</i> sp. B.K.A1	+	3.75	+	3.23	+	1.17
2	<i>Bacillus</i> sp. B.K.A3	+	3.20	+	3.73	+	1.27
3	<i>Bacillus</i> sp.PP.A5	+	5.00	+	4.40	+	1.46
Endophytic bacteria from the stem							
4	<i>Bacillus</i> sp.PPD.B2	+	3.00	+	3.90	+	2.60



Figure 2. Hydrolysis enzyme activity, (a) protease , (b) lipase and (c) phosphate solubilization.

The author should expand the discussion by looking at previous published studies and compare with current findings.

Formatted: English (South Africa)

CONCLUSION

Based on research carried out, it has been successfully isolated, morphologically and biochemically characterized four the endophytic bacteria that have the potential to be developed as biopesticides to control maize disease, especially *R. solani* and *Pantoea* sp. They can suppress the growth of *R.solani* by more than 50%, have a strong antagonistic index against *Pantoea* sp (> 4), and can produce protease and lipase enzyme, and phosphate solubilization.

ACKNOWLEDGEMENTS

This research was a part of the main study funded by Doctoral Dissertation Grant from the Indonesian Ministry of Research, Technology and Higher Education; for that, I deeply thank for the financial support

REFERENCES

Abed HN, Rouag, Mouatassem D, Rouabhi A. 2016. Screening for *Pseudomonas* and *Bacillus* antagonistic rhizobacteria strains for the biocontrol of Fusarium wilt of chickpea. Eurasian Journal of Soil Science 5(3):182–191

Abidin Z, Aini LQ, Abadi AL. 2015. Effect of *Bacillus* sp. and *Pseudomonas* sp. on the growth of the pathogenic fungus *Sclerotium rolfsii* Sacc. causes of seedling diseases in soybean plants. Jurnal HPT 3(1): 1-10.

Aini,LQ, Suryani L, Sugiharto AN,. Abadi A. 2013. Identification of bacterial wilt and leaf blight disease on maize (*Zea Mays*) found in Kediri, Indonesia. Agrivita 35 (1): 1-7.

Ammar, E, V.R. Correa, S.A. Hogenhout, and M.G. Redinbaugh. 2014. Immunofluorescence localization and ultrastructure of Stewart's wilt disease bacterium *Pantoea stewartii* in maize leaves and in its flea beetle vector *Chaetocnema pulicaria* (Coleoptera: Chrysomelidae). Journal of Microscopy and Ultrastructure 2: 28–33

Anderson LM, Stockwell VO, Loper JE. 2004. An extracellular protease of *Pseudomonas fluorescens* inactivates antibiotics of *Pantoea agglomerans*. Phytopathology 94:1228-1234

Arwiyanto T, Maryudani. YMS, Nurul N, Azizah. 2007. The phenotypic properties of *Pseudomonas fluorescens*, biological control agents for lincat disease in temanggung tobacco. Biodiversitas 8(2) : 147-151.

- Cavaglieri L, Orlando J, Etcheverry M. 2009. Rhizosphere microbial community structure at different maize plant growth stages and root locations. Microbiological
- Compant SB, Duffy, Nowak J, Clement C, Barka EA. 2005. Use of Plant Growth- Promoting Bacteria for Biocontrol of Plant Diseases: principles, Mechanisms of Action, and Future Prospects. Applied and Environmental Microbiology 71(9): 4951-4959.
- Coplin DL, Redinbaugh M G. 2012. The Bacterium *Pantoea stewartii* Uses Two Different Type III Secretion Systems To Colonize Its Plant Host and Insect Vector. Applied and Environmental Microbiology 78(17): 6327-6336.
- Costa FG, Zucchi TD, de Melo IS. 2013. Biological control of Phytopathogenic Fungi by Endophytic Actinomycetes Isolated from Maize (*Zea Mays* L.). Braz. Arch.Biol.Technol 56(6):948-955.
- Davis WW, Stout TR. 1971. Disc plate methods of microbiological antibiotic assay. Applied Microbiology 22(4):659-665.
- Djaenuddin N, Nonci N, Muis A. 2017. Effectiveness of the formula *Bacillus subtilis* TM4 for disease control in maize plants. Jurnal Fitopatologi Indonesia 13(4): 113-118
- Djatmiko HA, Arwiyanto T, Hadisutrisno B, Sunarminto BH. 2007. Potential of three bacterial genera from three plant rhizosphere as biological agents controlling linear disease. Jurnal ilmu-ilmu Pertanian 9(1):40-47.
- Djuric SA, Pavic, Jarak M, Pavlovic S, Starovic M, Pivic R, Josic D.. 2011. Selection of indigenous fluorescent pseudomonad isolates from maize rhizospheric soil in Vojvodina as possible PGPR. Romanian Biotechnological Letters 16(5): 6580-6590.DOI:10.1094/PHI-I-2004-0113-01.
- Farooq U, Bano A. 2013. Screening of indigenous bacteria from rhizosphere of maize (*Zea mays* L.) for their plant growth promotion ability and antagonism against fungal and bacterial pathogens. Journal of Animal and Plant Sciences, 23(6), pp.1642- 1652.
- Ganeshan G, Kumar AM. 2005. *Pseudomonas fluorescens*, a potential bacterial antagonist to control plant diseases. Journal of Plant Interactions 1(3): 123-134.
- Hastuti RD, Saraswati R, Sari AP. 2014. The effectiveness of the endophytic microbes in promoting plant growth and controlling leaf blight disease in the lowland rice. Jurnal Tanah dan Iklim 38(2) : 109-118.
- Heydari A, Pessarakhi M. 2010. A review on biological control of fungal plant pathogens using microbial antagonists. Journal of Biological Science 10(4): 273-290
- Motaa MS, Gomes CB, Júnior ITS, Moura AB. 2017. Bacterial selection for biological control of plant disease: criterion determination and validation. Brazilian Journal of Microbiology 48 : 62-70
- Muis A. 2007. Management of sheath blight disease (*Rhizoctonia solani* Kuhn.) in maize. Jurnal Litbang Pertanian 26(3) : 100-103
- Nasrun, Burhanudin. 2016. Evaluation of the efficacy of the formula *Pseudomonas fluorescens* for controlling bacterial wilt (*Ralstonia solanacearum*) patchouli. Buletin Penelitian Tanaman Rempah dan Obat, 27(1): 67-76.
- Nuryanto B, Priyatmojo A, Hadisutrisno B.. 2014. Pengaruh Tinggi Tempat dan Tipe Tanaman Padi terhadap Keparahan Penyakit Hawar Pelepah. Penelitian Pertanian Tanaman Pangan 33(1):1-8.
- Orole OO, Adejumo TO. 2011. Bacterial and fungal endophytes associated with grains and roots of maize. Journal of Ecology and the natural Environment 3(9):298-303.
- Pal KK, McSpadden Gardener B. 2006. Biological Control of Plant Pathogens.
- Pataky JK. 2004. Stewart's wilt of corn. The Plant Health Instructor.
- Rosenblueth M, Martinez-Romero E. 2006. Bacterial endophytes and their interaction with hosts (Review). MPMI 19 (8): 827-837.
- Santiago TR, Grabowski C, Rossato M, Romeiroa RS, Mizubuti ESG. 2015. Biological control of eucalyptus bacterial wilt with rhizobacteria. Biological Control 80:14-22. <http://dx.doi.org/10.1016/j.biocontrol.2014.09.007>.
- Shanti AT, Vittal RR. 2013. Biocontrol potentials of Plant Growth promoting Rhizobacteria Against Fusarium Wilt Disease of Cucurbit. Esci J. Plant Pathol 2(3): 155-161.
- Soesanto L, Mugiastuti E, Rahayuniati RF. 2010. Study of the antagonistic mechanism of *Pseudomonas fluorescens* P60 against *Fusarium oxysporum* f.sp. *lycopersici* in tomatoes in vivo. Jurnal HPT Tropika 10(2) : 108-115
- Soesanto L, Mugiastuti E, Rahayuniati RF. 2011. Utilization of some animal broths as a liquid formula for *Pseudomonas fluorescens* P60 to control *Sclerotium rolfsii* in cucumber plants. Jurnal Perlingungan Tanaman Indonesia, 17(1): 7-17.

1
2
3

4
5
6
7

Short Communication:
Isolation and characterization of the endophytic bacteria, and their potential as maize diseases control

Abstract. ~~The present was aimed to isolate and characterize the endophytic bacteria morphologically and biochemically and to study~~~~The research aimed to isolate and characterize morphologically and biochemically the endophytic bacteria, and their potential to control~~ maize diseases, especially sheat blight and bacterial wilt ~~causing pathogens~~. The study was conducted at the Plant Protection Laboratory, Faculty of Agriculture, Jenderal Soedirman University, from April to August 2019. The study consisted of four stages: isolation and characterization of endophytic bacteria, the antagonism test of the endophytic bacterial to *R. solani*, the antagonism test of the endophytic bacteria to *Pantoea* sp., and the mechanism test of the endophytic bacteria as biological control agents and plant growth-promoting bacteria. Based on the research, ~~four endophytic bacteria isolates~~ it has been successfully isolated, and characterized ~~morphologically and biochemically characterized four endophytic bacteria isolates~~ successfully and found that have the potential to be developed as biopesticides to control maize disease, especially *R. solani* and *Pantoea* sp. *Bacillus* sp. endophytic from the root (BK.A1; BK.A3; PP.A5) and *Bacillus* sp. endophytic from the stem (PPD.B2) can suppress the growth of *R. solani* by more than 50%, have a strong antagonistic index against *Pantoea* sp (> 4), and can produce protease and lipase enzyme, and phosphate solubilization.

Key words: *Bacillus* sp., Fluorescents *Pseudomonads*, *Pantoea* sp, *Rhizoctoni solani*

Running title: Isolation and characterization of the endophytic

INTRODUCTION

Maize is a strategic food commodity in the world. In Indonesia, the government seeks to achieve self-sufficiency in maize through increasing production of sustainable maize. However, these efforts have faced several ~~obstacles.obstacles~~; one of them is the presence of plant diseases such as ~~sheat~~~~sheath~~ blight–caused by *Rhizoctonia solani* Kuhn, and bacterial wilt caused by (*Pantoea* sp.). *R. solani* can infect up to the midrib of the cob (Djaenuddin et al. 2017), resulting in ~~up to 100% a decrease in the yield of up to 100%.~~ (Muis 2007). *Pantoea* sp. can attack all stages of the plant causing wilting and leaf blight, and is known as Stewart's wilt (Pataky 2004; Ammar et al. 2014). The pathogens can cause 40-100% yield loss.

Over the past 3 decades, the ~~concept of sustainable and environmentally friendly agriculture have~~~~concept of sustainable and environmentally friendly agriculture has~~ been carried out by minimizing the use of chemicals, both synthetic fertilizers and ~~synthetic~~–pesticides. In the management of pests and plant diseases, biological control is developed by applying biological ~~control~~–agents including the endophytic bacteria (Shanti and Vittal 2013). Many endophytic bacteria can pass the endodermic barrier across from the root cortex to the vascular system, and subsequently develop as endophytes in stems, leaves, tubers, and other organs (Compant et al. 2005). The use of endophytic bacteria as biological agents has an advantage compared to rhizosphere bacteria because endophytic bacteria live and survive in the plant tissue during plant development, thus protecting the plants.

Bacillus sp. and fluorescent *Pseudomonads* are reported to be able to live as endophytes and are widely used as biological control agents for soil-borne and air-borne diseases. The endophytic bacteria could control plant diseases through several mechanisms including competition, hyperparasit~~isme~~, producing microbial inhibiting compounds (antibiotics, lysis enzymes, other physical or chemical disorders), enhancing plant resistance, and promoting plant growth (Compant et al 2005, Pal and McSpadden 2006; Rosenblueth and Martinez Romero 2006:).

Based on the mechanisms, the use of endophytic bacteria isolated from maize, both upland and lowland, suggested potentially alternative control for sheath blight (*R. solani*) and bacterial wilt (*Pantoea* sp). The research aimed to isolate and characterize morphologically and biochemically the endophytic bacterial as well as their potential to control pathogens that cause disease in maize especially *R. solani* and *Pantoea* sp.

Commented [A1]: Add 1-2 line of introduction.

Commented [A2]: Check spelling? If genus, it should be in italics.

Commented [A3]: I think authors should write full name of pathogen as is available in literature. If possible, replace in whole manuscript.

Commented [A4]: Check spelling?

MATERIALS AND METHODS

This research was conducted at the Laboratory of Plant Protection, Faculty of Agriculture, Jenderal Soedirman University, Purwokerto, Central Java, Indonesia, from April to August 2019

Isolation *R. solani*

R. solani was isolated from maize with sheath blight symptoms and there was sclerotium as a resistant structure from the pathogenic fungi in Banyumas. Samples were isolated on PDA medium to obtain pure *R. solani* isolates.

Isolation *Pantoea* sp.

Pantoea sp. was isolated from diseased maize, which was samples taken from the maize growing area in Banyumas Regency. *Pantoea* sp. was isolated according to Coplin et al. (2012); Aini et al. (2013) and Desi et al. (2014). Diseased leaves or stems were washed with running water, then dried with a tissue. Diseased samples were cut 1.5 x 5 cm, then sterilized with ethanol 70% and rinsed with sterile water 3 times. Furthermore, the sample was crushed with 5 ml of sterile distilled water using a sterile mortar. The bacterial suspension was streaked on nutrient agar and incubated 3-5 days. Bacterial colonies that exhibit the character of *Pantoea* sp. are were yellow, shiny, slimy, flat or convex, then separated as pure cultures of *stewartii* of *stewartii* candidates. The culture was then tested by Gram Reaction (KOH test), Hugh-Leifson test, pigment production in YDC medium, oxidase test, hypersensitivity test, and pathogenicity test on maize.

Isolation and characterization of endophytic bacteria

Sampling for the isolation of endophytic bacteria was carried out in Banyumas and Purbalingga, Central Java, Indonesia, with purposive stratified random sampling. Samples were taken from two areas of altitude, i.e., low-moderate lands (0-500 m above sea level), and highlands (> 500 m above sea level) (Nuryanto et al., 2014). In each district, 2 locations were selected for the low-medium lands, and 1 location for the highlands. Age of maize plants was 20-30 days after planting, when the number of endophytic microbial populations that can be cultured is in the highest population (Cavaglieri et al. 2009).

The endophytic bacteria are were isolated from the roots and stems of healthy maize plants. Roots and stems are were washed, sterilized with 70% ethanol (1 minute), 20% natrium hypochlorite (5 minutes) and Ringer's thiosulfate solution (5 minutes). Separately, the roots and stems of 10 g each were crushed with 90 ml PBS on a sterile mortar. Subsequently, samples were plated on NA and Kings B media (Cavaglieri et al. 2009). To isolate *Bacillus* sp., the suspension is was heated for 10 minutes at 80 ° C, before plating on NA. Bacterial isolates were further purified and characterized, such as morphological characteristics, gram properties, catalase tests, and hypersensitivity tests

The antagonism test of endophytic bacterial to *R solani*

The antagonism test of endophytic bacteria on *R. solani* was carried out using the dual culture method. The level of inhibition of antagonist is calculated using the formula (Abidin et al., 2015).

$$I = \frac{C - T}{C} \times 100\%$$

I = The level of inhibition of antagonist (%)

C = The radius of pathogen colonies opposite antagonist

T = The radius of the colony of pathogens towards antagonist

The antagonism test of endophytic bacterial to bacterial pathogens

Antagonism testing was carried out using the double-layer test method (Santiago et al. 2015). Endophytic bacteria to be tested were grown on the NA medium, incubated at 28 ° C for 48 hours. In the upside-down position, 1 ml of chloroform was added to the cup lid and left for 2 hours. Next, add 5 mL so that 0.6% water containing 0.5 mL of *P. stewartii* bacterial suspension. The culture was re-incubated for 24 hours, and there were clear zones around the antagonistic bacterial colony. The antibiotic activity was assessed based on the diameter of the clear zone compared to the diameter of the colony. Characterization of the type of antibiosis can be divided into bactericidal and bacteriostatic types according to the method of Djatmiko et al. (2007).

The mechanism test of endophytic bacteria as controlling agents biological and plant growth-promoting microbial

The testing mechanism of endophytic bacteria is was carried out for bacteria that have the potential in testing the antagonism of the fungus *R. solani* and *Pantoea* sp.

Protease Test

The activity of the ability of antagonistic bacteria to produce extracellular protease enzymes was tested using Skim Milk Agar (SMA) medium. Each bacterium to be tested was grown in a medium SMA and incubated at 28 ° C for 24-48 hours. The presence of clear zones around the colony shows that positive bacteria produce protease enzymes (Abed et al.

Commented [A5]: Authors should write detailed protocol of isolation of *R. solani*.

Commented [A6]: Shocking towards authors, as heading is endophytic bacteria's antagonism to bacterial pathogens but mentioned here to *R. solani* too, which is fungi. Please check carefully.

Commented [A7]: Full abbreviation.

Commented [A8]: What this denote?

Formatted: Font: Italic

2016). The protease activity index is assessed based on the diameter of the clear zone compared to the diameter of the colony..

$$\text{Protease index} = \frac{(\text{clear zone diameter} - \text{colony diameter})}{\text{colony diameter}}$$

Lipase test

Detection of the ability of bacteria to produce the enzyme lipase was done by growing the antagonistic bacteria on a medium containing 1% Tween 80. The presence of lipase enzyme activity was demonstrated by milky white sediment around the bacterial colony, after incubating at 28 C for 4-7 days. The lipolytic index was measured using a formula Djuric et al. (201).

$$\text{Lipolytic index} = \frac{(\text{milky white diameter} - \text{colony diameter})}{\text{colony diameter}}$$

Uji fosfatase

Detection of the ability of bacteria to produce the enzyme phosphatase was done by growing bacterial isolates on Pikovkaya medium. After incubating for 7 days at 28 C, the presence of a clear zone around the bacterial colony shows that the bacteria has the ability to produce the phosphatase enzyme to dissolve phosphates. The solubility index is measured using a formula (Farooq and Bano 2013)

$$\text{Phosphatase Index} = \frac{\text{clear zone diameter} - \text{colony diameter}}{\text{colony diameter}}$$

RESULTS AND DISCUSSION

Isolation and characterization of endophytic bacteria

The results of the exploration, isolation, and characterization of endophytic bacteria obtained 23 isolates of endophytic bacteria, consisted of 9 isolates of the fluorescent *Pseudomonads* and 14 isolates of *Bacillus* sp. (Table 1). *Fluorescent Pseudomonads* colony on King's B ~~is was~~ round, with a flat edge, and yellowish-white, to greenish-yellow, gram-negative, rod-shaped, non-spore and fluorescent under ultraviolet light. According to Arwiyanto et al. (2007), *P. fluorescens* have round, flat-edged, fluidal and release greenish-yellow colony in the King's B. Individual rod-shaped bacteria with a size (0.5-1.0) - (1.5-4.0) µm. The *P. fluorescens* isolates is gram-negative, which can form catalase, a positive oxidase, needed to grow aerobes.

Bacillus sp. ~~has was observed with a~~ spherical colony, ~~having~~ cell rod-shaped, gram-positive, and endospores within cells (Table 1.). According to Slepecky and Hempill 2006; Amin et al. 2015, *Bacillus* sp. has the characteristics of a circular colony and punctiform (small round), variations in the entire margin and lobate, white dull, non-slimy, gram-positive, has endospores, flagellum and some are motile.

Based on its distribution, *fluorescent Pseudomonads* and *Bacillus* sp. found in all sampling locations, high or low-medium lands. This shows that *fluorescent Pseudomonads* and *Bacillus* sp. spread and can live in various altitudes, both high and low-medium land. According to Bacon and Hilton 2002 and Ganeshan and Kumar 2005, *P. fluoresscens* and *Bacillus* sp., are species of bacteria with a wide range of life and are very adaptive in various environments. Both types of bacteria are also found in the roots or corn stalks. According to Ganeshan and Kumar 2005; Orole and Adejumo 2011; Costa et al. 2013, *Bacillus* sp., and *Pseudomonas* sp. including a group of endophytic bacteria have a wide range of life and more isolated in maize.

Antagonism test between the endophytic bacteria against *R. solani*

Based on the results of *in vitro* tests (Table 2), 24 isolates of the endophytic bacteria were able to inhibit the growth of *R. solani*, with varying degrees of inhibition. The endophytic bacteria that have inhibition rates above 50%, i.e *Pseudomonas* Pf BK.A1 (51%), *Bacillus* sp. B.K.A1 (55.39%), *Bacillus* sp BK.A3 (51.52%), PP.A5 (50.66%), and PPD.B2 (50.8%). The effect of the endophytic bacteria in inhibiting the growth of *R. solani* is inversely proportional to the dry weight mycelium. The greater the percentage of inhibition of endophytic bacteria to the growth of *R. solani*, the smaller the dry weight mycelium (Table 2.)

The endophytic bacteria can inhibit the growth of *R. solani* shown by the inhibition zone around the bacterial colony (Fig. 1). The endophytic bacteria have anti-pathogenic properties and can produce antibiotic compounds. The ability of the endophytic bacteria to control plant pathogens occurs through the mechanism of antibiosis, competition, lysis, inducing resistance and producing growth substances. Bacteria capable of producing secondary metabolites that can inhibit growth or damage pathogens (Hastuti et al. 2014). These compounds, including alkaloids, phenols, flavonoids, glycosides, and phytoalexin (Soesanto et al. 2010).

Formatted: Font: Not Italic

Commented [A9]: Why to compare and discuss with *P. fluorescens*, when test bacteria was *Pseudomonads*?

Commented [A10]: References are not as per international referencing style. Please check?

Commented [A11]: Why it is in italics every time? If a genera, should be started with capital letter. Check in whole ms.

Commented [A12]: if authors are doing characterization, then why characterize *Bacillus* only up to genus level?

Commented [A13]: Wrong citation style.

Commented [A14]:

Commented [A15R14]: ?

Formatted: Font: Italic

Commented [A16]:

Commented [A17]: it is fungi not bacteria.

150
151

Table 1. Isolation and characterization of endophytic bacteria

Land	Sampling location	sample	Gram test	Catalase test	oxidase test	Colony morphology	colony pigment	Fluorescence on KB Medium	Cell morphology	Endospores	Isolat
Highland	1. Purbalingga, Pratin LS, 109.17'21" BT, TT 1.190 m dpl	Root	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PP.A1
		Root	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. PP.A3
		Root	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. PP.A5
		Stem	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PP. B4
Medium-Lowland	2.Banyumas, Baturaden 7.19"1" LS, 109.14'29" BT, TT 520 m dpl	Root	-	+	+	round	Greenish yellow	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> BB.A2
		Root	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. BB.A3
		Stem	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BB.B4
		Stem	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BB.B4
	1.Banyumas, Sumbang7.21'54" LS, 109.17'33"BT, TT 200 m dpl	Root	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> BS.A2
		Root	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BS.A1
		Root	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. BS.A3
		Stem	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. BS. B1
	2.Purbalingga, Bojongsari, 7.20'12" LS, 109.20'22" BT, TT 190 m dpl	Root	-	+	+	round	Greenish yellow	+	Small rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PB. A 4
		Stem	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. PB. B1
		Stem	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BB. B3
		Root	-	+	+	round	Greenish yellow	+	Small rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PPD A1
	3.Purbalingga, Padamara, 7.22'28" LS, 109.13'24" BT, TT 180 m dpl	Stem	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PPD. B1
		Stem	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> PPD. B5
		Stem	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. PPD. B2
		Stem	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. PPD. B4
	4.Banyumas, Kembaran 7.23'47" LS, 109.17'9" BT, TT 110 m dpl	Root	-	+	+	round	yellowish white	+	Medium rod	-	<i>Fluorescent Pseudomonads (Pf)</i> BK. A1
		Root	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BK.A1
		Root	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BK.A3
		Stem	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BK.B3

Commented [A18]: Check rules of binomial system of nomenclature???



Figure 1. Antagonism test between the endophytic bacteria against *R. solani* (a) and *Pantoea* sp..

Table 2. Inhibition of endophytic bacteria against *R. solani*

No	Isolate	Inhibition rate (%)	Dry weights Mycelium
1	Control	0	0,093
Endophytic bacteria from the root			
2	<i>Pseudomonas</i> Pf BB.A2	49,00	0,038
3	<i>Pseudomonas</i> Pf BS.A 2	45,00	0,027
4	<i>Pseudomonas</i> Pf BK.A1	51,00	0,017
5	<i>Pseudomonas</i> Pf PPD.A1	10,33	0,059
6	<i>Pseudomonas</i> Pf PP.A1	38,33	0,017
7	<i>Pseudomonas</i> Pf PB.A4	18,00	0,037
8	<i>Bacillus</i> sp.BB.A3	40,42	0,030
9	<i>Bacillus</i> sp.BS.A1	48,73	0,016
10	<i>Bacillus</i> sp. BSA3	37,42	0,039
11	<i>Bacillus</i> sp. B.K.A1	55,39	0,002
12	<i>Bacillus</i> sp. B.K.A3	51,52	0,003
13	<i>Bacillus</i> sp.PP.A3	46,65	0,019
14	<i>Bacillus</i> sp.PP.A5	50,66	0,009
Endophytic bacteria from the stem			
15	<i>Pseudomonas</i> Pf PPD.B1	27,00	0,020
16	<i>Pseudomonas</i> Pf PPD.B5	49,33	0,013
17	<i>Pseudomonas</i> Pf PP.B4	65,67	0,004
18	<i>Bacillus</i> sp. BB.B4	44,44	0,026
19	<i>Bacillus</i> sp. BS.B1	49,74	0,012
20	<i>Bacillus</i> sp. BK. B3	40,36	0,031
21	<i>Bacillus</i> sp.PPD.B2	50,8	0,007
22	<i>Bacillus</i> sp. PPD.B4	39,44	0,036
23	<i>Bacillus</i> sp. PB.B1	37,29	0,047
24	<i>Bacillus</i> sp. PB.B3	44,9	0,022

The endophytic bacterial antagonism test against *Pantoea* sp.

The results of antagonism between the endophytic bacteria and *Pantoea* sp. show varied results. The endophytic bacteria that can inhibit bacterial growth ~~are were~~ indicated by the presence of clear zones around the endophytic bacterial colonies (Fig.1). From the nine isolate *Pseudomonas* sp. were tested, only three isolates were able to inhibit the growth of the *Pantoea* sp., i.e Pf BS.A2, Pf BK.A1 Pf PPD.B5. While the isolates Pf BB.A2, Pf PPD.A1, Pf PP.A1, Pf PPD.B1, and Pf PP.B4 are not able to inhibit the growth of pathogenic bacteria. Meanwhile, all isolate *Bacillus* sp. tested (thirteen isolates) were able to inhibit the growth of *Pantoea* sp. (Table 3).

Commented [A19]:

The presence of clear zones around endophytic bacterial colonies ~~shows~~ showed the ability of endophytic bacteria to produce antibiotics to inhibit the growth of *Pantoea* sp. *P. fluorescens* P60 can produce antibiotics that inhibit the growth of pathogens (Soesanto 2011). *Pseudomonas fluorescens* is reported to produce phenazine-1-carboxylic acid (PCA) and other derivatives, 2,4 diacetyl phloroglucinol (DAPG), pyrrolnitrin (PRN) and or pyoluteorin (Plt) (Heydari and Pessarakli 2010). Nasrun and Burhanudin (2016) mention that *P. fluorescens* produce secondary metabolites, i.e. antimicrobial, cyanide acid and 2,4-diacetyl phloroglucinol phenazine, pyrrolnitrin, pyoluteorin antibiotics.

The level of bacteria's ability to inhibit growth can be shown by the large diameter of the clear zone. The results showed that the antagonism index ranged from 1.67 - 8.17. Based on this index, most endophytic bacteria have a strong antagonism (index of antagonism > 4) (Davis and Stout 1971). Furthermore, bacterial isolates that showed antagonistic activity were tested for types of antagonism based on Djatmiko (2007). Based on the type of antagonistic activity, ten isolates the endophytic bacteria were bacteriostatic and nine isolates the endophytic bacteria were bactericidal. Bacteriostatic activity, growth inhibition is temporary, it is shown that regrowth of bacteria after being transferred to a new medium, which is free from the influence of antagonistic bacteria. Bactericidal activity, inhibition is permanent. Bacteria were unable to grow even though they are transferred to new medium.

Table 3. Inhibition of endophytic bacteria against *Pantoea* sp.

No	Isolate	Antagonism	Antagonism index	Antagonism category*	Antagonism activity
Endophytic bacteria from the root					
1	<i>Pseudomonas</i> Pf BB.A2	-	0	-	-
2	<i>Pseudomonas</i> Pf BS.A 2	+	4,91	strong	bacteriostatic
3	<i>Pseudomonas</i> Pf BK.A1	+	4,42	strong	bacteriostatic
4	<i>Pseudomonas</i> Pf PPD.A1	-	0	-	-
5	<i>Pseudomonas</i> Pf PP.A1	-	0	-	-
6	<i>Pseudomonas</i> Pf PB.A4	+	5,29	strong	bactericidal
7	<i>Bacillus</i> sp.BB.A3	+	8,17	strong	bacteriostatic
8	<i>Bacillus</i> sp.BS.A1	+	4,00	strong	bacteriostatic
9	<i>Bacillus</i> sp. BSA3	+	5,07	strong	bactericidal
10	<i>Bacillus</i> sp. B.K.A1	+	4,01	strong	bakteriostatik
11	<i>Bacillus</i> sp. B.K.A3	+	4,91	strong	bacteriostatic
12	<i>Bacillus</i> sp.PP.A3	+	6,63	strong	bactericidal
13	<i>Bacillus</i> sp.PP.A5	+	6,56	strong	bactericidal
Endophytic bacteria from the stem					
14	<i>Pseudomonas</i> PPD.B1	-	0	-	-
15	<i>Pseudomonas</i> PPD.B5	+	5,86	strong	bactericidal
16	<i>Pseudomonas</i> PP.B4	-	0	-	-
17	<i>Bacillus</i> sp. BB.B4	+	7,80	strong	bactericidal
18	<i>Bacillus</i> sp. BS.B1	+	6,22	strong	bacteriostatic
19	<i>Bacillus</i> sp. BK. B3	+	5,33	strong	bacteriostatic
20	<i>Bacillus</i> sp.PPD.B2	+	5,00	strong	bacteriostatic
21	<i>Bacillus</i> sp. PPD.B4	+	8,75	strong	bacteriostatic
22	<i>Bacillus</i> sp. PB.B1	+	1,67	weak	bacteriostatic
23	<i>Bacillus</i> sp. PB.B3	+	5,67	strong	bactericidal

*Based on Davis and Stout, 1971

Test the mechanism of endophytic bacteria as biological control agents and plant growth-promoting microbes

The mechanism test ~~is was~~ carried out on endophytic bacteria that have the potential to control the fungus *R. solani* and *Pantoea* sp., i.e. *Bacillus* sp. B.K.A1, *Bacillus* sp. B.K.A3, *Bacillus* sp. PP.A5, *Bacillus* sp. PPD.B2. The production of compounds related to biocontrol of pathogens and promotion of plant growth in bacterial isolates was evaluated by measuring the production of antimicrobial compounds and hydrolytic enzymes (amylases, lipases, proteases, and chitinases) and phosphate solubilization. The results of enzyme activity tests are as shown in Table 4.

190 **Table 4.** Test results of proteases, lipases and phosphate solubilization.

No	Isolate	Protease Test activity	index	Lipase Test activity	index	Phosphate solubilization Activity	index
Endophytic bacteria from the root							
1	<i>Bacillus</i> sp. B.K.A1	+	3.75	+	3.23	+	1.17
2	<i>Bacillus</i> sp. B.K.A3	+	3.20	+	3.73	+	1.27
3	<i>Bacillus</i> sp. PP.A5	+	5.00	+	4.40	+	1.46
Endophytic bacteria from the stem							
4	<i>Bacillus</i> sp.PPD.B2	+	3.00	+	3.90	+	2.60

191 The four isolates *Bacillus* sp. tested were able to produce protease, lipase and phosphatase enzymes, with varied
 192 activity indexes. All isolates of *Bacillus* sp. those tested had a high index of protease and lipase enzymes (> 3) (Table 4.,
 193 Fig 2.). Protease and lipase enzymes, related to the ability of the endophytic bacteria as biological control agents. Based on
 194 the protease and lipase indexes, *Bacillus* sp. PP.A5 can produce the highest proteases and lipase enzymes compared to
 195 other isolates. The isolates that have high protein and fat hydrolysis enzymes have the potential as biological control
 196 agents because proteins and fats are constituents of pathogen cells (Mota et al 2016). Besides, the protease enzyme is
 197 thought to degrade antibiotics produced by fungal or bacterial pathogens. According to Anderson et al. (2014), the
 198 extracellular protease enzyme produced by *P. fluorescens* can inactivate antibiotic compounds produced by *Pantoea*
 199 *agglomerans*.

200 *Bacillus* sp. PPD.B2 has the highest phosphate solubility index. The phosphate solubilization is related to the ability of
 201 endophytic bacteria as a plant growth promoter, providing phosphates for plants. Microbes with high phosphate solubility
 202 activity are capable of producing and releasing metabolites such as organic acids that chelate cations that are bound to
 203 phosphate (especially calcium) and converting them into soluble forms. Solubilization of different forms of phosphate by
 204 microbes associated with plants, and increasing its availability for plants, will increase growth and production of the plant
 205 (Djuric et al., 2011)



209 **Figure 2.** Hydrolysis enzyme activity, (a) protease , (b) lipase and (c) phosphate solubilization.

212 CONCLUSION

213 Based on research carried out, it has been successfully isolated, morphologically and biochemically characterized four
 214 the endophytic bacteria that have the potential to be developed as biopesticides to control maize disease, especially *R.*
 215 *solani* and *Pantoea* sp. They can suppress the growth of *R.solani* by more than 50%, have a strong antagonistic index
 216 against *Pantoea* sp (> 4), and can produce protease and lipase enzyme, and phosphate solubilization.

217 ACKNOWLEDGEMENTS

218 This research was a part of the main study funded by Doctoral Dissertation Grant from the Indonesian Ministry of
 219 Research, Technology and Higher Education; for that, I deeply thank for the financial support

220 REFERENCES

221 Abed HN, Rouag.,Mouatasssem D, Rouabhi A. 2016. Screening for *Pseudomonas* and *Bacillus* antagonistic rhizobacteria strains for the biocontrol of
 222 Fusarium wilt of chickpea. Eurasian Journal of Soil Science 5(3):182–191

Abidin Z, Aini LQ, Abadi AL. 2015. Effect of *Bacillus* sp. and *Pseudomonas* sp. on the growth of the pathogenic fungus *Sclerotium rolfsii* Sacc. causes of seedling diseases in soybean plants. Jurnal HPT 3(1): 1-10.

Aini, LQ, Suryani L, Sugiharto AN., Abadi A. 2013. Identification of bacterial wilt and leaf blight disease on maize (*Zea Mays*) found in Kediri, Indonesia. Agrivita 35 (1): 1-7.

Amin M, Rakhisi Z, Ahmady AZ. 2015. Isolation and Identification of Bacillus Species From Soil and Evaluation of Their Antibacterial Properties. Avicenna J Clin Microb Infec. 2(1): e23233

Ammar, E, V.R. Correa, S.A. Hogenhout, and M.G. Redinbaugh. 2014. Immunofluorescence localization and ultrastructure of Stewart's wilt disease bacterium *Pantoea stewartii* in maize leaves and in its flea beetle vector *Chaetocnema pulicaria* (Coleoptera: Chrysomelidae). Journal of Microscopy and Ultrastructure 2: 28–33

Anderson LM, Stockwell VO, Loper JE. 2004. An extracellular protease of *Pseudomonas fluorescens* inactivates antibiotics of *Pantoea agglomerans*. Phytopathology 94:1228-1234

Arwiyanto T, Maryudani. YMS, Nurul N, Azizah. 2007. The phenotypic properties of *Pseudomonas fluorescens*, biological control agents for lincat disease in temanggung tobacco. Biodiversitas 8(2) : 147-151.

Cavaglieri L, Orlando J, Etcheverry M. 2009. Rhizosphere microbial community structure at different maize plant growth stages and root locations. Microbiological

Compant SB, Duffy, Nowak J, Clement C, Barka EA. 2005. Use of Plant Growth- Promoting Bacteria for Biocontrol of Plant Diseases: principles, Mechanisms of Action, and Future Prospects. Applied and Environmental Microbiology 71(9): 4951-4959.

Coplin DL, Redinbaugh M G. 2012. The Bacterium *Pantoea stewartii* Uses Two Different Type III Secretion Systems To Colonize Its Plant Host and Insect Vector. Applied and Environmental Microbiology 78(17): 6327-6336.

Costa FG, Zucchi TD, de Melo IS. 2013. Biological control of Phytopathogenic Fungi by Endophytic Actinomycetes Isolated from Maize (*Zea Mays* L.). Braz. Arch. Biol. Technol 56(6):948-955.

Davis WW, Stout TR. 1971. Disc plate methods of microbiological antibiotic assay. Applied Microbiology 22(4):659-665.

Djaenuddin N, Nonci N, Muis A. 2017. Effectiveness of the formula *Bacillus subtilis* TM4 for disease control in maize plants. Jurnal Fitopatologi Indonesia 13(4): 113-118

Djatmiko HA, Arwiyanto T, Hadisutrisno B, Sunarminto BH. 2007. Potential of three bacterial genera from three plant rhizosphere as biological agents controlling lincat disease. Jurnal ilmu-ilmu Pertanian 9(1):40-47.

Djuric SA, Pavic, Jarak M, Pavlovic S, Starovic M, Pivic R, Josic D.. 2011. Selection of indigenous fluorescent pseudomonad isolates from maize rhizospheric soil in Vojvodina as possible PGPR. Romanian Biotechnological Letters 16(5): 6580–6590. DOI:10.1094/PHI-I-2004-0113-01.

Farooq U, Bano A. 2013. Screening of indigenous bacteria from rhizosphere of maize (*Zea mays* L.) for their plant growth promotion ability and antagonism against fungal and bacterial pathogens. Journal of Animal and Plant Sciences, 23(6), pp.1642– 1652.

Ganeshan G, Kumar AM. 2005. *Pseudomonas fluorescens*, a potential bacterial antagonist to control plant diseases. Journal of Plant Interactions 1(3): 123-134.

Hastuti RD, Saraswati R, Sari AP. 2014. The effectiveness of the endophytic microbes in promoting plant growth and controlling leaf blight disease in the lowland rice. Jurnal Tanah dan Iklim 38(2) : 109-118.

Heydari A, Pessarakhi M. 2010. A review on biological control of fungal plant pathogens using microbial antagonists. Journal of Biological Science 10(4): 273-290

Motaa MS, Gomes CB, Júnior ITS, Moura AB. 2017. Bacterial selection for biological control of plant disease: criterion determination and validation. Brazilian Journal of Microbiology 48 : 62–70

Muis A. 2007. Mmanagement of sheath blight disease (*Rhizoctonia solani* Kuhn.) in maize. Jurnal Litbang Pertanian 26(3) : 100-103

Nasrun, Burhanudin. 2016. Evaluation of the efficacy of the formula *Pseudomonas fluorescens* for controlling bacterial wilt (*Ralstonia solanacearum*) patchouli. Buletin Penelitian Tanaman Rempah dan Obat, 27(1): 67-76.

Nuryanto B, Priyatmojo A, Hadisutrisno B.. 2014. Pengaruh Tinggi Tempat dan Tipe Tanaman Padi terhadap Keparahan Penyakit Hawar Pelepah. Penelitian Pertanian Tanaman Pangan 33(1):1–8.

Orole OO, Adejumo TO. 2011. Bacterial and fungal endophytes associated with grains and roots of maize. Journal of Ecology and the natural Environment 3(9):298-303.

Pal KK, McSpadden Gardener B. 2006. Biological Control of Plant Pathogens.

Pataky JK. 2004. Stewart's wilt of corn. The Plant Health Instructor.

Rosenblueth M, Martinez-Romero E. 2006. Bacterial endophytes and their interaction with hosts (Review). MPMI 19 (8): 827-837.

Santiago TR, Grabowski C, Rossato M, Romeiroa RS, Mizubuti ESG. 2015. Biological control of eucalyptus bacterial wilt with rhizobacteria. Biological Control 80:14–22. <http://dx.doi.org/10.1016/j.biocontrol.2014.09.007>.

Shanti AT, Vittal RR. 2013. Biocontrol potentials of Plant Growth promoting Rhizobacteria Against Fusarium Wilt Disease of Cucurbit. Esci J. Plant Pathol 2(3): 155-161.

Slepecky R.A, Hemphill H.E. 2006. The Genus Bacillus—Nonmedical. In: Dworkin M., Falkow S., Rosenberg E., Schleifer KH., Stackebrandt E. (eds) The Prokaryotes. Springer, New York, NY

Soesanto L, Mugiastuti E, Rahayuniati RF. 2010. Study of the antagonistic mechanism of *Pseudomonas fluorescens* P60 against *Fusarium oxysporum* f.sp. *lycopersici* in tomatoes in vivo. Jurnal HPT Tropika 10(2) : 108-115

Soesanto L, Mugiastuti E, Rahayuniati RF. 2011. Utilization of some animal broths as a liquid formula for *Pseudomonas fluorescens* P60 to control *Sclerotium rolfsii* in cucumber plants. Jurnal Perlindungan Tanaman Indonesia, 17(1): 7–17.

Short Communication:

Isolation and characterization of the endophytic bacteria, and their potential as maize diseases control

Abstract. Sheath blight and bacterial wilt are diseases that can reduce maize production. Biological control with the endophytic bacteria offers environmentally friendly control for these pathogens. The study was aimed to isolate and characterize the endophytic bacteria morphologically and biochemically and to study their potential to control maize diseases, especially sheath blight and bacterial wilt causing pathogens. The study was conducted at the Plant Protection Laboratory, Faculty of Agriculture, Jenderal Soedirman University, from April to August 2019. The study consisted of four stages: isolation and characterization of endophytic bacteria, the antagonism test of the endophytic bacteria to *Pantoea* sp., and the mechanism test of the endophytic bacteria as biological control agents and plant growth-promoting bacteria. Based on the research, four endophytic bacteria isolates have been successfully isolated, and characterized successfully and found have the potential to be developed as biopesticides to control maize disease, especially *R. solani* and *Pantoea* sp. *Bacillus* sp. endophytic from the root (BK.A1; BK.A3; PP.A5) and *Bacillus* sp. endophytic from the stem (PPD.B2) can suppress the growth of *R. solani* by more than 50%, have a strong antagonistic index against *Pantoea* sp. (> 4), and can produce protease and lipase enzyme, and phosphate solubilization.

Key words: *Bacillus* sp., fluorescent *Pseudomonas*, *Pantoea* sp., *Rhizoctonia solani*

Running title: Isolation and characterization of the endophytic

INTRODUCTION

Maize is a strategic food commodity in the world. In Indonesia, the government seeks to achieve self-sufficiency in maize through increasing production of sustainable maize. However, these efforts have faced several obstacles; one of them is the presence of plant diseases such as sheath blight caused by *Rhizoctonia solani* Kuhn, and bacterial wilt caused by (*Pantoea stewartii*). *R. solani* can infect up to the midrib of the cob (Djaenuddin et al. 2017), resulting in up to 100% decrease in the yield (Muis 2007). *Pantoea* sp. can attack all stages of the plant causing wilting and leaf blight, and is known as Stewart's wilt (Pataky 2004; Ammar et al. 2014). The pathogens can cause 40-100% yield loss.

Over the past 3 decades, the concept of sustainable and environmentally friendly agriculture has been carried out by minimizing the use of chemicals, both synthetic fertilizers and pesticides. In the management of pests and plant diseases, biological control is developed by applying biological agents including the endophytic bacteria (Shanti and Vittal 2013). Many endophytic bacteria can pass the endodermic barrier across from the root cortex to the vascular system, and subsequently develop as endophytes in stems, leaves, tubers, and other organs (Compant et al. 2005). The use of endophytic bacteria as biological agents has an advantage compared to rhizosphere bacteria because endophytic bacteria live and survive in the plant tissue during plant development, thus protecting the plants.

Bacillus sp. and fluorescent *Pseudomonas* are reported to be able to live as endophytes and are widely used as biological control agents for soil-borne and air-borne diseases. The endophytic bacteria could control plant diseases through several mechanisms including competition, hyperparasitism, producing microbial inhibiting compounds (antibiotics, lysis enzymes, other physical or chemical disorders), enhancing plant resistance, and promoting plant growth (Compant et al 2005, Pal and McSpadden 2006; Rosenblueth and Martinez Romero 2006).

Based on the mechanisms, the use of endophytic bacteria isolated from maize, both upland and lowland, suggested potentially alternative control for sheath blight (*R. solani*) and bacterial wilt (*Pantoea* sp). The research aimed to isolate and characterize morphologically and biochemically the endophytic bacteria as well as their potential to control pathogens that cause disease in maize especially *R. solani* and *Pantoea* sp.

MATERIALS AND METHODS

This research was conducted at the Laboratory of Plant Protection, Faculty of Agriculture, Jenderal Soedirman University, Purwokerto, Central Java, Indonesia, from April to August 2019

Isolation *R. solani*

R. solani was isolated from maize with sheath blight symptoms and there was sclerotium as a resistant structure from the pathogenic fungi in Banyumas. *R. solani* isolation was carried out based on Al-Fadhal et al. 2019. Disease samples were cut 0.5 x 0.5 cm, then sterilized with NaOCl (1%) for 2 min, and rinsed with sterile water 3 times. Disease samples pieces were then dried using sterile filter papers, and transferred to Petri dishes containing PDA medium to obtain pure *R. solani* isolates.

Isolation *Pantoea* sp.

Pantoea sp. was isolated from diseased maize samples taken from the maize growing area in Banyumas Regency according to Coplin et al. (2012); Aini et al. (2013) and Desi et al. (2014). Diseased leaves or stems were washed with running water, then dried with a tissue. Diseased samples were cut 1.5 x 5 cm, then sterilized with ethanol 70% and rinsed with sterile water 3 times. Furthermore, the sample was crushed with 5 ml of sterile distilled water using a sterile mortar. The bacterial suspension was streaked on nutrient agar and incubated 3-5 days. Bacterial colonies that exhibit the character of *Pantoea* sp. were yellow, shiny, slimy, flat or convex, then separated as pure cultures of *stewartii* candidates. The culture was then tested by Gram Reaction (KOH test), Hugh-Leifson test, pigment production in YDC medium, oxidase test, hypersensitivity test, and pathogenicity test on maize.

Isolation and characterization of endophytic bacteria

Sampling for the isolation of endophytic bacteria was carried out in Banyumas and Purbalingga, Central Java, Indonesia, with purposive stratified random sampling. Samples were taken from two areas of altitude, i.e., low-moderate lands (0-500 m above sea level), and highlands (> 500 m above sea level) (Nuryanto et al., 2014). In each district, 2 locations were selected for the low-medium lands, and 1 location for the highlands. Age of maize plants was 20-30 days after planting, when the number of endophytic microbial populations that can be cultured is in the highest population (Cavaglieri et al. 2009).

The endophytic bacteria were isolated from the roots and stems of healthy maize plants. Roots and stems were washed, sterilized with 70% ethanol (1 minute), 20% sodium hypochlorite (5 minutes) and Ringer's thiosulfate solution (5 minutes). Separately, the roots and stems of 10 g each were crushed with 90 ml PBS on a sterile mortar. Subsequently, samples were plated on NA and Kings B media (Cavaglieri et al. 2009). To isolate *Bacillus* sp., the suspension was heated for 10 minutes at 80 ° C, before plating on NA. Bacterial isolates were further purified and characterized, such as morphological characteristics, gram properties, catalase tests, and hypersensitivity tests

The antagonism test of endophytic bacterial to *R. solani*

The antagonism test of endophytic bacteria on *R. solani* was carried out using the dual culture method. The level of inhibition of antagonist is calculated using the formula (Abidin et al., 2015).

$$I = \frac{C - T}{C} \times 100\%$$

I = The level of inhibition of antagonist (%)

C = The radius of pathogen colonies opposite antagonist

T = The radius of the colony of pathogens towards antagonist

The antagonism test of endophytic bacterial to *Pantoea* sp.

Antagonism testing was carried out using the double-layer test method (Santiago et al. 2015). Endophytic bacteria to be tested were grown on the nutrient agar medium, incubated at 28 ° C for 48 hours. In the upside-down position, 1 ml of chloroform was added to the cup lid and left for 2 hours. Next, add 5 mL so that 0.6% water containing 0.5 mL of *Pantoea* sp. bacterial suspension. The culture was re-incubated for 24 hours, and there were clear zones around the antagonistic bacterial colony. The antibiotic activity was assessed based on the diameter of the clear zone compared to the diameter of the colony. Characterization of the type of antibiosis can be divided into bactericidal and bacteriostatic types according to the method of Djatmiko et al. (2007).

The mechanism test of endophytic bacteria as controlling agents biological and plant growth-promoting microbial

The testing mechanism of endophytic bacteria was carried out for bacteria that have the potential in testing the antagonism of the fungus *R. solani* and *Pantoea* sp.

Protease Test

The activity of the ability of antagonistic bacteria to produce extracellular protease enzymes was tested using Skim Milk Agar (SMA) medium. Each bacterium to be tested was grown in a medium SMA and incubated at 28 C for 24-48 hours. The presence of clear zones around the colony shows that positive bacteria produce protease enzymes (Abed et al. 2016). The protease activity index is assessed based on the diameter of the clear zone compared to the diameter of the colony..

$$\text{Protease index} = \frac{(\text{clear zone diameter} - \text{colony diameter})}{\text{colony diameter}}$$

Lipase test

Detection of the ability of bacteria to produce the enzyme lipase was done by growing the antagonistic bacteria on a medium containing 1% Tween 80. The presence of lipase enzyme activity was demonstrated by milky white sediment around the bacterial colony, after incubating at 28 C for 4-7 days. The lipolytic index was measured using a formula Djuric et al. (201).

$$\text{Lipolytic index} = \frac{(\text{milky white diameter} - \text{colony diameter})}{\text{colony diameter}}$$

Uji fosfatase

Detection of the ability of bacteria to produce the enzyme phosphatase was done by growing bacterial isolates on Pikovkaya medium. After incubating for 7 days at 28 C, the presence of a clear zone around the bacterial colony shows that the bacteria has the ability to produce the phosphatase enzyme to dissolve phosphates. The solubility index is measured using a formula (Farooq and Bano 2013)

$$\text{Phosphatase Index} = \frac{\text{clear zone diameter} - \text{colony diameter}}{\text{colony diameter}}$$

RESULTS AND DISCUSSION

Isolation and characterization of endophytic bacteria

The results of the exploration, isolation, and characterization of endophytic bacteria obtained 23 isolates of endophytic bacteria, consisted of 9 isolates of the fluorescent *Pseudomonas* and 14 isolates of *Bacillus* sp. (Table 1). *Fluorescent Pseudomonas* colony on King's B was round, with a flat edge, and yellowish-white, to greenish-yellow, gram-negative, rod-shaped, non-spore and fluorescent under ultraviolet light. Singh et al. 2017 reported fluorescent *Pseudomonas* showed light green, yellowish, creamy, circular, slimy, regular-irregular characteristics. Bacteria have short-long rod forms. The *Fluorescent Pseudomonas* isolates is gram-negative, which can form catalase, a positive oxidase, needed to grow aerobes.

Bacillus sp. was observed with spherical colony having cell rod-shaped, gram-positive, and endospores within cells (Table 1.). Slepecky and Hempill 2006; Amin et al. 2015 reported *Bacillus* sp. has the characteristics of a circular colony and punctiform (small round), variations in the entire margin and lobate, white dull, non-slimy, gram-positive, has endospores, flagellum and some are motile.

Based on its distribution, fluorescent *Pseudomonas* and *Bacillus* sp. found in all sampling locations, high or low-medium lands. This shows that fluorescent *Pseudomonas* and *Bacillus* sp. spread and can live in various altitudes, both high and low-medium land. Bacon and Hilton 2002; Ganeshan and Kumar 2005 reported *P. fluorescens* and *Bacillus* sp., are species of bacteria with a wide range of life and are very adaptive in various environments. Both types of bacteria are also found in the roots or corn stalks. *Bacillus* sp. and *Pseudomonas* sp. including a group of endophytic bacteria have a wide range of life and more isolated in maize (Ganeshan and Kumar 2005; Orole and Adejumo 2011; Costa et al. 2013)

Antagonism test between the endophytic bacteria against *R. solani*

Based on the results of *in vitro* tests (Table 2), 24 isolates of the endophytic bacteria were able to inhibit the growth of *R. solani*, with varying degrees of inhibition. The endophytic bacteria that have inhibition rates above 50%, i.e. fluorescent *Pseudomonas* BK.A1 (51%), *Bacillus* sp. B.K.A1 (55.39%), *Bacillus* sp BK.A3 (51.52%), PP.A5 (50.66%), and PPD.B2 (50.8%). The effect of the endophytic bacteria in inhibiting the growth of *R. solani* is inversely proportional to the dry weight mycelium. The greater the percentage of inhibition of endophytic bacteria to the growth of *R. solani*, the smaller the dry weight mycelium (Table 2.)

Endophytic bacteria can inhibit the growth of *R. solani*, which were shown by the inhibitory zone in the area bordering the bacterial streak (Fig. 1a). The endophytic bacteria have anti-pathogenic properties and can produce antibiotic compounds. The ability of the endophytic bacteria to control plant pathogens occurs through the mechanism of antibiosis, competition, lysis, inducing resistance and producing growth substances. Bacteria capable of producing secondary metabolites that can inhibit growth or damage pathogens (Hastuti et al. 2014). These compounds, including alkaloids, phenols, flavonoids, glycosides, and phytoalexin (Soesanto et al. 2010). *Fluorescent Pseudomonas* can produce various types of antibiotics including phenazine-1-carboxylic acid, pyocyanin, pyrrolnitrin, and pyoluteorin and

Table 1. Isolation and characterization of endophytic bacteria

Land	Sampling location	sample	Gram test	Catalase test	oxidase test	Colony morphology	colony pigment	Fluorescence on KB Medium	Cell morphology	Endospores	Isolat
Highland	1. Purbalingga, Pratin 7.13'33" LS, 109.17'21" BT, TT 1.190 m dpl	Root	-	+	+	round	yellowish white	+	Medium rod	-	fluorescent <i>Pseudomonas</i> PP.A1
		Root	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. PP.A3
		Root	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. PP.A5
		Stem	-	+	+	round	yellowish white	+	Medium rod	-	fluorescent <i>Pseudomonas</i> PP. B4
Medium-Lowland	2.Banyumas, Baturaden 7.19"1" LS, 109.14'29" BT, TT 520 m dpl	Root	-	+	+	round	Greenish yellow	+	Medium rod	-	fluorescent <i>Pseudomonas</i> BB.A2
		Root	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. BB.A3
		Stem	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BB.B4
		Stem	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BB.B4
	1.Banyumas, Sumbang 7.21'54" LS, 109.17'33"BT, TT 200 m dpl	Root	-	+	+	round	yellowish white	+	Medium rod	-	fluorescent <i>Pseudomonas</i> (Pf) BS.A2
		Root	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BS.A1
		Root	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. BS.A3
		Stem	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. BS. B1
	2.Purbalingga, Bojongsari, 7.20'12" LS, 109.20'22" BT, TT 190 m dpl	Root	-	+	+	round	Greenish yellow	+	Small rod	-	fluorescent <i>Pseudomonas</i> (Pf) PB. A4
		Stem	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. PB. B1
		Stem	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BB. B3
		Root	-	+	+	round	Greenish yellow	+	Small rod	-	fluorescent <i>Pseudomonas</i> (Pf) PPD A1
	3.Purbalingga, Padamara, 7.22'28" LS, 109.13'24" BT, TT 180 m dpl	Stem	-	+	+	round	yellowish white	+	Medium rod	-	fluorescent <i>Pseudomonas</i> (Pf) PPD. B1
		Stem	-	+	+	round	yellowish white	+	Medium rod	-	fluorescent <i>Pseudomonas</i> (Pf) PPD. B5
		Stem	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. PPD. B2
		Stem	+	+	+	round	white	-	Small rod	+	<i>Bacillus</i> sp. PPD. B4
	4.Banyumas, Kembaran 7.23'47" LS, 109.17'9" BT, TT 110 m dpl	Root	-	+	+	round	yellowish white	+	Medium rod	-	fluorescent <i>Pseudomonas</i> (Pf) BK. A1
		Root	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BK.A1
		Root	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BK.A3
		Stem	+	+	+	round	white	-	Medium rod	+	<i>Bacillus</i> sp. BK.B3

2,4-diacetylphloroglucinol (Phl). Phl is a phenolic metabolite with antibacterial and antifungal (Jain and Das 2016). *Bacillus* species can produce various kinds of volatile compounds and diffusible with strong inhibitory activity against plant pathogens (Lim et al. 2017).




Figure 1. Antagonism test between the endophytic bacteria against *R. solani* (a) and *Pantoea* sp (b)

Table 2. Inhibition of endophytic bacteria against *R. solani*

No	Isolate	Inhibition rate (%)	Dry weights mycelium
1	Control	0	0,093
Endophytic bacteria from the root			
2	fluorescent <i>Pseudomonas</i> BB.A2	49,00	0,038
3	fluorescent <i>Pseudomonas</i> BS.A 2	45,00	0,027
4	fluorescent <i>Pseudomonas</i> BK.A1	51,00	0,017
5	fluorescent <i>Pseudomonas</i> PPD.A1	10,33	0,059
6	fluorescent <i>Pseudomonas</i> PP.A1	38,33	0,017
7	fluorescent <i>Pseudomonas</i> PB.A4	18,00	0,037
8	<i>Bacillus</i> sp.BB.A3	40,42	0,030
9	<i>Bacillus</i> sp.BS.A1	48,73	0,016
10	<i>Bacillus</i> sp. BSA3	37,42	0,039
11	<i>Bacillus</i> sp. B.K.A1	55,39	0,002
12	<i>Bacillus</i> sp. B.K.A3	51,52	0,003
13	<i>Bacillus</i> sp.PP.A3	46,65	0,019
14	<i>Bacillus</i> sp.PP.A5	50,66	0,009
Endophytic bacteria from the stem			
15	fluorescent <i>Pseudomonas</i> PPD.B1	27,00	0,020
16	fluorescent <i>Pseudomonas</i> PPD.B5	49,33	0,013
17	fluorescent <i>Pseudomonas</i> PP.B4	65,67	0,004
18	<i>Bacillus</i> sp. BB.B4	44,44	0,026
19	<i>Bacillus</i> sp. BS.B1	49,74	0,012
20	<i>Bacillus</i> sp. BK. B3	40,36	0,031
21	<i>Bacillus</i> sp.PPD.B2	50,8	0,007
22	<i>Bacillus</i> sp. PPD.B4	39,44	0,036
23	<i>Bacillus</i> sp. PB.B1	37,29	0,047
24	<i>Bacillus</i> sp. PB.B3	44,9	0,022

The endophytic bacterial antagonism test against *Pantoea* sp.

The results of antagonism between the endophytic bacteria and *Pantoea* sp. show varied results. The endophytic bacteria that can inhibit bacterial growth were indicated by the presence of clear zones around the endophytic bacterial colonies (Fig.1). From the nine isolate fluorescent *Pseudomonas* were tested, only three isolates were able to inhibit the growth of the *Pantoea* sp., i.e Pf BS.A2, Pf BK.A1 Pf PPD.B5. While the isolates Pf BB.A2, Pf PPD.A1, Pf PP.A1, Pf

PPD.B1, and Pf PP.B4 are not able to inhibit the growth of pathogenic bacteria. Meanwhile, all isolate *Bacillus* sp. tested (thirteen isolates) were able to inhibit the growth of *Pantoea* sp. (Table 3).

The presence of clear zones around endophytic bacterial colonies showed the ability of endophytic bacteria to produce antibiotics to inhibit the growth of *Pantoea* sp. *P. fluorescens* P60 can produce antibiotics that inhibit the growth of pathogens (Soesanto 2011). *Pseudomonas fluorescens* is reported to produce phenazine-1-carboxylic acid (PCA) and other derivatives, 2,4 diacetyl phloroglucinol (DAPG), pyrrolnitrin (PRN) and or pyoluteorin (Plt) (Heydari and Pessarakli 2010). Nasrun and Burhanudin (2016) mention that *P. fluorescens* produce secondary metabolites, i.e. antimicrobial, cyanide acid and 2,4-diacetyl phloroglucinol phenazine, pyrrolnitrin, pyoluteorin antibiotics.

The level of bacteria's ability to inhibit growth can be shown by the large diameter of the clear zone. The results showed that the antagonism index ranged from 1.67 - 8.17. Based on this index, most endophytic bacteria have a strong antagonism (index of antagonism > 4) (Davis and Stout 1971). Furthermore, bacterial isolates that showed antagonistic activity were tested for types of antagonism based on Djatmiko (2007). Based on the type of antagonistic activity, ten isolates the endophytic bacteria were bacteriostatic and nine isolates the endophytic bacteria were bactericidal. Bacteriostatic activity, growth inhibition is temporary, it is shown that regrowth of bacteria after being transferred to a new medium, which is free from the influence of antagonistic bacteria. Bactericidal activity, inhibition is permanent. Bacteria were unable to grow even though they are transferred to new medium.

Test the mechanism of endophytic bacteria as biological control agents and plant growth-promoting microbes

The mechanism test was carried out on endophytic bacteria that have the potential to control the fungus *R. solani* and *Pantoea* sp., i.e. *Bacillus* sp. B.K.A1, *Bacillus* sp. B.K.A3, *Bacillus* sp. PP.A5, *Bacillus* sp. PPD.B2. The production of compounds related to biocontrol of pathogens and promotion of plant growth in bacterial isolates was evaluated by measuring the production of antimicrobial compounds and hydrolytic enzymes (amylases, lipases, proteases, and chitinases) and phosphate solubilization. The results of enzyme activity tests are as shown in Table 4. The four isolates *Bacillus* sp. tested were able to produce protease, lipase and phosphatase enzymes, with varied activity indexes. All isolates of *Bacillus* sp. those tested had a high index of protease and lipase enzymes (> 3) (Table 4., Fig 2.). Protease and lipase enzymes, related to the ability of the endophytic bacteria as biological control agents. Based on the protease and lipase indexes, *Bacillus* sp. PP.A5 can produce the highest proteases and lipase enzymes compared to other isolates. The isolates that have high protein and fat hydrolysis enzymes have the potential as biological control agents because proteins and fats are constituents of pathogen cells (Mota et al 2016). Besides, the protease enzyme is thought to degrade antibiotics produced by fungal or bacterial pathogens. According to Anderson et al. (2014), the extracellular protease enzyme produced by *P. fluorescens* can inactivate antibiotic compounds produced by *Pantoea agglomerans*.

Bacillus sp. PPD.B2 has the highest phosphate solubility index. The phosphate solubilization is related to the ability of endophytic bacteria as a plant growth promoter, providing phosphates for plants. Microbes with high phosphate solubility activity are capable of producing and releasing metabolites such as organic acids that chelate cations that are bound to phosphate (especially calcium) and converting them into soluble forms. Solubilization of different forms of phosphate by microbes associated with plants, and increasing its availability for plants, will increase growth and production of the plant (Djuric et al., 2011).



Figure 2. Hydrolysis enzyme activity, (a) protease , (b) lipase and (c) phosphate solubilization.

359

Table 3. Inhibition of endophytic bacteria against *Pantoea* sp.

No	Isolate	Antagonism	Antagonism index	Antagonism catagory*	Antagonism activity
Endophytic bacteria from the root					
1	fluorescent <i>Pseudomonas</i> BB.A2	-	0	-	-
2	fluorescent <i>Pseudomonas</i> BS.A 2	+	4,91	strong	bacteriostatic
3	fluorescent <i>Pseudomonas</i> BK.A1	+	4,42	strong	bacteriostatic
4	fluorescent <i>Pseudomonas</i> PPD.A1	-	0	-	-
5	fluorescent <i>Pseudomonas</i> PP.A1	-	0	-	-
6	fluorescent <i>Pseudomonas</i> PB.A4	+	5,29	strong	bactericidal
7	<i>Bacillus</i> sp.BB.A3	+	8,17	strong	bacteriostatic
8	<i>Bacillus</i> sp.BS.A1	+	4,00	strong	bacteriostatic
9	<i>Bacillus</i> sp. BSA3	+	5,07	strong	bactericidal
10	<i>Bacillus</i> sp. B.K.A1	+	4,01	strong	bakteriostatik
11	<i>Bacillus</i> sp. B.K.A3	+	4,91	strong	bacteriostatic
12	<i>Bacillus</i> sp.PP.A3	+	6,63	strong	bactericidal
13	<i>Bacillus</i> sp.PP.A5	+	6,56	strong	bactericidal
Endophytic bacteria from the stem					
14	fluorescent <i>Pseudomonas</i> PPD.B1	-	0	-	-
15	fluorescent <i>Pseudomonas</i> PPD.B5	+	5,86	strong	bactericidal
16	fluorescent <i>Pseudomonas</i> PP.B4	-	0	-	-
17	<i>Bacillus</i> sp. BB.B4	+	7,80	strong	bactericidal
18	<i>Bacillus</i> sp. BS.B1	+	6,22	strong	bacteriostatic
19	<i>Bacillus</i> sp. BK. B3	+	5,33	strong	bacteriostatic
20	<i>Bacillus</i> sp.PPD.B2	+	5,00	strong	bacteriostatic
21	<i>Bacillus</i> sp. PPD.B4	+	8,75	strong	bacteriostatic
22	<i>Bacillus</i> sp. PB.B1	+	1,67	weak	bacteriostatic
23	<i>Bacillus</i> sp. PB.B3	+	5,67	strong	bactericidal

•Based on Davis and Stout, 1971

360

361

362

Table 4. Test results of proteases, lipases and phosphate solubilization.

No	Isolate	Protease Test activity	index	Lipase Test activity	index	Phosphate solubilization Activity	index
Endophytic bacteria from the root							
1	<i>Bacillus</i> sp. B.K.A1	+	3.75	+	3.23	+	1.17
2	<i>Bacillus</i> sp. B.K.A3	+	3.20	+	3.73	+	1.27
3	<i>Bacillus</i> sp. PP.A5	+	5.00	+	4.40	+	1.46
Endophytic bacteria from the stem							
4	<i>Bacillus</i> sp.PPD.B2	+	3.00	+	3.90	+	2.60

363

364

CONCLUSION

Based on research carried out, it has been successfully isolated, morphologically and biochemically characterized four the endophytic bacteria that have the potential to be developed as biopesticides to control maize disease, especially *R. solani* and *Pantoea* sp. They can suppress the growth of *R. solani* by more than 50%, have a strong antagonistic index against *Pantoea* sp (> 4), and can produce protease and lipase enzyme, and phosphate solubilization.

ACKNOWLEDGEMENTS

This research was a part of the main study funded by Doctoral Dissertation Grant from the Indonesian Ministry of Research, Technology and Higher Education; for that, I deeply thank for the financial support

REFERENCES

- Abed HN, Rouag.,Mouatasse D, Rouabhi A. 2016. Screening for *Pseudomonas* and *Bacillus* antagonistic rhizobacteria strains for the biocontrol of Fusarium wilt of chickpea. Eurasian Journal of Soil Science 5(3):182–191
- Abidin Z, Aini LQ, Abadi AL. 2015. Effect of *Bacillus* sp. and *Pseudomonas* sp. on the growth of the pathogenic fungus *Sclerotium rolfsii* Sacc. causes of seedling diseases in soybean plants. Jurnal HPT 3(1): 1-10.
- Aini,LQ, Suryani L, Sugiharto AN,. Abadi A. 2013. Identification of bacterial wilt and leaf blight disease on maize (*Zea Mays*) found in Kediri, Indonesia. Agrivita 35 (1): 1-7.
- Al-Fadhal FA, AL-Abedy AN, Alkhafijel DA. 2019. Isolation and molecular identification of *Rhizoctonia solani* and *Fusarium solani* isolated from cucumber (*Cucumis sativus* L.) and their control feasibility by *Pseudomonas fluorescens* and *Bacillus subtilis*. Egyptian Journal of Biological Pest Control 29(47):1-11. <https://doi.org/10.1186/s41938-019-0145-5>
- Amin M, Rakhisi Z, Ahmady AZ. 2015. Isolation and Identification of *Bacillus* Species From Soil and Evaluation of Their Antibacterial Properties. Avicenna J Clin Microb Infec. 2(1): e23233
- Ammar, E, Correa VR, Hogenhout SA, Redinbaugh MG. 2014. Immunofluorescence localization and ultrastructure of Stewart's wilt disease bacterium *Pantoea stewartii* in maize leaves and in its flea beetle vector *Chaetocnema pulicaria* (Coleoptera: Chrysomelidae). Journal of Microscopy and Ultrastructure 2: 28–33
- Anderson LM, Stockwell VO, Loper JE. 2004. An extracellular protease of *Pseudomonas fluorescens* inactivates antibiotics of *Pantoea agglomerans*. Phytopathology 94:1228-1234
- Bacon CW, Hinton DM. 2002. Endophytic and biological control potential of *Bacillus mojavensis* and related species. Biological Control 23: 274-284.
- Cavaglieri L, Orlando J, Etcheverry M. 2009. Rhizosphere microbial community structure at different maize plant growth stages and root locations. Microbiological
- Compant SB, Duffy, Nowak J, Clement C, Barka EA. 2005. Use of plant growth-promoting bacteria for biocontrol of plant diseases: principles, mechanisms of action, and future prospects. Applied and Environmental Microbiology 71(9): 4951-4959.
- Coplin DL, Redinbaugh M G. 2012. The Bacterium *Pantoea stewartii* Uses Two Different Type III Secretion Systems To Colonize Its Plant Host and Insect Vector. Applied and Environmental Microbiology 78(17): 6327-6336.
- Costa FG, Zucchi TD, de Melo IS. 2013. Biological control of Phytopathogenic Fungi by Endophytic Actinomycetes Isolated from Maize (*Zea Mays* L.). Braz. Arch.Biol.Technol 56(6):948-955.
- Davis WW, Stout TR. 1971. Disc plate methods of microbiological antibiotic assay. Applied Microbiology 22(4):659-665.
- Desi Y, Habazar T, Agustian , Khairul U, Syamsuwirman, Novia P. 2014. Morphological dan physiological characterization of *Pantoea stewartii* subsp. *stewartii* from maize. Jurnal Fitopatologi Indonesia 10(2): 45–52. DOI: 10.14692/jfi.10.2.45
- Djaenuddin N, Nonci N, Muis A. 2017. Effectiveness of the formula *Bacillus subtilis* TM4 for disease control in maize plants. Jurnal Fitopatologi Indonesia 13(4): 113-118
- Djarmiko HA, Arwiyanto T, Hadisutrisno B, Sunarminto BH. 2007. Potential of three bacterial genera from three plant rhizosphere as biological agents controlling lincat disease. Jurnal ilmu-ilmu Pertanian 9(1):40-47.
- Djuric SA, Pavic, Jarak M, Pavlovic S, Starovic M, Pivic R, Josic D.. 2011. Selection of indigenous fluorescent pseudomonad isolates from maize rhizospheric soil in Vojvodina as possible PGPR. Romanian Biotechnological Letters 16(5): 6580–6590.DOI:10.1094/PHI-I-2004-0113-01.
- Farooq U, Bano A. 2013. Screening of indigenous bacteria from rhizosphere of maize (*Zea mays* L.) for their plant growth promotion ability and antagonism against fungal and bacterial pathogens. Journal of Animal and Plant Sciences, 23(6), pp.1642– 1652.
- Ganesan G, Kumar AM. 2005. *Pseudomonas fluorescens*, a potential bacterial antagonist to control plant diseases. Journal of Plant Interactions 1(3): 123_134.
- Hastuti RD, Saraswati R, Sari AP. 2014. The effectiveness of the endophytic microbes in promoting plant growth and controlling leaf blight disease in the lowland rice. Jurnal Tanah dan Iklim 38(2) : 109-118.
- Heydari A, Pessarakhi M. 2010. A review on biological control of fungal plant pathogens using microbial antagonists. Journal of Biological Science 10(4): 273-290
- Jain A, Das S. 2016. Insight into the Interaction between plants and associated fluorescent *Pseudomonas* spp. International Journal of Agronomy: 1-8. <http://dx.doi.org/10.1155/2016/4269010>
- Lim SM, Yoon M, Choi GJ, Choi YH, Jang KS, Shin TS, Park HW, Yu NH, Kim YH, Kim J. 2017. Diffusible and volatile zntifungal compounds produced by an antagonistic *Bacillus velezensis* G341 against various phytopathogenic fungi. Plant Pathol. J. 33(5) : 488-498. <https://doi.org/10.5423/PJ.OA.04.2017.0073>
- Motaa MS, Gomes CB, Júniora ITS, Moura AB. 2017. Bacterial selection for biological control of plant disease: criterion determination and validation. Brazilian Journal of Microbiology 48 : 62–70
- Muis A. 2007. Management of sheath blight disease (*Rhizoctonia solani* Kuhn.) in maize. Jurnal Litbang Pertanian 26(3) : 100-103Desi 2014
- Nasrun, Burhanudin. 2016. Evaluation of the efficacy of the formula *Pseudomonas fluorescens* for controlling bacterial wilt (*Ralstonia solanacearum*) patchouli. Buletin Penelitian Tanaman Rempah dan Obat, 27(1): 67-76.

427 Nuryanto B, Priyatmojo A, Hadisutrisno B.. 2014. Pengaruh Tinggi Tempat dan Tipe Tanaman Padi terhadap Keparahan Penyakit Hawar Pelepah.
 428 Penelitian Pertanian Tanaman Pangan 33(1):1–8.

429 Orole OO, Adejumo TO. 2011. Bacterial and fungal endophytes associated with grains and roots of maize. *Journal of Ecology and the natural*
 430 *Environment* 3(9):298-303.

431 Pal KK, McSpadden Gardener B. 2006. Biological Control of Plant Pathogens. *The Plant Health Instructor*. p1-25. DOI: 10.1094/PHI-A-2006-1117-02
 432 Pataky JK. 2004. Stewart's wilt of corn. *The Plant Health Instructor*. DOI:10.1094/PHI-I-2004-0113-01.;

433 Rosenblueth M, Martinez-Romero E. 2006. Bacterial endophytes and their interaction with hosts (Review). *MPMI* 19 (8): 827-837.

434 Santiago TR, Grabowski C, Rossato M, Romeiroa RS., Mizubuti ESG. 2015. Biological control of eucalyptus bacterial wilt with rhizobacteria. *Biological*
 435 *Control* 80:14–22. <http://dx.doi.org/10.1016/j.biocontrol.2014.09.007>.

436 Shanti AT, Vittal RR. 2013. Biocontrol potencies of Plant Growth promoting Rhizobacteria Against Fusarium Wilt Disease of Cucurbit. *Esci J. Plant*
 437 *Pathol* 2(3): 155-161.

438 Singh S, Dutta U, Bhat AK, Gupta S, Gupta V, Jamwal S. 2017. Morpho-cultural and biochemical identification of *Pseudomonas* sp. isolated from the
 439 rhizosphere of different vegetable crops and study its efficacy on *Solanum melongena* (Brinjal). *Journal of Pharmacognosy and Phytochemistry* 6(2):
 440 22-28

441 Slepecky R.A., Hemphill H.E. 2006. The Genus *Bacillus*—Nonmedical. In: Dworkin M., Falkow S., Rosenberg E., Schleifer KH., Stackebrandt E. (eds)
 442 *The Prokaryotes*. Springer, New York, NY

443 Soesanto L, Mugiastuti E, Rahayuniati RF. 2010. Study of the antagonistic mechanism of *Pseudomonas fluorescens* P60 against *Fusarium oxysporum*
 444 f.sp. *lycopersici* in tomatoes in vivo. *Jurnal HPT Tropika* 10(2) : 108-115

445 Soesanto L, Mugiastuti E, Rahayuniati RF. 2011. Utilization of some animal broths as a liquid formula for *Pseudomonas fluorescens* P60 to control
 446 *Sclerotium rolfsii* in cucumber plants. *Jurnal Perlindungan Tanaman Indonesia*, 17(1): 7–17.

Short Communication: Isolation and characterization of the endophytic bacteria, and their potential as maize diseases control

ENDANG MUGIASTUTI*, SUPRAYOGI, NUR PRIHATININGSIH, LOEKAS SOESANTO

Faculty of Agriculture, Universitas Jenderal Soedirman. Jl. Dr. Suparno, Karangwangkal, Purwokerto Utara, Banyumas 53123, Central Java, Indonesia.
Tel./fax.: +62-281-638791, *email: endangmugiaastuti@gmail.com.

Manuscript received: 13 November 2019. Revision accepted: xxx April 2020.

Abstract. Mugiaastuti E, Suprayogi, Prihatiningsih N, Soesanto L. 2020. Short Communication: Isolation And Characterization Of The Endophytic Bacteria, And Their Potential As Maize Diseases Control. *Biodiversitas* 21: xxxx. Sheath blight and bacterial wilt are diseases that can reduce maize production. Biological control with the endophytic bacteria offers environmentally friendly control for these pathogens. The study was aimed to isolate and characterize the endophytic bacteria morphologically and biochemically and to study their potential to control maize diseases, especially sheath blight and bacterial wilt causing pathogens. The study was conducted at the Plant Protection Laboratory, Faculty of Agriculture, Jenderal Soedirman University, from April to August 2019. The study consisted of four stages: isolation and characterization of endophytic bacteria, the antagonism test of the endophytic bacterial to *R. solani*, the antagonism test of the endophytic bacteria to *Pantoea* sp., and the mechanism test of the endophytic bacteria as biological control agents and plant growth-promoting bacteria. Based on the research, four endophytic bacteria isolates have been successfully isolated, and characterized successfully and found have the potential to be developed as biopesticides to control maize disease, especially *R. solani* and *Pantoea* sp. *Bacillus* sp. endophytic from the root (BK.A1; BK.A3; PP.A5) and *Bacillus* sp. endophytic from the stem (PPD.B2) can suppress the growth of *R. solani* by more than 50%, have a strong antagonistic index against *Pantoea* sp (> 4), and can produce protease and lipase enzyme, and phosphate solubilization.

Keywords: *Bacillus*, fluorescent *Pseudomonas*, *Pantoea*, *Rhizoctonia solani*

INTRODUCTION

Maize is a strategic food commodity in the world. In Indonesia, the government seeks to achieve self-sufficiency in maize through increasing production of sustainable maize. However, these efforts have faced several obstacles; one of them is the presence of plant diseases such as sheath blight caused by *Rhizoctonia solani* Kuhn, and bacterial wilt caused by *Pantoea stewartii*. *R. solani* can infect up to the midrib of the cob (Djaenuddin et al. 2017), resulting in up to 100% decrease in the yield (Muis 2007). *Pantoea* sp. can attack all stages of the plant causing wilting and leaf blight, and is known as Stewart's wilt (Pataky 2004; Ammar et al. 2014). The pathogens can cause 40-100% yield loss.

Over the past 3 decades, the concept of sustainable and environmentally friendly agriculture has been carried out by minimizing the use of chemicals, both synthetic fertilizers and pesticides. In the management of pests and plant diseases, biological control is developed by applying biological agents including the endophytic bacteria (Shanti and Vittal 2013). Many endophytic bacteria can pass the endodermic barrier across from the root cortex to the vascular system, and subsequently develop as endophytes in stems, leaves, tubers, and other organs (Compant et al. 2005). The use of endophytic bacteria as biological agents has an advantage compared to rhizosphere bacteria because

endophytic bacteria live and survive in the plant tissue during plant development, thus protecting the plants.

Bacillus sp. and fluorescent *Pseudomonas* are reported to be able to live as endophytes and are widely used as biological control agents for soil-borne and air-borne diseases. The endophytic bacteria could control plant diseases through several mechanisms including competition, hyperparasitism, producing microbial inhibiting compounds (antibiotics, lysis enzymes, other physical or chemical disorders), enhancing plant resistance, and promoting plant growth (Compant et al 2005, Pal and McSpadden 2006; Rosenblueth and Martinez-Romero 2006).

Based on the mechanisms, the use of endophytic bacteria isolated from maize, both upland and lowland, suggested potentially alternative control for sheath blight (*R. solani*) and bacterial wilt (*Pantoea* sp). The research aimed to isolate and characterize morphologically and biochemically the endophytic bacteria as well as their potential to control pathogens that cause disease in maize especially *R. solani* and *Pantoea* sp.

MATERIALS AND METHODS

This research was conducted at the Laboratory of Plant Protection, Faculty of Agriculture, Jenderal Soedirman

University, Purwokerto, Central Java, Indonesia, from April to August 2019

Isolation *R. solani*

R. solani was isolated from maize with sheath blight symptoms and there was sclerotium as a resistant structure from the pathogenic fungi in Banyumas. *R. solani* isolation was carried out based on Al-Fadhil et al. 2019. Disease samples were cut 0.5 x 0.5 cm, then sterilized with NaOCl (1%) for 2 min, and rinsed with sterile water 3 times. Disease samples pieces were then dried using sterile filter papers, and transferred to Petri dishes containing PDA medium to obtain pure *R. solani* isolates.

Isolation *Pantoea* sp.

Pantoea sp. was isolated from diseased maize samples taken from the maize growing area in Banyumas Regency according to Coplin et al. (2012); Aini et al. (2013) and Desi et al. (2014). Diseased leaves or stems were washed with running water, then dried with a tissue. Diseased samples were cut 1.5 x 5 cm, then sterilized with ethanol 70% and rinsed with sterile water 3 times. Furthermore, the sample was crushed with 5 ml of sterile distilled water using a sterile mortar. The bacterial suspension was streaked on nutrient agar and incubated 3-5 days. Bacterial colonies that exhibit the character of *Pantoea* sp. were yellow, shiny, slimy, flat or convex, then separated as pure cultures of *stewartii* candidates. The culture was then tested by Gram Reaction (KOH test), Hugh-Leifson test, pigment production in YDC medium, oxidase test, hypersensitivity test, and pathogenicity test on maize.

Isolation and characterization of endophytic bacteria

Sampling for the isolation of endophytic bacteria was carried out in Banyumas and Purbalingga, Central Java, Indonesia, with purposive stratified random sampling. Samples were taken from two areas of altitude, i.e., low-moderate lands (0-500 m above sea level), and highlands (> 500 m above sea level) (Nuryanto et al., 2014). In each district, 2 locations were selected for the low-medium lands, and 1 location for the highlands. Age of maize plants was 20-30 days after planting, when the number of endophytic microbial populations that can be cultured is in the highest population (Cavaglieri et al. 2009).

The endophytic bacteria were isolated from the roots and stems of healthy maize plants. Roots and stems were washed, sterilized with 70% ethanol (1 minute), 20% sodium hypochlorite (5 minutes) and Ringer's thiosulfate solution (5 minutes). Separately, the roots and stems of 10 g each were crushed with 90 ml PBS on a sterile mortar. Subsequently, samples were plated on NA and Kings B media (Cavaglieri et al. 2009). To isolate *Bacillus* sp., the suspension was heated for 10 minutes at 80 ° C, before plating on NA. Bacterial isolates were further purified and characterized, such as morphological characteristics, gram properties, catalase tests, and hypersensitivity tests

The antagonism test of endophytic bacterial to *R. solani*

The antagonism test of endophytic bacteria on *R. solani* was carried out using the dual culture method. The level of

inhibition of antagonist is calculated using the formula (Abidin et al., 2015).

$$I = \frac{C-T}{C} \times 100\%$$

Where:

I : The level of inhibition of antagonist (%)

C: The radius of pathogen colonies opposite antagonist

T: The radius of the colony of pathogens towards antagonist

The antagonism test of endophytic bacterial to *Pantoea* sp.

Antagonism testing was carried out using the double-layer test method (Santiago et al. 2015). Endophytic bacteria to be tested were grown on the nutrient agar medium, incubated at 28 °C for 48 hours. In the upside-down position, 1 ml of chloroform was added to the cup lid and left for 2 hours. Next, add 5 mL so that 0.6% water containing 0.5 mL of *Pantoea* sp. bacterial suspension. The culture was re-incubated for 24 hours, and there were clear zones around the antagonistic bacterial colony. The antibiotic activity was assessed based on the diameter of the clear zone compared to the diameter of the colony. Characterization of the type of antibiosis can be divided into bactericidal and bacteriostatic types according to the method of based on Djatmiko et al. (2007).

The mechanism test of endophytic bacteria as controlling agents biological and plant growth-promoting microbial.

The testing mechanism of endophytic bacteria was carried out for bacteria that have the potential in testing the antagonism of the fungus *R. solani* and *Pantoea* sp.

Protease Test

The activity of the ability of antagonistic bacteria to produce extracellular protease enzymes was tested using Skim Milk Agar (SMA) medium. Each bacterium to be tested was grown in a medium SMA and incubated at 28 °C for 24-48 hours. The presence of clear zones around the colony shows that positive bacteria produce protease enzymes (Abed et al. 2016). The protease activity index is assessed based on the diameter of the clear zone compared to the diameter of the colony.

$$\text{Protease index} = \frac{(\text{clear zone diameter} - \text{colony diameter})}{\text{colony diameter}}$$

Lipase test

Detection of the ability of bacteria to produce the enzyme lipase was done by growing the antagonistic bacteria on a medium containing 1% Tween 80. The presence of lipase enzyme activity was demonstrated by milky white sediment around the bacterial colony, after incubating at 28 °C for 4-7 days. The lipolytic index was measured using a formula Djuric et al. (201).

$$\text{Lipolytic index} = \frac{(\text{milky white diameter} - \text{colony diameter})}{\text{colony diameter}}$$

Formatted: Font: Bold

Formatted: Indent: First line: 0 cm

Commented [U1]: water agar

Formatted: Indent: First line: 0 cm

Formatted: Font: Bold

Phosphatase test

Detection of the ability of bacteria to produce the enzyme phosphatase was done by growing bacterial isolates on Pikovskaya medium. After incubating for 7 days at 28 °C, the presence of a clear zone around the bacterial colony shows that the bacteria has the ability to produce the phosphatase enzyme to dissolve phosphates. The solubility index is measured using a formula (Farooq and Bano 2013)

$$\text{Phosphatase Index} = \frac{(\text{clear zone diameter} - \text{colony diameter})}{\text{colony diameter}}$$

RESULTS AND DISCUSSION

Isolation and characterization of endophytic bacteria

The results of the exploration, isolation, and characterization of endophytic bacteria obtained 23 isolates of endophytic bacteria, consisted of 9 isolates of the fluorescent *Pseudomonas* and 14 isolates of *Bacillus* sp. (Table 1). Fluorescent *Pseudomonas* colony on King's B was round, with a flat edge, and yellowish-white, to greenish-yellow, gram-negative, rod-shaped, non-spore and fluorescent under ultraviolet light. Singh et al. 2017 reported fluorescent *Pseudomonas* showed light green, yellowish, creamy, circular, slimy, regular-irregular characteristics. Bacteria have short-long rod forms. The fluorescent *Pseudomonas* isolates is gram-negative, which can form catalase, a positive oxidase, needed to grow aerobes. *Bacillus* sp. was observed with a spherical colony having cell rod-shaped, gram-positive, and endospores within cells (Table 1.). Slepecky and Hempill 2006; Amin et al. 2015 reported *Bacillus* sp. has the characteristics of a circular colony and punctiform (small round), variations in the entire margin and lobate, white dull, non-slimy, gram-positive, has endospores, flagellum and some are motile.

Based on its distribution, fluorescent *Pseudomonas* and *Bacillus* sp. found in all sampling locations, high or low-medium lands. This shows that fluorescent *Pseudomonas* and *Bacillus* sp. spread and can live in various altitudes, both high and low-medium land. Bacon and Hilton 2002; Ganeshan and Kumar 2005 reported *P. fluorescens* and *Bacillus* sp., are species of bacteria with a wide range of life and are very adaptive in various environments. Both types of bacteria are also found in the roots or corn stalks. *Bacillus* sp. and *Pseudomonas* sp. including a group of endophytic bacteria have a wide range of life and more isolated in maize (Ganeshan and Kumar 2005; Orole and Adejumo 2011; Costa et al. 2013)

Antagonism test between the endophytic bacteria against *R. solani*

Based on the results of *in vitro* tests (Table 2), 24 isolates of the endophytic bacteria were able to inhibit the growth of *R. solani*, with varying degrees of inhibition. The endophytic bacteria that have inhibition rates above 50%, i.e. fluorescent *Pseudomonas* BK.A1 (51%), *Bacillus* sp. B.K.A1 (55.39%), *Bacillus* sp. BK.A3 (51.52%), PP.A5 (50.66%), and PPD.B2 (50.8%). The effect of the endophytic bacteria in inhibiting the growth of *R. solani* is inversely proportional to the dry weight mycelium. The greater the percentage of inhibition of endophytic bacteria to the growth of *R. solani*, the smaller the dry weight mycelium (Table 2.) Endophytic bacteria can inhibit the growth of *R. solani*, which were shown by the inhibitory zone in the area bordering the bacterial streak (Figure 1a). The endophytic bacteria have anti-pathogenic properties and can produce antibiotic compounds. The ability of the endophytic bacteria to control plant pathogens occurs through the mechanism of antibiosis, competition, lysis, inducing resistance and producing growth substances. Bacteria capable of producing secondary metabolites that can inhibit growth or damage pathogens (Hastuti et al. 2014). These compounds, including alkaloids, phenols, flavonoids, glycosides, and phytoalexin (Soesanto et al. 2010). Fluorescent *Pseudomonas* can produce various types of antibiotics including phenazine-1-carboxylic acid, pyocyanin, pyrrolnitrin, and pyoluteorin, 2,4-diacetylphloroglucinol (Phl). Phl is a phenolic metabolite with antibacterial and antifungal (Jain and Das 2016). *Bacillus* species can produce various kinds of volatile compounds and diffusible with strong inhibitory activity against plant pathogens (Lim et al. 2017).




Figure 1. Antagonism test between the endophytic bacteria against *R. solani* (a) and *Pantoea* sp (b)

Formatted: Underline

Formatted: Highlight

Formatted: No underline

Commented [U2]: move from bottom




Figure 2. Hydrolysis enzyme activity, (a) protease , (b) lipase and (c) phosphate solubilization.

Table 1. Isolation and characterization of endophytic bacteria.

Land	Sampling location	Sample	Gram test	Catalase test	Oxidase test	Colony morphology	Colony pigment	Fluorescence on KB medium	Cell morphology	Endo-spores	Isolate
Highland	Purbalingga, Pratin 7.13'33" S, 109.17'21" E, 1.190 m asl	Root	-	+	+	Round	yellowish white	+	Medium rod	-	fluorescent <i>Pseudomonas</i> PP.A1
		Root	+	+	+	Round	White	-	Small rod	+	<i>Bacillus</i> sp. PP.A3
		Root	+	+	+	Round	White	-	Medium rod	+	<i>Bacillus</i> sp. PP.A5
		Stem	-	+	+	Round	yellowish white	+	Medium rod	-	fluorescent <i>Pseudomonas</i> PP. B4
	Banyumas, Baturaden 7.19"1" S, 109.14'29" E, 520 m asl	Root	-	+	+	Round	Greenish yellow	+	Medium rod	-	fluorescent <i>Pseudomonas</i> BB.A2
		Root	+	+	+	Round	White	-	Small rod	+	<i>Bacillus</i> sp. BB.A3
		Stem	+	+	+	Round	White	-	Medium rod	+	<i>Bacillus</i> sp. BB.B4
	Medium-Lowland	Banyumas, Sumbang 7.21'54" S, 109.17'33"E, 200 m asl	Root	-	+	+	Round	yellowish white	+	Medium rod	-
Root			+	+	+	Round	White	-	Medium rod	+	<i>Bacillus</i> sp. BS.A1
Root			+	+	+	Round	White	-	Small rod	+	<i>Bacillus</i> sp. BS.A3
Stem			+	+	+	Round	White	-	Small rod	+	<i>Bacillus</i> sp. BS. B1
Purbalingga, Bojongsari 7.20'12" S, 109.20'22" E, 190 m asl		Root	-	+	+	Round	Greenish yellow	+	Small rod	-	fluorescent <i>Pseudomonas</i> (PP) PB. A 4
		Stem	+	+	+	Round	White	-	Small rod	+	<i>Bacillus</i> sp. PB. B1
		Stem	+	+	+	Round	White	-	Medium rod	+	<i>Bacillus</i> sp. BB. B3
Purbalingga, Padamara 7.22'28" S, 109.13'24" E, 180 m asl		Root	-	+	+	Round	Greenish yellow	+	Small rod	-	fluorescent <i>Pseudomonas</i> (PP) PPD A1
		Stem	-	+	+	Round	yellowish white	+	Medium rod	-	fluorescent <i>Pseudomonas</i> (PP) PPD. B1
		Stem	-	+	+	Round	yellowish white	+	Medium rod	-	fluorescent <i>Pseudomonas</i> (PP) PPD. B5
		Stem	+	+	+	Round	White	-	Small rod	+	<i>Bacillus</i> sp. PPD. B2
		Stem	+	+	+	Round	White	-	Small rod	+	<i>Bacillus</i> sp. PPD. B4
Banyumas, Kembaran 7.23'47" S, 109.17'9" E, 110 m asl		Root	-	+	+	Round	yellowish white	+	Medium rod	-	fluorescent <i>Pseudomonas</i> (PP) BK. A1
		Root	+	+	+	Round	White	-	Medium rod	+	<i>Bacillus</i> sp. BK.A1
		Root	+	+	+	Round	White	-	Medium rod	+	<i>Bacillus</i> sp. BK.A3
		Stem	+	+	+	Round	White	-	Medium rod	+	<i>Bacillus</i> sp. BK.B3

Table 2. Inhibition of endophytic bacteria against *R. solani*.

Isolate	Inhibition rate (%)	Dry weights mycelium
Control	0	0.093
Endophytic bacteria from the root		
fluorescent <i>Pseudomonas</i> BB.A2	49.00	0.038
fluorescent <i>Pseudomonas</i> BS.A 2	45.00	0.027
fluorescent <i>Pseudomonas</i> BK.A1	51.00	0.017
fluorescent <i>Pseudomonas</i> PPD.A1	10.33	0.059
fluorescent <i>Pseudomonas</i> PP.A1	38.33	0.017
fluorescent <i>Pseudomonas</i> PB.A4	18.00	0.037
<i>Bacillus</i> sp.BB.A3	40.42	0.030
<i>Bacillus</i> sp.BS.A1	48.73	0.016
<i>Bacillus</i> sp. BSA3	37.42	0.039
<i>Bacillus</i> sp. B.K.A1	55.39	0.002
<i>Bacillus</i> sp. B.K.A3	51.52	0.003
<i>Bacillus</i> sp.PP.A3	46.65	0.019
<i>Bacillus</i> sp.PP.A5	50.66	0.009
Endophytic bacteria from the stem		
fluorescent <i>Pseudomonas</i> PPD.B1	27.00	0.020
fluorescent <i>Pseudomonas</i> PPD.B5	49.33	0.013
fluorescent <i>Pseudomonas</i> PP.B4	65.67	0.004
<i>Bacillus</i> sp. BB.B4	44.44	0.026
<i>Bacillus</i> sp. BS.B1	49.74	0.012
<i>Bacillus</i> sp. BK. B3	40.36	0.031
<i>Bacillus</i> sp.PPD.B2	50.8	0.007
<i>Bacillus</i> sp. PPD.B4	39.44	0.036
<i>Bacillus</i> sp. PB.B1	37.29	0.047
<i>Bacillus</i> sp. PB.B3	44.9	0.022

2,4-diacetylphloroglucinol (Phl). Phl is a phenolic metabolite with antibacterial and antifungal (Jain and Das

Table 3. Inhibition of endophytic bacteria against *Pantoea* sp.

Isolate	Antagonism	Antagonism index	Antagonism catagory*	Antagonism activity
Endophytic bacteria from the root				
fluorescent <i>Pseudomonas</i> BB.A2	-	0	-	-
fluorescent <i>Pseudomonas</i> BS.A 2	+	4.91	Strong	Bacteriostatic
fluorescent <i>Pseudomonas</i> BK.A1	+	4.42	Strong	Bacteriostatic
fluorescent <i>Pseudomonas</i> PPD.A1	-	0	-	-
fluorescent <i>Pseudomonas</i> PP.A1	-	0	-	-
fluorescent <i>Pseudomonas</i> PB.A4	+	5.29	Strong	Bactericidal
<i>Bacillus</i> sp.BB.A3	+	8.17	Strong	Bacteriostatic
<i>Bacillus</i> sp.BS.A1	+	4.00	Strong	Bacteriostatic
<i>Bacillus</i> sp. BSA3	+	5.07	Strong	Bactericidal
<i>Bacillus</i> sp. B.K.A1	+	4.01	Strong	Bakteriostatik
<i>Bacillus</i> sp. B.K.A3	+	4.91	Strong	Bacteriostatic
<i>Bacillus</i> sp.PP.A3	+	6.63	Strong	Bactericidal
<i>Bacillus</i> sp.PP.A5	+	6.56	Strong	Bactericidal
Endophytic bacteria from the stem				
fluorescent <i>Pseudomonas</i> PPD.B1	-	0	-	-
fluorescent <i>Pseudomonas</i> PPD.B5	+	5.86	Strong	Bactericidal
fluorescent <i>Pseudomonas</i> PP.B4	-	0	-	-
<i>Bacillus</i> sp. BB.B4	+	7.80	Strong	Bactericidal
<i>Bacillus</i> sp. BS.B1	+	6.22	Strong	Bacteriostatic
<i>Bacillus</i> sp. BK. B3	+	5.33	Strong	Bacteriostatic
<i>Bacillus</i> sp.PPD.B2	+	5.00	Strong	Bacteriostatic
<i>Bacillus</i> sp. PPD.B4	+	8.75	Strong	Bacteriostatic
<i>Bacillus</i> sp. PB.B1	+	1.67	Weak	Bacteriostatic

2016). *Bacillus* species can produce various kinds of volatile compounds and diffusible with strong inhibitory activity against plant pathogens (Lim et al. 2017).

The endophytic bacterial antagonism test against *Pantoea* sp.

The results of antagonism between the endophytic bacteria and *Pantoea* sp. show varied results. The endophytic bacteria that can inhibit bacterial growth were indicated by the presence of clear zones around the endophytic bacterial colonies (Fig.1). From the nine isolate fluorescent *Pseudomonas* were tested, only three isolates were able to inhibit the growth of the *Pantoea* sp., i.e fluorescent *Pseudomonas* (Pf) BS.A2, Pf BK.A1 Pf PPD.B5. While the isolates Pf BB.A2, Pf PPD.A1, Pf PP.A1, Pf PPD.B1, and Pf PP.B4 are not able to inhibit the growth of pathogenic bacteria. Meanwhile, all isolate *Bacillus* sp. tested (thirteen isolates) were able to inhibit the growth of *Pantoea* sp. (Table 3).

The presence of clear zones around endophytic bacterial colonies showed the ability of endophytic bacteria to produce antibiotics to inhibit the growth of *Pantoea* sp. *P. fluorescens* P60 can produce antibiotics that inhibit the growth of pathogens (Soesanto 2011). *Pseudomonas fluorescens* is reported to produce phenazine-1-carboxylic acid (PCA) and other derivatives, 2,4 diacetyl phloroglucinol (Phl) (DAPG), pyrrolnitrin (PRN) and or pyoluteorin (Plt) (Heydari and Pessarakli 2010). Nasrun and Burhanudin (2016) mention that *P. fluorescens* produce secondary metabolites, i.e. antimicrobial, cyanide acid and 2,4-diacetyl phloroglucinol phenazine, pyrrolnitrin, pyoluteorin antibiotics.

Commented [U3]: move to the previous page

<i>Bacillus</i> sp. PB.B3	+	5,67	Strong	Bactericidal
---------------------------	---	------	--------	--------------

Note: •Based on Davis and Stout, 1971

Table 4. Test results of proteases, lipases and phosphate solubilization.

Isolate	Protease test		Lipase test		Phosphate solubilization	
	Activity	Index	Activity	Index	Activity	Index
Endophytic bacteria from the root						
<i>Bacillus</i> sp. B.K.A1	+	3.75	+	3.23	+	1.17
<i>Bacillus</i> sp. B.K.A3	+	3.20	+	3.73	+	1.27
<i>Bacillus</i> sp. PP.A5	+	5.00	+	4.40	+	1.46
Endophytic bacteria from the stem						
<i>Bacillus</i> sp.PPD.B2	+	3.00	+	3.90	+	2.60

The level of bacteria's ability to inhibit growth can be shown by the large diameter of the clear zone. The results showed that the antagonism index ranged from 1.67-8.17. Based on this index, most endophytic bacteria have a strong antagonism (index of antagonism > 4) (Davis and Stout 1971). Furthermore, bacterial isolates that showed antagonistic activity were tested for types of antagonism based on Djatmiko (2007). Based on the type of antagonistic activity, ten isolates the endophytic bacteria were bacteriostatic and nine isolates the endophytic bacteria were bactericidal. Bacteriostatic activity, growth inhibition is temporary, it is shown that regrowth of bacteria after being transferred to a new medium, which is free from the influence of antagonistic bacteria. Bactericidal activity, inhibition is permanent. Bacteria were unable to grow even though they are transferred to new medium.

Test the mechanism of endophytic bacteria as biological control agents and plant growth-promoting microbes

The mechanism test was carried out on endophytic bacteria that have the potential to control the fungus *R. solani* and *Pantoea* sp., i.e. *Bacillus* sp. B.K.A1, *Bacillus* sp. B.K.A3, *Bacillus* sp. PP.A5, *Bacillus* sp. PPD.B2. The production of compounds related to biocontrol of pathogens and promotion of plant growth in bacterial isolates was evaluated by measuring the production of antimicrobial compounds and hydrolytic enzymes (amylases, lipases, proteases, and chitinases) and phosphate solubilization. The results of enzyme activity tests are as shown in Table 4. The four isolates *Bacillus* sp. tested were able to produce protease, lipase and phosphatase enzymes, with varied activity indexes. All isolates of *Bacillus* sp. those tested had a high index of protease and lipase enzymes (> 3) (Table 4., Figure 2.). Protease and lipase enzymes, related to the ability of the endophytic bacteria as biological control agents. Based on the protease and lipase indexes, *Bacillus* sp. PP.A5 can produce the highest proteases and lipase enzymes compared to other isolates. The isolates that have high protein and fat hydrolysis enzymes have the potential as biological control agents because proteins and fats are constituents of pathogen cells (Mota et al 2016). Besides, the protease enzyme is thought to degrade antibiotics produced by fungal or bacterial pathogens. According to Anderson et al. (2014), the extracellular protease enzyme produced by *P. fluorescens*

can inactivate antibiotic compounds produced by *Pantoea agglomerans*.

Bacillus sp. PPD.B2 has the highest phosphate solubility index. The phosphate solubilization is related to the ability of endophytic bacteria as a plant growth promoter, providing phosphates for plants. Microbes with high phosphate solubility activity are capable of producing and releasing metabolites such as organic acids that chelate cations that are bound to phosphate (especially calcium) and converting them into soluble forms. Solubilization of different forms of phosphate by microbes associated with plants, and increasing its availability for plants, will increase growth and production of the plant (Djuric et al., 2011).

In conclusion, based on research carried out, it has been successfully isolated, morphologically and biochemically characterized four the endophytic bacteria that have the potential to be developed as biopesticides to control maize disease, especially *R. solani* and *Pantoea* sp. They can suppress the growth of *R.solani* by more than 50%, have a strong antagonistic index against *Pantoea* sp (> 4), and can produce protease and lipase enzyme, and phosphate solubilization.

ACKNOWLEDGEMENTS

This research was a part of the main study funded by Doctoral Dissertation Grant from the Indonesian Ministry of Research, Technology and Higher Education; for that, I deeply thank for the financial support.

REFERENCES

- Abed HN, Rouag, Mouatasssem D, Rouabhi A. 2016. Screening for *Pseudomonas* and *Bacillus* antagonistic rhizobacteria strains for the biocontrol of fusarium wilt of chickpea. *Eurasian J Soil Sci* 5 (3):182-191
- Abidin Z, Aini LQ, Abadi AL. 2015. Effect of *Bacillus* sp. and *Pseudomonas* sp. On the growth of the pathogenic fungus *Sclerotium rolfsii* Sacc. causes of seedling diseases in soybean plants. *Jurnal HPT* 3 (1): 1-10.
- Aini LQ, Suryani L, Sugiharto AN, Abadi A. 2013. Identification of bacterial wilt and leaf blight disease on maize (*Zea Mays*) found in Kediri, Indonesia. *Agrivita* 35 (1): 1-7.
- Al-Fadhal FA, AL-Abedy AN, Alkhafijel DA. 2019. Isolation and molecular identification of *Rhizoctonia solani* and *Fusarium solani* isolated from cucumber (*Cucumis sativus* L.) and their control

- feasibility by *Pseudomonas fluorescens* and *Bacillus subtilis*. Egypt J Biol Pest Control 29 (47):1-11. <https://doi.org/10.1186/s41938-019-0145-5>
- Amin M, Rakhisi Z, Ahmady AZ. 2015. Isolation and Identification of *Bacillus* Species From Soil and Evaluation of Their Antibacterial Properties. Avicenna J Clin Microb Infect. 2 (1): e23233
- Ammar, E, Correa VR, Hogenhout SA, Redinbaugh MG. 2014. Immunofluorescence localization and ultrastructure of Stewart's wilt disease bacterium *Pantoea stewartii* in maize leaves and in its flea beetle vector *Chaetocnema pulicaria* (Coleoptera: Chrysomelidae). J Microsc Ultrastructure 2: 28-33
- Anderson LM, Stockwell VO, Loper JE. 2004. An extracellular protease of *Pseudomonas fluorescens* inactivates antibiotics of *Pantoea agglomerans*. Phytopathology 94:1228-1234
- Bacon CW, Hinton DM. 2002. Endophytic and Biological Control Potential Of *Bacillus Mojavensis* And Related Species. Biol Control 23: 274-284.
- Cavaglieri L, Orlando J, Etcheverry M. 2009. Rhizosphere Microbial Community Structure At Different Maize Plant Growth Stages And Root Locations. Microbiological
- Compant SB, Duffy, Nowak J, Clement C, Barka EA. 2005. Use of plant growth-promoting bacteria for biocontrol of plant diseases: principles, mechanism of action, and future prospects. Appl Environmental Microbiol 71 (9): 4951-4959.
- Coplin DL, Redinbaugh MG. 2012. The Bacterium *Pantoea stewartii* Uses Two Different Type III Secretion Systems To Colonize Its Plant Host and Insect Vector. Appl Environmental Microbiol 78 (17): 6327-6336.
- Costa FG, Zucchi TD, de Melo IS. 2013. Biological control of phytopathogenic fungi by endophytic actinomycetes isolated from maize (*Zea mays* L.). Braz. Arch Biol Technol 56 (6):948-955.
- Davis WW, Stout TR. 1971. Disc plate methods of microbiological antibiotic assay. Appl Microbiol 22 (4):659-665.
- Desi Y, Habazar T, Agustian, Khairul U, Syamsuwirman, Novia P. 2014. Morphological dan Physiological Characterization of *Pantoea Stewartii* Subsp. *stewartii* from maize. Jurnal Fotopatologi Indonesia 10 (2): 45-52. DOI: 10.14692/jfi.10.2.45 [indonesian]
- Djaenuddin N, Nonci N, Muis A. 2017. Effectiveness of the formula *Bacillus subtilis* TM4 for disease control in maize plants. Jurnal Fitopatologi Indonesia 13 (4): 113-118 [Indonesian]
- Djamtiko HA, Arwiyanto T, Hadisutrisno B, Sunarminto BH. 2007. Potential of three bacterial genera from three plant rhizosphere as biological agents controlling linat disease. Jurnal ilmu-ilmu Pertanian 9 (1):40-47. [Indonesia]
- Djuric SA, Pavic, Jarak M, Pavlovic S, Starovic M, Pivic R, Josic D. 2011. Selection of indigenous fluorescent pseudomonad isolates from maize rhizospheric soil in Vojvodina as possible PGPR. Romanian Biotechnological Letters 16 (5): 6580-6590. DOI:10.1094/PHI-I-2004-0113-01.
- Farooq U, Bano A. 2013. Screening of indigenous bacteria from rhizosphere of maize (*Zea mays* L.) for their plant growth promotion ability and antagonism against fungal and bacterial pathogens. J Animal Plant Sci, 23 (6):1642-1652.
- Ganeshan G, Kumar AM. 2005. *Pseudomonas fluorescens*, a potential bacterial antagonist to control plant diseases. J Plant Interactions 1 (3): 123_134.
- Hastuti RD, Saraswati R, Sari AP. 2014. The Effectiveness Of The Endophytic Microbes In Promoting Plant Growth And Controlling Leaf Blight Disease In The Lowland Rice. Jurnal Tanah dan Iklim 38 (2): 109-118. [indonesian]
- Heydari A, Pessarakhi M. 2010. A review on biological control of fungal plant pathogens using microbial antagonists. J Biol Sci 10 (4): 273-290
- Jain A, Das S. 2016. Insight Into The Interaction Between Plants And Associated Fluorescent *Pseudomonas* spp. Int J Agron: 1-8. <http://dx.doi.org/10.1155/2016/4269010>
- Lim SM, Yoon M, Choi GJ, Choi YH, Jang KS, Shin TS, Park HW, Yu NH, Kim YH, Kim J. 2017. Diffusible and volatile zntifungal compounds produced by an antagonistic *Bacillus velezensis* G341 against various phytopathogenic fungi. Plant Pathol J 33 (5): 488-498. <https://doi.org/10.5423/PPJ.OA.04.2017.0073>
- Motaa MS, Gomes CB, Jüniora ITS, Moura AB. 2017. Bacterial Selection For Biological Control Of Plant Disease: Criterion Determination And Validation. Braz J Microbiol 48: 62-70
- Muis A. 2007. Management of sheath blight disease (*Rhizoctonia solani* Kuhn.) in maize. Jurnal Litbang Pertanian 26 (3): 100-103 Desi 2014 [indonesian]
- Nasrun, Burhanudin. 2016. Evaluation of the efficacy of the formula *Pseudomonas fluorescens* for controlling bacterial wilt (*Ralstonia solanacearum*) patchouli. Buletin Penelitian Tanaman Rempah dan Obat, 27 (1): 67-76. [indonesian]
- Nuryanto B, Priyatmojo A, Hadisutrisno B. 2014. Pengaruh Tinggi Tempat dan Tipe Tanaman Padi terhadap Keparahan Penyakit Hawar Pelepah. Penelitian Pertanian Tanaman Pangan 33 (1):1-8. [indonesian]
- Orole OO, Adejumo TO. 2011. Bacterial and fungal endophytes associated with grains and roots of maize. J Ecol Nat Environment 3 (9):298-303.
- Pal KK, McSpadden Gardener B. 2006. Biological Control of Plant Pathogens. The Plant Health Instructor. p1-25. DOI: 10.1094/PHI-A-2006-1117-02
- Pataky JK. 2004. Stewart's wilt of corn. The Plant Health Instructor. DOI:10.1094/PHI-I-2004-0113-01.
- Rosenblueth M, Martínez-Romero E. 2006. Bacterial endophytes and their interaction with hosts (Review). MPMI 19 (8): 827-837.
- Santiago TR, Grabowski C, Rossato M, Romeiroa RS., Mizubuti ESG. 2015. Biological control of eucalyptus bacterial wilt with rhizobacteria. Biological Control 80:14-22. <http://dx.doi.org/10.1016/j.biocontrol.2014.09.007>.
- Shanti AT, Vittal RR. 2013. Biocontrol potenciales of Plant Growth promoting Rhizobacteria Against Fusarium Wilt Disease of Cucurbit. Esci J. Plant Pathol 2 (3): 155-161.
- Singh S, Dutta U, Bhat AK, Gupta S, Gupta V, Jamwal S. 2017. Morpho Cultural And Biochemical Identification Of *Pseudomonas* Sp. Isolated From The Rhizosphere Of Different Vegetable Crops And Study Its Efficacy On *Solanum melongena* (Brinjal). J Pharmacognosy Phytochem 6 (2): 22-28
- Slepecky R.A., Hemphill H.E. 2006. The Genus *Bacillus*—Nonmedical. In: Dworkin M., Falkow S., Rosenberg E., Schleifer KH., Stackebrandt E. (eds) The Prokaryotes. Springer, New York, NY
- Soesanto L, Mugiastuti E, Rahayuniati RF. 2010. Study Of The Antagonistic Mechanism Of *Pseudomonas Fluorescens* P60 Against *Fusarium Oxysporum* F.Sp. *Lycopersici* In Tomatoes In Vivo. Jurnal HPT Tropika 10 (2): 108-115
- Soesanto L, Mugiastuti E, Rahayuniati RF. 2011. Utilization of some animal broths as a liquid formula for *Pseudomonas fluorescens* P60 to control *Sclerotium rolfsii* in cucumber plants. Jurnal Perlindungan Tanaman Indonesia, 17 (1): 7-17.

Short Communication: Isolation and characterization of the endophytic bacteria, and their potential as maize diseases control

ENDANG MUGIASTUTI[✉], SUPRAYOGI, NUR PRIHATININGSIH, LOEKAS SOESANTO

Faculty of Agriculture, Universitas Jenderal Soedirman. Jl. Dr. Suparno, Karangwangkal, Purwokerto Utara, Banyumas 53123, Central Java, Indonesia.
Tel./fax.: +62-281-638791, ✉email: endangmugiastuti@gmail.com.

Manuscript received: 13 November 2019. Revision accepted: 6 April 2020.

Abstract. Mugiastuti E, Suprayogi, Prihatiningsih N, Soesanto L. 2020. Short Communication: Isolation And Characterization Of The Endophytic Bacteria, And Their Potential As Maize Diseases Control. *Biodiversitas* 21: 1809-1815. Sheath blight and bacterial wilt are diseases that can reduce maize production. Biological control with the endophytic bacteria offers environmentally friendly control for these pathogens. The study aimed to isolate and characterize the endophytic bacteria morphologically and biochemically and to study their potential to control maize diseases, especially sheath blight and bacterial wilt causing pathogens. The study was conducted at the Plant Protection Laboratory, Faculty of Agriculture, Jenderal Soedirman University, from April to August 2019. The study consisted of four stages: isolation and characterization of endophytic bacteria, the antagonism test of the endophytic bacteria to *R. solani*, the antagonism test of the endophytic bacteria to *Pantoea* sp., and the mechanical test of the endophytic bacteria as biological control agents and plant growth-promoting bacteria. Based on the research, four endophytic bacteria isolates have been successfully isolated, and characterized successfully and found have the potential to be developed as biopesticides to control maize disease, especially *R. solani* and *Pantoea* sp. *Bacillus* sp. endophytic from the root (BK.A1; BK.A3; PP.A5) and *Bacillus* sp. endophytic from the stem (PPD.B2) can suppress the growth of *R. solani* by more than 50%, have a strong antagonistic index against *Pantoea* sp (> 4), and can produce protease and lipase enzyme, and phosphate solubilization.

Keywords: *Bacillus*, fluorescent *Pseudomonas*, *Pantoea*, *Rhizoctonia solani*

INTRODUCTION

Maize is a strategic food commodity in the world. In Indonesia, the government seeks to achieve self-sufficiency in maize through increasing production of sustainable maize. However, these efforts have faced several obstacles; one of them is the presence of plant diseases such as sheath blight caused by *Rhizoctonia solani* Kuhn, and bacterial wilt caused by *Pantoea stewartii*. *R. solani* can infect up to the midrib of the cob (Djaenuddin et al. 2017), resulting in up to 100% decrease in the yield (Muis 2007). *Pantoea* sp. can attack all stages of the plant causing wilting and leaf blight, and is known as Stewart's wilt (Pataky 2004; Ammar et al. 2014). The pathogens can cause 40-100% yield loss.

Over the past 3 decades, the concept of sustainable and environmentally friendly agriculture has been carried out by minimizing the use of chemicals, both synthetic fertilizers, and pesticides. In the management of pests and plant diseases, biological control is developed by applying biological agents including endophytic bacteria (Shanti and Vittal 2013). Many endophytic bacteria can pass the endodermic barrier across from the root cortex to the vascular system, and subsequently develop as endophytes in stems, leaves, tubers, and other organs (Compant et al. 2005). The use of endophytic bacteria as biological agents has an advantage compared to rhizosphere bacteria because

endophytic bacteria live and survive in the plant tissue during plant development, thus protecting the plants.

Bacillus sp. and fluorescent *Pseudomonas* are reported to be able to live as endophytes and are widely used as biological control agents for soil-borne and air-borne diseases. The endophytic bacteria could control plant diseases through several mechanisms including competition, hyperparasitism, producing microbial inhibiting compounds (antibiotics, lysis enzymes, other physical or chemical disorders), enhancing plant resistance, and promoting plant growth (Compant et al 2005, Pal and McSpadden-Gardener 2006; Rosenblueth and Martinez-Romero 2006).

Based on the mechanisms, the use of endophytic bacteria isolated from maize, both upland and lowland, suggested potentially alternative control for sheath blight (*R. solani*) and bacterial wilt (*Pantoea* sp). The research aimed to isolate and characterize morphologically and biochemically the endophytic bacteria as well as their potential to control pathogens that cause disease in maize especially *R. solani* and *Pantoea* sp.

MATERIALS AND METHODS

This research was conducted at the Laboratory of Plant Protection, Faculty of Agriculture, Jenderal Soedirman

University, Purwokerto, Central Java, Indonesia, from April to August 2019

Isolation *Rhizoctonia solani*

Rhizoctonia solani was isolated from maize with sheath blight symptoms and there was sclerotium as a resistant structure from the pathogenic fungi in Banyumas. *R. solani* isolation was carried out based on Al-Fadhal et al. 2019. Disease samples were cut 0.5 x 0.5 cm, then sterilized with NaOCl (1%) for 2 min, and rinsed with sterile water 3 times. Disease samples pieces were then dried using sterile filter papers, and transferred to Petri dishes containing PDA medium to obtain pure *R. solani* isolates.

Isolation *Pantoea* sp.

Pantoea sp. was isolated from diseased maize samples taken from the maize growing area in Banyumas Regency according to Coplin et al. (2012); Aini et al. (2013) and Desi et al. (2014). Diseased leaves or stems were washed with running water, then dried with a tissue. Diseased samples were cut 1.5 x 5 cm, then sterilized with ethanol 70% and rinsed with sterile water 3 times. Furthermore, the sample was crushed with 5 ml of sterile distilled water using a sterile mortar. The bacterial suspension was streaked on nutrient agar and incubated 3-5 days. Bacterial colonies that exhibit the character of *Pantoea* sp. were yellow, shiny, slimy, flat or convex, then separated as pure cultures of *P. stewartii* candidates. The culture was then tested by Gram Reaction (KOH test), Hugh-Leifson test, pigment production in YDC medium, oxidase test, hypersensitivity test, and pathogenicity test on maize.

Isolation and characterization of endophytic bacteria

Sampling for the isolation of endophytic bacteria was carried out in Banyumas and Purbalingga, Central Java, Indonesia, with purposive stratified random sampling. Samples were taken from two areas of altitude, i.e., low-moderate lands (0-500 m above sea level), and highlands (> 500 m above sea level) (Nuryanto et al., 2014). In each district, 2 locations were selected for the low-medium lands, and 1 location for the highlands. Age of maize plants was 20-30 days after planting, when the number of endophytic microbial populations that can be cultured is in the highest population (Cavaglieri et al. 2009).

The endophytic bacteria were isolated from the roots and stems of healthy maize plants. Roots and stems were washed, sterilized with 70% ethanol (1 minute), 20% sodium hypochlorite (5 minutes) and Ringer's thiosulfate solution (5 minutes). Separately, the roots and stems of 10 g each were crushed with 90 ml PBS on a sterile mortar. Subsequently, samples were plated on NA and Kings B media (Cavaglieri et al. 2009). To isolate *Bacillus* sp., the suspension was heated for 10 minutes at 80 ° C, before plating on NA. Bacterial isolates were further purified and characterized, such as morphological characteristics, gram properties, catalase tests, and hypersensitivity tests

The antagonism test of endophytic bacterial to *Rhizoctonia solani*

The antagonism test of endophytic bacteria on *R. solani* was carried out using the dual culture method. The level of inhibition of antagonist is calculated using the formula (Abidin et al. 2015).

$$I = \frac{C-T}{C} \times 100\%$$

Where:

I : The level of inhibition of antagonist (%)

C: The radius of pathogen colonies opposite antagonist

T: The radius of the colony of pathogens towards antagonist

The antagonism test of endophytic bacterial to *Pantoea* sp.

Antagonism testing was carried out using the double-layer test method (Santiago et al. 2015). Endophytic bacteria to be tested were grown on the nutrient agar medium, incubated at 28 °C for 48 hours. In the upside-down position, 1 ml of chloroform was added to the cup lid and left for 2 hours. Next, add 5 mL so that 0.6% water containing 0.5 mL of *Pantoea* sp. bacterial suspension. The culture was re-incubated for 24 hours, and there were clear zones around the antagonistic bacterial colony. The antibiotic activity was assessed based on the diameter of the clear zone compared to the diameter of the colony. Characterization of the type of antibiosis can be divided into bactericidal and bacteriostatic types based on Djatmiko et al. (2007).

The mechanism test of endophytic bacteria as controlling agents biological and plant growth-promoting microbial

The testing mechanism of endophytic bacteria was carried out for bacteria that have the potential in testing the antagonism of the fungus *R. solani* and *Pantoea* sp.

Protease test

The activity of the ability of antagonistic bacteria to produce extracellular protease enzymes was tested using Skim Milk Agar (SMA) medium. Each bacterium to be tested was grown in a medium SMA and incubated at 28 °C for 24-48 hours. The presence of clear zones around the colony shows that positive bacteria produce protease enzymes (Abed et al. 2016). The protease activity index is assessed based on the diameter of the clear zone compared to the diameter of the colony.

$$\text{Protease index} = \frac{(\text{clear zone diameter} - \text{colony diameter})}{\text{Colony diameter}}$$

Lipase test

Detection of the ability of bacteria to produce the enzyme lipase was done by growing the antagonistic bacteria on a medium containing 1% Tween 80. The presence of lipase enzyme activity was demonstrated by milky white sediment around the bacterial colony, after incubating at 28 °C for 4-7 days. The lipolytic index was measured using a formula Djuric et al. (2011).

$$\text{Lipolytic index} = \frac{(\text{Milky white diameter-colony diameter})}{\text{Colony diameter}}$$

Phosphatase test

Detection of the ability of bacteria to produce the enzyme phosphatase was done by growing bacterial isolates on Pikovskaya medium. After incubating for 7 days at 28 °C, the presence of a clear zone around the bacterial colony shows that the bacteria has the ability to produce the phosphatase enzyme to dissolve phosphates. The solubility index is measured using a formula (Farooq and Bano 2013)

$$\text{Phosphatase Index} = \frac{(\text{Clear zone diameter-Colony diameter})}{\text{Colony diameter}}$$

RESULTS AND DISCUSSION

Isolation and characterization of endophytic bacteria

The results of the exploration, isolation, and characterization of endophytic bacteria obtained 23 isolates of endophytic bacteria, consisted of 9 isolates of the fluorescent *Pseudomonas* and 14 isolates of *Bacillus* sp. (Table 1). Fluorescent *Pseudomonas* colony on King's B was round, with a flat edge, and yellowish-white, to greenish-yellow, gram-negative, rod-shaped, non-spore and fluorescent under ultraviolet light. Singh et al. 2017 reported fluorescent *Pseudomonas* showed light green, yellowish, creamy, circular, slimy, regular-irregular characteristics. Bacteria have short-long rod forms. The Fluorescent *Pseudomonas* isolates is gram-negative, which can form catalase, a positive oxidase, needed to grow aerobes. *Bacillus* sp. was observed with a spherical colony having cell rod-shaped, gram-positive, and endospores within cells (Table 1.). Slepecky and Hempill (2006); Amin et al. (2015) reported *Bacillus* sp. has the characteristics of a circular colony and punctiform (small round), variations in the entire margin and lobate, white dull, non-slimy, gram-positive, has endospores, flagellum and some are motile.

Based on its distribution, fluorescent *Pseudomonas* and *Bacillus* sp. found in all sampling locations, high or low-medium lands. This shows that fluorescent *Pseudomonas* and *Bacillus* sp. spread and can live in various altitudes, both high and low-medium land. Bacon and Hilton 2002; Ganeshan and Kumar 2005 reported *P. fluorescens* and *Bacillus* sp., are species of bacteria with a wide range of life and are very adaptive in various environments. Both types of bacteria are also found in the roots or corn stalks. *Bacillus* sp. and *Pseudomonas* sp. including a group of endophytic bacteria have a wide range of life and more

isolated in maize (Ganeshan and Kumar 2005; Orole and Adejumo 2011; Costa et al. 2013)

Antagonism test between the endophytic bacteria against *R. solani*

Based on the results of *in vitro* tests (Table 2), 24 isolates of the endophytic bacteria were able to inhibit the growth of *R. solani*, with varying degrees of inhibition. The endophytic bacteria that have inhibition rates above 50%, i.e. fluorescent *Pseudomonas* BK.A1 (51%), *Bacillus* sp. B.K.A1 (55.39%), *Bacillus* sp. BK.A3 (51.52%), PP.A5 (50.66%), and PPD.B2 (50.8%). The effect of the endophytic bacteria in inhibiting the growth of *R. solani* is inversely proportional to the dry weight mycelium. The greater the percentage of inhibition of endophytic bacteria to the growth of *R. solani*, the smaller the dry weight mycelium (Table 2) Endophytic bacteria can inhibit the growth of *R. solani*, which were shown by the inhibitory zone in the area bordering the bacterial streak (Figure 1.A).

The endophytic bacteria have anti-pathogenic properties and can produce antibiotic compounds. The ability of the endophytic bacteria to control plant pathogens occurs through the mechanism of antibiosis, competition, lysis, inducing resistance and producing growth substances. Bacteria capable of producing secondary metabolites that can inhibit growth or damage pathogens (Hastuti et al. 2014). These compounds, including alkaloids, phenols, flavonoids, glycosides, and phytoalexin (Soesanto et al. 2010). Fluorescent *Pseudomonas* can produce various types of antibiotics including phenazine-1-carboxylic acid, pyocyanin, pyrrolnitrin, and pyoluteorin, 2,4-diacetyl phloroglucinol (Phl). Phl is a phenolic metabolite with antibacterial and antifungal (Jain and Das 2016). *Bacillus* species can produce various kinds of volatile compounds and diffusible with strong inhibitory activity against plant pathogens (Lim et al. 2017).




Figure 1. Antagonism test between the endophytic bacteria against *R. solani* (A) and *Pantoea* sp. (B)

Table 1. Isolation and characterization of endophytic bacteria.

Land	Sampling location	Sample	Gram test	Catalase test	Oxidase test	Colony morphology	Colony pigment	Fluorescence on KB medium	Cell morphology	Endospores	Isolate
Highland	Purbalingga, Pratin 7.13'33" S, 109.17'21" E, 1.190 m asl	Root	-	+	+	Round	yellowish-white	+	Medium rod	-	fluorescent <i>Pseudomonas</i> PP.A1
		Root	+	+	+	Round	White	-	Small rod	+	<i>Bacillus</i> sp. PP.A3
		Root	+	+	+	Round	White	-	Medium rod	+	<i>Bacillus</i> sp. PP.A5
		Stem	-	+	+	Round	yellowish-white	+	Medium rod	-	fluorescent <i>Pseudomonas</i> PP. B4
	Banyumas, Baturaden 7.19"1" S, 109.14'29" E, 520 m asl	Root	-	+	+	Round	Greenish-yellow	+	Medium rod	-	fluorescent <i>Pseudomonas</i> BB.A2
		Root	+	+	+	Round	White	-	Small rod	+	<i>Bacillus</i> sp. BB.A3
		Stem	+	+	+	Round	White	-	Medium rod	+	<i>Bacillus</i> sp. BB.B4
Medium-Lowland	Banyumas, Sumbang 7.21'54" S, 109.17'33"E, 200 m asl	Root	-	+	+	Round	yellowish-white	+	Medium rod	-	fluorescent <i>Pseudomonas</i> BS.A2
		Root	+	+	+	Round	White	-	Medium rod	+	<i>Bacillus</i> sp. BS.A1
		Root	+	+	+	Round	White	-	Small rod	+	<i>Bacillus</i> sp. BS.A3
		Stem	+	+	+	Round	White	-	Small rod	+	<i>Bacillus</i> sp. BS. B1
	Purbalingga, Bojongsari 7.20'12" S, 109.20'22" E, 190 m asl	Root	-	+	+	Round	Greenish-yellow	+	Small rod	-	fluorescent <i>Pseudomonas</i> PB. A 4
		Stem	+	+	+	Round	White	-	Small rod	+	<i>Bacillus</i> sp. PB. B1
		Stem	+	+	+	Round	White	-	Medium rod	+	<i>Bacillus</i> sp. BB. B3
	Purbalingga, Padamara 7.22'28" S, 109.13'24" E, 180 m asl	Root	-	+	+	Round	Greenish-yellow	+	Small rod	-	fluorescent <i>Pseudomonas</i> PPD A1
		Stem	-	+	+	Round	yellowish white	+	Medium rod	-	fluorescent <i>Pseudomonas</i> PPD. B1
		Stem	-	+	+	Round	yellowish-white	+	Medium rod	-	fluorescent <i>Pseudomonas</i> PPD. B5
		Stem	+	+	+	Round	White	-	Small rod	+	<i>Bacillus</i> sp. PPD. B2
		Stem	+	+	+	Round	White	-	Small rod	+	<i>Bacillus</i> sp. PPD. B4
	Banyumas, Kembaran 7.23'47" S, 109.17'9" E, 110 m asl	Root	-	+	+	Round	yellowish-white	+	Medium rod	-	fluorescent <i>Pseudomonas</i> BK. A1
		Root	+	+	+	Round	White	-	Medium rod	+	<i>Bacillus</i> sp. BK.A1
		Root	+	+	+	Round	White	-	Medium rod	+	<i>Bacillus</i> sp. BK.A3
		Stem	+	+	+	Round	White	-	Medium rod	+	<i>Bacillus</i> sp. BK.B3

Table 2. Inhibition of endophytic bacteria against *R. solani*.

Isolate	Inhibition rate (%)	Dry weight mycelium
Control	0	0.093
Endophytic bacteria from the root		
fluorescent <i>Pseudomonas</i> BB.A2	49.00	0.038
fluorescent <i>Pseudomonas</i> BS.A 2	45.00	0.027
fluorescent <i>Pseudomonas</i> BK.A1	51.00	0.017
fluorescent <i>Pseudomonas</i> PPD.A1	10.33	0.059
fluorescent <i>Pseudomonas</i> PP.A1	38.33	0.017
fluorescent <i>Pseudomonas</i> PB.A4	18.00	0.037
<i>Bacillus</i> sp.BB.A3	40.42	0.030
<i>Bacillus</i> sp.BS.A1	48.73	0.016
<i>Bacillus</i> sp. BSA3	37.42	0.039
<i>Bacillus</i> sp. B.K.A1	55.39	0.002
<i>Bacillus</i> sp. B.K.A3	51.52	0.003
<i>Bacillus</i> sp.PP.A3	46.65	0.019
<i>Bacillus</i> sp.PP.A5	50.66	0.009
Endophytic bacteria from the stem		
fluorescent <i>Pseudomonas</i> PPD.B1	27.00	0.020
fluorescent <i>Pseudomonas</i> PPD.B5	49.33	0.013
fluorescent <i>Pseudomonas</i> PP.B4	65.67	0.004
<i>Bacillus</i> sp. BB.B4	44.44	0.026
<i>Bacillus</i> sp. BS.B1	49.74	0.012
<i>Bacillus</i> sp. BK. B3	40.36	0.031
<i>Bacillus</i> sp.PPD.B2	50.8	0.007
<i>Bacillus</i> sp. PPD.B4	39.44	0.036
<i>Bacillus</i> sp. PB.B1	37.29	0.047
<i>Bacillus</i> sp. PB.B3	44.9	0.022

The endophytic bacterial antagonism test against *Pantoea* sp.

The results of antagonism between the endophytic bacteria and *Pantoea* sp. show varied results. The endophytic bacteria that can inhibit bacterial growth were indicated by the presence of clear zones around the endophytic bacterial colonies (Figure1). From the nine isolate fluorescent, *Pseudomonas* were tested, only three isolates were able to inhibit the growth of the *Pantoea* sp., i.e fluorescent *Pseudomonas* (Pf) BS.A2, Pf BK.A1 Pf PPD.B5. While the isolates Pf BB.A2, Pf PPD.A1, Pf PP.A1, Pf PPD.B1, and Pf PP.B4 are not able to inhibit the growth of pathogenic bacteria. Meanwhile, all isolate *Bacillus* sp. tested (thirteen isolates) were able to inhibit the growth of *Pantoea* sp. (Table 3).

The presence of clear zones around endophytic bacterial colonies showed the ability of endophytic bacteria to produce antibiotics to inhibit the growth of *Pantoea* sp. *P. fluorescens* P60 can produce antibiotics that inhibit the growth of pathogens (Soesanto 2011). *Pseudomonas fluorescens* is reported to produce phenazine-1-carboxylic acid (PCA) and other derivatives, 2,4 diacetyl phloroglucinol (Phl), pyrrolnitrin (PRN) and or pyoluteorin (Plt) (Heydari and Pessarakli 2010). Nasrun and Burhanudin (2016) mention that *P. fluorescens* produce secondary metabolites, i.e. antimicrobial, cyanide acid and 2,4-diacetyl phloroglucinol phenazine, pyrrolnitrin, pyoluteorin antibiotics.


Table 3. Inhibition of endophytic bacteria against *Pantoea* sp.

Isolate	Antagonism	Antagonism index	Antagonism category*	Antagonism activity
Endophytic bacteria from the root				
fluorescent <i>Pseudomonas</i> BB.A2	-	0	-	-
fluorescent <i>Pseudomonas</i> BS.A 2	+	4.91	Strong	Bacteriostatic
fluorescent <i>Pseudomonas</i> BK.A1	+	4.42	Strong	Bacteriostatic
fluorescent <i>Pseudomonas</i> PPD.A1	-	0	-	-
fluorescent <i>Pseudomonas</i> PP.A1	-	0	-	-
fluorescent <i>Pseudomonas</i> PB.A4	+	5.29	Strong	Bactericidal
<i>Bacillus</i> sp.BB.A3	+	8.17	Strong	Bacteriostatic
<i>Bacillus</i> sp.BS.A1	+	4.00	Strong	Bacteriostatic
<i>Bacillus</i> sp. BSA3	+	5.07	Strong	Bactericidal
<i>Bacillus</i> sp. B.K.A1	+	4.01	Strong	Bacteriostatic
<i>Bacillus</i> sp. B.K.A3	+	4.91	Strong	Bacteriostatic
<i>Bacillus</i> sp.PP.A3	+	6.63	Strong	Bactericidal
<i>Bacillus</i> sp.PP.A5	+	6.56	Strong	Bactericidal
Endophytic bacteria from the stem				
fluorescent <i>Pseudomonas</i> PPD.B1	-	0	-	-
fluorescent <i>Pseudomonas</i> PPD.B5	+	5.86	Strong	Bactericidal
fluorescent <i>Pseudomonas</i> PP.B4	-	0	-	-
<i>Bacillus</i> sp. BB.B4	+	7.80	Strong	Bactericidal
<i>Bacillus</i> sp. BS.B1	+	6.22	Strong	Bacteriostatic
<i>Bacillus</i> sp. BK. B3	+	5.33	Strong	Bacteriostatic
<i>Bacillus</i> sp.PPD.B2	+	5.00	Strong	Bacteriostatic
<i>Bacillus</i> sp. PPD.B4	+	8.75	Strong	Bacteriostatic
<i>Bacillus</i> sp. PB.B1	+	1.67	Weak	Bacteriostatic
<i>Bacillus</i> sp. PB.B3	+	5.67	Strong	Bactericidal

Note: •Based on Davis and Stout (1971)

Table 4. Test results of proteases, lipases and phosphate solubilization

Isolate	Protease test		Lipase test		Phosphate solubilization	
	Activity	Index	Activity	Index	Activity	Index
Endophytic bacteria from the root						
<i>Bacillus</i> sp. B.K.A1	+	3.75	+	3.23	+	1.17
<i>Bacillus</i> sp. B.K.A3	+	3.20	+	3.73	+	1.27
<i>Bacillus</i> sp. PP.A5	+	5.00	+	4.40	+	1.46
Endophytic bacteria from the stem						
<i>Bacillus</i> sp.PPD.B2	+	3.00	+	3.90	+	2.60

**Figure 2.** Hydrolysis enzyme activity, (A) protease, (B) lipase and (C) phosphate solubilization

The level of bacteria's ability to inhibit growth can be shown by the large diameter of the clear zone. The results showed that the antagonism index ranged from 1.67-8.17. Based on this index, most endophytic bacteria have a strong antagonism (index of antagonism > 4) (Davis and Stout 1971). Furthermore, bacterial isolates that showed antagonistic activity were tested for types of antagonism based on Djatmiko (2007). Based on the type of antagonistic activity, ten isolates the endophytic bacteria were bacteriostatic and nine isolates the endophytic bacteria were bactericidal. Bacteriostatic activity, growth inhibition is temporary, it is shown that regrowth of bacteria after being transferred to a new medium, which is free from the influence of antagonistic bacteria. Bactericidal activity, inhibition is permanent. Bacteria were unable to grow even though they are transferred to new medium.

Test the mechanism of endophytic bacteria as biological control agents and plant growth-promoting microbes

The mechanism test was carried out on endophytic bacteria that have the potential to control the fungus *R. solani* and *Pantoea* sp., i.e. *Bacillus* sp. B.K.A1, *Bacillus* sp. B.K.A3, *Bacillus* sp. PP.A5, *Bacillus* sp. PPD.B2. The production of compounds related to biocontrol of pathogens and promotion of plant growth in bacterial isolates was evaluated by measuring the production of antimicrobial compounds and hydrolytic enzymes (amylases, lipases, proteases, and chitinases) and phosphate solubilization. The results of enzyme activity tests are as shown in Table 4. The four isolates *Bacillus* sp. tested were able to produce protease, lipase and phosphatase enzymes, with varied activity indexes. All isolates of *Bacillus* sp.

those tested had a high index of protease and lipase enzymes (> 3) (Table 4, Figure 2). Protease and lipase enzymes, related to the ability of the endophytic bacteria as biological control agents. Based on the protease and lipase indexes, *Bacillus* sp. PP.A5 can produce the highest proteases and lipase enzymes compared to other isolates. The isolates that have high protein and fat hydrolysis enzymes have the potential as biological control agents because proteins and fats are constituents of pathogen cells (Mota et al 2016). Besides, the protease enzyme is thought to degrade antibiotics produced by fungal or bacterial pathogens. According to Anderson et al. (2014), the extracellular protease enzyme produced by *P. fluorescens* can inactivate antibiotic compounds produced by *Pantoea agglomerans*.

Bacillus sp. PPD.B2 has the highest phosphate solubility index. The phosphate solubilization is related to the ability of endophytic bacteria as a plant growth promoter, providing phosphates for plants. Microbes with high phosphate solubility activity are capable of producing and releasing metabolites such as organic acids that chelate cations that are bound to phosphate (especially calcium) and converting them into soluble forms. Solubilization of different forms of phosphate by microbes associated with plants, and increasing its availability for plants, will increase growth and production of the plant (Djuric et al., 2011).

In conclusion, based on research carried out, it has been successfully isolated, morphologically and biochemically characterized four the endophytic bacteria that have the potential to be developed as biopesticides to control maize disease, especially *R. solani* and *Pantoea* sp. They can suppress the growth of *R.solani* by more than 50%, have a

strong antagonistic index against *Pantoea* sp (> 4), and can produce protease and lipase enzyme, and phosphate solubilization.

ACKNOWLEDGEMENTS

This research was a part of the main study funded by Doctoral Dissertation Grant from the Indonesian Ministry of Research, Technology and Higher Education; for that, I deeply thank for the financial support.

REFERENCES

- Abed HN, Rouag N, Mouatassef D, Rouabhi A. 2016. Screening for *Pseudomonas* and *Bacillus* antagonistic rhizobacteria strains for the biocontrol of fusarium wilt of chickpea. *Eurasian J Soil Sci* 5 (3): 182-191.
- Abidin Z, Aini LQ, Abadi AL. 2015. Effect of *Bacillus* sp. and *Pseudomonas* sp. On the growth of the pathogenic fungus *Sclerotium rolfsii* Sacc. causes of seedling diseases in soybean plants. *Jurnal HPT* 3 (1): 1-10.
- Aini LQ, Suryani L, Sugiharto AN, Abadi A. 2013. Identification of bacterial wilt and leaf blight disease on maize (*Zea Mays*) found in Kediri, Indonesia. *Agrivita* 35 (1): 1-7.
- Al-Fadhal FA, AL-Abedy AN, Alkhafijel DA. 2019. Isolation and molecular identification of *Rhizoctonia solani* and *Fusarium solani* isolated from cucumber (*Cucumis sativus* L.) and their control feasibility by *Pseudomonas fluorescens* and *Bacillus subtilis*. *Egypt J Biol Pest Control* 29 (47): 1-11. DOI: 10.1186/s41938-019-0145-5
- Amin M, Rakhisi Z, Ahmady AZ. 2015. Isolation and Identification of *Bacillus* Species From Soil and Evaluation of Their Antibacterial Properties. *Avicenna J Clin Microb Infec* 2 (1): e23233. DOI: 10.17795/ajcmi-23233.
- Ammar, E, Correa VR, Hogenhout SA, Redinbaugh MG. 2014. Immunofluorescence localization and ultrastructure of Stewart's wilt disease bacterium *Pantoea stewartii* in maize leaves and in its flea beetle vector *Chaetocnema pulicaria* (Coleoptera: Chrysomelidae). *J Microsc Ultrastructure* 2: 28-33.
- Anderson LM, Stockwell VO, Loper JE. 2004. An extracellular protease of *Pseudomonas fluorescens* inactivates antibiotics of *Pantoea agglomerans*. *Phytopathology* 94: 1228-1234
- Bacon CW, Hinton DM. 2002. Endophytic and Biological Control Potential Of *Bacillus Mojavensis* And Related Species. *Biol Control* 23: 274-284.
- Cavaglieri L, Orlando J, Etcheverry M. 2009. Rhizosphere microbial community structure at different maize plant growth stages and root locations. *Microbiol Res* 164 (4): 391-399.
- Compant SB, Duffy, Nowak J, Clement C, Barka EA. 2005. Use of plant growth-promoting bacteria for biocontrol of plant diseases: principles, mechanism of action, and future prospects. *Appl Environ Microbiol* 71 (9): 4951-4959.
- Coplin DL, Redinbaugh MG. 2012. The bacterium *Pantoea stewartii* uses two different Type III secretion systems to colonize its plant host and insect vector. *Appl Environ Microbiol* 78 (17): 6327-6336.
- Costa FG, Zucchi TD, de Melo IS. 2013. Biological control of phytopathogenic fungi by endophytic actinomycetes isolated from maize (*Zea mays* L.). *Braz. Arch Biol Technol* 56 (6): 948-955.
- Davis WW, Stout TR. 1971. Disc plate methods of microbiological antibiotic assay. *Appl Microbiol* 22 (4): 659-665.
- Desi Y, Habazar T, Agustian, Khairul U, Syamsuwirman, Novia P. 2014. Morphological and physiological characterization of *Pantoea stewartii* subsp. *stewartii* from maize. *Jurnal Fotopatologi Indonesia* 10 (2): 45-52. DOI: 10.14692/jfi.10.2.45. [Indonesian]
- Djaenuddin N, Nonci N, Muis A. 2017. Effectiveness of the formula *Bacillus subtilis* TM4 for disease control in maize plants. *Jurnal Fitopatologi Indonesia* 13 (4): 113-118. [Indonesian]
- Djarmiko HA, Arwiyanto T, Hadisutrisno B, Sunarminto BH. 2007. Potential of three bacterial genera from three plant rhizosphere as biological agents controlling Lincat disease. *Jurnal ilmu-ilmu Pertanian* 9 (1): 40-47. [Indonesian]
- Djuric S, Pavic A, Jarak M, Pavlovic S, Starovic M, Pivic R, Josic D. 2011. Selection of indigenous fluorescent pseudomonad isolates from maize rhizospheric soil in Vojvodina as possible PGPR. *Romanian Biotechnol Lett* 16 (5): 6580-6590. DOI: 10.1094/PHI-I-2004-0113-01.
- Farooq U, Bano A. 2013. Screening of indigenous bacteria from rhizosphere of maize (*Zea mays* L.) for their plant growth promotion ability and antagonism against fungal and bacterial pathogens. *J Anim Plant Sci* 23 (6): 1642-1652.
- Ganeshan G, Kumar AM. 2005. *Pseudomonas fluorescens*, a potential bacterial antagonist to control plant diseases. *J Plant Interact* 1 (3): 123-134.
- Hastuti RD, Saraswati R, Sari AP. 2014. The effectiveness of the endophytic microbes in promoting plant growth and controlling leaf blight disease in the lowland rice. *Jurnal Tanah dan Iklim* 38 (2): 109-118. [Indonesian]
- Heydari A, Pessarakhi M. 2010. A review on biological control of fungal plant pathogens using microbial antagonists. *J Biol Sci* 10 (4): 273-290
- Jain A, Das S. 2016. Insight into the interaction between plants and associated fluorescent *Pseudomonas* spp. *Intl J Agron*. DOI: 10.1155/2016/4269010
- Lim SM, Yoon M, Choi GJ, Choi YH, Jang KS, Shin TS, Park HW, Yu NH, Kim YH, Kim J. 2017. Diffusible and volatile antifungal compounds produced by an antagonistic *Bacillus velezensis* G341 against various phytopathogenic fungi. *Plant Pathol J* 33 (5): 488-498. DOI: 10.5423/PPJ.OA.04.2017.0073
- Motaa MS, Gomes CB, Júniora ITS, Moura AB. 2017. Bacterial selection for biological control of plant disease: criterion determination and validation. *Braz J Microbiol* 48: 62-70
- Muis A. 2007. Management of sheath blight disease (*Rhizoctonia solani* Kuhn.) in maize. *Jurnal Litbang Pertanian* 26 (3): 100-103 Desi 2014 [Indonesian]
- Nasrun, Burhanudin. 2016. Evaluation of the efficacy of the formula *Pseudomonas fluorescens* for controlling bacterial wilt (*Ralstonia solanacearum*) patchouli. *Buletin Penelitian Tanaman Rempah dan Obat*, 27 (1): 67-76. [Indonesian]
- Nuryanto B, Priyatmojo A, Hadisutrisno B. 2014. Pengaruh tinggi tempat dan tipe tanaman padi terhadap keparahan penyakit hawar pelepah. *Penelitian Pertanian Tanaman Pangan* 33 (1): 1-8. [Indonesian]
- Orole OO, Adejumo TO. 2011. Bacterial and fungal endophytes associated with grains and roots of maize. *J Ecol Nat Environ* 3 (9): 298-303.
- Pal KK, McSpadden-Gardener B. 2006. Biological Control of Plant Pathogens. *The Plant Health Instructor*. DOI: 10.1094/PHI-A-2006-1117-02
- Pataky JK. 2004. Stewart's wilt of corn. *The Plant Health Instructor*. DOI: 10.1094/PHI-I-2004-0113-01.
- Rosenbluth M, Martinez-Romero E. 2006. Bacterial endophytes and their interaction with hosts (Review). *Mol Plant Microbe Interact* 19 (8): 827-837.
- Santiago TR, Grabowski C, Rossato M, Romeiroa RS, Mizubuti ESG. 2015. Biological control of eucalyptus bacterial wilt with rhizobacteria. *Biol Contr* 80: 14-22. DOI: 10.1016/j.biocontrol.2014.09.007.
- Shanti AT, Vittal RR. 2013. Biocontrol potencies of plant growth promoting rhizobacteria against *Fusarium* wilt disease of cucurbit. *J Plant Pathol* 2 (3): 155-161.
- Singh S, Dutta U, Bhat AK, Gupta S, Gupta V, Jamwal S. 2017. Morphological and biochemical identification of *Pseudomonas* sp. isolated from the rhizosphere of different vegetable crops and study its efficacy on *Solanum melongena* (Brinjal). *J Pharmacogn Phytochem* 6 (2): 22-28
- Slepecky R.A., Hemphill H.E. 2006. The Genus *Bacillus*—Nonmedical. In: Dworkin M, Falkow S, Rosenberg E, Schleifer KH, Stackebrandt E. (eds.). *The Prokaryotes*. Springer, New York.
- Soesanto L, Mugiausti E, Rahayuniati RF. 2010. Study of the antagonistic mechanism of *Pseudomonas fluorescens* P60 against *Fusarium oxysporum* f.sp. *Lycopersici* in tomatoes in vivo. *Jurnal HPT Tropika* 10 (2): 108-115
- Soesanto L, Mugiausti E, Rahayuniati RF. 2011. Utilization of some animal broths as a liquid formula for *Pseudomonas fluorescens* P60 to control *Sclerotium rolfsii* in cucumber plants. *Jurnal Perlindungan Tanaman Indonesia* 17 (1): 7-17.



endang mugiasuti <endangmugiasuti@gmail.com>

[biodiv] Submission Acknowledgement

1 message

Ahmad Dwi Setyawan <smujo.id@gmail.com>

Thu, Nov 14, 2019 at 12:06 AM

Reply-To: Ahmad Dwi Setyawan <editors@smujo.id>

To: endang mugiasuti <endangmugiasuti@gmail.com>

endang mugiasuti:

Thank you for submitting the manuscript, "Isolation and Characterization of The Endophytic Bacteria, And Their Potential As Maize Diseases Control" to Biodiversitas Journal of Biological Diversity. With the online journal management system that we are using, you will be able to track its progress through the editorial process by logging in to the journal web site:

Submission URL: <https://smujo.id/biodiv/authorDashboard/submission/4844>

Username: endang_mugi

If you have any questions, please contact me. Thank you for considering this journal as a venue for your work.

Ahmad Dwi Setyawan

[Biodiversitas Journal of Biological Diversity](#)

Notifications

**[biodiv] Editor Decision**

2020-01-10 07:29 AM

endang mugiaastuti:

We have reached a decision regarding your submission to Biodiversitas Journal of Biological Diversity, "Isolation and characterization of the endophytic bacteria, and their potential as maize diseases control".

Our decision is: Revisions Required

Smujo Editors
editors@smujo.id

[Biodiversitas Journal of Biological Diversity](#)

Notifications

**[biodiv] Editor Decision**

2020-03-11 12:14 AM

endang mugiaastuti:

We have reached a decision regarding your submission to Biodiversitas Journal of Biological Diversity, "Isolation and characterization of the endophytic bacteria, and their potential as maize diseases control".

Our decision is: Revisions Required

Smujo Editors
editors@smujo.id

Reviewer A:
Recommendation: See Comments

[Biodiversitas Journal of Biological Diversity](#)

Notifications

**[biodiv] Editor Decision**

2020-04-07 05:40 AM

ENDANG MUGIASTUTI, SUPRAYOGI, NUR PRIHATININGSIH, LOEKAS SOESANTO :

We have reached a decision regarding your submission to Biodiversitas Journal of Biological Diversity, "Short Communication: Isolation and characterization of the endophytic bacteria, and their potential as maize diseases control".

Our decision is to: Accept Submission

Smujo Editors
editors@smujo.id

[Biodiversitas Journal of Biological Diversity](#)

Notifications

**[biodiv] Editor Decision**

2020-04-07 05:54 AM

ENDANG MUGIASTUTI, SUPRAYOGI, NUR PRIHATININGSIH, LOEKAS SOESANTO :

The editing of your submission, "Short Communication: Isolation and characterization of the endophytic bacteria, and their potential as maize diseases control," is complete. We are now sending it to production.

Submission URL: <https://smujo.id/biodiv/authorDashboard/submission/4844>

Smujo Editors
editors@smujo.id

[Biodiversitas Journal of Biological Diversity](#)