



FR-2021-533



Compose

Inbox

Snoozed

Sent

Drafts

Categories

More

Labels

Boxbe Waiting List

Manuscript ID: FR-2021-533



Food Research <foodresearch.my@outlook.com>
to me

Sat, Jul 17, 2021, 11:52 PM

Dear Professor Dr. Hery Winarsi,

This message is to acknowledge receipt of the above manuscript that you submitted via email to Food manuscript has been successfully checked-in. Please refer to the assigned manuscript ID number in any c with the Food Research Editorial Office or with the editor.

Your paper will be reviewed by three or more reviewers assigned by the Food Research editorial board ar made by the editor will be informed by email in due course. Reviewers' suggestions and editor's comme made available via email attached file. You can monitor the review process for your paper by emailing us c my manuscript".



FR-2021-533



Compose

Inbox

Snoozed

Sent

Drafts

Categories

More

Labels

Boxbe Waiting List

Please refer to the attachment for the galley proof of your manuscript FR-2021-533 entitled 'Imprc of immune and antioxidant status of COVID-19 vulnerable groups using mung bean sprout yoghu Please check the content of the galley proof. If there are any mistakes, please comment and high the PDF itself and revert to us within two (2) days of receipt. Once we have finalized the PDF vers manuscript will be published online for early viewing.

Please see the attachment for the invoice INV22043. We hope that you can make the payment as possible before 15 March 2022 for us to complete the publication of your manuscript. The manus information e.g. volume, issue, page numbers and DOI, will be provided once we have received th payment.

Thanks & Regards,
Vivian New
Editor



FR-2021-533



Compose

Inbox

Snoozed

Sent

Drafts

Categories

More

Labels

Boxbe Waiting List

Re: FR-2021-533 - Article Production

Inbox x



Food Research <foodresearch.my@outlook.com>
to me

Sun, Feb 20, 2022, 1:50 PM

Dear Dr Winarsi,

Manuscript ID: FR-2021-533

Manuscript Title: Improvement of immune and antioxidant status of Covid-19 vulnerable groups using mung
yoghurt

Before we can proceed with the article production, I would like to clarify a few points that I have commented
manuscript. Please refer to the attachment. Please address the issues raised in the comments.

Please use the attached copy to make your revisions as it has been corrected to the Journal's format. Once



FR-2021-533



Compose

Inbox

Snoozed

Sent

Drafts

Categories

More

Labels

Boxbe Waiting List

--
Prof. Dr. Ir. Hery Winarsi, MS
Department of Nutrition Science, Faculty of Health Sciences,
University of Jenderal Soedirman, Purwokerto, 53123.
Central Java, Indonesia.

Reply

Forward

19th February 2022

Dear Dr Winarsi,

ACCEPTANCE LETTER

Food Research is pleased to inform you that the following manuscript has been accepted for publication in Food Research journal.

Manuscript Title : Improvement of immune and antioxidant status of Covid-19 vulnerable groups using mung bean sprout yoghurt

Authors : Winarsi, H., Erminawati, E. and Andreas, A.

We thank you for your fine contribution to the Food Research journal and encourage you to submit other articles to the Journal.

Yours sincerely,



Professor Dr. Son Radu
Chief Editor
Food Research

Improvement of immune and antioxidant status of COVID-19 vulnerable groups using mung bean sprout yoghurt

^{1,*}Winarsi, H., ²Erminawati, E. and ³Andreas, A.

¹Department of Nutrition Science, Faculty of Health Sciences, Universitas Jenderal Soedirman, Purwokerto, 53123, Central Java, Indonesia

²Department of Food Technology, Faculty of Agricultural, Universitas Jenderal Soedirman, Purwokerto, 53123, Central Java, Indonesia

³Sinar Kasih General Hospital, Purwokerto, Indonesia

Article history:

Received: 18 September

2021

Received in revised form: 20

October 2021

Accepted: 19 February 2022

Available Online: 16 March

2022

Keywords:

Raise immune level,

CRP,

Mung bean sprout yoghurt,

SOD

DOI:

[https://doi.org/10.26656/fr.2017.6\(2\).533](https://doi.org/10.26656/fr.2017.6(2).533)

Abstract

This study aimed to determine the effect of mung bean sprout yoghurt (MungbeS-Yo) on plasma CRP and SOD levels of healthcare workers. MungbeS-Yo was prepared by mixing green bean sprout milk and skim milk, sucrose, and Lactic Acid Bacteria (LAB). The subjects of this study were 20 female healthcare workers, randomly divided into 2 groups. Group-1 was given MungbeS-Yo, while group-2 was given Skim-Yo (as control), as much as 200 mL/day for 2 months. MungbeS-Yo is prepared from 70 parts of mung bean sprouted milk, 30 parts skim, 0.5% LAB, and 10% sugar. Blood samples were taken before and after the intervention. Blood plasma was tested for CRP and SOD levels. CRP levels decreased from 5.41 to 3.61 mg/L ($P = 0.037$), while SOD levels increased from 88.86 pg/mL to 176.05 pg/mL ($P < 0.044$). MungbeS-Yo has the potential to improve the immune status and enzymatic antioxidant status of plasma susceptible groups infected with Covid-19. Thus, MungbeS-Yo can help raise the immune level against infection.

1. Introduction

Healthcare workers are the front line in handling COVID-19 patients. Therefore, healthcare workers are required to have excellent physical and psychological conditions. However, the high workload and the anxiety of being infected, create a big stressor. Stress is closely related to a decrease in immune status, a self-defence response against viruses. Nakata (2012) and Maydych (2019) added that stress affects the inflammatory response, characterized by increased levels of C-Reactive Protein (CRP). Shivpuri *et al.* (2012) reported that interpersonal stress is greater in women than men are. Therefore, it is very reasonable if female healthcare workers are one of the vulnerable groups for COVID-19. It has been reported that 32.9% of healthcare workers were infected with Covid-19 due to direct contact with patients and 22.7% were infected due to direct contact with co-workers of healthcare workers (Huang *et al.*, 2020; Vandercam *et al.*, 2020). Considering the number of Covid-19 sufferers is still increasing, it is important to make efforts to improve the immune status of healthcare workers, they can continue their services to patients.

Currently, the right medicine has not been found to

defend the body during a pandemic like today. Possibly, going back to nature is a wise step to overcome this. Some experts say that immune status is closely related to antioxidant status. Therefore, the role of antioxidant-rich food products being able to overcome oxidative stress by increasing the activity of antioxidant enzymes and improving the performance of the immune system by suppressing inflammatory markers (Middleton *et al.*, 2000; Winarsi *et al.*, 2010; Winarsi *et al.*, 2012; Winarsi *et al.*, 2013; Winarsi *et al.*, 2016).

One type of legume that is rich in antioxidants is green beans. Besides being known to be rich in nutrients, green beans also contain oligosaccharides, which often cause flatulence. To minimize the levels, it is necessary to soak and germinate. Germination reported Winarsi *et al.* (2019) can remove beany flavour, increase antioxidant content, digestibility (Sokrab *et al.*, 2012) and total phenolic content (Guo *et al.*, 2012; Winarsi *et al.*, 2020) whose antioxidant activity can protect the body from oxidative stress (Winarsi *et al.*, 2016). The activity of protease, lipase, and carbohydrase increases, causing the content of amino acids, fatty acids, and glucose in sprouts to be higher than in ungerminated

*Corresponding author.

Email: winarsi12@gmail.com

beans. Simple compounds such as glucose, amino acids, fatty acids, vitamins, and minerals are important nutrients for the growth of lactic acid bacteria (LAB), a bacterial species in the manufacture of yoghurt. Mung bean sprouted yoghurt prepared from mung bean sprouts and skim milk, which is fermented using LAB, contains 1,273.11 mg GAE/L, 41.40% protein, and 12.17% fibre, and *L. acidophilus* in MungbeS-Yo as much as 6.56×10^7 CFU/mL, it is called rich functional drink antioxidants and probiotics (Winarsi et al., 2021).

The problem is, can MungbeS-Yo improve the immune status and antioxidant status of healthcare workers? This study aimed to explore the benefits of MungbeS-Yo on improving the immune status and antioxidant status of healthcare workers.

2. Materials and methods

This study used a true experimental design. This study has gained ethical approval from the Komisi Etik Penelitian Kesehatan (KEPK) of the Faculty of Health Sciences, Jenderal Soedirman University, with a reference number: 253/EC/KEPK/I/2021.

2.1 Production of MungbeS-Yo

Mung beans were germinated for 10 hrs, blended until a smooth liquid was obtained called mung bean sprout milk. Mung bean sprout yoghurt was prepared by mixing green bean sprout milk and skim milk in a ratio of 70:30. As much as 10% of sucrose was added and pasteurized at 80°C for 10 mins, cooled to 45°C, then inoculated with 0.5% LAB starter of the total volume, and incubated at room temperature of 27-35°C for 24 hrs (Winarsi et al., 2021).

2.2 Research subjects

A total of 20 female healthcare workers became the research subjects with the criteria as followed aged 30-50 years old, had central obesity, inflammation, and oxidative stress did not leave the research site for two months and were willing to sign informed consent. The subjects were randomly divided into two groups, with a total of ten people in each group.

2.3 MungbeS-Yo intervention to research subjects

The first group received a MungbeS-Yo as much as 200 mL/d of regularly for two months, while the second group was given a placebo (Skim-Yo).

2.4 Blood sampling

Blood samples were taken twice (0 and 2 months after the intervention), with a total of 3 mL samples for each blood sampling. The blood sample was centrifuged

at 3,000 rpm for 10 mins, and the plasma portion was separated from the erythrocytes. Blood plasma was examined to determine the SOD levels (using Abbkine SOD ELISA) and CRP levels (using Calbiotech High Sensitivity C-Reactive Protein CRP ELISA).

2.5 Data analysis

Data were analyzed with the help of the SPSS Statistics 17.0 program. All data were tested for Shapiro-wilk normality. Differences in plasma CRP and SOD levels before and after the intervention were analyzed using the Paired t-test.

3. Results and discussion

Subject characteristics include a description of age, BMI, CRP and SOD levels initial (Table 1). Healthcare workers have an age range between 30-50 years, are obese, and have CRP levels above normal. According to Kratz et al. (2012), the normal threshold for CRP levels is 0.2 – 3.0 mg/L. This condition proves that healthcare workers experience inflammation, and it is possible that the levels will continue to increase, due to frequent contact with various patients, including Covid-19 sufferers in their work area. CRP levels are generally not associated with viral infections, but in fact, high CRP levels are closely related to the length of hospitalization of patients (Jeon et al., 2017). This proves the important role of CRP levels in respiratory viral infections, such as the coronavirus that attacks the respiratory tract.

Table 1. Subject characteristics

Variable	MungbeS-Yo	Skim-Yo	p*
	(n = 10)	(n = 10)	
	mean±SD	mean±SD	
Age (y)	38.86±3.49	38.14±4.88	0.758
BMI (kg/m ²)	30.96±3.87	28.61±4.16	0.297
CRP level (mg/L)	5.41±1.93	6.31±2.99	0.559
SOD level (pg/mL)	88.87±16.19	96.15±25.42	0.239

*Independent t-test. MungbeS-Yo: the group that was given mung bean sprout yogurt, Skim-Yo: the group given skim yogurt as a control

CRP levels are also used as a prognostic marker for pneumonia in COVID-19. The plasma CRP concentration increased to 11.47 mg/L in the mild infection group and even increased to 23.4 mg/L in the moderate to severe infection group. Elevated CRP levels are used as an early indicator of nosocomial infection and help detect the severity of COVID-19 regardless of age (Chen et al., 2020). CRP is considered a biomarker that easy and inexpensive to measure, compared to other markers (Armani and Becker, 2005). A decrease in CRP levels indicates an improvement in immune status.

Among inflammatory markers, CRP is the most

sensitive marker that can predict future cardiovascular disease events (Chen *et al.*, 2010). Various factors influence inflammatory markers, including nutrition, antioxidants, physical fitness and exercise. Physical activity has been reported to reduce the inflammatory marker CRP (Meyer *et al.*, 2007; Pyne and Pyne, 2013). According to Sproston and Ashworth (2018), CRP is the most sensitive marker of acute inflammation, because its levels can increase up to 1000 times during infection. CRP synthesized by hepatocytes, in response to increased levels of IL-6, IL-1, and TNF- released by inflamed tissue. CRP plays a key role in activating the complement system, inducing the expression of adhesion molecules, increasing phagocytosis and macrophages, as well as leukocyte activation and opsonization (Armani and Becker, 2005).

CRP levels in blood plasma will increase more than normal levels, after 6 hrs of infection, and even reach their highest levels at 50 hrs after trauma, infection or inflammation. CRP levels decreased exponentially after 18-20 hrs later (Ridker, 2003; Kaski, 2010). High serum CRP levels are common in viral infections, as well as SARS, which is the closest human disease to COVID-19. IL-6 production is also high, even IL-6 production due to SARS is higher than other viruses in general, such as influenza (Okabayashi *et al.*, 2006). Therefore, blockade of cytokines or pro-inflammatory receptors is therapeutic in COVID-19 (Shakoory *et al.*, 2016). Some researchers mention that functional foods containing antioxidants can reduce plasma CRP levels (Winarsi *et al.*, 2016). It was also reported that soybean sprouted milk can reduce CRP levels in pregnant women to reach 74.34%, because of the antioxidant role of isoflavones in it.

3.1 Effect of MungbeS-Yo on healthcare workers plasma CRP levels

After 2 months of intervention, CRP levels in the MungbeS-Yo group decreased from 5.41 to 3.61 mg/L ($P = 0.037$) (Figure 1), compared to the control ($P = 0.13$). The same finding was reported by Salehzadeh (2015) that yoghurt given to athletes could reduce CRP levels. Winarsi *et al.* (2021) reported that MungbeS-Yo contained 1,273.11 mg GAE/L phenolic, 41.40% protein, and 12.17% fibre.

As antioxidant compounds, phenolics reduce the activity of reactive compounds (Amaretti *et al.*, 2013), chelate transition metal ions, inhibit several enzymes associated with ascorbate autoxidation (Talwalkar and Kailasapathy, 2003), reduce oxidant compounds in the intestine (Azcarate-Peril *et al.*, 2011), induces the transcription of genes involved in glutathione biosynthesis in the intestinal mucosa (Lutgendorff *et al.*, 2009), increases glutathione synthesis in pancreatic cells,

and stimulates the action of enzymes in the intestinal microflora (Lutgendorff *et al.*, 2008). MungbeS-Yo with its protein and fiber contents considered useful as a product to support the immune system during the Covid-19 pandemic. Phenolic acids are simple compounds that play a role in reducing the risk of chronic diseases, various types of cancer (Reboredo-Rodríguez *et al.*, 2018), and various viruses (Kaila *et al.*, 1994).




Figure 1. Effect of mung beans sprout yogurt on CRP plasma levels of healthcare workers. Values followed by the same superscript are statistically not significant with $\alpha = 5\%$. MungbeS-Yo: the group that was given mung bean sprout yogurt, Skim-Yo: the group given skim yogurt as a control.

Generally, a product is considered a probiotic, if it contains good bacteria as much as 10^6 - 10^7 CFU/mL. In this study, MungbeS-Yo contained LAB as much as 6.56×10^7 CFU/mL (Winarsi, 2021). Therefore, MungbeS-Yo deserves to be considered as a probiotic drink. Probiotics are antimicrobials, which can increase lactose absorption, stimulate the immune system, treat and prevent allergies, prevent bacterial infections in preterm birth, inflammatory bowel disease, bladder and ear infections, tooth decay, chronic diarrhoea and travel diarrhoea. Beneficial bacteria of *Lactobacillus*, *Streptococcus thermophilus* and *Vulgaris*, which are often found in yoghurt, can improve the digestive system and have a major impact on health (Salarkia *et al.*, 2013). These bacteria also inhibit the growth and multiplication of pathogenic bacteria.

The decrease in plasma CRP levels may be related to the role of phenolic compounds in MungbeS-Yo. According to Winarsi *et al.* (2016), isoflavones (a form of phenolic) in sprouted soybean milk is capable to improve the CRP of pregnant women after 2 months of intervention. Isoflavones can suppress plasma IL-6 production in lactating mothers who are given sprouted soybean milk. IL-6 is a pro-inflammatory cytokine, which is secreted by immune cells and other cells that experience inflammation in response to the body's defence against foreign substances. IL-6 stimulates hepatocyte CRP production (Tanaka and Kishimoto, 2014). Once produced, CRP is then released into the inflamed tissue or organ. At the site of inflammation,

CRP performs a defence mechanism by activating the classical complement pathway, bacterial opsonization and complement-mediated phagocytosis (Sproston and Ashworth, 2018). Phenolics reduce CRP levels by inhibiting IL-6 regulation. The inhibition of IL-6 regulation has an impact on suppressing CRP production, and the levels decrease.

In addition to phenolics, MungbeS-Yo probiotics also have the effect of improving the level of inflammation. According to Duncan and Flint (2013), probiotics are live microbes which when given in adequate amounts can be beneficial for the health of the digestive tract (Antoine, 2010). The probiotic bacterial species in yoghurt are *Lactobacillus bulgaricus*, *Lactobacillus acidophilus* and *Streptococcus thermophilus*, used as a starter in the yoghurt fermentation process. It was reported that the amount of LAB in MungbeS-Yo was 6.56×10^7 CFU/mL (Winarsi et al., 2021), it is suspected that these bacteria contributed to suppressing CRP levels.

Giving probiotics early in life accelerates the formation of a balance of intestinal microflora (Kabir, 2009), thus supporting the formation of a good immune system in the intestinal lumen (Jeppsson et al., 2004). Probiotics are also capable to modulate physiological characteristics in the digestive tract, such as mucosal immunity, mucosal trophicity, and intestinal permeability (Fioramonti et al., 2003). Supplementation of probiotics can stimulate the production of short-chain fatty acids (SCFA), such as butyric acid, which plays an important role in the proliferation of intestinal epithelial cells. Increased production of butyric acid can help expand the absorption area in the intestinal lumen, so that nutrient absorption is more optimal (Ohashi and Ushida, 2009).

SCFA production requires cooperation between organic acid-producing bacteria and other bacteria in the gut, through a cross-feeding mechanism (Van Immerseel et al., 2006). These bacteria hydrolyze undigested carbohydrates into monosaccharides, through anaerobic fermentation in the intestine. The monosaccharides formed are then converted into phosphoenolpyruvate (PEP) via the Embden-Meyerhof-Parnas pathway (glycolysis) and pentose-phosphate, which are then converted into organic acids (Besten et al., 2013). The absorbed SCFA is the main energy source for enterocytes, so it can help meet the energy needs of the host.

According to Hegazy and El-Bedewy (2010), SCFA produced by probiotics can reduce plasma CRP levels, by blocking enzyme synthesis from liver CRP. CRP synthesized by the liver is a response to release factors

by fat cells, such as IL-6 (Kinoshita et al., 2015). In their study, Hegazy and El-Bedewy (2010) observed that the consumption of probiotics in patients with ulcerative colitis for 8 weeks, can significantly improve inflammation, by reducing IL-6 levels, TNF- expression, and NF- κ B. The decrease in the concentration of IL-6 can indirectly lead to the reduction of CRP production. Decreased CRP concentration is associated with reduced expression of inflammatory factors, in other words, an improvement in immune status.

The plasma C-RP is a marker of systemic inflammation, whose levels are elevated by chronic conditions, including obesity and high waist circumference. In this study, subjects were obese, as indicated by a large BMI > 30 kg/m² and had a waist circumference of 93.72 cm. However, BMI decreased significantly from 32.99 to 28.95 kg/m² (P = 0.043), and waist size also decreased from 93.72 to 89.43 Cm (P = 0.044). Thus, the decrease in plasma CRP was supported by a decrease in the subjects' BMI and waist size.

The anti-inflammatory properties of some probiotic strains are thought to work by reducing mucosal inflammation by modulating levels of cytokines and other inflammatory mediators (Choi et al., 2006). Studies on the consumption of probiotic yoghurt containing *L. acidophilus* and *Bifidobacterium* in pregnant women for nine weeks led to a decrease in CRP sensitivity. This reduction in inflammation due to the probiotic MungbeS-Yo may be related to its effect on increasing levels of SOD, which scavenges superoxide and hydroxyl radicals, thereby suppressing plasma CRP expression.

3.2 Effect of MungbeS-Yo on plasma SOD levels of healthcare workers

Subjects in this study were obese if their BMI was 25 kg/m² (WHO, 2000). According to Winarsi et al. (2012), obese women experience high oxidative stress in their bodies. In obese conditions, the wider adipose tissue can lead to hypoxic conditions (O₂ deficiency). Chronic hypoxia increases oxidative stress by producing excessive free radicals without compensating for the activity of antioxidant enzymes. The higher the level of oxidative stress in obese patients, the lower the activity of the SOD enzyme. During a pandemic, healthcare workers are at risk of being exposed to the virus due to direct contact with various kinds of patients. Baseline data showed that plasma SOD levels were low. Several researchers reported that antioxidant-rich products could improve the antioxidant activity of SOD (Harun et al., 2017; Winarsi et al., 2016; Setiawan et al., 2016). According to Winarsi et al. (2021), MungbeS-Yo contains phenolic antioxidants of 1,107.80 mg GAE/L, which is expected to increase plasma SOD levels of

healthcare workers.

After two months of administration of MungbeS-Yo, there was a significant increase in SOD levels from 88.86 pg/mL to 176.05 pg/mL ($p = 0.044$), as did the Skim-Yo group but not significantly ($p = 0.590$) (Figure 2). The increase in SOD levels in this study concurred with the findings of Setiawan *et al.* (2016), who had given sprout soybean juice at a dose of 2.12 g/d for 28 days in rats. This is due to the presence of phenolic compounds that have antioxidant abilities. In this study, subjects were given 200 mL of yocamio, which contains 221.56 mg of phenolic GAE (Winarsi *et al.*, 2021). Rahmawati *et al.* (2014) proved that the administration of a product with a flavonoid content of 21.46 mg/g in 14 days could increase the SOD activity of rats. Winarsi *et al.* (2016) also reported that an increase in plasma SOD also occurred in the provision of sprouted soybean milk for 2 months containing 39.1 ppm isoflavones. Isoflavones are able to increase SOD levels by increasing SOD activity, suppressing free radical activity by binding to reactive oxygen species (ROS) and converting them into isoflavone radicals (Winarsi *et al.*, 2016). Thus, the phenolic action of MungbeS-Yo in increasing plasma SOD levels is the same as that of isoflavones.




Figure 2. Effect of Mung beans sprout yogurt on SOD plasma levels of healthcare workers. Values followed by the same superscript are statistically not significant with $\alpha = 5\%$.

MungbeS-Yo: the group that was given mung bean sprout yogurt, Skim-Yo: the group given skim yogurt as a control.

Mung bean sprout yoghurt is a probiotic drink that has many health benefits. The *L. acidophilus* content in MungbeS-Yo is 6.56×10^7 CFU/mL (Winarsi *et al.*, 2021). This is in accordance with the SNI standard (2009) that fermented milk is said to be a probiotic drink if the number of LAB reaches 10^7 (Utami *et al.*, 2017). Wang *et al.* (2017) reported that athletes under conditions of high physical stress and oxidative stress can be neutralized by administering *Lactobacillus rhamnosus* because probiotic bacteria can suppress the production of free radicals, and this is reflected by increasing antioxidant levels in the body. Utami *et al.* (2017), reported that the administration of probiotic bacteria *Lactobacillus casei* at a dose of 10.9×10^7 cells/

kgBW/d for 2 weeks increased SOD activity by 50.7%. Probiotics are reported to reduce oxidative stress by neutralizing oxidants in the gastrointestinal tract expressed in antioxidant enzyme activity, stimulating the immune system by suppressing cytokine-induced inflammation and preventing oxidative stress and reducing intestinal pathogenic bacteria (Kleniewska *et al.*, 2016).

MungbeS-Yo contains 17.26% Vitamin C (Winarsi *et al.*, 2021). Based on the age and gender of the subjects, the amount of Vitamin C needed in a day is 75 mg. Vitamin C is a natural antioxidant that works synergistically to neutralize free radicals (Suhail *et al.*, 2011). Several studies have shown that vitamin C supplementation can improve the activity of the SOD enzyme (Suhail *et al.*, 2011; Boudouris *et al.*, 2013; Rafighi *et al.*, 2013), it is possible that the Vitamin C contained in MungbeS-Yo also plays a role in increasing SOD levels of healthcare worker plasma. Mung bean sprouts are the raw material for making yoghurt. Thus, the product obtained by LAB fermentation makes food products have therapeutic benefits, including antioxidant and immunostimulant activity, prophylaxis (preventing infection) against several types of intestinal infections, increasing tolerance to foods containing lactose, and preventing cancer.

4. Conclusion

MungbeS-Yo is a food product fermented by LAB which has the potential as an immune-stimulant by suppressing the level of inflammation, reducing CRP levels by 33.27%. MungbeS-Yo is rich in phenolic antioxidants; which have been shown to increase SOD antioxidant levels by 98.01%. Thus, it is believed that MungbeS-Yo is a probiotic drink product that is useful for improving the immune system of healthcare workers who are susceptible to exposure to COVID-19. Therefore, MungbeS-Yo can help raise the immune level against infection.

Acknowledgement

The author offered the deepest gratitude towards the Education Fund Management Institution from the Ministry of Finance of the Republic of Indonesia, through *Lembaga Penelitian dan Pengabdian Masyarakat* (LPPM) Chairman for funding this research in the skim of covid prevention, as well as the facilities. To the four students who helped with this research, namely Kristina Citra Palimirma, Rosyalya Alhanifa, Nur Amalia Lael, and Injih Prastiwi.

References

- Armani, A. and Becker, R.C. (2005). The Biology, Utilization, and Attenuation of C-Reactive Protein in Cardiovascular Disease. *American Heart Journal*, 149(6), 971-976. <https://doi.org/10.1016/j.ahj.2004.12.031>
- Amaretti, A., di Nunzio, M., Pompei, A., Raimondi, S., Rossi, M. and Bordoni, A. (2013). Antioxidant properties of potentially probiotic bacteria: *in vitro* and *in vivo* activities. *Applied Microbiology and Biotechnology*, 97(2), 809-817. <https://doi.org/10.1007/s00253-012-4241-7>
- Antoine, J.M. (2010). Probiotics in the defence and metabolic balance of the organism Probiotics: beneficial factors of the defence system. The 3rd International Immunonutrition Workshop. *Proceedings of the Nutrition Society*, 69(3), 429-433. <https://doi.org/10.1017/S0029665110001692>
- Azcarate-Peril, M.A., Sikes, M. and Bruno-Barcena, J.M. (2011). The intestinal microbiota, gastrointestinal environment and colorectal cancer: a putative role for probiotics in prevention of colorectal cancer? *American Journal of Physiology - Gastrointestinal and Liver Physiology*, 301(3), G401-G424. <https://doi.org/10.1152/ajpgi.00110.2011>
- Besten, G.D., Eunen, K.V., Groen, A.K., Venema, K., Reijngoud, D.J. and Bakker, B.M. (2013). The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism. *Journal of Lipid Research*, 54(9), 2325- 2340. <https://doi.org/10.1194/jlr.R036012>
- Boudouris, G., Verginadis I.I. and Simos, Y.V. (2013). Oxidative stress in patients treated with continuous ambulatory peritoneal dialysis (CAPD) and the significant role of vitamin C and E supplementation. *International Urology and Nephrology*, 45(4), 1137-1144. <https://doi.org/10.1007/s11255-012-0334-6>
- Chen, W., Zheng, K. I., Liu, S., Yan, Z., Xu, C. and Qiao, Z. (2020). Plasma CRP level is positively associated with the severity of COVID-19. *Annals of Clinical Microbiology and Antimicrobials*, 19(18), 1-7. <https://doi.org/10.1186/s12941-020-00362-2>
- Chen, G. and Nuñez, G. (2010). Sterile inflammation: sensing and reacting to damage. *Nature Reviews Immunology*, 10(12), 826-837. <https://doi.org/10.1038/nri2873>
- Choi, S.S., Kim, Y., Han, K.S., You, S., Oh, S. and Kim, S.H. (2006). Effects of Lactobacillus strains on cancer cell proliferation and oxidative stress *in vitro*. *Letters in Applied Microbiology*, 42(5), 452-458. doi:10.1111/j.1472-765x.2006.01913.x
- Duncan, S.H. and Flint, H.J. (2013). Probiotics and prebiotics and health in ageing populations. *Maturitas*, 75(1), 44-50. <https://doi.org/10.1016/j.maturitas.2013.02.004>
- Fioramonti, J., Theodorou, V. and Bueno, L. (2003). Probiotics: what are they? What are their effects on gut physiology? *Best Practice and Research Clinical Gastroenterology*, 17, 711-724. [https://doi.org/10.1016/S1521-6918\(03\)00075-1](https://doi.org/10.1016/S1521-6918(03)00075-1)
- Guo, X., Li, T., Tang, K. and Liu, R.H. (2012). Effect of germination on phytochemical profiles and antioxidant activity of mung bean sprouts (*Vigna radiata*). *Journal of Agricultural and Food Chemistry*, 60(44), 11050-11055. <https://doi.org/10.1021/jf304443u>
- Harun, I., Susanto, H. and Rosidi, A. (2017). Pemberian tempeh menurunkan kadar malondialdehide (MDA) dan meningkatkan aktivitas enzim superoxide dismutase (SOD) pada tikus dengan aktivitas fisik tinggi. *Jurnal Gizi Pangan*, 12(3), 211-216. <https://doi.org/10.25182/jgp.2017.12.3.211-216> [In Bahasa Indonesia].
- Hegazy, S.K. and El-Bedewy, M.M. (2010). Effect of probiotics on pro-inflammatory cytokines and NFkappaB activation in ulcerative colitis. *World Journal of Gastroenterology*, 16(33), 4145-4151. <https://doi.org/10.3748/wjg.v16.i33.4145>
- Huang, L., Lin, G., Tang, Li., Yu, L. and Zhou, Z. (2020). Special attention to nurse protection during the COVID-19 epidemic. *Critical Care*, 24(120), 1-3. <https://doi.org/10.1186/s13054-020-2841-7>
- Jeon, J. S., Rheem, I. and Kim, J.K. (2017). C-Reactive Protein and respiratory viral infection. *Korean Journal of Clinical Laboratory Science*, 49(1), 15-21. <https://doi.org/10.15324/kjcls.2017.49.1.15>
- Jeppsson, B., Mongell, P., Adawi, D. and Molin, G. (2004). Bacterial translocation: impact of probiotics. *Scandinavian Journal of Social Medicine*, 48(1), 37-41. <https://doi.org/10.1080/11026480410026816>
- Kabir, S.M.L. (2009). The role of probiotics in the poultry industry. *International Journal of Molecular Sciences*, 10(8), 3531-3546. <https://doi.org/10.3390/ijms10083531>
- Kaila, M., Arvilommi, H., Soppi, E., Laine, S. and Isolauri, E. (1994). A prospective study of humoral immune responses to cow milk antigens in the first year of life. *Pediatric Allergy and Immunology*, 5 (3), 164-169. <https://doi.org/10.1111/j.1399-3038.1994.tb00233.x>
- Kaski, J.C. (2010). C-Reactive Protein improves risk prediction in patients with acute coronary syndrome, or does it? *European Heart Journal*, 31, 274-277.

- <https://doi.org/10.1093/eurheartj/ehp435>
- Kinoshita, A., Onoda, H., Imai, N., Nishino, H. and Tajiri, H. (2015). C-Reactive Protein as a prognostic marker in patients with hepatocellular carcinoma. *Hepatology*, 62(140), 966-970.
- Kleniewska, P., Hoffmann, A., Pniewska, E. and Pawliczak, R. (2016). The influence of probiotic *Lactobacillus casei* in combination with prebiotic inulin on the antioxidant capacity of human plasma. *Oxidative Medicine and Cellular Longevity*, 2016, 1340903. <https://doi.org/10.1155/2016/1340903>
- Kratz, A., Pesce, M.A., Basner, R.C. and Einstein, A.J. (2012). Appendix: Laboratory Values of Clinical Importance. 19th ed. New York, USA: The McGraw Hill Companies Inc.
- Lutgendorff, F., Akkermans, L.M. and Soderholm, J.D. (2008). The role of microbiota and probiotics in stress-induced gastro-intestinal damage. *Current Molecular Medicine*, 8(4), 282-298. <https://doi.org/10.2174/156652408784533779>
- Lutgendorff, F., Nijmeijer, R.M., Sandstrom, P.A., Trulsson, L.M., Magnusson, K.E., Timmerman, H.M., van Minnen, L.P., Rijkers, G.T., Gooszen, H.G., Akkermans, L.M. and Soderholm, J.D. (2009). Probiotics prevent intestinal barrier dysfunction in acute pancreatitis in rats via induction of ileal mucosal glutathione biosynthesis. *PloS One*, 4(2), e4512. <https://doi.org/10.1371/journal.pone.0004512>
- Maydych, V. (2019). The interplay between stress, inflammation, and emotional attention: relevance for depression. *Frontiers in Neuroscience*, 13, 384. <https://doi.org/10.3389/fnins.2019.00384>
- Meyer, T., Stockfleth, E. and Christophers, E. (2007). Immune response profiles in human skin. *British Journal of Dermatology*, 157(Suppl. 2), 1-7. doi:10.1111/j.1365-2133.2007.08264.x
- Middleton, E., Kandaswamy, C. and Theoharides, T.C. (2000). The effects of plant flavonoids on mammalian cells: Implications for inflammation, heart disease, and cancer. *Pharmacological Reviews*, 52(4), 673-751.
- Nakata, A. (2012). Psychosocial job stress and immunity: A systematic review. In Yan, Q. (Ed.) *Psychoneuroimmunology: Methods and Protocols*, p. 39-75. Totowa, New Jersey, USA: Humana Press. https://doi.org/10.1007/978-1-62703-071-7_3
- Ohashi, Y. and Ushida, K. (2009). Health beneficial effects of probiotics: Its mode of action. *Journal of Animal Science*, 80(4), 361-371. <https://doi.org/10.1111/j.1740-0929.2009.00645.x>
- Okabayashi, T., Kariwa, H., Yokota, S., Iki, S., Indoh, T., Yokosawa, N. and Fujii, N. (2006). Cytokine regulation in SARS coronavirus infection compared to other respiratory virus infections. *Journal of Medical Virology*, 78(4), 417-424. <https://doi.org/10.1002/jmv.20556>
- Pyne, N.J. and Pyne, S. (2013). Sphingosine 1-phosphate is a missing link between chronic inflammation and colon cancer. *Cancer Cell*, 23(1), 5-7. <https://doi.org/10.1016/j.ccr.2012.12.005>
- Rafiqi, Z., Shiva, A., Arab, S. and Mohd Yousof, R. (2013). Association of dietary vitamin C and E intake and antioxidant enzymes in type 2 diabetes mellitus patients. *Global Journal of Health Science*, 5(3), 183-187. <https://doi.org/10.5539/gjhs.v5n3p183>
- Rahmawati, G., Rachmawati, F. and Winarsi, H. (2014). Aktivitas superoksida dismutase tikus diabetes yang diberi ekstrak batang kapulaga dan glibenklamid. *Scripta Biologica*, 1(3), 197-201. <https://doi.org/10.20884/1.sb.2014.1.3.42> [In Bahasa Indonesia].
- Reboredo-Rodríguez, P., Varela-López, A., Forbes-Hernández, T.Y., Gasparini, M., Afrin, S., Cianciosi, D. and Giampieri, F. (2018). Phenolic compounds isolated from olive oil as nutraceutical tools for the prevention and management of cancer and cardiovascular diseases. *International Journal of Molecular Sciences*, 19(8), 2305. <https://doi.org/10.3390/ijms19082305>
- Ridker, P.M. (2003). Clinical application of c-reactive protein for cardiovascular disease detection and prevention. *Circulation*, 107(3), 363-369. <https://doi.org/10.1161/01.CIR.0000053730.47739.3C>
- Salarkia, N., Ghadamli, L., Zaeri, F. and Sabaghian Rad, L. (2013). Effects of probiotic yogurt on performance, respiratory and digestive systems of young adult female endurance swimmers: a randomized controlled trial. *Medical Journal of the Islamic Republic of Iran*, 27(3), 141-146.
- Salehzadeh, K. (2015). The effects of probiotic yogurt drink on lipid profile, CRP and record changes in aerobic athletes. *International Journal of Life Sciences*, 9, 32-37. <https://doi.org/10.3126/ijls.v9i4.12672>
- Setiawan, D.I., Tjahyono, K. and Afifah, D.N. (2016). Pemberian kacang kedelai terhadap kadar malondialdehid (MDA) dan superoxide dismutase (SOD) tikus Sprague Dawley hiperkolesterolemia. *Jurnal Gizi Klinik Indonesia*, 13(1), 20-26. <https://doi.org/10.22146/ijcn.22815>
- Shivpuri, S., Gallo, L.C., Crouse, J.R. and Allison, M.A. (2012). The association between chronic stress type and C-Reactive Protein in the multi-ethnic study of atherosclerosis (MESA): Does gender make a

- difference? *Journal of Behavioral Medicine*, 35(1), 74-85. <https://doi.org/10.1007/s10865-011-9345-5>
- Sokrab, A.M., Ahmed, I.A.M. and Babiker, E.E. (2012). Effect of germination on antinutritional factors, total, and extractable minerals of high and low phytate corn (*Zea mays* L.) genotypes. *Journal of the Saudi Society of Agricultural Sciences*, 11(2), 123-128. <https://doi.org/10.1016/j.jssas.2012.02.002>
- Sproston, N.R. and Ashworth, J.J. (2018). Role of C-Reactive Protein at sites of inflammation and infection. *Frontiers in Immunology*, 9, 754. <https://doi.org/10.3389/fimmu.2018.00754>
- Suhail, N., Bilal, N., Khan, H., Hasan, S., Sharma, S., Khan, F., Mansoor, T. and Banu, N. (2012). Effect of Vitamins C and E on antioxidant status of breast-cancer patients undergoing chemotherapy. *Journal of Clinical Pharmacy and Therapeutics*, 37(1), 22-26. <https://doi.org/10.1111/j.1365-2710.2010.01237.x>
- Talwalkar, A. and Kailasapathy, K. (2003). Metabolic and biochemical responses of probiotic bacteria to oxygen. *Journal of Dairy Science*, 86(8), 2537-2546. [https://doi.org/10.3168/jds.S0022-0302\(03\)73848-X](https://doi.org/10.3168/jds.S0022-0302(03)73848-X)
- Tanaka, T. and Kishimoto, T. (2014). The biology and medical implications of Interleukin-6. *Cancer Immunology Research*, 2(4), 288-294. <https://doi.org/10.1158/2326-6066.CIR-14-0022>
- Utami, K., Mahdi, C. and Aulanni'am, A. (2017). Potential of *Lactobacillus casei* Shirota strain probiotic toward total cholesterol levels and SOD activity in rat with high cholesterol diet. *Molekul*, 12(2), 153-158. <https://doi.org/10.20884/1.jm.2017.12.2.364>
- Van Immerseel, F., Russel, J., Flythe, M., Gantois, I., Timbermont, J., Pasmans, F., Haesebrouck, F. and Ductalle, R. (2006). The use organic acids to combat *Salmonella* in poultry: a mechanistic explanation of the efficacy. *Avian Pathology*, 35, 182-188. <https://doi.org/10.1080/03079450600711045>
- Vandercam, G., Simon, A., Scohy, A., Belkhir, L., Kabamba, B., Rodriguez-Villalobos, H. and Yombi, J.C. (2020). Clinical characteristics and humoral immune response in healthcare workers with COVID-19 in a teaching hospital in Belgium. *Journal of Hospital Infection*, 106(4), 713-720. <https://doi.org/10.1016/j.jhin.2020.09.018>
- Wang, Y., Wu, Y., Wang, Y., Xu, H., Mei, X., Yu, D., Wang, Y. and Li, W. (2017). Antioxidant properties of probiotic bacteria. *Nutrients*, 9(5), 521. <https://doi.org/10.3390/nu9050521>
- Winarsi, H., Erminawati, E., Andreas, A., and Nuraeni, I. (2021). Mung bean sprouts yogurt rich in antioxidants as a functional drink during pandemic. *Food Research*. [Accepted].
- Winarsi, H., Purwanto, A. and Dwiyantri, H. (2010). Aktivitas glutation peroksidase (GSH-PX) wanita penderita diabetes melitus tipe-2 yang mendapat suplementasi susu kaya protein kecambah kedelai plus Zn. Prosiding Seminar Nasional Hari Lingkungan Hidup se Dunia 2010. 12 June 2010. Purwokerto, Indonesia. [In Bahasa Indonesia].
- Winarsi, H., Sasongko, N.D., Purwanto, A., Arinton, I.G. and Nuraeni, I. (2012). In vitro antioxidant activity of the stem and leaves *Amomum cardamomum* extracts, presented at the International Conference on Medicinal Plants. 11-13 Oct 2012. Purwokerto, Indonesia.
- Winarsi, H., Sasongko, N.D. and Purwanto, A. (2016). Germinated-Soy milk in suppressing inflammation and oxidative stress in blood plasma and breast milk of lactating mothers. *International Food Research Journal*, 23(2), 646-652.
- Winarsi, H., Sasongko, N.D., Purwanto, A. and Nuraeni, I. (2013). Ekstrak daun kapulaga menurunkan indeks atherogenik dan kadar gula darah tikus diabetes induksi Alloxan. *Agritech*, 33(3), 273-280. [In Bahasa Indonesia].
- Winarsi, H., Septiana, A.T., Kartini dan Hanifah, I.N. (2019). Fermentasi bakteri asam laktat meningkatkan kandungan fenolik dan serat yogurt susu kecambah kacang merah (*Phaseolus vulgaris* L.), minuman fungsional untuk obesitas. *Jurnal Gizi dan Pangan Soedirman*, 3(1), 64-75. <https://doi.org/10.20884/1.jgpps.2019.3.1.1526>
- Winarsi, H., Septiana, A.T. dan Wulandari, S.P. (2020). Germination improves sensory, phenolic, protein content and anti-inflammatory properties of red kidney bean (*Phaseolus vulgaris* L.) sprouts milk. *Food Research*, 4(6), 1921-1928. [https://doi.org/10.26656/fr.2017.4\(6\).188](https://doi.org/10.26656/fr.2017.4(6).188)
- Winarsi, H., Siwi P., Wijayanti, M. and Purwanto, A. (2012). Aktivitas enzim superoksida dismutase, katalase, dan glutation peroksidase wanita penderita sindrom metabolik. *Majalah Kedokteran Bandung*, 44(1), 7-12. <https://doi.org/10.15395/mkb.v44n1.75> [In Bahasa Indonesia].