

JOURNAL CONTENT

Search

Search Scope

Browse

- » [By Issue](#)
- » [By Author](#)
- » [By Title](#)

JOURNAL TEMPLATE



VISITORS

00331615

[View My Stats](#)

USER

You are logged in as...

nugrohoaras

- » [My Profile](#)
- » [Log Out](#)
- » [Log Out as User](#)

[Home](#) » [User](#) » [Author](#) » [Submissions](#) » [#747](#) » [Summary](#)

#747 Summary

[SUMMARY](#) [REVIEW](#) [EDITING](#)

Submission

Authors	Mas Yedi Sumaryadi, Dadang Mulyadi Saleh, Aras Prasetyo Nugroho, Nu'man Hidayat, Chomsiatun Nurul Hidayah
Title	Oxytocin Hormone Induction on Milk Production in Relation to Dairy Kid Performance
Original file	747-2845-1-SM.DOC 2019-11-25
Supp. files	None
Submitter	Mr Mas Yedi Sumaryadi
Date submitted	November 25, 2019 - 07:30 AM
Section	Articles
Editor	Agus Susanto
Abstract Views	1937

Status

Status	Published Vol 21, No 3 (2019)
Initiated	2020-03-25
Last modified	2020-12-14

Submission Metadata

Authors

Name	Mas Yedi Sumaryadi
Affiliation	Faculty of Animal Science Jenderal Soedirman University
Country	Indonesia
Bio Statement	—
Name	Dadang Mulyadi Saleh
Affiliation	Faculty of Animal Science Jenderal Soedirman University
Country	Indonesia
Bio Statement	—
Name	Aras Prasetyo Nugroho
Affiliation	Faculty of Animal Science Jenderal Soedirman University
Country	Indonesia
Bio Statement	—
Principal contact for editorial correspondence.	
Name	Nu'man Hidayat
Affiliation	Faculty of Animal Science Jenderal Soedirman University
Country	Indonesia
Bio Statement	—
Name	Chomsiatun Nurul Hidayah
Affiliation	Faculty of Animal Science Jenderal Soedirman University
Country	Indonesia
Bio Statement	—

Title and Abstract

Title	Oxytocin Hormone Induction on Milk Production in Relation to Dairy Kid Performance
Abstract	The application of oxytocin induction to milk production in relation to the performance of kids in dairy goats had been carried out in the "PEGUMAS" dairy goat farmer group Gumelar District, Banyumas. This activity aimed to increase milk production and kid performance through the induction of the hormone oxytocin in dairy goats. The approach method was designed in two stages: the first stage carried out the socialization activities and demonstration of techniques for applying IPTEKS, and the second stage applied hormonal technology in the form of oxytocin hormone induction in dairy goats. The material used was dairy goat parity 1 aged 1.5 - 2 years allocated into 2 groups of 10 individuals each. Group I was dairy goat experiment not induced with the oxytocin hormone as the control. Group II was dairy goat experiment induced with the oxytocin hormone at a dose of 1 ml (10 IU) intramuscularly before milking. Milk production was measured morning and evening as daily milk production in milliliters (ml). Milk was given to the kid of each parent. The kid's performance was measured based on the daily weight gain of the kid during the study. Overall, the application of science and technology could be adopted by farmers, and oxytocin-induced dairy goat milk production (503.2 ml) was significantly higher (P<0.05) than control (305.4 ml) with an increase of 64.77%, with a positive level of correlation (r = 0.45) and contributed 19.83% to the kids daily body weight gain.

Indexing

Keywords	Milk yield, oxytocin, daily body weight gain, dairy goat
Language	en

Supporting Agencies

Agencies	Research Centre and Community Service, Jenderal Soedirman University, Widi Utami Prahayuningtyas, PEGUMAS dairy goat farmers
----------	--

References

References	Adriani. 2013. Pengaruh suplementasi mineral seng terhadap kualitas susu kambing Peranakan Etawah. Jurnal Penelitian Seri Humaniora. 15(1):1 – 8. Adriani, S Novianti and Fatati. 2014. Usaha IbiKK kambing Perah. Jurnal Pengabdian pada Masyarakat. 29(4):1 – 10. Akhtar MS, LA Lodhi, AA Farooq, MM Ayaz, M Hussain, MH Lashari and ZI Chaudhary. 2012. Effect of oxytocin administration before milking on milk production, somatic cells count and fat contents in milk
------------	--

Editorial Board

Focus & Scope

Author Guidelines

Publication Ethics

Reviewer Acknowledgement

List of Reviewers

Online Submission

Open Access Policy

Plagiarism Policy

Submission and Publication Charge

TOOLS



KEYWORDS

Chloramphenicol, rabbit doe, toxicity, teratogenicity, fertility Dairy waste water sludge, broiler, microflora, cecum Essential oil, red ginger, rumen fermentation, in vitro Polymorphism, Association, MSTN gene and Batur sheep Production Quran animal breed conception rate dairy diluents type duration of estrus intact plasma membrane l-threonine l-tryptophan mastitis, microorganisms motility native chicken pregnancy rate productivity, pedet, rice straw the onset of estrus

AUTHOR

Submissions

- » [Active \(0\)](#)
- » [Archive \(1\)](#)
- » [New Submission](#)

- oxytocin administration before milking on milk production, somatic cells count and fat contents in milk of Nili-Ravi Buffaloes. *Pakistan Veterinary Journal*. 32(3):435-437.
- Antonic J, V Tancin, M Uhrincat, L Macuhova, J Macuhova and L Jackuliakova. 2013. The effect of exogenous oxytocin on milkability and milk composition in ewes differed in milk flow pattern. *Small Ruminant Research*. 113:254-257.
- Astuti P, H Surti and NE Sukarini. 2017. Produksi dan komposisi susu kambing Peranakan Ettawa melalui pemberian ekstrak meniran. *Jurnal Ilmu-ilmu Pertanian*. 1(2):82 – 87.
- Belo CJ and RM Bruckmaier. 2010. Sutability of low-dosage oxytocin treatment to induce milk ejection in dairy cows. *Journal of Dairy Science*. 93:63-69.
- Bhimte A, NS Thakur, VP Maurya and G Singh. 2018. Neurohormonal control of lactation and milk let-down in dairy animals. *International Journal of Current Microbiology and Applied Sciences*. 7(7):970-977.
- Elkhair AAM, IA Abdalla and GH El Obied. 2017. Effect of a home-made milk replacer on pre-weaning performance of Nubian male kids. *Gezira J. of Agric. Sci*. 15(2):107-118.
- Ginting SP. 2009. Pedoman Teknis Pemeliharaan Induk dan Anak Kambing Pra-Sapih. Pusat Penelitian dan Pengembangan Peternakan, Loka Penelitian Kambing Potong. Sei Putih, Sumatera Utara. p:33-44.
- Heppelmann M, J Volland, C Pfarrer, M Kietzmann, W Baumer, S Merbach, HA Schoon, O Wellnitz, M Schmicke, M Hoedemarker and H Bollwein. 2018. Effects of oxytocin and PGF2a on uterine contractility in cows with and without metritis – An in-vitro study. *Animal Reproduction*. 188:144-154.
- Hossain SMJ, MF Afroz, AKFH Bhuiyan and MA Habib. 2018. Milk Production Performance of Red Chittagong Cattle Based on Phenotypic and Genetic Parameters. *Glo. Adv. Res. J. Agric. Sci*. 7(4):110-117.
- Htoo NN, B Zeshan, AT Khaing, T Kyaw, EA Woldegiorgis and MA Khan. 2018. Creep Feeding Supplemented with Roughages Improve Rumen Morphology in Pre-Weaning Goat Kids. *Pakistan Journal of Zoology*. 50(2):703-709.
- Kaunang D, Suyadi and S Wahyuningsih. 2012. Analisis litter size, bobot lahir dan bobot sapih hasil perkawinan kawin alam dan inseminasi buatan kambing Boer dan Peranakan Etawah. *Jurnal Ilmu-ilmu Peternakan*. 23(3):41– 46.
- Kim SW and G Wu. 2009. Regulatory role for amino acids in mammary gland growth and milk synthesis. *Amino Acids*. 37: 89-95.
- Kurniasih NN, AM Fuah and R Priyanto. 2013. Karakteristik reproduksi dan perkembangan populasi kambing Peranakan Etawah di lahan pasca galian pasir. *Jurnal Ilmu Produksi dan Teknologi Peternakan*. 1(3):132 –137.
- La Ode Nafiu, W Kurniawan, PN Kusuma, and M Akramullah. 2017. Produktivitas dan Kualitas Susu Berdasarkan Bangsa dan Paritas kambing di Kabupaten Kolaka. *Seminar Nasional Peternakan 3*. P:19 – 28.
- Marwah MP, YY Suranindiyah and TW Murti. 2010. Produksi dan komposisi susu kambing Peranakan Etawa yang diberi suplemen daun katuk (*Sauropus androgynous* L. Merr) pada awal masa laktasi. *Buletin Peternakan*. 54(2):94 – 102.
- Masedunskas A, Y Chen, R Stussman, R Weigert, and IH Mather. 2017. Kinetics of milk lipid droplet transport, growth, and secretion revealed by intravital imaging: lipid droplet release is intermittently stimulated by oxytocin. *MBoC*. 28:935-946.
- Maylinda. 2010. Pengantar Pemuliaan Ternak. Universitas Brawijaya Press. Malang.
- Nwosu EU, IJ James, O Olowofeso, TJ Williams, and MN Bemji. 2019. Effects of oxytocin administration on milk yield and milk composition of West African dwarf does. *Small Ruminant Research*. 181:45-50.
- Ramadhan BG, TH Suprayogi and A Sustiyah. 2013. Tampilan produksi susu dan kadar lemak susu kambing Peranakan Etawa akibat pemberian pakan dengan imbalanced hijauan dan konsentrat yang berbeda. *Animal Agricultural Journal*. 2(1):351 – 356.
- Rizzo A, M Mutinati, G Minoia, M Spedicato, M Panteleo, and RL Sciorsci. 2012. The impact of oxytocin on the hemodynamic features of the milk vein in dairy cows: A color Doppler investigation. *Research in Veterinary Science*. 93:983-988.
- Sudewo ATA, SA Santosa and A Susanto. 2012. Produktivitas kambing peranakan etawah berdasarkan litter size, tipe kelahiran dan mortalitas di village breeding centre kabupaten banyumas. *Prosiding Seminar Nasional: Pengembangan Sumber Daya Pedesaan dan Kearifan Lokal Berkelanjutan II*. p:1-7.
- Yustendi D. 2017. Pemanfaatan tanaman katuk (*Sauropus androgynous* L. Merr) dalam ransum untuk meningkatkan produksi susu kambing betina Peranakan Etawah. *Jurnal Biology Education*. 6(1):21 – 29.

Animal Production Indexed In



This work is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-sa/4.0/).

OXITOCYNE HORMONE INDUCTION ON MILK PRODUCTION IN RELATION TO DAIRY KID PERFORMANCE

(APLIKASI INDUKSI HORMON OKSITOSIN TERHADAP PRODUKSI SUSU KAITANNYA DENGAN KINERJA ANAK PADA KAMBING PERAH)

Mas Yedi Sumaryadi, Dadang Mulyadi Saleh, Aras Prasetyo Nugroho, Nu'man Hidayat, Chomsiatun Nurul Hidayah, and Widi Utami Prahayuningtyas
Faculty of Animal Science, Jenderal Soedirman University – Purwokerto

Corresponding email: yedi.sumaryadi@yahoo.com, nugrohoaras@yahoo.com

ABSTRACT

The application of oxytocin induction to milk production in relation to the performance of kids in dairy goats has been carried out in the "PEGUMAS" dairy goat farmer group Gumelar District, Banyumas. This activity aims to increase milk production and kid performance through the induction of the hormone oxytocin in dairy goats. The approach method used was designed into two stages. The first stage was carried out the socialization activities and demonstration of techniques for applying IPTEKS. The second stage applies hormonal technology in the form of oxytocin hormone induction in dairy goats. The material used was dairy goat parity I with age 1.5 - 2 years which was allocated into 2 groups of 10 individuals each. Group I: dairy goat experiment was not induced by the hormone oxytocin which is used as a control. Group II: dairy goat experiment was induced by the hormone oxytocin at a dose of 1 ml (10 IU) intramuscularly before milking. Milk production is measured morning and evening as daily milk production in milliliters (ml). Milking milk is given to the kid of each parent. The kids performance is measured based on the daily weight gain of the kid during the activity. Overall the application of science and technology can be adopted by farmers, and oxytocin-induced dairy goat milk production (503.2 ml) is significantly higher ($P < 0.05$) than control (305.4 ml) with an increase of 64.77%, and has a positive level of correlation ($r = 0.45$) and contributed 19.83% to the kids daily body weight gain.

Key words : Milk yield, oxytocine, daily body weight gain, dairy goat

ABSTRAK

Aplikasi induksi hormon oksitosin terhadap produksi susu kaitannya dengan kinerja anak pada kambing perah telah dilaksanakan di kelompok peternak kambing perah "PEGUMAS" Kecamatan Gumelar, Banyumas. Kegiatan ini bertujuan untuk meningkatkan produksi susu dan kinerja anak melalui induksi hormon oksitosin pada kambing perah. Metode pendekatan yang digunakan dirancang menjadi dua tahap. Tahap pertama dilakukan kegiatan sosialisasi dan peragaan teknik penerapan IPTEKS. Tahap kedua mengaplikasikan teknologi hormonal berupa induksi hormon oksitosin pada kambing perah. Materi yang digunakan adalah kambing perah paritas I dengan umur 1.5 – 2 tahun yang dialokasikan menjadi 2 kelompok masing-masing 10 ekor. Kelompok I: kambing perah percobaan tidak diinduksi hormon oksitosin yang digunakan sebagai kontrol. Kelompok II: kambing perah percobaan diinduksi hormon oksitosin dengan dosis 1 ml (10 IU) secara intramuskulair sebelum dilakukan pemerahan. Produksi susu diukur pagi dan sore hari sebagai produksi susu harian dalam satuan milliliter (ml). Air susu hasil pemerahan diberikan kepada anak dari masing-masing induk. Kinerja anak diukur berdasarkan pertambahan bobot badan anak harian selama kegiatan. Secara keseluruhan penerapan IPTEKS ini dapat diadopsi peternak dengan baik, dan produksi susu kambing perah yang diinduksi oksitosin (503.2 ml) nyata lebih tinggi ($P < 0.05$) daripada kontrol (305.4 ml) dengan peningkatan sebesar 64.77 %, dan memiliki tingkat keeratan hubungan yang positif ($r = 0.45$) serta berkontribusi sebesar 19.83% terhadap pertambahan bobot badan harian anak.

Kata Kunci : Produksi susu, oksitosin, pertambahan bobot badan harian, kambing perah

INTRODUCTION

Improving population and productivity of dairy goat depends on calving process to breed healthy, weaned kids in one reproduction cycle. The potential improvement is attributed to the prolific nature of goat breeding – goat can breed one to four kids per delivery (Kaunang, et al., 2012; Kurniasih et al., 2013). However, the success to raise the kids post-weaning is related to milk availability because milk is the primary source of nutrient for kids since delivery to pre-weaning, and it is essential for kids' growth and development. Malnutrition (milk) in pre-weaned kids would inhibit the growth and decrease immunity against disease, which eventually leads to mortality (Lu, 2002; Maylinda; 2010). This incident is prevalent – the more kids delivered, the lower the body immune throughout the critical pre-weaning period. Therefore, only a few kids made it through the weaning period.

The increase of dairy goat's milk production for human consumption or the kids is below expectation. Previous studies reported that Etawah goat milk production varied around 0.67L or 756.8 gram/head/day (Adriani, 2013; Adriani et al., 2014), in a range of 0.20 – 1,3L/head/day (Taofik and Depison, 2008; ; Marwah et al., 2010; Ramadhan et al., 2013; Astuti et al., 2017; Yustendi, 2017). The current problem is the high mortality rate in kids at an early age or post-weaning time which inhibits the increase of cattle population and productivity. The mortality rate could reach 12.5 – 30.0% (Kaunang et al., 2012; Adriani et al., 2014), up to 28.57 – 57.14% (Kurniasih et al., 2013)

Milk production (milk let down) is naturally controlled by the oxytocin hormone from anterior hypophysis (McDonald, 1980). However, environment-generated stress or milking disorder could inhibit the secretion of endogenous oxytocin through the circulation system; consequently, milk synthesis products in alveoli cells of milk gland is not entirely discharged and potentially causes mastitis disease. Previous studies on sheep showed that exogenous oxytocin injection before milking could improve milk production in sheep (Sumaryadi and Manalu, 1995a,b; Sumaryadi and Manalu, 1998; Manalu and Sumaryadi, 1998; Sumaryadi and Manalu, 1999a,b). Furthermore, it could increase milk production by 18,64% in oxytocin-induced dairy cows (Sumaryadi et al., 2002). It provides an opportunity for further application to goat cattle, particularly dairy goat.

The result of this research focused on increasing milk production as well as improving the performance of weaned dairy kids. Additionally, milk is a potential food product for human consumption.

RESEARCH METHODS

The research method applied a two-stage design. The first stage included socialization and exhibition of science-technology application technique. The second stage was hormonal technology application through oxytocin induction into dairy goats.

The materials used were dairy goats at parity I aged 1.5 – 2 years old allocated into two groups, ten goats each. Group 1 was an experimental dairy goat without oxytocin induction as control groups, and Group II was experimental dairy goats induced with 1ml (10IU) oxytocin hormone intramuscularly before milking.

Milk production was measured in the morning and afternoon in ml, then provided to kids in a separate cage from each mother. Kid performance was measured by body weight per time unit to obtain daily body weight gain throughout the study.

Success evaluation was conducted at the end of activity in two waves. The first wave was the farmers' level of adoption to apply science-technology application, and the second wave was testing oxytocin-induction application on milk production and kid performance.

RESULT AND DISCUSSION

Oxytocin induction to dairy goats was started with socialization to disseminate information regarding the activity materials and the exhibition of science-technology application for the dairy goat farmers. The result of socialization showed that 90% of the farmers could adopt the oxytocin induction to improve milk production and kid performance. Total milk production at the initial phase from Etawah (PE), Saanen and Jawa Randu goats parity I aged 1.5 – 2 years old (Figure 1).




Figure 1. Average daily milk production of dairy goats

The statistical analysis result showed that milk production was not significantly different ($P>0.05$) across goat breeds (PE, Saanen and Jawa Randu). It indicated that milk production across breeds or genus was relatively similar. The average milk production of dairy goat in this study was 435.37 ± 248.09 ml per head. The contributing factors may include similar age, parity, lactating period and breeding management in addition to the long adaptation to the local environment. However, the average milk production was higher than that of Etawah crossbred, Etawah-Kacang crossbred (PEK) and Etawah-Boer crossbred, namely 272.58 ± 55.73 , 153.77 ± 24.76 , and 174.61 ± 35.22 ml, respectively (La Ode Nafiu et al., 2017). According to NUgroho et al. (2000), cattle potential productivity are attributed to genetic factors, environmental factors and genetic-environment interaction. Also, goat productivity is influenced by climate, parity, litter size and lactating period (Sudewo et al., 2012). The results of the milk production of dairy goats before and after oxytocin hormone induction measured in the morning and afternoon, as well as total milk production, are presented in Table 1.

Table 1. The average milk production of dairy goats induced with Oxitocyn hormone

Variables	Control	Oxitocyn induction
Milk production (ml)		
Morning	358.0 ± 274.9	658.0 ± 320.6
Afternoon	147.5 ± 99.5	348.3 ± 200.8
Total per day (ml/day)	305.4 ± 229.6^a	503.2 ± 249.6^b

a,b) Values bearing different superscript within rows show significant difference ($P<0.05$)

Table 1 shows that oxytocin-induced dairy goats produced more milk in the morning and afternoon milking by 83.80 and 136.14 %, respectively, compared to non-induced dairy goats (control group). Statistical analysis result (see appendix) showed that oxytocin hormone significantly affected ($P<0.05$) total daily milk production. Therefore, the average daily milk production of oxytocin-induced dairy goats (503.2ml) was significantly higher than that of the control group (305.4 ml) with a 64.77% increase in milk production. The current study showed a higher milk production in oxytocin-induced dairy goats than that in dairy cows (Sumaryadi et al., 2002). Milk production is affected by total secretory cells in udder tissue, secretory cell activity to synthesize milk, and substrate availability for milk synthesis. Milk synthesis is conducted by secretory cells in milk gland using the nutrition from feed (Manalu et al., 2000). However, the increased milk production of dairy goats in this study was not due to milk synthesis, but rather the

clear milk secreted from alveoli cell after oxytocin induction on the cell contraction (Linda et al., 1993). Stimulation from milking is forwarded to the hypothalamus that produces oxytocin hormone. Furthermore, oxytocin would trigger soft muscle around alveoli to contract and secrete milk (Rizqi, 2018). Similarly, Caldwell and Young (2006) stated that oxytocin stimulates contraction in the uterus wall to ease calving and secretes milk by stimulating the ductus lactiferous contraction of milk gland (udder) to secrete milk.

The average daily milk production of dairy goats in this study was 435.37 ± 248.09 ml per head. It was higher than that of Etawah dairy goats (PE), Etawah-Kacang crossbred (PEK), and Etawah-Boer crossbred (PEB) namely 272.58 ± 55.73 , 153.77 ± 24.76 , and 174.61 ± 35.22 ml, respectively (La Ode Nafiu et al., 2017). Daily body weight gain (DWG) of kid consuming the milk of each mother was 191.25 ± 91.60 g/day. The DWG was higher than that reported by Ginting (2009) on Boerka goats (118g/day) and Kacang goats (52–70g/day).

There is a significant ($P < 0.05$) correlation between daily milk production and DWG with 0.45 coefficient correlation. The regression analysis result showed that daily milk production of dairy goats was significantly correlated ($P < 0.05$) with kid's daily weight gain (DWG) using an equation $Y = 0.24 + 0.000217 X$ and 0.1983 coefficient of determination. It showed that kid's DWG was 19.83% affected by daily milk production offered to the kids from each mother. Post-calving growth is significantly affected by milk consumption sufficiency before taking forage. According to Surama et al. (1998), kid's growth from the first month of age was solely dependent on the mother's milk – the higher the dependency, the lower milk production when the kids start eating solid food. Similarly, Devandra and Burns (1994) reported that kids were dependent on their mother's milk up to 7-8 weeks post-calving before their rumens start to function and forage intake ensue. Pre-weaning growth was determined by birth weight, milk production, mother's welfare, total calves and their conditions, sex, genetic and breeding management (Lu, 2002; Sutama, 2007; Maylinda, 2010).

CONCLUSIONS

Farmers well adopted Oxytocin hormone induction to improve milk production and kid performance in PEGUMAS association of dairy goat farmers. Oxytocin induction could increase milk production by 64.77%, showed a close relationship ($r = 0.45$), and contributed 19.83% to kid's daily weight gain. However, this science-technology application needs further investigation in the economic aspect and utilization of goat milk for human consumption.

ACKNOWLEDGEMENT

The authors express their gratitude to the Chief of Research Centre and Community Service, Jenderal Soedirman University for the opportunity and research grant through DIPA funding Jenderal Soedirman University 2019 according to the Letter of Assignment to Community Service for implementing Science and Technology No.:P/916/UN 23/ 14/PM/2019. We also thank Widi Utami Prahayuningtyas, a student of Magister Program for assisting community service, as well as the chief and members of PEGUMAS dairy goat farmers community for the contribution to this study.

REFERENCES

- Adiati, U., Hastono, R.S.G. Sianturi, T.D. Chaniago, dan I.K. Utama. 1998. Sinkronisasi birahi secara biologis pada kambing Peranakan Etawah. *Prosiding Seminar Nasional Peternakan dan Veteriner*. Jilid II : 411 – 416.
- Adriani. 2013. Pengaruh suplementasi mineral seng terhadap kualitas susu kambing Peranakan Etawah. *Jurnal Penelitian Seri Humaniora*. Vol. 15 (1): 1 – 8.
- Adriani., S. Novianti, dan Fatati. 2014. Usaha IbKK kambing Perah. *Jurnal Pengabdian pada Masyarakat*. Vol. 29 (4): 1 – 10.
- Astuti, P., H. Surti, dan N.E. Sukarini. 2017. Produksi dan komposisi susu kambing Peranakan Etawah melalui pemberian ekstrak meniran. *Jurnal Ilmu-ilmu Pertanian*. Vol. 1 (2): 82 – 87.
- Caldwell, H.K. and Young, W.S., 2006. Oxytocin and Vasopressin: Genetics and Behavioral Implications in Lim, R. (ed.) *Handbook of Neurochemistry and Molecular Neurobiology*, 3rd edition, Springer, New York, pp. 573-607.
- Davendra C. dan M. Burns. 1994. *Produksi Kambing di Daerah Tropis*. Penerbit Institut Teknologi Bandung, Bandung. (Diterjemahkan oleh I.D.K Harya Putra).
- Edey, T.N. 1983. *A Course Manual in Tropical Sheep and Goat*. Published by AUIDP. Canberra.
- Febriana, D.N., D.W. Harjanti, dan P. Sambodho. 2018. Korelasi ukuran badan, volume ambung, dan produksi susu kambing Peranakan Etawah di kecamatan Turi, kabupaten Sleman Yogyakarta. *Jurnal Ilmu-ilmu Peternakan*. Vol. 28 (2): 134 – 140.
- Ginting, S. P. 2009. *Pedoman Teknis Pemeliharaan Induk dan Anak Kambing Pra-Sapih*. Pusat Penelitian dan Pengembangan Peternakan, Loka Penelitian Kambing Potong. Sei Putih, Sumatera Utara. hal 33-44.
- Kaunang, D., Suyadi, dan S. Wahyuningsih. 2012. Analisis litter size, bobot lahir dan bobot sapih hasil perkawinan kawin alam dan inseminasi buatan kambing Boer dan Peranakan Etawah. *Jurnal Ilmu-ilmu Peternakan*. Vol. 23 (3): 41– 46.
- Kurniasih, N.N., A.M. Fuah, dan R. Priyanto. 2013. Karakteristik reproduksi dan perkembangan populasi kambing Peranakan Etawah di lahan pasca galian pasir. *Jurnal Ilmu Produksi dan Teknologi Peternakan*. Vol. 01 (3): 132 –137.
- La Ode Nafiu, Widhi Kurniawan, Putu Nara Kusuma, dan Muh Akramullah. 2017. Produktivitas dan Kualitas Susu Berdasarkan Bangsa dan Paritas kambing di Kabupaten Kolaka. *Seminar Nasional Peternakan 3*. Pp. 19 – 28.

- Linda, V.B., J.L. Bleek, and R.D. Bremel. 1993. The effect of dairy oxytocin injection before and after milking on milk production, milk plasmin, and milk composition. *J. Dairy Sci.* 76: 315-320.
- Lu, C.D. 2002. *Boer Goat Production: Progress and Prespective*. Vice Chancellor of Academic Affairs. University of HAWAI'I Hilo. Hawaii
- Manalu, W., dan M.Y. Sumaryadi, 1998a. Tantangan dan kesempatan dalam bidang endokrinologi dalam penelitian ilmu-ilmu peternakan dan peningkatan produksi ternak di Indonesia. *Il. Pert. Indon.* Vol. 7 (1):20 – 42.
- Manalu W., M.Y. Sumaryadi, Sudjatmogo and A.S. Satyaningtjas. 2000. Effect of superovulation prior to mating on milk production performance during lactation in ewes. *J. Dairy Sci.* 83:477 –83
- Marwah, M. P., Y.Y. Suranindyah, dan T.W. Murti. 2010. Produksi dan komposisi susu kambing Peranakan Etawa yang diberi suplemen daun katuk (*Sauropus androgynous L. Merr*) pada awal masa laktasi. *Buletin Peternakan.* Vol. 54 (2): 94 – 102.
- Maylinda. 2010. *Pengantar Pemuliaan Ternak*. Universitas Brawijaya Press. Malang.
- McDonald, L.E. 1980. *Veterinary Endocrinology and Reproduction*. 3rd Ed. Lea and Febiger. Philadelphia.
- Nugroho, A. T., P. Surjowardojo dan M. N. Ihsan. Penampilan produksi sapi perah friesian holstein (FH) pada berbagai paritas dan bulan laktasi di ketinggian tempat yang berbeda. *JIPB*, 20(1): 55-64.
- Ramadhan, B.G., T.H. Suprayogi dan A. Sustiyah. 2013. Tampilan produksi susu dan kadar lemak susu kambing Peranakan Etawa akibat pemberian pakan denganimbangan hijauan dan konsentrat yang berbeda. *Animal Agricultural Journal*. Vol. 2 (1): 351 – 356.
- Ranjhan, S.K. 1980. *Animal Nutrition in Tropics*. Villeos Publishing House. New Delhi.
- Rizqi, V.L.P. 2018. Ini-Fungsi-Prolaktin-dan-Oksitosin-Untuk-Kelancaran-ASI Diakses 31 Oktober 2019 pada laman <https://www.motherandbaby.co.id/article/2018/7/38/10272/>
- Setiadi, B. dan I. Inounu 1991. Perkembangbiakan. *Dalam* Beternak Kambing-Domba Sebagai Ternak Potong. Pusat Penelitian dan Pengembangan Peternakan. Bogor. pp. 1-28.
- Stephenson, R.G.A. 1982. On Property Services to Improve Reproductive Rates of Sheep. *Qd. Agric.J.* 108:329 – 332.
- Subandriyo 1990. Ewe productivity in villages in the distric of Garut West Java. *Ilmu dan Peternakan* 4: 307 – 310.
- Sudewo, A. A., S. A. Santosa dan A. Susanto. 2012. Produktivitas kambing peranakan etawah berdasarkanlitter size, tipe kelahiran dan mortalitas di village breeding centre kabupaten banyumas. *Prosiding*,3(1):1-7.
- Sumaryadi, M.Y. dan W. Manalu. 1995a. Peranan konsentrasi progesteron dalam serum induk selama periode kebuntingan dalam mendukung keberhasilan reproduksi: Pengaruh pada pertumbuhan fetus dan kelenjar susu, produksi susu induk, serta berat sapih anak. Makalah Lima Terbaik pada Lomba Karya Ilmiah Boedi Rahardjani Award di *Prosiding Seminar Nasional Ikatan Ahli Ilmu Faal Indonesia (IAIFI)*, Bandung, Semarang.
- Sumaryadi, M.Y. dan W. Manalu. 1995b. Contributions of Maternal Serum Progesterone and Estradiol Concentrations or Corpora Luteal and Fetal Number to Mammary Growth and Development of Ewes During Pregnancy. *Bulletin of Animal Science*. Special Edition : 243 – 247.
- Sumaryadi, M.Y. dan W. Manalu. 1998. Pemanfaatan beberapa profil hormon dan metabolit darah induk domba selama kebuntingan sebagai prediktor untuk menunjang keberhasilan reproduksi: Prediksi jumlah anak, bobot lahir, pertumbuhan kelenjar susu dan produksi susu. *Prosiding Seminar Nasional Peternakan dan Veteriner*. Jilid II : 425 – 436.

- 270 Sumaryadi, M.Y. dan W. Manalu. 1999a. Prediksi pertumbuhan kelenjar susu produksi susu, dan
 271 involusi kelenjar susu berdasarkan konsentrasi beberapa hormon dan metabolit darah
 272 induk selama kebuntingan pada domba ekor tipis Jawa Barat. *Buletin Peternakan*.
 273 Vol.23 (3):103-126
- 274 Sumaryadi, M.Y. dan W. Manalu. 1999a. Prediksi jumlah anak yang akan dilahirkan dan bobot
 275 lahir anak berdasarkan konsentrasi hormon dan metabolit darah induk selama
 276 kebuntingan pada domba ekor tipis Jawa Barat *Buletin Peternakan*. Vol.23 (2):46-63
- 277 Sumaryadi, M.Y., S. Utami. dan B. Hartoyo. 2002. Peningkatan produksi susu melalui induksi
 278 hormone oksitosin pada sapi perah di tingkat peternakan rakyat. *Jurnal Pembangunan*
 279 *Pedesaan*. Vol. 11 (2): 19 – 25.
- 280 Sutama, I.K. 1992. Reproductive development and performance of small ruminants in Indonesia.
 281 *In: New Technology for Small Ruminant Production in Indonesia*. P. Ludgar and S. Scolz
 282 Ed. Winrock International Institute for Agriculture Development, Morritton, Arkansas.
 283 pp. 7 – 14.
- 284 Sutama, I.K., B. Setiadi, I.G.M. Buadiarsana, dan U. Adiati. 1998. Aktivitas seksual setelah
 285 beranak dari kambing perah Peranakan Etawah dengan tingkat produksi susu yang
 286 berbeda. *Prosiding Seminar Nasional Peternakan dan Veteriner*. Jilid II : 401 – 409.
- 287 Sutama, I.K. dan I.G.M. Budiarsana. 2007. *Panduan Lengkap Kambing dan Domba*. Penebar
 288 Swadaya. Jakarta.
- 289 Sutama, I-K. 2007. Tantangan dan peluang peningkatan produktivitas kambing melalui inovasi
 290 teknologi reproduksi. *Prosiding Lokakarya Nasional Kambing Potong*. Pp. 51-60.
- 291 Taofik, A. dan Depison. 2008. Hubungan antar lingkaran perut dan volume ambung dengan
 292 kemampuan produksi susu kambing Peranakan Etawa. *Jurnal Ilmiah Ilmu-ilmu*
 293 *Peternakan*. Vol. 11 (2): 59 – 74.
- 294 Tiesnamurti, B., 1992. Reducing the preweaning mortality rate of Javanese thin-tail sheep. *In:*
 295 *New Technology for Small Ruminant Production in Indonesia*. P. Ludgar and S. Scolz Ed.
 296 Winrock International Institute for Agriculture Development, Morritton, Arkansas. pp.
 297 71 – 80.
- 298 Yustendi, D. 2017. Pemanfaatan tanaman katuk (*Sauropus androgynous* L. Merr) dalam ransum
 299 untuk meningkatkan produksi susu kambing betina Peranakan Etawah. *Jurnal Biology*
 300 *Education*. Vol. 6 (1): 21 – 29.

JOURNAL CONTENT

Search

Search Scope

All

Search

Browse

» By Issue

» By Author

» By Title

JOURNAL TEMPLATE

Journal Template

VISITORS

00331616

View My Stats

USER

You are logged in as...
nugrohoaras

» My Profile

» Log Out

» Log Out as User

Home > User > Author > Submissions > #747 > Review

#747 Review

SUMMARYREVIEWEDITING

Submission

Authors

Mas Yedi Sumaryadi, Dadang Mulyadi Saleh, Aras Prasetyo Nugroho, Nu'man Hidayat, Chomsiatun Nurul Hidayah

Title

Oxytocin Hormone Induction on Milk Production in Relation to Dairy Kid Performance

Section

Articles

Editor

Agus Susanto

Peer Review

Round 1

Review Version

747-2846-1-RV.DOC

2019-11-25

Initiated

2020-01-24

Last modified

2020-01-29

Uploaded file

Reviewer A 747-2896-1-RV.DOC

2020-01-29

Editor Decision

Decision

Accept Submission

2020-02-27

Notify Editor

Editor/Author Email Record

2020-03-19

Editor Version

None

Author Version

747-2895-1-ED.DOC

2020-02-18

DELETE

Upload Author Version

Choose File

No file chosen

Upload

Animal Production Indexed In

Google Scholar

ISJD

Crossref member

MENDELEY

citeulike

Academic Resource Index ResearchBib

EZ3 Electronic Journals Library

BASE

OAJ Open Academic Journals Index

WorldCat

Academic Journals Database

one search

PKP INDEX

Sinta

DOAJ

IPI

CC BY SA

This work is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](#).

Editorial Board

Focus & Scope

Author Guidelines

Publication Ethics

Reviewer Acknowledgement

List of Reviewers

Online Submission

Open Access Policy

Plagiarism Policy

Submission and Publication Charge

TOOLS

MENDELEY

grammarly

KEYWORDS

Chloramphenicol, rabbit doe, toxicity, teratogenicity, fertility Dairy waste water sludge, broiler, microflora, cecum Essential oil, red ginger, rumen fermentation, in vitro Polymorphism, Association, MSTN gene and Batur sheep Production Quran animal breed conception rate dairy diluents type duration of estrus intact plasma membrane l-threonine l-tryptophan mastitis, microorganisms motility native chicken pregnancy rate productivity, pedet, rice straw the onset of estrus

AUTHOR

Submissions

» Active (0)

» Archive (1)

» New Submission

Powered by OJS | Design by ThemeOJS

Back to Top

OXITOCYNE HORMONE INDUCTION ON MILK PRODUCTION IN RELATION TO DAIRY KID PERFORMANCE

(APLIKASI INDUKSI HORMON OKSITOSIN TERHADAP PRODUKSI SUSU KAITANNYA DENGAN KINERJA ANAK PADA KAMBING PERAH)

Mas Yedi Sumaryadi, Dadang Mulyadi Saleh, Aras Prasetyo Nugroho, Nu'man Hidayat, Chomsiatun Nurul Hidayah, and Widi Utami Prahayuningtyas
Faculty of Animal Science, Jenderal Soedirman University – Purwokerto

Corresponding email: yedi.sumaryadi@yahoo.com, nugrohoaras@yahoo.com

ABSTRACT

The application of oxytocin induction to milk production in relation to the performance of kids in dairy goats has been carried out in the "PEGUMAS" dairy goat farmer group Gumelar District, Banyumas. This activity aims to increase milk production and kid performance through the induction of the hormone oxytocin in dairy goats. The approach method used was designed into two stages. The first stage was carried out the socialization activities and demonstration of techniques for applying IPTEKS. The second stage applies hormonal technology in the form of oxytocin hormone induction in dairy goats. The material used was dairy goat parity I with age 1.5 - 2 years which was allocated into 2 groups of 10 individuals each. Group I: dairy goat experiment was not induced by the hormone oxytocin which is used as a control. Group II: dairy goat experiment was induced by the hormone oxytocin at a dose of 1 ml (10 IU) intramuscularly before milking. Milk production is measured morning and evening as daily milk production in milliliters (ml). Milking milk is given to the kid of each parent. The kids performance is measured based on the daily weight gain of the kid during the activity. Overall the application of science and technology can be adopted by farmers, and oxytocin-induced dairy goat milk production (503.2 ml) is significantly higher ($P < 0.05$) than control (305.4 ml) with an increase of 64.77%, and has a positive level of correlation ($r = 0.45$) and contributed 19.83% to the kids daily body weight gain.

Key words : Milk yield, oxytocine, daily body weight gain, dairy goat

ABSTRAK

Aplikasi induksi hormon oksitosin terhadap produksi susu kaitannya dengan kinerja anak pada kambing perah telah dilaksanakan di kelompok peternak kambing perah "PEGUMAS" Kecamatan Gumelar, Banyumas. Kegiatan ini bertujuan untuk meningkatkan produksi susu dan kinerja anak melalui induksi hormon oksitosin pada kambing perah. Metode pendekatan yang digunakan dirancang menjadi dua tahap. Tahap pertama dilakukan kegiatan sosialisasi dan peragaan teknik penerapan IPTEKS. Tahap kedua mengaplikasikan teknologi hormonal berupa induksi hormon oksitosin pada kambing perah. Materi yang digunakan adalah kambing perah paritas I dengan umur 1.5 – 2 tahun yang dialokasikan menjadi 2 kelompok masing-masing 10 ekor. Kelompok I: kambing perah percobaan tidak diinduksi hormon oksitosin yang digunakan sebagai kontrol. Kelompok II: kambing perah percobaan diinduksi hormon oksitosin dengan dosis 1 ml (10 IU) secara intramuskulair sebelum dilakukan pemerahan. Produksi susu diukur pagi dan sore hari sebagai produksi susu harian dalam satuan milliliter (ml). Air susu hasil pemerahan diberikan kepada anak dari masing-masing induk. Kinerja anak diukur berdasarkan pertambahan bobot badan anak harian selama kegiatan. Secara keseluruhan penerapan IPTEKS ini dapat diadopsi peternak dengan baik, dan produksi susu kambing perah yang diinduksi oksitosin (503.2 ml) nyata lebih tinggi ($P < 0.05$) daripada kontrol (305.4 ml) dengan peningkatan sebesar 64.77 %, dan memiliki tingkat keeratan hubungan yang positif ($r = 0.45$) serta berkontribusi sebesar 19.83% terhadap pertambahan bobot badan harian anak.

Kata Kunci : Produksi susu, oksitosin, pertambahan bobot badan harian, kambing perah

Commented [I1]: See the guidelines instruction for author

Commented [I2]: It is better to add the measured variable

Commented [I3]: Adapt to english please

Commented [I4]: How many lambs are used. How old

Commented [I5]: Too much space

INTRODUCTION

Improving population and productivity of dairy goat depends on calving process to breed healthy, weaned kids in one reproduction cycle. The potential improvement is attributed to the prolific nature of goat breeding – goat can breed one to four kids per delivery (Kaunang, et al., 2012; Kurniasih et al., 2013). However, the success to raise the kids post-weaning is related to milk availability because milk is the primary source of nutrient for kids since delivery to pre-weaning, and it is essential for kids' growth and development. Malnutrition (milk) in pre-weaned kids would inhibit the growth and decrease immunity against disease, which eventually leads to mortality (Lu, 2002; Maylinda; 2010). This incident is prevalent – the more kids delivered, the lower the body immune throughout the critical pre-weaning period. Therefore, only a few kids made it through the weaning period.

The increase of dairy goat's milk production for human consumption or the kids is below expectation. Previous studies reported that Etawah goat milk production varied around 0.67L or 756.8 gram/head/day (Adriani, 2013; Adriani et al., 2014), in a range of 0.20 – 1.3L/head/day (Taofik and Depison, 2008; ; Marwah et al., 2010; Ramadhan et al., 2013; Astuti et al., 2017; Yustendi, 2017). The current problem is the high mortality rate in kids at an early age or post-weaning time which inhibits the increase of cattle population and productivity. The mortality rate could reach 12.5 – 30.0% (Kaunang et al., 2012; Adriani et al., 2014), up to 28.57 – 57.14% (Kurniasih et al., 2013)

Milk production (milk let down) is naturally controlled by the oxytocin hormone from anterior hypophysis (McDonald, 1980). However, environment-generated stress or milking disorder could inhibit the secretion of endogenous oxytocin through the circulation system; consequently, milk synthesis products in alveoli cells of milk gland is not entirely discharged and potentially causes mastitis disease. Previous studies on sheep showed that exogenous oxytocin injection before milking could improve milk production in sheep (Sumaryadi and Manalu, 1995a,b; Sumaryadi and Manalu, 1998; Manalu and Sumaryadi, 1998; Sumaryadi and Manalu, 1999a,b). Furthermore, it could increase milk production by 18,64% in oxytocin-induced dairy cows (Sumaryadi et al., 2002). It provides an opportunity for further application to goat cattle, particularly dairy goat.

The result of this research focused on increasing milk production as well as improving the performance of weaned dairy kids. Additionally, milk is a potential food product for human consumption.

Commented [I6]: So...
Add background to be clear

RESEARCH METHODS

The research method applied a two-stage design. The first stage included socialization and exhibition of science-technology application technique. The second stage was hormonal technology application through oxytocin induction into dairy goats.

The materials used were dairy goats at parity I aged 1.5 – 2 years old allocated into two groups, ten goats each. Group 1 was an experimental dairy goat without oxytocin induction as control groups, and Group II was experimental dairy goats induced with 1ml (10IU) oxytocin hormone intramuscularly before milking.

Milk production was measured in the morning and afternoon in ml, then provided to kids in a separate cage from each mother. Kid performance was measured by body weight per time unit to obtain daily body weight gain throughout the study.

Success evaluation was conducted at the end of activity in two waves. The first wave was the farmers' level of adoption to apply science-technology application, and the second wave was testing oxytocin-induction application on milk production and kid performance.

RESULT AND DISCUSSION

Oxytocin induction to dairy goats was started with socialization to disseminate information regarding the activity materials and the exhibition of science-technology application for the dairy goat farmers. The result of socialization showed that 90% of the farmers could adopt the oxytocin induction to improve milk production and kid performance. Total milk production at the initial phase from Etawah (PE), Saanen and Jawa Randu goats parity I aged 1.5 – 2 years old (Figure 1).




Figure 1. Average daily milk production of dairy goats

Commented [17]: Please write in detail. It is better to add a variable that is measured
Research design?
Analysis?

Commented [18]: ???
Detail please

Commented [19]: Breed not explained in the method?

The statistical analysis result showed that milk production was not significantly different ($P>0.05$) across goat breeds (PE, Saanen and Jawa Randu). It indicated that milk production across breeds or genus was relatively similar. The average milk production of dairy goat in this study was 435.37 ± 248.09 ml per head. The contributing factors may include similar age, parity, lactating period and breeding management in addition to the long adaptation to the local environment. However, the average milk production was higher than that of Etawah crossbred, Etawah-Kacang crossbred (PEK) and Etawah-Boer crossbred, namely 272.58 ± 55.73 , 153.77 ± 24.76 , and 174.61 ± 35.22 ml, respectively (La Ode Nafiu et al., 2017). According to Nugroho et al. (2000), cattle potential productivity are attributed to genetic factors, environmental factors and genetic-environment interaction. Also, goat productivity is influenced by climate, parity, litter size and lactating period (Sudewo et al., 2012). The results of the milk production of dairy goats before and after oxytocin hormone induction measured in the morning and afternoon, as well as total milk production, are presented in Table 1.

Table 1. The average milk production of dairy goats induced with Oxitocyn hormone

Variables	Control	Oxitocyn induction
Milk production (ml)		
Morning	358.0 ± 274.9	658.0 ± 320.6
Afternoon	147.5 ± 99.5	348.3 ± 200.8
Total per day (ml/day)	305.4 ± 229.6^a	503.2 ± 249.6^b

a,b) Values bearing different superscript within rows show significant difference ($P<0.05$)

Table 1 shows that oxytocin-induced dairy goats produced more milk in the morning and afternoon milking by 83.80 and 136.14 %, respectively, compared to non-induced dairy goats (control group). Statistical analysis result (see appendix) showed that oxytocin hormone significantly affected ($P<0.05$) total daily milk production. Therefore, the average daily milk production of oxytocin-induced dairy goats (503.2ml) was significantly higher than that of the control group (305.4 ml) with a 64.77% increase in milk production. The current study showed a higher milk production in oxytocin-induced dairy goats than that in dairy cows (Sumaryadi et al., 2002). Milk production is affected by total secretory cells in udder tissue, secretory cell activity to synthesize milk, and substrate availability for milk synthesis. Milk synthesis is conducted by secretory cells in milk gland using the nutrition from feed (Manalu et al., 2000). However, the increased milk production of dairy goats in this study was not due to milk synthesis, but rather the

clear milk secreted from alveoli cell after oxytocin induction on the cell contraction (Linda et al., 1993). Stimulation from milking is forwarded to the hypothalamus that produces oxytocin hormone. Furthermore, oxytocin would trigger soft muscle around alveoli to contract and secrete milk (Rizqi, 2018). Similarly, Caldwell and Young (2006) stated that oxytocin stimulates contraction in the uterus wall to ease calving and secretes milk by stimulating the ductus lactiferous contraction of milk gland (udder) to secrete milk.

The average daily milk production of dairy goats in this study was 435.37 ± 248.09 ml per head. It was higher than that of Etawah dairy goats (PE), Etawah-Kacang crossbred (PEK), and Etawah-Boer crossbred (PEB) namely 272.58 ± 55.73 , 153.77 ± 24.76 , and 174.61 ± 35.22 ml, respectively (La Ode Nafiu et al., 2017). Daily body weight gain (DWG) of kid consuming the milk of each mother was 191.25 ± 91.60 g/day. The DWG was higher than that reported by Ginting (2009) on Boerka goats (118g/day) and Kacang goats (52–70g/day).

There is a significant ($P < 0.05$) correlation between daily milk production and DWG with 0.45 coefficient correlation. The regression analysis result showed that daily milk production of dairy goats was significantly correlated ($P < 0.05$) with kid's daily weight gain (DWG) using an equation $Y = 0.24 + 0.000217 X$ and 0.1983 coefficient of determination. It showed that kid's DWG was 19.83% affected by daily milk production offered to the kids from each mother. Post-calving growth is significantly affected by milk consumption sufficiency before taking forage. According to Surama et al. (1998), kid's growth from the first month of age was solely dependent on the mother's milk – the higher the dependency, the lower milk production when the kids start eating solid food. Similarly, Devandra and Burns (1994) reported that kids were dependent on their mother's milk up to 7-8 weeks post-calving before their rumens start to function and forage intake ensue. Pre-weaning growth was determined by birth weight, milk production, mother's welfare, total calves and their conditions, sex, genetic and breeding management (Lu, 2002; Sutama, 2007; Maylinda, 2010).

CONCLUSIONS

Farmers well adopted Oxytocin hormone induction to improve milk production and kid performance in PEGUMAS association of dairy goat farmers. Oxytocin induction could increase milk production by 64.77%, showed a close relationship ($r=0.45$), and contributed 19.83% to kid's daily weight gain. However, this science-technology application needs further investigation in the economic aspect and utilization of goat milk for human consumption.

ACKNOWLEDGEMENT

The authors express their gratitude to the Chief of Research Centre and Community Service, Jenderal Soedirman University for the opportunity and research grant through DIPA funding Jenderal Soedirman University 2019 according to the Letter of Assignment to Community Service for implementing Science and Technology No.:P/916/UN 23/ 14/PM/2019. We also thank Widi Utami Prahayuningtyas, a student of Magister Program for assisting community service, as well as the chief and members of PEGUMAS dairy goat farmers community for the contribution to this study.

REFERENCES

- Adiati, U., Hastono, R.S.G. Sianturi, T.D. Chaniago, dan I.K. Utama. 1998. Sinkronisasi birahi secara biologis pada kambing Peranakan Etawah. *Prosiding Seminar Nasional Peternakan dan Veteriner*. Jilid II : 411 – 416.
- Adriani. 2013. Pengaruh suplementasi mineral seng terhadap kualitas susu kambing Peranakan Etawah. *Jurnal Penelitian Seri Humaniora*. Vol. 15 (1): 1 – 8.
- Adriani., S. Novianti, dan Fatati. 2014. Usaha IBKK kambing Perah. *Jurnal Pengabdian pada Masyarakat*. Vol. 29 (4): 1 – 10.
- Astuti, P., H. Surti, dan N.E. Sukarini. 2017. Produksi dan komposisi susu kambing Peranakan Etawah melalui pemberian ekstrak meniran. *Jurnal Ilmu-ilmu Pertanian*. Vol. 1 (2): 82 – 87.
- Caldwell, H.K. and Young, W.S., 2006. Oxytocin and Vasopressin: Genetics and Behavioral Implications in Lim, R. (ed.) *Handbook of Neurochemistry and Molecular Neurobiology*, 3rd edition, Springer, New York, pp. 573-607.
- Davendra C. dan M. Burns. 1994. *Produksi Kambing di Daerah Tropis*. Penerbit Intitut Teknologi Bandung, Bandung. (Diterjemahkan oleh I.D.K Harya Putra).
- Edey, T.N. 1983. *A Course Mannual in Tropical Sheep and Goat*. Published by AUIDP. Canberra.
- Febriana, D.N., D.W. Harjanti, dan P. Sambodho. 2018. Korelai ukuran badan, volume ambing, dan produksi susu kambing Preranakan Etawah di kecamatan Turi, kabupaten Sleman Yogyakarta. *Jurnal Ilmu-ilmu Peternakan*. Vol. 28 (2): 134 – 140.
- Ginting, S. P. 2009. *Pedoman Teknis Pemeliharaan Induk dan Anak Kambing Pra-Sapih*. Pusat Penelitian dan Pengembangan Peternakan, Loka Penelitian Kambing Potong. Sei Putih, Sumatera Utara. hal 33-44.
- Kaunang, D., Suyadi, dan S. Wahyuningsih. 2012. Analisis littersize, bobot lahir dan bobot sapih hasil perkawinan kawin alam dan inseminasi buatan kambing Boer dan Peranakan Etawah. *Jurnal Ilmu-ilmu Peternakan*. Vol. 23 (3): 41– 46.
- Kurniasih, N.N., A.M. Fuah, dan R. Priyanto. 2013. Karakteristik reproduksi dan perkembangan populasi kambing Peranakan Etawah di lahan pasca galian pasir. *Jurnal Ilmu Produksi dan Teknologi Peternakan*. Vol. 01 (3): 132 –137.
- La Ode Nafiu, Widhi Kurniawan, Putu Nara Kusuma, dan Muh Akramullah. 2017. Produktivitas dan Kualitas Susu Berdasarkan Bangsa dan Paritas kambing di Kabupaten Kolaka. *Seminar Nasional Peternakan 3*. Pp. 19 – 28.

Commented [I10]: Update references please

- Linda, V.B., J.L. Bleek, and R.D. Bremel. 1993. The effect of dairy oxytocin injection before and after milking on milk production, milk plasmin, and milk composition. *J. Dairy Sci.* 76: 315-320.
- Lu, C.D. 2002. *Boer Goat Production: Progress and Prespective*. Vice Chancellor of Academic Affairs. University of HAWAI'I Hilo. Hawai
- Manalu, W., dan M.Y. Sumaryadi, 1998a. Tantangan dan kesempatan dalam bidang endokrinologi dalam penelitian ilmu-ilmu peternakan dan peningkatan produksi ternak di Indonesia. *Il. Pert. Indon.* Vol. 7 (1):20 – 42.
- Manalu W., M.Y. Sumaryadi, Sudjatmogo and A.S. Satyaningtijas. 2000. Effect of superovulation prior to mating on milk production performance during lactation in ewes. *J. Dairy Sci.* 83:477 –83
- Marwah, M. P., Y.Y. Suranindyah, dan T.W. Murti. 2010. Produksi dan komposisi susu kambing Peranakan Etawa yang diberi suplemen daun katuk (*Sauropus androgynous L. Merr*) pada awal masa laktasi. *Buletin Peternakan.* Vol. 54 (2): 94 – 102.
- Maylinda. 2010. *Pengantar Pemuliaan Ternak*. Universitas Brawijaya Press. Malang.
- McDonald, L.E. 1980. *Veterinary Endocrinology and Reproduction*. 3rd Ed. Lea and Febiger. Philadelphia.
- Nugroho, A. T., P. Surjowardojo dan M. N. Ihsan. Penampilan produksi sapi perah friesland holstein (FH) pada berbagai paritas dan bulan laktasi di ketinggian tempat yang berbeda. *JIPB*, 20(1): 55-64.
- Ramadhan, B.G., T.H. Suprayogi dan A. Sustiyah. 2013. Tampilan produksi susu dan kadar lemak susu kambing Peranakan Etawa akibat pemberian pakan dengan imbalanced hijauan dan konsentrat yang berbeda. *Animal Agricultural Journal.* Vol. 2 (1): 351 – 356.
- Ranjhan, S.K. 1980. *Animal Nutrition in Tropics*. Vileos Publishing House. New Delhi.
- Rizqi, V.L.P. 2018. Ini-Fungsi-Prolaktin-dan-Oksitosin-Untuk-Kelancaran-ASI Diakses 31 Oktober 2019 pada laman <https://www.motherandbaby.co.id/article/2018/7/38/10272/>
- Setiadi, B. dan I. Inounu 1991. Perkembangbiakan. *Dalam* Beternak Kambing-Domba Sebagai Ternak Potong. Pusat Penelitian dan Pengembangan Peternakan. Bogor. pp. 1-28.
- Stephenson, R.G.A. 1982. On Property Services to Improve Reproductive Rates of Sheep. *Qd. Agric.J.* 108:329 – 332.
- Subandriyo 1990. Ewe productivity in villages in the distric of Garut West Java. *Ilmu dan Peternakan* 4: 307 – 310.
- Sudewo, A. A., S. A. Santosa dan A. Susanto. 2012. Produktivitas kambing peranakan etawah berdasarkan litter size, tipe kelahiran dan mortalitas di village breeding centre kabupaten banyumas. *Prosiding*, 3(1):1-7.
- Sumaryadi, M.Y. dan W. Manalu. 1995a. Peranan konsentrasi progesteron dalam serum induk selama periode kebuntingan dalam mendukung keberhasilan reproduksi: Pengaruh pada pertumbuhan fetus dan kelenjar susu, produksi susu induk, serta berat sapih anak. Makalah Lima Terbaik pada Lomba Karya Ilmiah Boedi Rahardjani Award di *Prosiding Seminar Nasional Ikatan Ahli Ilmu Faal Indonesia (IAIFI)*, Bandung, Semarang.
- Sumaryadi, M.Y. dan W. Manalu. 1995b. Contributions of Maternal Serum Progesterone and Estradiol Concentrations or Corpora Luteal and Fetal Number to Mammary Growth and Development of Ewes During Pregnancy. *Bulletin of Animal Science*. Special Edition : 243 – 247.
- Sumaryadi, M.Y. dan W. Manalu. 1998. Pemanfaatan beberapa profil hormon dan metabolit darah induk domba selama kebuntingan sebagai prediktor untuk menunjang keberhasilan reproduksi: Prediksi jumlah anak, bobot lahir, pertumbuhan kelenjar susu dan produksi susu. *Prosiding Seminar Nasional Peternakan dan Veteriner*. Jilid II : 425 – 436.

- Sumaryadi, M.Y. dan W. Manalu. 1999a. Prediksi pertumbuhan kelenjar susu produksi susu, dan involusi kelenjar susu berdasarkan konsentrasi beberapa hormon dan metabolit darah induk selama kebuntingan pada domba ekor tipis Jawa Barat. *Buletin Peternakan*. Vol.23 (3):103-126
- Sumaryadi, M.Y. dan W. Manalu. 1999a. Prediksi jumlah anak yang akan dilahirkan dan bobot lahir anak berdasarkan konsentrasi hormon dan metabolit darah induk selama kebuntingan pada domba ekor tipis Jawa Barat *Buletin Peternakan*. Vol.23 (2):46-63
- Sumaryadi, M.Y., S. Utami. dan B. Hartoyo. 2002. Peningkatan produksi susu melalui induksi hormone oksitosin pada sapi perah di tingkat peternakan rakyat. *Jurnal Pembangunan Pedesaan*. Vol. 11 (2): 19 – 25.
- Sutama, I.K. 1992. Reproductive development and performance of small ruminants in Indonesia. In: New Technology for Small Ruminant Production in Indonesia. P. Ludgar and S. Scolz Ed. Winrock International Institute for Agriculture Development, Morritton, Arkansas. pp. 7 – 14.
- Sutama, I.K., B. Setiadi, I.G.M. Buadiarsana, dan U. Adiati. 1998. Aktivitas seksual setelah beranak dari kambing perah Peranakan Etawah dengan tingkat produksi susu yang berbeda. *Prosiding Seminar Nasional Peternakan dan Veteriner*. Jilid II : 401 – 409.
- Sutama, I.K. dan I.G.M. Budiarsana. 2007. *Panduan Lengkap Kambing dan Domba*. Penebar Swadaya. Jakarta.
- Sutama, I.K. 2007. Tantangan dan peluang peningkatan produktivitas kambing melalui inovasi teknologi reproduksi. Prosiding Lokakarya Nasional Kambing Potong. Pp. 51-60.
- Taofik, A. dan Depison. 2008. Hubungan antar lingkaran perut dan volume ambing dengan kemampuan produksi susu kambing Peranakan Etawa. *Jurnal Ilmiah Ilmu-ilmu Peternakan*. Vol. 11 (2): 59 – 74.
- Tiesnamurti, B., 1992. Reducing the preweaning mortality rate of Javanese thin-tail sheep. In: New Technology for Small Ruminant Production in Indonesia. P. Ludgar and S. Scolz Ed. Winrock International Institute for Agriculture Development, Morritton, Arkansas. pp. 71 – 80.
- Yustendi, D. 2017. Pemanfaatan tanaman katuk (*Sauropus androgynous L. Merr*) dalam ransum untuk meningkatkan produksi susu kambing betina Peranakan Etawah. *Jurnal Biology Education*. Vol. 6 (1): 21 – 29.

OXITOCYN HORMONE INDUCTION ON MILK PRODUCTION IN RELATION TO DAILY BODY WEIGHT GAIN OF DAIRY KID

(APLIKASI INDUKSI HORMON OKSITOSIN TERHADAP PRODUKSI SUSU KAITANNYA DENGAN PERTAMBAHAN BOBOT BADAN HARIAN ANAK KAMBING PERAH)

Mas Yedi Sumaryadi, Dadang Mulyadi Saleh, Aras Prasetyo Nugroho, Nu'man Hidayat, and Chomsiatun Nurul Hidayah

Faculty of Animal Science, Jenderal Soedirman University - Purwokerto 53123, Indonesia
Corresponding email: yedi.sumaryadi@yahoo.com

ABSTRACT

The application of oxytocin induction to milk production in relation to the daily weight gain of the kid in dairy goats have been carried out in the "PEGUMAS" dairy goat farmer group Gumelar District, Banyumas. This activity aims to increase milk production and kid performance through the induction of the hormone oxytocin in dairy goats. The approach method used was designed into two stages. The first stage was carried out the socialization activities and demonstration of techniques for applying IPTEKS. The second stage applies hormonal technology in the form of oxytocin hormone induction in dairy goats. The material used was dairy goat parity I (twins, two dairy kid, and one month old) with age 1.5 - 2 years which was allocated into 2 groups of 10 individuals each. Group I: dairy goat experiment was not induced by the hormone oxytocin which is used as a control. Group II: dairy goat experiment was induced by the hormone oxytocin at a dose of 1 mL (10 IU) intramuscularly before milking. Milk production is measured morning and evening as daily milk production in milliliters (mL). Milking milk is given to the kid of each parent. The kid's performance is measured based on the daily weight gain of the kid during the activity. Overall the application of science and technology can be adopted by farmers, and oxytocin-induced dairy goat milk production (503.2 mL) is significantly higher ($P < 0.05$) than control (305.4 mL) with an increase of 64.77%, and has a positive level of correlation ($r = 0.45$) with equation $Y = 0.24 + 0.000217 X$ and contributed 12.90% to the kids daily body weight gain.

Keywords: Milk yield, oxytocin, daily body weight gain, dairy goat

ABSTRAK

Aplikasi induksi hormon oksitosin terhadap produksi susu kaitannya dengan pertambahan bobot badan harian anak kambing perah telah dilaksanakan di kelompok peternak kambing perah "PEGUMAS" Kecamatan Gumelar, Banyumas. Kegiatan ini bertujuan untuk meningkatkan produksi susu dan kinerja anak melalui induksi hormon oksitosin pada kambing perah. Metode pendekatan yang digunakan dirancang menjadi dua tahap. Tahap pertama dilakukan kegiatan sosialisasi dan peragaan teknik penerapan IPTEKS. Tahap kedua mengaplikasikan teknologi hormonal berupa induksi hormon oksitosin pada kambing perah. Materi yang digunakan adalah kambing perah paritas I (kembar, dua anak, umur satu bulan) dengan umur 1,5 – 2 tahun, dialokasikan menjadi 2 kelompok masing-masing 10 ekor. Kelompok I: kambing perah percobaan tidak diinduksi hormon oksitosin yang digunakan sebagai kontrol. Kelompok II: kambing perah percobaan diinduksi hormon oksitosin dengan dosis 1 mL (10 IU) secara intramuskuler sebelum dilakukan pemerahan. Produksi susu diukur pagi dan sore hari sebagai produksi susu harian dalam satuan mililiter (mL). Air susu hasil pemerahan diberikan kepada anak dari masing-masing induk. Kinerja anak diukur berdasarkan pertambahan bobot badan anak harian selama kegiatan. Secara keseluruhan penerapan IPTEKS ini dapat diadopsi peternak dengan baik, dan produksi susu kambing perah yang diinduksi oksitosin (503,2 mL) nyata lebih tinggi ($P < 0,05$) daripada kontrol (305,4 mL) dengan peningkatan sebesar 64,77%, dan memiliki tingkat keeratan hubungan yang positif ($r = 0,45$) dengan persamaan $Y = 0.24 + 0.000217 X$ serta berkontribusi sebesar 12,90% terhadap pertambahan bobot badan harian anak.

Kata Kunci: Produksi susu, oksitosin, pertambahan bobot badan harian, kambing perah

Commented [I1]: See the guidelines instruction for author

Commented [a2R1]: Oke

Commented [I3]: It is better to add the measured variable

Commented [a4R3]: Oke

Commented [I5]: Adapt to english please

Commented [a6R5]: Oke

Commented [I7]: How many lambs are used. How old

Commented [a8R7]: Oke

Commented [I9]: Too much space

Commented [a10R9]: Oke

Introduction

Improving population and productivity of dairy goat depends on calving process to breed healthy, weaned kids in one reproduction cycle. The potential improvement is attributed to the prolific nature of goat breeding – goat can breed one to four kids per delivery (Kaunang et al., 2012; Kurniasih et al., 2013). However, the success to raise the kids post-weaning is related to milk availability because milk is the primary source of nutrient for kids since delivery to pre-weaning, and it is essential for kids' growth and development. Malnutrition (milk) in pre-weaned kids would inhibit the growth and decrease immunity against the disease, which eventually leads to mortality (Maylinda, 2010). This incident is prevalent – the more kids delivered, the lower the body immune throughout the critical pre-weaning period. Therefore, only a few kids made it through the weaning period.

The increase of dairy goat's milk production for human consumption or the kids is below expectation. Previous studies reported that Etawah goat milk production varied around 0.67 L or 756.8 gram/head/day (Adriani, 2013; Adriani et al., 2014), in a range of 0.20 – 1,3 L/head/day (Marwah et al., 2010; Ramadhan et al., 2013; Astuti et al., 2017; Yustendi, 2017). The current problem is the high mortality rate in kids at an early age or post-weaning time which inhibits the increase of cattle population and productivity. The mortality rate could reach 12.5 – 30.0% (Kaunang et al., 2012; Adriani et al., 2014), up to 28.57 – 57.14% (Kurniasih et al., 2013)

Milk production (milk let down) is naturally controlled by the oxytocin hormone from anterior hypophysis (Bhimte et al., 2018). However, environment-generated stress or milking disorder could inhibit the secretion of endogenous oxytocin through the circulation system; consequently, milk synthesis products in alveoli cells of milk gland is not entirely discharged and potentially causes mastitis disease. Previous studies showed that exogenous oxytocin injection before milking could improve milk production in sheep (Sumaryadi and Manalu, 1995; Sumaryadi and Manalu, 1999a; Sumaryadi and Manalu, 1999b), cow (Belo and Bruckmaier, 2010; Rizzo et al., 2012) and buffaloes (Akhtar et al., 2012). Furthermore, it could increase milk production by 18,64% (Sumaryadi et al., 2002) and 15,32% (Antonic et al., 2013) in oxytocin-induced sheep. It provides an opportunity for further application to goat, particularly dairy goat.

From the results of previous studies, researchers suspect that if there is an increase in milk production, it will be related to the daily weight gain of pre-weaning. Therefore researchers are very interested in researching giving oxytocin to dairy goats.

Commented [I11]: So...
Add background to be clear

Commented [a12R11]: Oke

Materials and Methods

The materials used are dairy goats of three species, namely Etawah (PE), Saanen, and Jawa Randu on parity I (twins, two dairy kid, and one month old) aged 1.5 – 2 years, oxytocin, injection syringe, gloves, milking bottle, scale, mask, Vaseline, measuring cup, and office stationery. The Study have been carried out in the "PEGUMAS" dairy goat farmer group Gumelar District, Banyumas.

The research method applied a two-stage design. The first stage included socialization and exhibition of science-technology application technique. The second stage was hormonal technology application through oxytocin induction into dairy goats. The study was designed using a completely randomized designed (CRD) which was allocated into two treatment groups, with each repetition being carried out three times. Group 1 was an experimental dairy goat without oxytocin induction as control groups, and Group II was experimental dairy goats induced with 1 mL (10 IU) oxytocin hormone intramuscularly before milking. Milk production was measured in the morning and afternoon in mL, then provided to dairy kids in a separate cage from each mother. Dairy Kid performance was measured by body weight per time unit to obtain daily body weight gain throughout the study. Milk production of each species will be analyzed descriptively, while the relationship between milk production and dairy kid daily body weight gain (DWG) will be analyzed by correlation and regression.

Success evaluation was conducted at the end of activity in two waves. The first wave was the farmers' level of adoption to apply science-technology application, and the second wave was testing an oxytocin-induction application on milk production and kid performance.

Result and Discussion

Oxytocin induction to dairy goats was started with socialization to disseminate information regarding the activity materials and the exhibition of science-technology application for the dairy goat farmers. The result of socialization showed that 90% of the farmers could adopt the oxytocin induction to improve milk production and kid performance. Total milk production at the initial phase from Etawah (PE), Saanen and Jawa Randu goats parity I aged 1.5 – 2 years old (Figure 1).

Commented [I13]: ???
Detail please

Commented [I14]: Breed not explained in the method?

Commented [a15R14]: Oke..., already mentioned




Figure 1. Average daily milk production of dairy goats

The statistical analysis result showed that milk production was not significantly different ($P>0.05$) across goat breeds (PE, Saanen and Jawa Randu). It indicated that milk production across breeds or genus was relatively similar. The average milk production of dairy goat in this study was 435.37 ± 248.09 mL/head. The contributing factors may include similar age, parity, lactating period and breeding management in addition to the long adaptation to the local environment. However, the average milk production was higher than that of Etawah crossbred, Etawah-Kacang crossbred (PEK) and Etawah-Boer crossbred, namely 272.58 ± 55.73 , 153.77 ± 24.76 , and 174.61 ± 35.22 mL, respectively (La Ode Nafiu et al., 2017). According to Hossain et al. (2018), cattle potential productivity are attributed to genetic factors, environmental factors and genetic-environment interaction. Also, goat productivity is influenced by climate, parity, litter size and lactating period (Sudewo et al., 2012). The results of the milk production of dairy goats before and after oxytocin hormone induction measured in the morning and afternoon, as well as total milk production, are presented in Table 1.

Table 1. The average milk production of dairy goats induced with Oxytocin hormone

Variables	Control	Oxytocin induction
Milk production (mL)		
Morning	358.0 ± 274.9	658.0 ± 320.6
Afternoon	147.5 ± 99.5	348.3 ± 200.8
Total (mL/day)	305.4 ± 229.6^a	503.2 ± 249.6^b

(a,b) Values bearing different superscript within rows show a significant difference ($P<0.05$)

Table 1 shows that oxytocin-induced dairy goats produced more milk in the morning and afternoon milking by 83.80 and 136.14%, respectively, compared to non-induced dairy goats (control group). Statistical analysis result (see appendix) showed that oxytocin hormone significantly affected ($P<0.05$) total daily milk production. Therefore, the average daily milk production of oxytocin-induced dairy goats (503.2 mL) was significantly higher than that of the control group (305.4 mL) with a 64.77% increase in milk production. In line with the previous study, oxytocin injection in goat can increase milk production, especially milk fat (Nwosu et al., 2019). The current study showed a higher milk production in oxytocin-induced dairy goats than that in sheep (Antonic et al., 2013). Milk production is affected by total secretory cells in udder tissue, secretory cell activity to synthesize milk, and substrate availability for milk synthesis. Milk synthesis is conducted by secretory cells in milk gland using nutrition from feed (Kim and Wu, 2009). Stimulation from milking is forwarded to the hypothalamus that produces oxytocin hormone. Furthermore, oxytocin stimulates contraction in the uterus wall (Heppelmann et al., 2018) and would trigger soft muscle around alveoli to contract and secretes milk by stimulating the ductus lactiferous contraction of milk gland (udder) to secrete milk (Masedunskas et al., 2017).

The average daily milk production of dairy goats in this study was 435.37 ± 248.09 ml per head. It was higher than that of Etawah dairy goats (PE), Etawah-Kacang crossbred (PEK), and Etawah-Boer crossbred (PEB) namely 272.58 ± 55.73 , 153.77 ± 24.76 , and 174.61 ± 35.22 mL, respectively (La Ode Nafiu et al., 2017). Daily body weight gain (DWG) of dairy kid consuming the milk of each mother was 0.35 ± 0.13 Kg/day. The DWG was higher than that reported by Ginting (2009) on Boer goats (118 g/day) and Kacang goats (52–70 g/day).

Based on Table 2 it turns out that the average daily body weight gain (DWG) of dairy kid fed with milk from oxytocin-induced mothers was 0.35 ± 0.13 Kg significantly higher ($P<0.05$) than that of Kid dairy DWG from control was 0.31 ± 0.12 Kg. This means that the DWG of kid dairy from oxytocin-induced parents is 12.90% higher than DWG of the parent of the control.

Table 2. Milk production and daily body weight gain (DWG)

	Daily weight gain (Control) (Kg/day)	Daily weight gain with oxytocin (kg/day)
Mean	0.31 ± 0.12^a	0.35 ± 0.13^b
Correlation (r)	0.45	
Regression	$Y = 0.24 + 0.000217 X$	

(a,b) Values bearing different superscript within rows show a significant difference ($P<0.05$)

There is a significant ($P < 0.05$) correlation between daily milk production and DWG with 0.45 coefficient correlation. The regression analysis result showed that daily milk production of dairy goats was significantly correlated ($P < 0.05$) with kid's daily weight gain (DWG) using an equation $Y = 0.24 + 0.000217 X$ and 0.1290 coefficient of determination. It showed that kid's DWG was 12.90% affected by daily milk production offered to the kids from each mother. Post-calving growth is significantly affected by milk consumption sufficiency before taking forage. According to Elkhair et al. (2017), kid's growth from the first month of age was solely dependent on the mother's milk – the higher the dependency, the lower milk production when the kids start eating solid food. Similarly, Htoo et al. (2018) reported that kids were dependent on their mother's milk up to 12 weeks post-calving before their rumens start to function and forage intake ensue. Pre-weaning growth was determined by birth weight, milk production, mother's welfare, total calves and their conditions, sex, genetic and breeding management (Maylinda, 2010).

Conclusions

Farmers well adopted Oxytocin hormone induction to improve milk production and kid performance in PEGUMAS association of dairy goat farmers. Oxytocin induction could increase milk production by 64.77%, showed a close relationship ($r = 0.45$), and contributed 12.90% to dairy kid's daily weight gain. However, this science-technology application needs further investigation in the economic aspect and utilization of goat milk for human consumption.

Acknowledgement

The authors express their gratitude to the Chief of Research Centre and Community Service, Jenderal Soedirman University for the opportunity and research grant through DIPA funding Jenderal Soedirman University 2019 according to the Letter of Assignment to Community Service for implementing Science and Technology No.: P/916/UN 23/ 14/PM/2019. We also thank Widi Utami Prahayuningtyas, a student of Magister Program for assisting community service, as well as the chief and members of PEGUMAS dairy goat farmer's community for the contribution to this study.

References

- Adriani. 2013. Pengaruh suplementasi mineral seng terhadap kualitas susu kambing Peranakan Etawah. *Jurnal Penelitian Seri Humaniora*. 15(1):1 – 8.
- Adriani, S Novianti and Fatati. 2014. Usaha IBKK kambing Perah. *Jurnal Pengabdian pada Masyarakat*. 29(4):1 – 10.

Commented [I16]: Update references please

Commented [a17R16]: OKe

- 204 Akhtar MS, LA Lodhi, AA Farooq, MM Ayaz, M Hussain, MH Lashari and ZI Chaudhary. 2012. Effect
205 of oxytocin administration before milking on milk production, somatic cells count and
206 fat contents in milk of Nili-Ravi Buffaloes. *Pakistan Veterinary Journal*. 32(3):435-437.
- 207 Antonic J, V Tancin, M Uhrincat, L Macuhova, J Macuhova and L Jackuliakova. 2013. The effect of
208 exogenous oxytocin on milkability and milk composition in ewes differed in milk flow
209 pattern. *Small Ruminant Research*. 113:254-257.
- 210 Astuti P, H Sripita and NE Sukarini. 2017. Produksi dan komposisi susu kambing Peranakan Ettawa
211 melalui pemberian ekstrak meniran. *Jurnal Ilmu-ilmu Pertanian*. 1(2):82 – 87.
- 212 Belo CJ and RM Bruckmaier. 2010. Sutability of low-dosage oxytocin treatment to induce milk
213 ejection in dairy cows. *Journal of Dairy Science*. 93:63-69.
- 214 Bhimte A, NS Thakur, VP Maurya and G Singh. 2018. Neurohormonal control of lactation and milk
215 let-down in dairy animals. *International Journal of Current Microbiology and Applied
216 Sciences*. 7(7):970-977.
- 217 Elkhair AAM, IA Abdalla and GH El Obied. 2017. Effect of a home-made milk replacer on pre-
218 weaning performance of Nubian male kids. *Gezira J. of Agric. Sci*. 15(2):107-118.
- 219 Ginting SP. 2009. Pedoman Teknis Pemeliharaan Induk dan Anak Kambing Pra-Sapih. Pusat
220 Penelitian dan Pengembangan Peternakan, Loka Penelitian Kambing Potong. Sei Putih,
221 Sumatera Utara. p:33-44.
- 222 Heppelmann M, J Volland, C Pfarrer, M Kietzmann, W Baumer, S Merbach, HA Schoon, O Wellnitz,
223 M Schmicke, M Hoedemarker and H Bollwein. 2018. Effects of oxytocin and PGF2a on
224 uterine contractility in cows with and without metritis – An in-vitro study. *Animal
225 Reproduction*. 188:144-154.
- 226 Hossain SMJ, MF Afroz, AKFH Bhuiyan and MA Habib. 2018. Milk Production Performance of Red
227 Chittagong Cattle Based on Phenotypic and Genetic Parameters. *Glo. Adv. Res. J. Agric.
228 Sci*. 7(4):110-117.
- 229 Htoo NN, B Zeshan, AT Khaing, T Kyaw, EA Woldegiorgis and MA Khan. 2018. Creep Feeding
230 Supplemented with Roughages Improve Rumen Morphology in Pre-Weaning Goat Kids.
231 *Pakistan Journal of Zoology*. 50(2):703-709.
- 232 Kaunang D, Suyadi and S Wahyuningsih. 2012. Analisis litter size, bobot lahir dan bobot sapih hasil
233 perkawinan kawin alam dan inseminasi buatan kambing Boer dan Peranakan Etawah.
234 *Jurnal Ilmu-ilmu Peternakan*. 23(3):41– 46.
- 235 Kim SW and G Wu. 2009. Regulatory role for amino acids in mammary gland growth and milk
236 synthesis. *Amino Acids*. 37: 89-95.
- 237 Kurniasih NN, AM Fuah and R Priyanto. 2013. Karakteristik reproduksi dan perkembangan
238 populasi kambing Peranakan Etawah di lahan pasca galian pasir. *Jurnal Ilmu Produksi
239 dan Teknologi Peternakan*. 1(3):132 –137.
- 240 La Ode Nafiu, W Kurniawan, PN Kusuma, and M Akramullah. 2017. Produktivitas dan Kualitas Susu
241 Berdasarkan Bangsa dan Paritas kambing di Kabupaten Kolaka. *Seminar Nasional
242 Peternakan* 3. P:19 – 28.
- 243 Marwah MP, YY Suranindyah and TW Murti. 2010. Produksi dan komposisi susu kambing
244 Peranakan Etawa yang diberi suplemen daun katuk (*Sauropus androgynous* L. Merr)
245 pada awal masa laktasi. *Buletin Peternakan*. 54(2):94 – 102.
- 246 Masedunskas A, Y Chen, R Stussman, R Weigert, and IH Mather. 2017. Kinetics of milk lipid droplet
247 transport, growth, and secretion revealed by intravital imaging: lipid droplet release is
248 intermittently stimulated by oxytocin. *MBoC*. 28:935-946.
- 249 Maylinda. 2010. Pengantar Pemuliaan Ternak. Universitas Brawijaya Press. Malang.
- 250 Nwosu EU, IJ James, O Olowofeso, TJ Williams, and MN Bemji. 2019. Effects of oxytocin
251 administration on milk yield and milk composition of West African dwarf does. *Small
252 Ruminant Research*. 181:45-50.
- 253 Ramadhan BG, TH Suprayogi and A Sustiyah. 2013. Tampilan produksi susu dan kadar lemak susu
254 kambing Peranakan Etawa akibat pemberian pakan dengan imbalanced hijauan dan
255 konsentrat yang berbeda. *Animal Agricultural Journal*. 2(1):351 – 356.

- 256 Rizzo A, M Mutinati, G Minoia, M Spedicato, M Panteleo, and RL Sciorsci. 2012. The impact of
 257 oxytocin on the hemodynamic features of the milk vein in dairy cows: A color Doppler
 258 investigation. *Research in Veterinary Science*. 93:983-988.
- 259 Sudewo ATA, SA Santosa and A Susanto. 2012. Produktivitas kambing peranakan etawah
 260 berdasarkan litter size, tipe kelahiran dan mortalitas di village breeding centre
 261 kabupaten banyumas. *Prosiding Seminar Nasional: Pengembangan Sumber Daya*
 262 *Pedesaan dan Kearifan Lokal Berkelanjutan II*. p:1-7.
- 263 Sumaryadi, M.Y. and W. Manalu. 1995. Contributions of Maternal Serum Progesterone and
 264 Estradiol Concentrations or Corpora Luteal and Fetal Number to Mammary Growth
 265 and Development of Ewes During Pregnancy. *Bulletin of Animal Science*. Special
 266 Edition: 243 – 247.
- 267 Sumaryadi, M.Y. dan W. Manalu. 1999a. Prediksi pertumbuhan kelenjar susu produksi susu, dan
 268 involusi kelenjar susu berdasarkan konsentrasi beberapa hormon dan metabolit darah
 269 induk selama kebuntingan pada domba ekor tipis Jawa Barat. *Buletin Peternakan*.
 270 Vol.23 (3):103-126
- 271 Sumaryadi, M.Y. dan W. Manalu. 1999b. Prediksi jumlah anak yang akan dilahirkan dan bobot
 272 lahir anak berdasarkan konsentrasi hormon dan metabolit darah induk selama
 273 kebuntingan pada domba ekor tipis Jawa Barat. *Buletin Peternakan*. Vol. 23 (2):46-63.
- 274 Sumaryadi, M.Y., S. Utami. dan B. Hartoyo. 2002. Peningkatan produksi susu melalui induksi
 275 hormone oksitosin pada sapi perah di tingkat peternakan rakyat. *Jurnal Pembangunan*
 276 *Pedesaan*. Vol. 11 (2): 19 – 25.
- 277 Yustendi D. 2017. Pemanfaatan tanaman katuk (*Sauropus androgynous L. Merr*) dalam ransum
 278 untuk meningkatkan produksi susu kambing betina Peranakan Etawah. *Jurnal Biology*
 279 *Education*. 6(1):21 – 29.

JOURNAL CONTENT

Search

Search Scope

All

Search

Browse

» By Issue

» By Author

» By Title

JOURNAL TEMPLATE

Journal Template

VISITORS

00331617

View My Stats

USER

You are logged in as...

nugrohoaras

» My Profile

» Log Out

» Log Out as User

Home > User > Author > Submissions > #747 > Editing

#747 Editing

SUMMARYREVIEWEDITING

Submission

Authors

Mas Yedi Sumaryadi, Dadang Mulyadi Saleh, Aras Prasetyo Nugroho, Nu'man Hidayat, Chomsiatun Nurul Hidayah

Title

Oxytocin Hormone Induction on Milk Production in Relation to Dairy Kid Performance

Section

Articles

Editor

Agus Susanto

Copyediting

COPYEDIT INSTRUCTIONS

REVIEW METADATA

REQUEST

UNDERWAY

COMPLETE

1.

Initial Copyedit

File: None

2.

Author Copyedit

File: None

Choose File

No file chosen

Upload

3.

Final Copyedit

File: None

Copyedit Comments

No Comments

Layout

Layout Editor

None

Layout Version

REQUEST

UNDERWAY

COMPLETE

VIEWS

None

Galley Format

FILE

1.

PDF

VIEW PROOF

747-3034-1-PB.PDF

2020-03-25

580

Supplementary Files

FILE

None

Layout Comments

No Comments

Proofreading

REVIEW METADATA

REQUEST

UNDERWAY

COMPLETE

1.

Author

2.

Proofreader

3.

Layout Editor

Proofreading Corrections

No Comments

PROOFING INSTRUCTIONS

Animal Production Indexed In

Google Scholar

ISJD

Crossref

MEMDELEY

citeulike

Academic Resource Index Researchbib

E3B Electronic Journals Library

BASE Biofield Academic Search Engine

OAJ Open Academic Journals Index

WORLD Cat

Academic Journals Database

indonesianone search

PKP INDEX

Sinta

DOAJ DIRECTORY OF OPEN ACCESS JOURNALS

IPI

CC BY SA

This work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License.

Editorial Board

Focus & Scope

Author Guidelines

Publication Ethics

Reviewer Acknowledgement

List of Reviewers

Online Submission

Open Access Policy

Plagiarism Policy

Submission and Publication Charge

TOOLS

MEMDELEY

grammarly

KEYWORDS

Chloramphenicol, rabbit doe, toxicity, teratogenicity, fertility Dairy waste water sludge, broiler, microflora, cecum Essential oil, red ginger, rumen fermentation, in vitro Polymorphism, Association, MSTN gene and Batur sheep Production Quran animal breed conception rate dairy diluents type duration of estrus intact plasma membrane l-threonine l-tryptophan mastitis, microorganisms motility native chicken pregnancy rate productivity, pedet, rice straw the onset of estrus

AUTHOR

Submissions

» Active (0)

» Archive (1)

» New Submission

Powered by OJS | Design by ThemeOJS

Back to Top

AWAL

MAIL

BERITA

KEUANGAN

OLAHRAGA

SELEB

LIFESTYLE

LAINNYA...

yahoo!mail PRO

Tingkatkan Sekarang

yahoo!mail

Temukan pesan, dokumen, foto, atau orang

Lanjutan

Awal

Tulis

Email Masuk 916

Belum Dibaca

Berbintang

Draft 44

Terkirim

Arsip

Spam

Sampah

Lebih sedikit

Tampilan Sembunyikan

Foto

Dokumen

Langganan

Folder Sembunyikan

Folder Baru

Kembali

Arsipkan

Pindahkan

Hapus

Spam

[JAP] Editor Decision

Yahoo/Email M... ☆

Dr. Novie Andri Setianto <nsetianto@animalprod>

Kepada: Mr Mas Yedi Sumaryadi

Cc: redaksijap.as@gmail.com

Kam, 19 Mar 2020 jam 15.39

☆

Dear Mr Mas Yedi Sumaryadi:

This is an official notification that your paper titled "Oxytocin Hormone Induction on Milk Production in Relation to Dairy Kid Performance". We have reached a decision regarding your submission to ANIMAL PRODUCTION.

We are pleased to inform you that your paper has been reviewed, and accepted for publication.

Thank you for making the journal a vehicle for your research interests

Sincerely,

Dr. Novie Andri Setianto
Department of Animal Production, Faculty of Animal Science, Jenderal Soedirman University
Phone +62 281 638792
Fax +62 281 626080
nsetianto@animalproduction.net
Editor in Chief
Animal Production
www.animalproduction.net

Animal Production
<http://www.animalproduction.net/index.php/JAP>

Balas, Balas ke Semua atau Teruskan

WAKTUNYA
EASY
BIKIN
PILIHAN

17:00

GO AHEAD

PERINGATAN:

KARENA MEROKOK, SAYA TERKENA
KANKER TENGGOROKAN.
LAYANAN BERHENTI MEROKOK (0800-177-6565)

<https://mail.yahoo.com/d/search/keyword=novie/messages/7755>

1/1

Oxytocin Hormone Induction on Milk Production in Relation to Dairy Kid Performance

Mas Yedi Sumaryadi*, Dadang Mulyadi Saleh, Aras Prasetyo Nugroho,
Nu'man Hidayat and Chomsiatun Nurul Hidayah

Faculty of Animal Science, Jenderal Soedirman University, Purwokerto 53123, Jawa Tengah, Indonesia
*Corresponding author email: yedi.sumaryadi@yahoo.com

Abstract. The application of oxytocin induction to milk production in relation to the performance of kids in dairy goats had been carried out in the "PEGUMAS" dairy goat farmer group Gumelar District, Banyumas. This activity aimed to increase milk production and kid performance through the induction of the hormone oxytocin in dairy goats. The approach method was designed in two stages: the first stage carried out the socialization activities and demonstration of techniques for applying IPTEKS, and the second stage applied hormonal technology in the form of oxytocin hormone induction in dairy goats. The material used was dairy goat parity I aged 1.5 - 2 years allocated into 2 groups of 10 individuals each. Group I was dairy goat experiment not induced with the oxytocin hormone as the control. Group II was dairy goat experiment induced with the oxytocin hormone at a dose of 1 ml (10 IU) intramuscularly before milking. Milk production was measured morning and evening as daily milk production in milliliters (ml). Milk was given to the kid of each parent. The kid's performance was measured based on the daily weight gain of the kid during the study. Overall, the application of science and technology could be adopted by farmers, and oxytocin-induced dairy goat milk production (503.2 ml) was significantly higher than control (305.4 ml) with an increase of 64.77%, with a positive level of correlation ($r = 0.45$) and contributed 19.83% to the kids daily body weight gain.

Keywords: milk yield, oxytocin, daily body weight gain, dairy goat

Abstrak. Aplikasi induksi hormon oksitosin terhadap produksi susu kaitannya dengan kinerja anak pada kambing perah telah dilaksanakan di kelompok peternak kambing perah "PEGUMAS" Kecamatan Gumelar, Banyumas. Kegiatan ini bertujuan untuk meningkatkan produksi susu dan kinerja anak melalui induksi hormon oksitosin pada kambing perah. Metode pendekatan yang digunakan dirancang menjadi dua tahap. Tahap pertama dilakukan kegiatan sosialisasi dan peragaan teknik penerapan IPTEKS. Tahap kedua mengaplikasikan teknologi hormonal berupa induksi hormon oksitosin pada kambing perah. Materi yang digunakan adalah kambing perah paritas I dengan umur 1,5 – 2 tahun yang dialokasikan menjadi 2 kelompok masing-masing 10 ekor. Kelompok I: kambing perah percobaan tidak diinduksi hormon oksitosin yang digunakan sebagai kontrol. Kelompok II: kambing perah percobaan diinduksi hormon oksitosin dengan dosis 1 ml (10 IU) secara intramuskuler sebelum dilakukan pemerahan. Produksi susu diukur pagi dan sore hari sebagai produksi susu harian dalam satuan milliliter (ml). Air susu hasil pemerahan diberikan kepada anak dari masing-masing induk. Kinerja anak diukur berdasarkan pertambahan bobot badan anak harian selama kegiatan. Secara keseluruhan penerapan IPTEKS ini dapat diadopsi peternak dengan baik, dan produksi susu kambing perah yang diinduksi oksitosin (503,2 ml) nyata lebih tinggi dibandingkan kontrol (305,4 ml) dengan peningkatan sebesar 64,77 %, dan memiliki tingkat keeratan hubungan yang positif ($r = 0,45$) serta berkontribusi sebesar 19,83% terhadap pertambahan bobot badan harian anak.

Kata kunci: produksi susu, oksitosin, pertambahan bobot badan harian, kambing perah

Introduction

Improving population and productivity of dairy goat depends on calving process to breed healthy, weaned kids in one reproduction cycle. The potential improvement is attributed to the prolific nature of goat breeding – goat can breed one to four kids per delivery (Kaunang et al., 2012; Kurniasih et al., 2013). However, the

success to raise the kids post-weaning is related to milk availability because milk is the primary source of nutrient for kids since delivery to pre-weaning, and it is essential for kids' growth and development. Malnutrition (milk) in pre-weaned kids would inhibit the growth and decrease immunity against disease, which eventually leads to mortality (Maylinda, 2010).

This incident is prevalent the more kids delivered, the lower the body immune throughout the critical pre-weaning period. Therefore, only a few kids made it through the weaning period.

The increase of dairy goat's milk production for human consumption or the kids is below expectation. Previous studies reported that Etawah goat milk production varied around 0.67 L or 756.8 gram/head/day (Adriani et al., 2014), in a range of 0.20 – 1.3 l/head/day (Marwah et al., 2010; Ramadhan et al., 2013; Astuti et al., 2017; Yustendi, 2017). The current problem is the high mortality rate in kids at an early age or post-weaning time which inhibits the increase of cattle population and productivity. The mortality rate could reach 12.5 – 30.0% (Kaunang et al., 2012; Adriani et al., 2014), up to 28.57 – 57.14% (Kurniasih et al., 2013)

Milk production (milk let down) is naturally controlled by the oxytocin hormone from anterior hypophysis (Bhimte et al., 2018). However, environment-generated stress or milking disorder could inhibit the secretion of endogenous oxytocin through the circulation system; consequently, milk synthesis products in alveoli cells of milk gland is not entirely discharged and potentially causes mastitis disease. Previous studies showed that exogenous oxytocin injection before milking could improve milk production in cow (Belo and Bruckmaier, 2010; Rizzo et al., 2012) and buffaloes (Akhtar et al., 2012). Furthermore, it could increase milk production by 15.32% in oxytocin-induced sheep (Antonic et al., 2013). It provides an opportunity for further application to goat cattle, particularly dairy goat.

The result of this research focused on increasing milk production as well as improving the performance of weaned dairy kids. Additionally, milk is a potential food product for human consumption.

Materials and Methods

The research method applied a two-stage design. The first stage included socialization and exhibition of science-technology application technique. The second stage was hormonal technology application through oxytocin induction into dairy goats.

The materials used were dairy goats at parity I aged 1.5 – 2 years old allocated into two groups, ten goats each. Group 1 was an experimental dairy goat without oxytocin induction as control groups, and Group II was experimental dairy goats induced with 1 mL (10 IU) oxytocin hormone intramuscularly before milking.

Milk production was measured in the morning and afternoon in ml, then provided to kids in a separate cage from each mother. Kid performance was measured by body weight per time unit to obtain daily body weight gain throughout the study.

Success evaluation was conducted at the end of activity in two waves. The first wave was the farmers' level of adoption to apply science-technology application, and the second wave was testing an oxytocin-induction application on milk production and kid performance.

Results and Discussions

Oxytocin induction to dairy goats was started with socialization to disseminate information regarding the activity materials and the exhibition of science-technology application for the dairy goat farmers. The result of socialization showed that 90% of the farmers could adopt the oxytocin induction to improve milk production and kid performance. Total milk production at the initial phase from Peranakan Etawah (PE), Saanen and Jawa Randu goats parity I aged 1.5 – 2 years old is presented in Figure 1.




Figure 1. Average daily milk production of dairy goats

The statistical analysis result showed that milk production was not significantly different ($P>0.05$) across goat breeds (PE, Saanen and Jawa Randu). The average milk production of dairy goat in this study was 435.37 ± 248.09 ml per head. The contributing factors may include similar age, parity, lactating period and breeding management in addition to the long adaptation to the local environment. However, the average milk production was higher than that of Etawah crossbred, Etawah-Kacang crossbred (PEK) and Etawah-Boer crossbred, namely 272.58 ± 55.73 , 153.77 ± 24.76 , and 174.61 ± 35.22 ml, respectively (La Ode Nafiu et al., 2017). According to Hossain et al. (2018), cattle potential productivity is attributed to genetic factors, environmental factors and genetic-environment interaction. Also, goat productivity is influenced by climate, parity, litter size and lactating period (Sudewo et al., 2012). Milk production of dairy goats before and after oxytocin hormone induction measured in the morning and afternoon, as well as total milk production presented in Table 1.

Table 1 shows that oxytocin-induced dairy goats produced more milk in the morning and afternoon milking by 83.80 and 136.14%, respectively, compared to non-induced dairy

goats (control group). Statistical analysis result (see appendix) showed that oxytocin hormone significantly affected ($P<0.05$) total daily milk production. Therefore, the average daily milk production of oxytocin-induced dairy goats (503.2 ml) was significantly higher than that of the control group (305.4 ml) with a 64.77% increase in milk production. In line with the previous study, oxytocin injection in goat can increase milk production, especially milk fat (Nwosu et al., 2019). The current study showed a higher milk production in oxytocin-induced dairy goats than that in sheep (Antonic et al., 2013). Milk production is affected by total secretory cells in udder tissue, secretory cell activity to synthesize milk, and substrate availability for milk synthesis. Milk synthesis is conducted by secretory cells in milk gland using nutrition from feed (Kim and Wu, 2009). Stimulation from milking is forwarded to the hypothalamus that produces oxytocin hormone. Furthermore, oxytocin stimulates contraction in the uterus wall (Heppelmann et al., 2018) and would trigger soft muscle around alveoli to contract and secretes milk by stimulating the ductus lactiferous contraction of milk gland (udder) to secrete milk (Masedunskas et al., 2017).

Table 1. The average milk production of dairy goats induced with oxytocin hormone

Variables	Control	Oxitocyn induction
Milk production (ml)		
Morning	358.0 ± 274.9	658.0 ± 320.6
Afternoon	147.5 ± 99.5	348.3 ± 200.8
Total per day (ml/day)	305.4 ± 229.6 ^a	503.2 ± 249.6 ^b

(a,b) Values bearing different superscript within rows show a significant difference (P<0.05)

The average daily milk production of dairy goats in this study was 435.37 ± 248.09 ml per head. It was higher than that of Etawah dairy goats (PE), Etawah-Kacang crossbred (PEK), and Etawah-Boer crossbred (PEB) namely 272.58 ± 55.73 , 153.77 ± 24.76 , and 174.61 ± 35.22 ml, respectively (La Ode Nafiu et al., 2017). Daily body weight gain (DWG) of kid consuming the milk of each mother was 191.25 ± 91.60 g/day. The DWG was higher than that reported by Ginting (2009) on Boer goats (118 g/day) and Kacang goats (52 – 70 g/day).

There is a significant (P<0.05) correlation between daily milk production and DWG with 0.45 coefficient correlation. The regression analysis result showed that daily milk production of dairy goats was significantly correlated (P<0.05) with kid's daily weight gain (DWG) using an equation $Y = 0.24 + 0.000217x$ and 0.1983 coefficient of determination. It showed that kid's DWG was 19.83% affected by daily milk production offered to the kids from each mother.

Post-calving growth is significantly affected by milk consumption sufficiency before taking forage. According to Elkhair et al. (2017), kid's growth from the first month of age was solely dependent on the mother's milk – the higher the dependency, the lower milk production when the kids start eating solid food. Similarly, Htoo et al. (2018) reported that kids were dependent on their mother's milk up to 12 weeks post-calving before their rumens start to function and forage intake ensue. Pre-weaning growth was determined by birth weight, milk production, mother's welfare, total calves and their conditions, sex, genetic and breeding management (Maylinda, 2010).

Conclusions

Farmers well adopted Oxytocin hormone induction to improve milk production and kid performance in PEGUMAS association of dairy goat farmers. Oxytocin induction could increase milk production by 64.77%, showed a close relationship ($r=0.45$) and contributed 19.83% to kid's daily weight gain. However, this science-technology application needs further investigation in the economic aspect and utilization of goat milk for human consumption.

Acknowledgement

The authors express their gratitude to the Chief of Research Centre and Community Service, Jenderal Soedirman University for the opportunity and research grant through DIPA funding Jenderal Soedirman University 2019 according to the Letter of Assignment to Community Service for implementing Science and Technology No: P/916/UN 23/14/PM/2019. We also thank Widi Utami Prahayuningtyas, a student of Magister Program for assisting community service, as well as the chief and members of PEGUMAS dairy goat farmer's community for the contribution to this study.

References

- Adriani, S. N. and Fatati. 2014. Usaha IBKK kambing Perah. Jurnal Pengabdian pada Masyarakat. 29(4):1 – 10.
- Akhtar, M. S., L. A. Lodhi, A. A. Farooq, M. M. Ayaz, M. Hussain, M. H. Lashari and Z. I. Chaudhary. 2012. Effect of oxytocin administration before milking on milk production, somatic cells count and fat contents in milk of Nili-Ravi Buffaloes. Pakistan Veterinary Journal. 32(3):435-437.
- Antonic, J., V. Tancin, M. Uhrincat, L. Macuhova, J. Macuhova and L. Jackuliakova. 2013. The effect of exogenous oxytocin on milkability and milk

- composition in ewes differed in milk flow pattern. *Small Ruminant Research*. 113:254-257.
- Astuti, P., H. Surti and N. E. Sukarini. 2017. Produksi dan komposisi susu kambing Peranakan Ettawa melalui pemberian ekstrak meniran. *Jurnal Ilmu-ilmu Pertanian*. 1(2):82 – 87.
- Belo, C. J. and R. M. Bruckmaier. 2010. Sutability of low-dosage oxytocin treatment to induce milk ejection in dairy cows. *Journal of Dairy Science*. 93:63-69.
- Bhimte, A., N. S. Thakur, V. P. Maurya and G. Singh. 2018. Neurohormonal control of lactation and milk let-down in dairy animals. *International Journal of Current Microbiology and Applied Sciences*. 7(7):970-977.
- Elkhair, A. A. M., I. A. Abdalla and G. H. El Obied. 2017. Effect of a home-made milk replacer on pre-weaning performance of Nubian male kids. *Gezira J. of Agric. Sci*. 15(2):107-118.
- Ginting, S. P. 2009. Pedoman Teknis Pemeliharaan Induk dan Anak Kambing Pra-Sapih. Pusat Penelitian dan Pengembangan Peternakan, Loka Penelitian Kambing Potong. Sei Putih, Sumatera Utara. p:33-44.
- Heppelmann, M., J. Volland, C. Pfarrer, M. Kietzmann, W. Baumer, S. Merbach, H. A. Schoon, O. Wellnitz, M. Schmicke, M. Hoedemarker and H. Bollwein. 2018. Effects of oxytocin and PGF2a on uterine contractility in cows with and without metritis – An in-vitro study. *Animal Reproduction*. 188:144-154.
- Hossain, S. M. J., M. F. Afroz, A. K. F. H. Bhuiyan and M. A. Habib. 2018. Milk Production Performance of Red Chittagong Cattle Based on Phenotypic and Genetic Parameters. *Glo. Adv. Res. J. Agric. Sci*. 7(4):110-117.
- Htoo, N. N., B. Zeshan, A. T. Khaing, T. Kyaw, E. A. Woldegiorgis and M. A. Khan. 2018. Creep Feeding Supplemented with Roughages Improve Rumen Morphology in Pre-Weaning Goat Kids. *Pakistan Journal of Zoology*. 50(2):703-709.
- Kaunang, D., Suyadi and S. Wahyuningsih. 2012. Analisis litter size, bobot lahir dan bobot sapih hasil perkawinan kawin alam dan inseminasi buatan kambing Boer dan Peranakan Etawah. *Jurnal Ilmu-ilmu Peternakan*. 23(3):41– 46.
- Kim, S. W. and G. Wu. 2009. Regulatory role for amino acids in mammary gland growth and milk synthesis. *Amino Acids*. 37: 89-95.
- Kurniasih, N. N., A. M. Fuah and R. Priyanto. 2013. Karakteristik reproduksi dan perkembangan populasi kambing Peranakan Etawah di lahan pasca galian pasir. *Jurnal Ilmu Produksi dan Teknologi Peternakan*. 1(3):132 –137.
- La Ode Nafiu, W. Kurniawan, P. N. Kusuma and M. Akramullah. 2017. Produktivitas dan Kualitas Susu Berdasarkan Bangsa dan Paritas kambing di Kabupaten Kolaka. *Seminar Nasional Peternakan* 3. P:19 – 28.
- Marwah, M. P., Y. Y. Suranindiyah and T. W. Murti. 2010. Produksi dan komposisi susu kambing Peranakan Etawa yang diberi suplemen daun katuk (*Sauropus androgynous L. Merr*) pada awal masa laktasi. *Buletin Peternakan*. 54(2):94 – 102.
- Masedunskas, A., Y. Chen, R. Stussman, R. Weigert, and I. H. Mather. 2017. Kinetics of milk lipid droplet transport, growth, and secretion revealed by intravital imaging: lipid droplet release is intermittently stimulated by oxytocin. *MBOC*. 28:935-946.
- Maylinda. 2010. Pengantar Pemuliaan Ternak. Universitas Brawijaya Press. Malang.
- Nwosu, E. U., I. J. James, O. Olowofeso, T. J. Williams and M. N. Bemji. 2019. Effects of oxytocin administration on milk yield and milk composition of West African dwarf does. *Small Ruminant Research*. 181:45-50.
- Ramadhan, B. G., T. H. Suprayogi and A. Sustiyah. 2013. Tampilan produksi susu dan kadar lemak susu kambing Peranakan Etawa akibat pemberian pakan denganimbangan hijauan dan konsentrat yang berbeda. *Animal Agricultural Journal*. 2(1):351 – 356.
- Rizzo, A., M. Mutinati, G. Minoia, M. Spedicato, M. Panteleo, and R. L. Sciorsci. 2012. The impact of oxytocin on the hemodynamic features of the milk vein in dairy cows: A color Doppler investigation. *Research in Veterinary Science*. 93:983-988.
- Sudewo, A. T. A., S. A. Santosa and A. Susanto. 2012. Produktivitas kambing peranakan etawah berdasarkanlitter size, tipe kelahiran dan mortalitas di village breeding centre kabupaten banyumas. *Prosiding Seminar Nasional: Pengembangan Sumber Daya Pedesaan dan Kearifan Lokal Berkelanjutan II*. p:1-7.
- Yustendi, D. 2017. Pemanfaatan tanaman katuk (*Sauropus androgynous L. Merr*) dalam ransum untuk meningkatkan produksi susu kambing betina Peranakan Etawah. *Jurnal Biology Education*. 6(1):21 – 29.