

IOP Proceeding 2020

by Ardhini Maharning

Submission date: 13-May-2023 08:40PM (UTC+0700)

Submission ID: 2092102612

File name: IOP_EES_Ekowati_Maharning_593_2020.pdf (748.57K)

Word count: 2911


Character count: 15578

PAPER • OPEN ACCESS**Effects of Ethyl Acetate Extract of Jew's Ear Mushrooms (*Auricularia auricula*) on Cytotoxic and Apoptosis of Cervical Cancer Cells (HeLa)****1**To cite this article: Nuraeni Ekowati *et al* 2020 *IOP Conf. Ser.: Earth Environ. Sci.* **593** 012011View the [article online](#) for updates and enhancements.**The Electrochemical Society**
Advancing solid state & electrochemical science & technology

The ECS is seeking candidates to serve as the
Founding Editor-in-Chief (EIC) of ECS Sensors Plus,
a journal in the process of being launched in 2021

The goal of ECS Sensors Plus, as a one-stop shop journal for sensors, is to advance the fundamental science and understanding of sensors and detection technologies for efficient monitoring and control of industrial processes and the environment, and improving quality of life and human health.

Nomination submission begins: May 18, 2021



Effects of Ethyl Acetate Extract of Jew's Ear Mushrooms (*Auricularia auricula*) on Cytotoxic and Apoptosis of Cervical Cancer Cells (HeLa)

Nuraeni Ekowati, Ardhini R Maharning, Nuniek I Ratnaningtyas, Aris Mumpuni, and Arif R Hikam

Faculty of Biology, Universitas Jenderal Soedirman, Purwokerto, Indonesia

E-mail: nuraeniekowati@yahoo.com

Abstract. The increasing number of people living with cervical cancer encourages the search for bioactive compounds from natural ingredients such as macroscopic fungi that have the potential to induce apoptosis. Jew's ear fungus (*Auricularia auricula*) is an edible mushroom that is also commonly used by society for medicinal purposes including curing cancer. The study aimed to examine the cytotoxic, antiproliferative, and apoptotic effects of ethyl acetate extract from the mushroom on cervical cancer cells (HeLa) *in vitro*. The research consisted of experiments. Cytotoxic and antiproliferative tests were carried out using MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide) assay, while apoptosis test was the double staining method using acridine orange/ethidium bromide. The data of cytotoxic and antiproliferative effects were analyzed with linear regression, whereas apoptosis test result was analyzed descriptively. The results showed that the best cytotoxic effect was ethyl acetate extract of mushroom fruit body with IC₅₀ of 538 µg/ml. Antiproliferative analysis of HeLa cells showed the best inhibitory effect of the mushroom fruit body extract at 24-hour incubation time. The apoptosis test showed that HeLa cell death through apoptosis mechanisms was higher than by necrosis.

1. Introduction

Auricularia auricula (Bull.) J. Schröt. is an edible mushroom found throughout the world. It grows on wood with ear-like shape, and brown fruiting body. The fungus is useful as a medicine due to its bioactive compounds. It has been used as a traditional medicine in China, but further scientific evidence to support it is still required⁴. The dichloromethane extract of *A. auricula* has the potential to be anti-inflammatory and inhibits the production of Nitric Oxide (NO), Interleukin 6 (IL-6), Tumor Necrosis Factor- α (TNF- α), and Interleukin-1 β (IL-1 β) in macrophages [1].

A. auricula has been extensively studied for its use as a medicine and a source of new chemicals with potential therapeutic effects. In a study of water extract from seven edible fungal species tested for antitumor activity against Sarcoma 180 transplantation in Swiss albino mice, the results showed that *A. auricula* significantly inhibited the growth of transplanted tumors [2]. Furthermore, studies of the hypoglycemic effect of water-soluble polysaccharides from *A. auricula* fruit bodies were investigated in genetic diabetic rats (KK-Ay) ages 10-14 weeks. The rats showed a decrease in plasma glucose, insulin, and urine glucose [3]. Other studies about the effects of *A. auricula* on anticoagulant aggregation have been investigated [4]. No results, however, concerning the anticancer effects of *A. auricula* ethyl acetate extract from Indonesia have been reported.



² Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Published under licence by IOP Publishing Ltd

The potential of bioactive compounds of *A. auricula* as an anticancer can be detected through testing the cancer cells directly [5, 6]. One type of cancer cell used for *in vitro* testing is cervical cancer cells (HeLa cells). Cervical cancer is the first incidence of cancer in women, and no treatment method is yet available [7].

The most widely used cancer treatment is chemotherapy. This method, however, has side effects, one of which is the high damage of healthy cells. Thus, through this study, it is expected to reveal the bioactive metabolites potential of *A. auricula* in suppressing HeLa cell growth. Bioactive metabolites of *A. auricula* can be isolated from the fruiting bodies, which are obtained by cultivation of the fungus on solid medium. The metabolites can also be isolated from mycelium biomass and culture filtrate by liquid medium cultivation [8]. The β -glucan is the fungus bioactive metabolite which has been studied for its therapeutic potential [5], however, other bioactive metabolites require further studies. Thus, it is necessary to isolate and identify the classes of bioactive metabolites produced by *A. auricula* and test their potential. The study aimed to examine the cytotoxic, antiproliferative, and apoptotic effects of ethyl acetate extract from ear mushroom on cervical cancer cells (HeLa) *in vitro*.

2. Methods

The method used was *in vitro* experimental research. Four treatments were used in the experiments, i.e., fruit body, mycelium, and culture filtrate of *A. auricula*, and Doxorubicin. Anticancer activity test was applied to cervical cancer cells (HeLa). Cytotoxic and antiproliferative tests were carried out using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) assay. A double staining method using acridine orange and ethidium bromide dyes was applied for apoptosis test. Cytotoxic and antiproliferative data were analyzed by linear regression whereas descriptive analysis was for apoptosis test. Cytotoxic tests were used to treat extracts of secondary metabolites with concentrations of 62.50, 125, 250, 500 and 1,000 $\mu\text{g/ml}$. The variables observed were living cells, and Inhibition Concentration 50 (IC_{50}) values, cervical cancer cell (HeLa) morphology, cancer cell proliferation kinetics, and cancer cell death mechanisms.

The experiments included (1) making the Roswell Park Memorial Institute (RPMI) medium for cancer cells, (2) preparing and multiplying cervical cancer cells (HeLa), (3) diluting the bioactive compound extract for treatments with the given concentration series, and (4) testing the cytotoxic activity of each cancer cell.

Bioactive metabolite testing procedure [9]

A total of 100 μl of the extract was taken from the treated medium prepared with the varied metabolite concentrations, and transferred to a 96-well plate containing 100 μL of HeLa cells aged 24 hours. Each 96-well plate was used to test four treatments with each using five series of extract concentrations (three replicates). The wells with no extract were used for cell and medium controls (three replicates each). For medium control, 200 μl of complete medium was filled into well, while for cell control, 100 μl of complete medium was filled with 100 μl of HeLa cells (10^4 cells). Furthermore, the 96-well plates that already contained cells and extracts were incubated in an incubator with a flow of 5% carbon dioxide at 37° C for 24 hours.

The cells in each extract concentration were observed and photographed under an inverted microscope after incubation. By reversing the plate, the media in the 96-well plate were removed. The remnants in the well were washed with PBS 1x. In each well, 10 μl (with a concentration of 0.5 mg/ml in PBS) MTT reagents were added, and 100 μl complete medium for each well including the control medium, were incubated for 4 hours in a 5% CO_2 incubator. After 4 hours, it was observed under an inverted microscope. When formazan was formed, the observation was stopped by adding 100 μl of SDS 10% stopper reagent in 0.1 N HCl. The enzymes of the living cells reacted with MTT demonstrating purple or blue color in the cell. The plate was wrapped in paper and incubated overnight at room temperature. The following procedure was observing the plate with ELISA reader. The plate was opened and placed on the ELISA reader with a wavelength of 595 nm ($\lambda = 550\text{-}600$

nm). The readable absorbance was converted into the percentage of HeLa living cell. Data were analyzed by regression analysis to obtain Inhibition Concentration 50 (IC₅₀) values.

Data analysis

The living cells were counted for cytotoxicity analysis. The absorption responses of the cytotoxicity test were converted into percent of cancer cell life with the following formula:

$$\text{Living cell} = \frac{\text{absorbance of treatment} - \text{absorbance of control medium}}{\text{absorbance of control cell} - \text{absorbance of control medium}} \times 100\%$$

Based on the results of living cells, the IC₅₀ values were calculated using the Statistics for Window program.

3. Results

IC₅₀ values of anticancer tests showed various results. The best value was obtained from ethyl acetate extract of the fruit body (IC₅₀ value of 538 µg/mL), followed by ethyl acetate extract of mycelium (IC₅₀ value of 579 µg/mL) whereas ethyl acetate extract of culture filtrate reached IC₅₀ value of 2,246 µg/mL. Table 1 presented the IC₅₀ values of *A. auricula* extract.

Table1. IC₅₀ values of *A. auricula* extract

No	Treatments	IC ₅₀ (µg/ml)
1	<i>A. auricula</i> ethyl acetate extract of the fruitbody	53
2	<i>A. auricula</i> ethyl acetate extract of the mycelium	579
3	<i>A. auricula</i> ethyl acetate extract of the culture filtrate	2,246
4	Doxorubicin	15,80

The linear regression analysis correlating the concentration of *A. auricula* extract to HeLa cell viability indicated the inhibition of HeLa cell growth. Ethyl acetate extract of fruitbody (Figure 1), and mycelium (Figure 2) demonstrated sharper decreasing slope than culture filtrate (Figure 3), i.e., 0.092, 0.099, and 0.02 respectively. The antiproliferative test revealed an increasing trend over time except for Doxorubicin (positive control) (Figure 4). Figure 5 represented the micrograph of HeLa morphological cells showing the damaged and healthy cells.




Figure 1. Linear regression curve for ethyl acetate extract of *A. auricula* fruit body




Figure 2. Linear regression curve for ethyl acetate extract of *A. auricula* mycelium




Figure 3. Linear regression curve for ethyl acetate extract of *A. auricula* culture filtrate




Figure 4. Antiproliferative test for ethyl acetate extract of *A. auricula* (EeTbAa: fruit body, EeMiAa: mycelium).




Figure 5. HeLa cell morphology from cytotoxic tests (left: damaged HeLa cells, right: healthy HeLa cells).

4. Discussion

The IC_{50} values indicate that the two types of extract (fruit body and mycelium) have potential anticancer compounds for HeLa cells. The ethyl acetate extract of culture filtrate showed value higher than 1,000 $\mu\text{g/mL}$ suggesting no potential anticancer compound. An IC_{50} value of 45.3 $\mu\text{g/mL}$ from ethyl acetate extract of *A. auricula* fruit body cultivated in Egypt was able to inhibit HCT116 colon cancer cells [10]. Testing of ethyl acetate extract carried out in Korea reported that ethyl acetate extract (IC_{50} of 133 $\mu\text{g/mL}$) inhibited Sarcoma 180 cells [11]. The IC_{50} value ≤ 1000 $\mu\text{g/mL}$ suggests a good anticancer activity and that the extract is potential for further testing [12].

The fruit body and mycelium extract of *A. auricula* showed to inhibit HeLa cell viability. The viability of HeLa cells decreased sharply as the concentration of *A. auricula* extract increased. At the extract concentration of 1,000 $\mu\text{g/mL}$, the HeLa cell viability reached a zero value indicating death. This differed from culture filtrate extract that demonstrated a slow decrease of HeLa cell viability. This condition was possibly due to the capacity of bioactive compounds in the extract was low, thus unable to suppress HeLa cell growth. Even in a concentration of 1,000 $\mu\text{g/mL}$ the extract was not able to suppress HeLa cell growth by 50%. Thus, the extract of culture filtrate is not potential as a HeLa cell anticancer agent.

The subsequent tests were antiproliferative test and apoptosis test. The tests were only used for extracts of fruit body and mycelium of *A. auricula*. The fruit body extract of *A. auricula* mushroom has been studied and claimed that *A. auricula* contains alkaloid compounds, flavonoids, saponins, steroids, and carbohydrates [13]. With the various contents of the bioactive compounds, *A. auricula* is very potential as an anticancer compound.

Observation of apoptosis test on HeLa cells showed that the ethyl acetate extract of *A. auricula* fruit body and mycelium induced apoptosis. Red marks of the DNA showed the apoptosis as a result of ethidium bromide dye penetrating the cells through the damaged membrane. The apoptosis was characterized by fragmented, damaged cells. There were visible circles in cells that were still covered by cell membranes. A small number of cells also experienced necrosis characterized by the cell membrane rupture so that the cell cytoplasm spilled out and toxic for the surrounding cells. *A. auricula* extract might damage cancer cells through chromatin disintegration mechanism that induces apoptosis [14].

5. Conclusion

The results showed that the best cytotoxic effect of Jew's Ear Mushroom extract was the mushroom fruit body with an IC₅₀ value of 538 µg/ml (medium level). Antiproliferative analysis toward HeLa cells showed the best inhibitory effect in ethyl acetate extract of mushroom fruit body and the best part-time effect was at 24-hour incubation. The apoptosis test showed that cell death through apoptosis mechanisms was more than necrosis.

References

- [1] Damte D, Reza A, Lee S J, Jo W S and Park S C 2011 Anti-inflammatory activity of Dichloromethane extract of *Auricularia auricula-judae* in RAW264.7 cells *Toxicol. Res.* **27** 11–4
- [2] Misaki A, Kakuta M, Sasaki T, Tanaka M and Miyaji H 1981 Studies on interrelation of structure and antitumor effects of polysaccharides: antitumor action of periodate-modified, branched (1→3)-β-D-glucan of *Auricularia auricula-judae*, and other polysaccharides containing (1→3)-glycosidic linkages *Carbohydrate Research* **92** 115–29
- [3] Yuan Z, He P, Cui J and Takeuchi H 1998 Hypoglycemic effect of water-soluble polysaccharide from *Auricularia auricula-judae* Quel on genetically diabetic KK-A^y mice *Biosci. Biotechnol. Biochem.* **62** 1898–903
- [4] Yoon S J, Yu M A, Pyun Y R, Hwang J K, Chu D C, Juneja L R and Mourão P A S 2003 The nontoxic mushroom *Auricularia auricula* contains a polysaccharide with anticoagulant activity mediated by antithrombin *Thromb. Res.* **112** 151–8
- [5] Smith J E, Rowan N J and Sullivan R 2002 *Medicinal Mushrooms: Their Therapeutic Properties and Current Medical Usage with Special Emphasis on Cancer Treatments* (London, UK: Cancer Research UK, University of Strathclyde)
- [6] Israilides C, Kletsas D, Arapoglou D, Philippoussis A, Pratsinis H, Ebringerová A, Hribalová V and Harding S E 2008 In vitro cytostatic and immunomodulatory properties of the medicinal mushroom *Lentinula edodes* *Phytomedicine* **15** 512–9
- [7] Tjindarbumi D and Mangunkusumo R 2002 Cancer in Indonesia, present and future *Jpn. J. Clin. Oncol.* **32** 17–21
- [8] Ekowati N, Aris M and Muljowati J S 2017 Effectiveness of *Pleurotus ostreatus* extract through cytotoxic test and apoptosis mechanism of cervical cancer cells *Biosaintifika* **9** 148–55
- [9] Langdon S P 2003 *Methods in Molecular Medicine: Cancer Cell Culture, Methods and Protocols* (Totowa, NJ: Humana Press Inc.)
- [10] Elkhateeb W A, El-Hagrassi A M, Fayad W, El-Manawaty M A, Zaghlol G M, Daba G M and Ahmed E F 2018 Cytotoxicity and hypoglycemic effect of the Japanese jelly mushroom *Auricularia auricula-judae* *Chemistry Research Journal* **3** 123–33
- [11] Reza A, Choi M J, Damte D, Jo W S, Lee S J, Lee J S and Park S C 2011 Comparative antitumor activity of different solvent fractions from an *Auricularia auricula-judae* ethanol extract in P388D1 and sarcoma 180 cells *Toxicol. Res.* **27** 77–83
- [12] Meiyanto E, Susidarti R A, Handayani S and Rahmi F 2008 Ethanol extract of areca nut seeds (*Areca catechu* L.) is able to inhibit proliferation and stimulate MCF-7 cell apoptosis *Pharmacy Magazine Indonesia* **19** 12–9
- [13] Essien E E, Mkpene V N and Akpan S M 2015 Phytochemical and mineral elements composition of *Bondazewia berkeleyi*, *Auricularia auricula* and *Ganoderma lucidum* fruiting bodies *Res. J. Pharm. Biol. Chem. Sci.* **6** 200–4

- [14] Gurusamy R and Arthe R 2012 Effect of medicinal mushroom, *Auricularia auriculajudae* polysaccharides against EAC cell lines *Res. J. Biotech.* **7** 14–7

ORIGINALITY REPORT

14%

SIMILARITY INDEX

12%

INTERNET SOURCES

13%

PUBLICATIONS

11%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Universitas Jenderal Soedirman Student Paper	8%
2	backend.orbit.dtu.dk Internet Source	2%
3	Vartika Mishra, Sarika Tomar, Priyanka Yadav, M.P. Singh. "Promising anticancer activity of polysaccharides and other macromolecules derived from oyster mushroom (<i>Pleurotus</i> sp.): An updated review", International Journal of Biological Macromolecules, 2021 Publication	1%
4	referencecitationanalysis.com Internet Source	1%
5	dlibrary.univ-boumerdes.dz:8080 Internet Source	1%
6	moam.info Internet Source	<1%
7	mail.scialert.net Internet Source	<1%

8

Yupeng Chen, Yuanyuan Wu.
"Hsa_circ_0070661 inhibits cancer progression
through miR-556-5p/TEK axis in lung
adenocarcinoma", Cancer Biomarkers, 2023

Publication

<1 %

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On