

Determinant Factors Stroke Prevention Behavior among Hypertension Patient in Indonesia

by Yunita Sari

Submission date: 09-Oct-2022 03:42PM (UTC+0700)

Submission ID: 1920389438

File name: oamjms-9e-336.pdf (211.58K)

Word count: 2708

Character count: 14310



Determinant Factors Stroke Prevention Behavior among Hypertension Patient in Indonesia

Arif Setyo Upoyo*, Atyanti Isworo, Yunita Sari, Agis Taufik, Annas Sumeru, Akhyarul Anam

Department of Nursing, Faculty of Health Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia

Abstract

Stroke is the leading cause of death in the world and in Indonesia. Hypertension is a major risk of stroke. Modified stroke risk factors are behavioral factors. The purpose of this study is to identify the factors that influence stroke prevention behavior in hypertensive patients. The research method uses cross sectional. Total participants were 461 hypertensive patients who came to Community Health Center. Sampling technique used consecutive sampling. The questionnaire used contains demographics, knowledge, self-efficacy, awareness of stroke risk, and stroke prevention behaviors which include diet, exercise, smoking, and drinking alcohol. Data analysis used Chi-square and logistic regression. The results showed that 65.7% of respondents showed poor stroke prevention behavior. Factors related to stroke prevention behavior include self-efficacy ($p = 0.043$; 95%CI = 1.01–2.22; OR = 1.5), perception ($p = 0.038$; 95%CI = 0.45–0.98; OR = 0.663), and awareness of stroke risk ($p \leq 0.001$; 95%CI = 2.255–6.333; OR = 3.779). The dominant factor influencing stroke prevention behavior in hypertensive patients is awareness of stroke risk.

Edited by: Sasho Stoleski
Citation: Upoyo AS, Isworo A, Sari Y, Taufik A, Sumeru A, Anam A. Determinant Factors Stroke Prevention Behavior among Hypertension Patient in Indonesia. Open Access Maced J Med Sci. 2021 May 07; 9(E):336-339. <https://doi.org/10.3889/oamjms.2021.6040>
Keywords: Behavior; Stroke prevention; Hypertension; Awareness
***Correspondence:** Arif Setyo Upoyo, Department of Nursing, Faculty of Health Sciences, Universitas Jenderal Soedirman, E-mail: adholi@gmail.com
Received: 17-Mar-2021
Revised: 24-Apr-2021
Accepted: 28-Apr-2021
Copyright: © 2021 Arif Setyo Upoyo, Atyanti Isworo, Yunita Sari, Agis Taufik, Annas Sumeru, Akhyarul Anam
Funding: This study was supported by the Research Community Service Institute of Universitas Jenderal Soedirman, Indonesia
Competing Interest: The authors have declared that no competing interest exists
Open Access: This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

Introduction

Stroke is a major health-care problem in South, East, and Southeast Asia. With a majority of the world's population living in the developing countries of these regions, the global burden of stroke will have the largest contribution from Asia [1]. The prevalence of stroke in Indonesia increases every year. Based on the 2013 Basic Health Research, the prevalence of stroke was 7 per thousand population, while in 2018, the prevalence of stroke was 10.9 per thousand population [2].

Risk factors for stroke include factors that cannot be modified and factors that can be modified. Factors that cannot be modified include age, sex, race/ethnicity, and genetics, while factors that can be modified include hypertension, diabetes, atrial fibrillation, dyslipidemia, diet, physical activity, obesity, metabolic syndrome, alcohol consumption, and smoking [3], [4], [5]. Risk factors for stroke that can be modified are influenced by patient behavior.

Hypertension is one of the main factors of stroke. Primary hypertension is a major cause of cardiovascular disease morbidity which has a significant influence on the incidence of stroke [6], [7]. Goldstein states that the prevalence of stroke in hypertensive patients aged 50 years is 20% with a risk ratio of 4 and the prevalence

continues to increase with increasing age [8], whereas according to Ghani *et al.*, hypertension sufferers have a risk of 2.87 times getting a stroke [7].

The purpose of this study was to determine the factors that influence stroke prevention behavior in hypertensive patients.

Methodology

Design

The research method used cross-sectional study. The research was conducted in April–August 2019.

Sample

The subjects of the study were 461 participants of hypertension. Sampling technique used consecutive sampling from five public health centers in Banyumas Regency, Central of Java. Inclusion criteria of sample were hypertensive patient, >35 years old and visit to public health center. Exclusion criteria were hypertensive patients with complications and refused as a respondent.

Instrument

The questionnaire used contained demographics, knowledge, self-efficacy, awareness of stroke risk, and stroke prevention behaviors. Knowledge instrument modified from Hypertension Knowledge-Level Scale by Erkok *et al.* [9] that divided into three categories: Low, middle, and high. Self-efficacy and awareness of stroke risk used numeric score 1–10, high score if above average, and low score if below or equal to average. Stroke prevention behaviors which included low salt and cholesterol diet, exercise, smoking, and drinking alcohol used Likert score 1–4, total score categorized good if the total score is more than average and bad if the total score is below or equal to average.

Data analysis

Data analysis used Chi-square for bivariate analysis and logistic regression for multivariate analysis with SPSS 16.

Results

Based on Table 1, it is known that the respondents are mostly female (80.9%), over 60 years old (56.8%), still married (71.4%), living together with family (49.9%), with income below the regional minimum wage (65.7%), and low education (70.9%). Respondent characteristics that significantly influence behavior include gender ($p=0.000$; 95%CI=7.12–22.28; OR=12.59), age ($p = 0.000$; 95%CI = 0.24–0.56; OR = 0.369), marital status ($p = 0.008$; 95%CI = 1.17–2.89; OR = 1.83), and income ($p = 0.008$; 95%CI = 0.39–0.87; OR = 0.58). Female, old age, married, and low income have a tendency to behave badly in stroke prevention behavior.

Based on Table 2, it is known that the majority showed poor stroke prevention behavior (65.7%), high

19
Table 1: Characteristics of respondents

14 Characteristics of respondents	n	%	OR	95%CI	p
Sex					
Male	88	19.1	12.59	7.12–22.28	<0.001
Female	373	80.9			
Age					
<60 years old	199	43.2	0.369	0.24–0.56	<0.001
>60 years old	262	56.8			
Marital status					
Married	329	71.4	1.83	1.17–2.89	0.008
Not yet	132	28.6			
Living together					
Living alone	12	2.6	1.27	0.86–1.87	0.227
With couple	230	49.9			
With family	201	43.6			
Other	18	3.9			
Income					
Under minimum standard payment	303	65.7	0.58	0.39–0.87	0.008
Standard payment	24	5.2			
17, per minimum standard payment	134	29.1			
Level education					
No	40	8.7	0.807	0.39–1.63	0.551
Elementary	327	70.9			
Junior high school	57	12.4			
Senior high school	34	7.4			
High education	3	0.7			

knowledge of hypertension (89.4%), low self-efficacy (56.2%), good perception (57.3%), information about low stroke (41.4%), and low stroke risk awareness (60.1%).

Factors related to stroke prevention behavior include self-efficacy ($p = 0.043$; 95%CI = 1.01–2.22; OR = 1.50), perception ($p = 0.038$; 95%CI = 0.45–0.98; OR = 0.663), and awareness of stroke risk ($p = 0.000$; 95%CI = 0.24–0.53; OR = 0.354).

The results of multivariate analysis showed that factors related to stroke prevention behavior were age, sex, efficacy, and awareness of stroke risk. The dominant factor influencing behavior is awareness of stroke risk with OR 3779 (Table 3).

Discussion

Most respondents are female, the elderly, have low education, and earn below the regional minimum wage standard. The risk of hypertension increases with age. This is related to vascular changes that cause an increased peripheral resistance. Postmenopausal women increase the risk of developing hypertension due to hormonal changes. Low education and income can affect health behavior. 24

The result shows that the majority (65.7%) of hypertension patients had a poor stroke prevention behavior. The poor behavior is high consumption of sodium and fat and also lack of regular exercise. The factors that influence stroke prevention behavior in this study include perception, self-efficacy, and stroke risk awareness.

The majority of the respondents have poor perception and low self-efficacy in stroke prevention. Perception and self-efficacy can influence behavior. Imprecise perception in stroke prevention and low self-efficacy will inhibit stroke prevention behavior [10].

One of the factors that influence perception is information. In this study, we obtained bad perception

Table 2: Stroke prevention behavior and associated factors

Variable	n	%	OR	95% CI	p
Stroke prevention behavior					
Good	158	34.3	-	-	-
Bad	303	65.7			
Knowledge					
Low	0	0	0.830	0.44–1.58	0.568
Middle	49	10.6			
High	412	89.4			
Self-efficacy					
Low	259	56.2	1.50	1.01–2.22	0.043
High	202	43.8			
Perception					
Bad	264	57.3	0.663	0.45–0.98	0.038
Good	197	42.7			
Information about stroke					
Low	191	41.4	1.19	0.811–1.766	0.366
Middle	126	27.3			
Good	144	31.2			
Stroke risk awareness					
Low	277	60.1	0.354	0.24–0.53	<0.001
High	184	39.9			

Table 3: Multivariate analysis results (n = 461)

Variable	SE	Sig.	OR	CI 95%	
				Min.	Max.
Sex	0.339	<0.001	0.072	0.037	0.139
Age	0.261	<0.001	2.598	1.557	4.335
Marital status	0.327	0.218	0.669	0.352	1.269
Income	0.260	0.832	0.946	0.569	1.574
Stroke risk awareness	0.263	<0.001	3.779	2.255	6.333
Perception	0.253	0.272	1.321	0.804	2.169
Self-efficacy	0.261	<0.001	0.370	0.222	0.617
Living together	0.278	0.893	1.038	0.602	1.792

and information about stroke (41.4%). Plenty of information can affect or increase someone's knowledge and with knowledge creates awareness that eventually someone will behave in accordance with the knowledge they have.

Self-efficacy is an important precondition for successful self-management. Many studies have shown that self-efficacy is the strongest determinant of smoking cessation, increasing physical activity, and healthy diet [11], [12], [13]. Thus, increasing self-efficacy is an effective method to support healthy behavioral change in patients with stroke.

Respondents also indicated that the majority has a low stroke risk awareness. Many patients are not aware that hypertension is a risk of stroke so they ignore stroke prevention behavior. This is in line with the research of Hertz *et al.* which stated that 7.6% had the chance to become stroke, 46.4% had no chance to become a stroke, and 46.0 were unsure [14]. These findings indicate that self-awareness about stroke risk is poor. This disease attitude can inhibit efforts to raise awareness about stroke in society.

Most people think that stroke is more serious than heart attack. However, because stroke symptoms appear in variety of ways, they are not easily recognized. People with stroke report that they are confused by the symptoms such as migraine-like, headache, ear disease, or food poisoning. This kind of confusion can be one of the causes of delays to the hospital. Studies suggest that myocardial infarction patients with atypical symptoms lead to increased delay in many patients [15], [16]. Dracup *et al.* found the people who thought that they were simply having gas discomfort or indigestion significantly delayed seeing a doctor [17]. This study also revealed that people who had a low stroke self-awareness had a 3.78 times risk of having poor stroke prevention behavior. Therefore, increasing public awareness about the risk of stroke is an important factor in the stroke risk patients management.

Conclusion

Most respondents stroke prevention behaviors are bad. Respondent characteristics that significantly

influence behavior include gender, age, marital status, and income. Factors related to stroke prevention behavior include self-efficacy, perception, and awareness of stroke risk. The dominant factor influencing behavior is awareness of stroke risk. It is necessary to intervene to increase stroke prevention behavior in risk groups by increasing self-efficacy, perception, and awareness of stroke risk.

Ethical Aspect and Declaration of Interest Statement

The study has been declared ethical by the Dr. Moewardi Hospital's health research ethics commission number: 789/VI/HREC/2019. Informed consent was given to the participants before study.

Authors' Contributions

Concept and study design (Upoyo, Sari, Isworo), data collection, data analysis and interpretations (Taufik, Sumeru, Anam), processing the draft of the manuscript (Upoyo, Isworo), critical revision of the manuscript (Sari, Upoyo, Isworo), and article finalization (Upoyo, Isworo).

References

1. Venketasubramanian N, Yoon BW, Pandian J, Navarrod JC. Stroke epidemiology in South, East, and South-East Asia: A review. *J Stroke*. 2017;19(3):286-94. <https://doi.org/10.5853/jos.2017.00234>
PMid:29037005
2. Ministry of Health of the Republic of Indonesia. The Reports of the Results of Basic Health Research; 2019. Available from: <https://www.litbang.kemkes.go.id/laporan-riiset-kesehatan-dasar-risikesdas>. [Last accessed on 2020 Mar 17].
3. Slark J, Sharma P. Risk awareness in secondary stroke prevention: A review of the literature. *JRSM Cardiovasc Dis*. 2014;3:1-6. DOI: 10.1177/2048004013514737.
PMid:24570838
4. Boehme AK, Esenwa C, Elkind MS. Stroke risk factors, genetics, and prevention. *Circ Res*. 2017;120(3):472-95. <https://doi.org/10.1161/circresaha.116.308398>
PMid:28154098
5. Upoyo AS, Setyopranoto I, Pangastuti HS. The modifiable risk factors of uncontrolled hypertension in stroke: A systematic review and meta-analysis. *Stroke Res Treat*. 2021;2021:6683256. <https://doi.org/10.1155/2021/6683256>
PMid:33680423
6. Hanchaiphiboolkul S, Pongvarin N, Nidhinandana S,

- Suwanwela NC, Puthkhao P, Towanabut S, et al. Prevalence of stroke and stroke risk factors in thailand: Thai epidemiologic stroke (TES) study. *J Med Assoc Thai*. 2011;94(4):427-36. <https://doi.org/10.1016/j.jstrokecerebrovasdis.2012.05.013>
PMid:21591527
7. Ghani L, Mihardja LK, Delima D. Dominant risk factors of stroke in Indonesia. *Buletin Penelitian Kesehatan*. 2016;44(1):49-58. <https://doi.org/10.22435/bpk.v44i1.4949.49-58>
 8. Goldstein LB, Adams R, Albert MJ, Appel LJ, Brass LM, Bushnell CD, et al. Primary prevention of ischemic stroke. *Stroke*. 2006;37:1583-633.
 9. Erkoc SB, Isikli B, Metintas S, Kalyoncu C. Hypertension knowledge-level scale (HK-LS): A study on development, validity and reliability. *Int J Environ Res Public Health*. 2012;9(3):1018-29. <https://doi.org/10.3390/ijerph9031018>
PMid:22690180
 10. Michie S, Atkins L, West R. *The Behaviour Change Wheel: A Guide to Designing Interventions*. London: Silverback Publishing; 2014.
 11. Garcia K, Mann T. From "i wish" to "i will": Social-cognitive predictors of behavioral intentions. *J Health Psychol*. 2003;8(3):347-60.
PMid:14670213
 12. Sniehotta FF, Scholz U, Schwarzer R. Bridging the intention-behaviour gap: Planning, self-efficacy, and action control in the adoption and maintenance of physical exercise. *Psychol Health*. 2005;20(2):143-60. <https://doi.org/10.1080/08870440512331317670>
 13. Sol BG, van der Bijl JJ, Banga JD, Visseren FL. Vascular risk management through nurse-led self-management programs. *J Vasc Nurs*. 2005;23(1):20-4. <https://doi.org/10.1016/j.jvn.2004.12.003>
PMid:15741961
 14. Hertz JT, Madut DB, William G, Maro VP, Crump JA, Rubach MP. Perceptions of stroke and associated health-care-seeking behavior in Northern Tanzania: A community-based study. *Neuroepidemiology*. 2019;53(1-2):41-7. <https://doi.org/10.1159/000499069>
PMid:30986785
 15. Moser DK, Dracup K. Gender differences in treatment-seeking delay in acute myocardial infarction. *Prog Cardiovasc Nurs*. 1993;8(1):6-12.
PMid:8372090
 16. Schwarz B, Schoberberger R, Rieder A, Kunze M. Factors delaying treatment of acute myocardial infarction. *Eur Heart J*. 1994;15(12):1595-8. <https://doi.org/10.1093/oxfordjournals.eurheartj.a060439>
PMid:7698126
 17. Dracup K, McKinley SM, Moser DK. Australian patients' delay in response to heart attack symptoms. *Med J Aust*. 1997;166(5):233-6. <https://doi.org/10.5694/j.1326-5377.1997.tb140101.x>
PMid:9076265

Determinant Factors Stroke Prevention Behavior among Hypertension Patient in Indonesia

ORIGINALITY REPORT

17%

SIMILARITY INDEX

13%

INTERNET SOURCES

13%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1	pdfcoffee.com Internet Source	2%
2	www2.j-stroke.org Internet Source	2%
3	bmccresnotes.biomedcentral.com Internet Source	2%
4	www.pubmedcentral.nih.gov Internet Source	1%
5	www.cancerindex.org Internet Source	1%
6	Lis Mukti Lestari, Dwi Pudjonarko, Fitria Handayani Handayani. "Characteristics of stroke patients: An analytical description of outpatient at the hospital in Semarang Indonesia", Jurnal Aisyah : Jurnal Ilmu Kesehatan, 2020 Publication	1%
7	Vivin Nur Hafifah. "Influence of Hypertension and Lifestyle History on Stroke Occurrence	1%

Rates in Probolinggo District", Journal for Quality in Public Health, 2020

Publication

8	jgmds.org.pk Internet Source	1 %
9	knepublishing.com Internet Source	1 %
10	pure.eur.nl Internet Source	1 %
11	Submitted to Clemson University Student Paper	1 %
12	Shu-xia Li, Ling Zhang. "Health behavior of hypertensive elderly patients and influencing factors", Aging Clinical and Experimental Research, 2013 Publication	<1 %
13	Submitted to Sultan Agung Islamic University Student Paper	<1 %
14	cardiab.biomedcentral.com Internet Source	<1 %
15	tnsroindia.org.in Internet Source	<1 %
16	Submitted to Saint Leo University Student Paper	<1 %
17	repository.uin-suska.ac.id Internet Source	

<1 %

18

trialsjournal.biomedcentral.com

Internet Source

<1 %

19

Anggraini Dwi Kurnia, Nur Melizza, Faqih Ruhyanudin, Nur Lailatul Masruroh et al. "The Effect of Educational Program on Hypertension Management Toward Knowledge and Attitude Among Uncontrolled Hypertension Patients in Rural Area of Indonesia", *Community Health Equity Research & Policy*, 2020

Publication

<1 %

20

Devy Putri Nursanti, Eri Puji Kumalasari, Aan Setyaningsih, Julia Siekmans, Vera Wabeke. "ANALYSIS OF ADOLESCENT REPRODUCTIVE HEALTH KNOWLEDGE ON PREMARITAL SEXUAL ATTITUDE", *Journal for Quality in Women's Health*, 2022

Publication

<1 %

21

Falko F. Sniehotta, Urte Scholz, Ralf Schwarzer. "Bridging the intention-behaviour gap: Planning, self-efficacy, and action control in the adoption and maintenance of physical exercise", *Psychology & Health*, 2005

Publication

<1 %

22 Md. Abdur Rafi, Md. Zahidus Sayeed, Papia Sultana, Saw Aik, Golam Hossain. "Pre-hospital delay in patients with myocardial infarction: an observational study in a tertiary care hospital of northern Bangladesh", Research Square Platform LLC, 2020
Publication <1 %

23 Wittawat Chantkran, Janeyuth Chaisakul, Ram Rangsin, Mathirut Mungthin, Boonsub Sakboonyarat. "Prevalence of and factors associated with stroke in hypertensive patients in Thailand from 2014 to 2018: A nationwide cross-sectional study", Scientific Reports, 2021
Publication <1 %

24 proceeding.tenjic.org
Internet Source <1 %

25 www.coursehero.com
Internet Source <1 %

26 Jörg Ederle, Martin M. Brown. "Schlaganfallprävention", Herz Kardiovaskuläre Erkrankungen, 2008
Publication <1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

