

Turnitin Originality Report

Processed on: 05-May-2023 13:00 WIB

ID: 2084834046

Word Count: 2657

Submitted: 1

Oligocene-Pleistocene
Paleogeography within B... By
Eko Bayu Purwasatriya

Similarity Index

19%

Similarity by Source

Internet Sources: 18%
Publications: 9%
Student Papers: 3%

exclude quoted

include bibliography

exclude small matches

mode:

quickview (classic) report



print

refresh

download

7% match (Internet from 29-Jun-2021)

<https://icoesmb.febi.umkendari.ac.id/downloads/ICST2017-abstract-book.pdf>

1% match (Internet from 01-Jan-2019)

<https://docplayer.net/49133481-Welcoming-remarks-from-the-3-rd-icst-chairman.html>

1% match (Internet from 23-Feb-2020)

<https://aip.scitation.org/doi/pdf/10.1063/1.5097478>

1% match (Internet from 09-Apr-2021)

<https://ugrg.ft.ugm.ac.id/conference-proceedings/>

1% match ()

[Satria, Eko Bayu, Surjono, Sugeng Sapto, Amijaya, D. Hendra. "Sejarah Geologi Pembentukan Cekungan Banyumas Serta Implikasinya Terhadap Sistem Minyak dan Gas Bumi", 'Universitas Jenderal Soedirman', 2019](#)

1% match (Firman Herdiansyah, Muhammad Burhannudinnur, Dewi Syavitri. "An unique of the late Miocene turbidite succession in Brebes and Banyumas Central Java", AIP Publishing, 2021)

[Firman Herdiansyah, Muhammad Burhannudinnur, Dewi Syavitri. "An unique of the late Miocene turbidite succession in Brebes and Banyumas Central Java", AIP Publishing, 2021](#)

1% match (Internet from 21-Jan-2023)

<https://www.mdpi.com/2077-0375/13/1/86>

1% match (student papers from 24-Nov-2014)

[Submitted to University of Leeds on 2014-11-24](#)

1% match (student papers from 16-Jul-2020)

[Submitted to MiraCosta College on 2020-07-16](#)

1% match (student papers from 11-Dec-2012)

[Submitted to National University on 2012-12-11](#)

1% match (student papers from 02-Nov-2018)

[Submitted to University Der Es Salaam on 2018-11-02](#)

1% match (Ahmad Setiawan, Zulfakriza Zulfakriza, Andri Dian Nugraha, Shindy Rosalia et al. "Delineation of sedimentary basin structure beneath the Banyumas Basin, Central Java, Indonesia, using ambient seismic noise tomography", Geoscience Letters, 2021)

[Ahmad Setiawan, Zulfakriza Zulfakriza, Andri Dian Nugraha, Shindy Rosalia et al. "Delineation of sedimentary basin structure beneath the Banyumas Basin, Central Java, Indonesia, using ambient seismic noise tomography", Geoscience Letters, 2021](#) ✕

<1% match (Internet from 24-Jul-2020)

<http://dinarek.unsoed.ac.id> ✕

<1% match (Kent C. Condie. "Tectonic Settings", Earth as an Evolving Planetary System, 2011)

[Kent C. Condie. "Tectonic Settings", Earth as an Evolving Planetary System, 2011](#) ✕

<1% match (Internet from 25-Mar-2023)

<https://geoscienceletters.springeropen.com/articles/10.1186/s40562-021-00202-x> ✕

<1% match (Internet from 17-Jun-2022)

https://webofproceedings.org/proceedings_series/ACE/ICCEME%202019/ICCEME19011.pdf ✕

<1% match (Ebrahim Mohammadi, Mahboubeh Hasanzadeh-Dastgerdi, Mehdi Ghaedi, Rezvan Dehghan et al. "The Tethyan Seaway Iranian Plate Oligo-Miocene deposits (the Qom Formation): distribution of Rupelian (Early Oligocene) and evaporate deposits as evidences for timing and trending of opening and closure of the Tethyan Seaway", Carbonates and Evaporites, 2012)

[Ebrahim Mohammadi, Mahboubeh Hasanzadeh-Dastgerdi, Mehdi Ghaedi, Rezvan Dehghan et al. "The Tethyan Seaway Iranian Plate Oligo-Miocene deposits \(the Qom Formation\): distribution of Rupelian \(Early Oligocene\) and evaporate deposits as evidences for timing and trending of opening and closure of the Tethyan Seaway", Carbonates and Evaporites, 2012](#) ✕

<1% match (Kent C. Condie. "Tectonic Settings", Elsevier BV, 2005)

[Kent C. Condie. "Tectonic Settings", Elsevier BV, 2005](#) ✕

<1% match (Internet from 22-Jun-2022)

<https://www.livescience.com/56930-prehistoric-mammals-in-images.html> ✕

<1% match (Internet from 14-Jan-2020)

https://www.scilit.net/articles/10/130/Newest?Search=doi%3A10.29037%2F%2A&has_free_version=&open-access=&preprints= ✕

UGM Digital Press Physical Sciences and Engineering (2018)1: 45–51 [The 3rd International Conference on Science and Technology Oligocene-Pleistocene Paleogeography within Banyumas Basin and implication to petroleum potential Eko Bayu Purwasatriya^{1,2*}, Sugeng Sapto Surjono^{3,,} and D. Hendra Amijaya³](#)
¹Doctoral Program Student, Geological Engineering Department, Faculty of Engineering, Universitas Gadjah Mada, Jl. Grafika No.2, Yogyakarta, 55281 2
 Geological Engineering, Faculty of Engineering, Universitas Jenderal Soedirman, Jl. Mayjen Sungkono, Km.5, Blater, Purbalingga, 53371 3 Geological Engineering Department, Faculty of Engineering, Universitas Gadjah Mada, Jl. Grafika No.2, Yogyakarta, 55281 *e-mail: bayusatriya@yahoo.com Abstract [This study attempts to reconstruct paleogeography of Banyumas Basin in association with magmatic arc evolution and its implication to petroleum potential. Based on the volcanic rocks distribution, their association and relatives age, there are three alignments of a magmatic arc, that are: \(1\) Oligo-Miocene arc in the south \(2\) Mio- Pliocene arc in the middle \(3\) Plio-Pleistocene arc in the north. The consequences of the magmatic arc movement were tectonic setting changing during Oligocene to Pleistocene, as well as their paleogeography. During Oligo-Miocene where magmatic arc existed in the southern part, the Banyumas tectonic setting was a back-arc basin. This tectonic setting was changing to intra-arc basin during Mio-Pliocene and subsequently to fore-arc basin since Plio-Pleistocene until today. Back-arc basin is the most suitable paleogeography to create a depositional](#)

environment for potential source rocks. Exploration activity to prove the existence of source rocks during Oligo-Miocene is needed to reveal petroleum potential in Banyumas Basin. Keywords Back-arc Basin, Banyumas Basin, Magmatic arc, Paleogeography, Petroleum potential Citation: PurwasatriyaE. B., Surjono S. S., D. H. Amijaya.: Oligocene-Pleistocene paleogeography within Banyumas basin and implication to petroleum potential. In: I.W. Mustika, I. Kartini. (eds.): Proceeding of the 3rd International Conference on Science and Technology, Vol. 1, pp. 45–51.

UGM Digital Press Physical Science and Engineering, Yogyakarta (2018).

Published: October, 2018 1 Introduction Banyumas basin is one of a sedimentary basin in the southern coast of Central Java, Indonesia. Banyumas also categorized as frontier basin because although several oil and gas seeps found, but, not yet economic petroleum revealed in this area. A previous researcher called Banyumas Basin as "Terra Incognita," a place that not yet fully understood its petroleum systems [1]. It is important to know how is the petroleum systems works in Banyumas basin and one step to know the petroleum systems better is by study the paleogeography of Banyumas basin through time by time. Paleogeography means in this topic is related to its position to a magmatic arc. If the basin's position in front of a magmatic arc and directly face opening sea, it is called fore-arc basin. If the basin's position is among the mountains on the 10.29037/digitalpress.11244 magmatic arc, it is called intra-arc basin. If the position of the basin is in the back of a magmatic arc, it is called back-arc basin. Fore-arc basins are marine depositional basins on the trench side of arcs, and back-arc basins occur over descending slabs behind arc systems [2] (Fig. 1). In petroleum systems, back-arc basin and intra-arc basin are the paleogeography which has important role to create a petroleum source rock. Petroleum source rock is usually a fine grain rock, contain sufficient organic matter and will be mature by thermal conduct. In back-arc and intra-arc basin, the depositional environment is quiet of current and wave, so, the fine grain sediments can be deposited rapidly, and the organic matter also can be preserved in a reduction environment. Banyumas basin indicates changing its paleogeography in different geological time. The indication is visibly by the distribution of volcanic rock with different age and made an alignment. The geological study is needed to find the evidence in the surface, and geophysical data such as gravity data is needed to support the interpretation by providing the subsurface data. Integration of geology and geophysical data will increase the confidence of the interpretation. Fig. 1 Diagrammatic cross sections of (a) continental margin and (b) island arc, showing fore-arc basin, intra-arc basin and back-arc basin [2]. 2 Methods The methods divide into two parts, first is the geological method and the second is a geophysical method. The geological method is conducted to get the surface data, such as lithology, stratigraphy and structural geology and geophysical method is to provide the subsurface data. The geological method comprises fieldwork and studio analysis. Fieldwork means in this study is a general geological surface mapping to identify the type of lithology that indicates the volcanic product and also the location, is it made alignment of magmatic arc or not. Studio analysis is comprised of making the general geological map from the fieldwork and analysis the type of rock samples picked from the field. The relative age of the rock is taken from the previous researcher that already have the age data for Banyumas area. The data needs for the geological method are : - Regional geological maps for Banyumas and surrounding area, scale 1:100,000 - Topographic maps for Banyumas and surrounding area, scale 1:25,000 - Literature from the previous researcher for references The tools needed for geological fieldwork comprises of Geological Compass, Geological Hammer, Loupe, GPS, Camera, Stationery, Field books, medicine, and other personal needs. The second method is the geophysical method. Geophysical method means in this study is the gravity method. Gravity data is taken from the topex website http://topex.ucsd.edu/cgi-bin/get_data.cgi [3]. It is a satellite gravity data and can be download for free of charge. Around 1,800 point of gravity data was download for Banyumas and surrounding area and need to process to get the gravity anomaly map. The step of gravity processing data is provided on (Fig. 2). The figure 2 shows the steps from acquisition data to regional and residual anomaly maps, but, if we download the data from topex's website, the data already a free-air anomaly, so the step is shorter than the flow diagram. Gravity data will support the analysis and interpretation of magmatic arc from subsurface information. Gravity works based on the contrast density of rocks in the subsurface and volcanic rocks havea higher density than other sedimentary rocks. Regional anomaly gravity usually used to identify the deep anomaly such as the border of a paleobasin and the residual anomaly gravity usually used to

identify the shallow and local anomaly. This study used the residual anomaly to detect the alignment of the paleo-magmatic arc and matching the interpretation with the geological surface data. The integration of surface data from the geological study and subsurface data from gravity interpretation should give better results to define the alignment of the paleo-magmatic arc and then interpret the paleogeography of the basin through geological time. Reading value at gravimeter - Instrument scale factor - Tidal correction Observed Gravity -Latitude Correction Normal Gravity - [Free-air Correction](#) [Free-air Anomaly](#) - [Bouguer Correction](#) [Bouguer Anomaly](#) - [Terrain Correction](#) Complete [Bouguer Anomaly](#) Regional Anomaly Residual Anomaly Fig. 2. Flow diagram of gravity processing to get the regional and residual anomaly maps 3. Results and Discussions Geological fieldworks and studio analysis had been conducted in Banyumas area. Some association of volcanic rocks found and made an East-West alignment based on its relative age. Association of volcanic rocks related to the central zone and proximal zone of a magmatic arc are: breccia, lava, intrusion dike or sill. Association of volcanic rocks related to the distal zone of magmatic arcs such as tuff, marl, tuffaceous sandstone, siltstone, and claystone. Figure 3 shows the example of volcanic rocks in the field. Figure 3(A) is a very rare kind of breccia; it is called limestone breccia. It is very rare because usually limestone is not associated with breccia and in this case, the fragment of breccia is andesite, and the matrix of breccia is limestone. It probably happens when volcanic activity is shut off, so carbonate reef can build up, but there is a small volcanic activity like diatreme made explosive and mixed with reworked carbonate reef. Figure 3(B) is a "common" breccia, called andesite breccia, composed of andesite as fragments and sandstone as a matrix. Andesite breccia is distributed almost at all magmatic arc alignment but with different geological age. Figure 3(C) shows the magmatic intrusion in a limestone. The intrusion indicates magmatic activity near the central zone of a magmatic arc. Figure 3(D) shows the volcanoclastic rocks, intercalation of tuffaceous sandstone and siltstone. The volcanoclastic rocks usually indicate the distal zone of the magmatic arc. Fig. 3 (A) Limestone Breccia, (B) Andesite Breccia, (C) Magmatic intrusion on limestone and (D) intercalating of tuffaceous sandstone and siltstone After doing fieldwork and studio analysis, a regional geological map of Banyumas and the surrounding area has made. Geological map of Banyumas and surrounding area is shown in (Figure 4). The picture in figure 4 a little bit complex because the area is large, but the legend gives the explanation of the figure to understand the meaning. The various range of color from red to brown is showing the association of volcanic rocks such as breccia, lava and intrusion, means the area is near with magmatic arc. The various range of color from yellow to green is showing the volcanoclastic rocks such as tuff, marl, tuffaceous sandstone, siltstone, and claystone, means the area probably a distal zone or far zone of a magmatic arc. The blue color means limestone and grey color means alluvium. The brown line crossing the map is structural geology such as a fault or fold axis, and the blue curve is a delineation of Banyumas Basin taken from [4]. This geological map is modified from the previous regional geological map such as Banyumas area [5], Purwokerto and Tegal area [6], Majenang area [7] and Pangandaran area [8]. There are three alignment of magmatic arc in Banyumas basin and bordered by [the black dashed line](#) in [the](#) geological map. [The](#) first alignment is [in the south of Banyumas basin](#), called Oligo-Miocene magmatic arc. The second alignment is [in the middle of the basin](#), called [Mio-Pliocene](#) magmatic [arc](#) and [the](#) third alignment is Plio-Pleistocene magmatic arc in the north of the basin. Oligocene was around 33.9 [million years ago](#), [Miocene](#) is around [23 million years ago](#), Pliocene is around [5.3 million years ago](#), and Pleistocene is around [2,6 million years ago](#) [9]. Based [on the](#) position [of](#) a magmatic arc, [the](#) paleogeography of Banyumas basin change through geological time. At Oligocene-Miocene, the paleogeography probably a back-arc basin if the basement is continent or an island arc if the basement is the oceanic crust. At Miocene-Pliocene time, the paleogeography probably an intra-arc basin and at Pliocene-Pleistocene the paleogeography becomes a fore-arc basin until now. Moreover, Banyumas basin still keep the petroleum potential, in 2010 a water well drilling found oil at depth 90 meter, [in Tipar Kidul village, Ajibarang Sub- district](#), Banyumas regency [10], [this](#) proved [the](#) interesting challenge of Banyumas basin and also petroleum potential. Fig. 4 Geological map of Banyumas and surrounding area shows basin delineation and three alignments of a magmatic arc. JAVA Fig. 5 Gravity residual map showing the three alignments of magmatic arc indicated by low anomaly (green) among high anomaly (orange) The geophysical method creates a residual gravity map and regional gravity map, but in this study, only

residual gravity map used to support the interpretation of alignment of magmatic arc from surface data. Gravity residual map (Fig. 5) also shown the alignment of three magmatic arcs, same with geological map. The low anomaly with green color among high anomaly with orange color interpreted as a mountainous arc. This interpretation based on Slamet mountain in the north of Banyumas basin that gives the same response, a low anomaly among high anomaly. It can be understood because gravity works based on the density of medium and mountains will have a hole in the middle of it is cone filled with soft sediments and give low anomaly respond in gravity. The flank of the cone usually composes of breccia, lava, and intrusion which are high densities and give high anomaly response in gravity.

4 Conclusions There is three magmatic arcs within Banyumas basin, first is [Oligo-Miocene magmatic arc in the southern part of the basin](#), second is [Mio-Pliocene magmatic arc in the middle of the basin](#), and third is Plio- Pleistocene magmatic arc in the northern part of the basin. The movement of the magmatic arc through geological time, giving implication to the paleogeography of Banyumas basin. At Oligo-Miocene, the paleogeography probably back-arc basin or island arc, at Mio-Pliocene the paleogeography probably intra-arc basin and at Plio-Pleistocene the paleogeography should be fore-[arc basin](#). [Back-arc basin](#) and [the intra-arc basin is the](#) most suitable paleogeography for petroleum potential because the environment is quite of current and wave, so, the fine grain sediment and organic matter can be deposited and preserved. By burial for million years, the temperature will increase, and the organic matter should be mature and convert to petroleum. A petroleum source rock that had been mature and full of petroleum will expel to carrier bed like sandstone and migrate to a trap consist of the good reservoir. Need more investigation about the petroleum systems in this area to making a prospect of petroleum.

Acknowledgments I would like to thank to [Geological Engineering Department, Faculty of Engineering, Universitas Gadjah Mada](#) and Geological Engineering, Faculty of Engineering, Universitas Jenderal Soedirman.

References [1]. Satyana, A.H., Central java indonesia-a'terra incognita' in petroleum exploration : new consideration on the tectonic evolution and petroleum implication, Proceedings Indonesian Petroleum Association, 31st annual convention Jakarta, pp. 105-126 (2007). DOI: 10.29118/IPA.1211.07.G.085 [2]. Kent C. Condie, Plate Tectonics and Crustal Evolution, Fourth Edition, Butterworth-Heinemann, Oxford (1997). <https://doi.org/10.1016/B978-0-7506-3386-4.X5000-9> [3]. Topex (2014), Extract XYZ Grid – Topography or Gravity, http://topex.ucsd.edu/cgi-bin/get_data.cgi, Accessed: May, 2014 [4]. Purwasatriya, E.B., Tinjauan kembali potensi hidrokarbon Cekungan Banyumas berdasarkan data geologi dan geofisika, Proceeding Seminar Kebumihan Ke-7, Yogyakarta (2014). https://www.academia.edu/29740002/TINJAUAN_KEMBALI_POTENSI_HIDROKARBON_CEKUNGAN_BANYUMAS_BERDASARKAN_DATA_GEOLOGI_DAN_DATA_GEOFISIKA [5]. Asikin, S., Handoyo, A., Prastistho, B., dan Gafoer, S., Peta Geologi Lembar Banyumas, Jawa, Pusat Penelitian dan Pengembangan Geologi, Bandung, Indonesia (1992) http://library.vsi.esdm.go.id/katalog/slims7_cendana-master//index.php?p=show_detail&id=11478 [6]. Djuri, M., Samodra, H., Amin, T.C., dan Gafoer, S., Peta Geologi Lembar Purwokerto Dan Tegal, Jawa, Pusat Penelitian dan Pengembangan Geologi, Bandung, Indonesia (1996) [https://www.dropbox.com/s/q1n6fgqy11bv29/22-1308-6-1309-3-Purwokerto dan Tegal.jpg?dl=0](https://www.dropbox.com/s/q1n6fgqy11bv29/22-1308-6-1309-3-Purwokerto%20dan%20Tegal.jpg?dl=0) [7]. Kastowo, Peta Geologi Lembar Majenang, Jawa, Direktorat Geologi, Bandung, Indonesia (1975). <https://www.dropbox.com/s/lxyl9vlymsyxvti/20-1308-5-Majenang.jpg?dl=0> [8]. Simandjuntak, T.O., dan Surono, Peta Geologi Lembar Pangandaran, Jawa, Pusat Penelitian dan Pengembangan Geologi, Bandung, Indonesia (1992). <https://www.dropbox.com/s/dyeglqt4ek3m8ig/21-1308-2-Pangandaran.jpg?dl=0> [9]. Walker J.D., Geissman, J.W., Bowring, S.A., Babcock L.A., GSA Geologic Time Scale v. 4.0, Geological Society of America, America (2012). https://www.geosociety.org/GSA/Education_Careers/Geologic_Time_Scale/GSA/timescale/home.asp [10]. Purwasatriya, E.B., and Waluyo, G., Studi Potensi Minyak Dangkal dengan Pendekatan Metode Statistik Berdasar Data Geologi Permukaan Di Cekungan Banyumas, Jurnal Dinamika Rekayasa, Vol.8 No.2 (2012). DOI: <http://dx.doi.org/10.20884/1.dr.2012.8.2.103> Proceeding of the ICST 2017 Physical Sciences and Engineering UGM Digital Press (2018) Vol. 1: 45–51 Proceeding of the ICST 2017 Oligocene-Pleistocene Paleogeography within Banyumas Basin.... Physical Sciences and Engineering UGM Digital Press (2018) Vol. 1: 45–51 Proceeding of the ICST 2017 Oligocene-Pleistocene Paleogeography within Banyumas Basin.... Physical Sciences and Engineering UGM Digital Press

(2018) Vol. 1: 45–51 Proceeding of the ICST 2017 Oligocene-Pleistocene Paleogeography within Banyumas Basin.... 45 46 [E. B. Purwasatriya, S. S. Surjono & D. H. Amijaya](#) 10.29037/digitalpress.11244 47 48 [E. B. Purwasatriya, S. S. Surjono & D. H. Amijaya](#) 10.29037/digitalpress.11244 49 50 E. B. Purwasatriya, S. S. Surjono & D. H. Amijaya 10.29037/digitalpress.11244 51